

# Effect of simultaneous exercise and cognitive training on executive functions, baroreflex sensitivity and pre-frontal cortex oxygenation in healthy older adults: a pilot study Manon Pellegrini-Laplagne, Olivier Dupuy, Philippe Sosner, Laurent Bosquet

# ▶ To cite this version:

Manon Pellegrini-Laplagne, Olivier Dupuy, Philippe Sosner, Laurent Bosquet. Effect of simultaneous exercise and cognitive training on executive functions, baroreflex sensitivity and pre-frontal cortex oxygenation in healthy older adults: a pilot study. GeroScience, 2022, 45, pp.119-140. 10.1007/s11357-022-00595-3. hal-04538515

# HAL Id: hal-04538515 https://hal.science/hal-04538515

Submitted on 9 Apr 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. **Title**: Effect of simultaneous exercise and cognitive training on executive functions, baroreflex sensitivity and pre-frontal cortex oxygenation in healthy older adults: a pilot study.

Running title: Simultaneous exercise/cognitive training and cognition

Authors : Manon Pellegrini-Laplagne<sup>1</sup>, Olivier Dupuy<sup>1,2\*</sup>, Phillipe Sosner<sup>1,3</sup>, Laurent Bosquet<sup>1,2</sup>

#### Affiliations:

- 1. Laboratoire MOVE (EA6314), Université de Poitiers, Faculté des sciences du sport, 8 allée Jean Monnet – TSA 31113 – 96073 Poitiers cedex 9
- 2. School of Kinesiology and Physical Activity Sciences (EKSAP), Faculty of Medicine, University of Montreal, Montreal, CANADA
- 3. Mon Stade, Paris, FRANCE

Corresponding author: Dr Olivier Dupuy 2: +33 (0)5 49 45 33 43 : olivier.dupuy@univ-poitiers.fr

Laboratoire MOVE (EA6314), Université de Poitiers, Faculté des sciences du sport, 8 allée Jean Monnet – TSA 31113 – 96073 Poitiers cedex 9

#### Abstract (270 words):

Aging is characterized by cognitive decline affecting daily functioning. To manage this socioeconomic challenge, several non-pharmacological methods such as physical, cognitive, and combined training are proposed. Although there is an important interest in this subject, the literature is still heterogeneous. The superiority of simultaneous training compared to passive control and physical training alone seems clear but very few studies compared simultaneous training to cognitive training alone. The aim of this pilot study was to investigate the effect of simultaneous exercise and cognitive training on several cognitive domains in healthy older adults, in comparison with either training alone. Thirty-five healthy older adults were randomized into one of three experimental groups: exercise training, cognitive training, and simultaneous exercise and cognitive training. The protocol involved two 30-min sessions per week for 24 weeks. Cognitive performance in several domains, pre-frontal cortex oxygenation, and baroreflex sensitivity were assessed before and after the intervention. All groups improved executive performance, including flexibility or working memory. We found a group by time interaction for inhibition cost ( $F_{(2,28)}$  = 6,44; p < 0,01) and baroreflex sensitivity during controlled breathing (F<sub>(2,25)</sub> = 4,22; p = 0,01), the magnitude of improvement of each variable being positively associated (r = 0.39; p = 0.03). We also found a decrease in left and right pre-frontal cortex oxygenation in all groups during the trail making test B. A simultaneous exercise and cognitive training are more efficient than either training alone to improve executive function and baroreflex sensitivity. The results of this study may have important clinical repercussions by allowing to optimize the interventions designed to maintain the physical and cognitive health of older adults.

**Keywords:** Exercise training, Cognitive training, Simultaneous training, Motor-cognitive dual-task training, Healthy aging, Cognition.

## **INTRODUCTION**

Aging process is associated with a normal cognitive decline, which may affect well-being, quality of life and ultimately autonomy and life expectancy [1]. Cognition covers a wide range of mental capacities necessary to perceive, process, and interact with our environment [2]. Normal aging is associated with a decline in cognitive functions such as processing speed, memory, and executive functions [3]. In addition to some specific neuro-anatomical and functional changes in the brain [4], age-related changes in the cardiovascular system are also involved in this cognitive decline [5]–[8]. In fact, an alteration of cardiac autonomic control, and more specifically of baroreflex sensitivity (BRS), as well as decrease in cerebral perfusion and cerebral oxygenation, have been shown to increase the risk of cognitive impairment [9].

Cognitive functions are influenced by many physiological factors and to a large extent by cardiovascular mechanisms. During the ageing process, the development of cognitive dysfunction and dementia can be attributed to cerebral and vascular disorders (Sahathevan et al, 2012). In particular, many chronic diseases affect cerebral blood flow and explain an accentuated cognitive decline. It is well known that our cognitive functions are influenced by CBF as shown by numerous studies modulating CBF and affecting cognitive function (de Aquino Lemos V et al, 2012, Liu et al, 2013, Sun et al, 2012, Virtues-Ortega et al, 2004). One would expect that CBF and its regulation would be a key factor in the modification of cognitive function. Furthermore, cerebral metabolism and in particular cerebral oxygenation may be as important in influencing cognitive function, rather than changes in CBF (Ogoh, 2017). Indeed, several studies in exercise or at rest have shown that executive functions are influenced by cerebral oxygenation (Mekari et al, 2015; Dupuy et al, 2015). All these mechanisms are themselves under the direct influence of arterial baroreflex (Ogoh and Tarumi, 2019). A direct association between the arterial baroreflex and cerebral circulation, via the autonomic nervous system, has been demonstrated in some animal models (Nakai, 1985, Nakai et al, 1982; Iadecola et al, 1983). In human models, several studies have shown a direct association between baroreflex activity and cognitive functioning (Saint Martin, 2013), and the perturbation of baroreflex activity during aging process explains the perturbation of memory functions (Saint Martin, 2015 ). Indeed, many brain structures involved in many cognitive functions are also responsible for the regulation of the baroreflex (Ding, et al, 2020; Kimmerly et al, 2017) and could explain the association between cognitive functioning and baroreflex activity. For example, an original article incorporating a wide range of these physiological mechanisms (Tarumi et al, 2015), showed that baroreflex activity in older subjects, is associated to better aortic stiffness and white matter integrity, all this combined with better cognition in the Trail test.

The demographic transition toward an older society is not the only cause of these alterations. Life habits also play a major role, since physical inactivity and sedentary behaviors have been associated with reduced cognitive performances [10]. Yet, the level of physical activity in western countries decreased continuously over the past decades, while the duration of sedentary behaviors increased dramatically [11]. The consequence is an increase in the incidence of cognitive impairment in the near future, that is expected to double within the next 20 years [12]. In this context, one of the main challenges consists in developing efficient strategies to strengthen the "cognitive reserve" of the elderly [13]. Fortunately, interventions aimed at mitigating cognitive decline are attracting great research and societal interest. To date, two strategies have been used to prevent cognitive decline and of dementia: (i) cognitive training and (ii) exercise training [14], [15].

There is growing epidemiological evidence showing the importance of cognitive stimulation in the prevention of cognitive decline and dementia [1]. From an ecological point of view, La Rue

(2010) showed a link between the frequency of activities like lecture, puzzles, knitting, or gardening and the 4-year risk of neurogenerative diseases [15]. Furthermore, several cross-sectional studies have also shown that cognitive stimulation was negatively associated with brain beta-amyloid plaque (a sign associated with Alzheimer's disease) [16] or atrophy of the brain hippocampus [17]. Lastly, recent meta-analyses and reviews have reported that cognitive training was also effective in improving cognitive function in older people [18]. On the other hand, there is an important bulk of knowledge showing a close relationship between physical activity and performance in several cognitive domains in the elderly [4], [19]–[22], or the risk of dementia [23]. Several mechanisms have been proposed to understand this beneficial effect of physical activity and/or exercise training, including a reduction in inflammation or the promotion of neurovascular integrity and brain plasticity through hormones and growth factors release [24]–[26]

Considering the positive impact of exercise training or cognitive training on the cognitive health of the elderly when they are implemented separately, we could question the interest of combining these two types of training within the same intervention. In fact, synergistic effects between both strategies can be expected, thus inducing greater benefits than those obtained by each intervention alone [27]. For example, exercise training may improve brain plasticity by increasing cell proliferation and synaptic plasticity, while cognitive training may orientate this plasticity toward the brain areas solicited during cognitive exercises by increasing the number of neurons in pre-existing neural networks [27], [28]. This is at least what suggests a study [29] who showed in mices that exercise combined with a sequence of enriched environment yielded an approximately 30% greater increase in new neurons than either stimulus alone. On their side, Smith and colleagues [30] reported that exercise combined with cognitive training resulted in a greater cognitive improvement in mices than either stimulus alone, thus suggesting that adaptations reported by Fabel et al. [29] may have functional consequences on these animals. Results are more equivocal in humans, since recent narrative reviews [31]–[33], systematic reviews [34]–[36], and meta-analyses [37]–[41] have reported conflicting results. The positive benefits of combined exercise and cognitive training on cognitive performances in the elderly is unanimous. What is debated is rather the superiority of this combined training when compared with either training alone, since the difference between conditions is not systematic. One of the reasons for this heterogeneity is the diversity of protocols used to achieve this double stimulation. As underscored by Herold et al. [32], combined exercise and cognitive training can take several forms. The combination of both stimulations can be sequential (i.e., at different moments), or simultaneous (i.e., at the same time). Regarding the simultaneous combination, physical and cognitive exercises can belong to the same task, as is the case in exergaming, or be independent [32]. The very recent metaanalysis from 2021 [38] shows that the simultaneous combination is more efficient to improve cognitive performance, whenever people suffer from mild cognitive impairment or not. The simultaneous combination is also less time-consuming and probably represent a lower training load than the sequential combination [39], which potentially prevents from excessive stress, less engagement in home- or community-based activities, or other changes that could inhibit rather than promote neural plasticity and cognitive improvement [42].

To date, only one experimental study compared the cognitive benefits of simultaneous combined training with either training in healthy older people [9]. The authors reported a greater improvement in divided attention, executive control, and working memory following combined training when compared with either training [9]. These promising results need to be confirmed, and to be associated to neurophysiological measures in under to understand the mechanisms that underpin this better efficiency. Thus, the main purpose of this pilot study was to assess the impact of a simultaneous combination of exercise and cognitive training on several cognitive

domains in healthy older adults, in comparison with exercise and cognitive training alone. Based on Kraft theory [27], we hypothesized that simultaneous combined training would be effective in improving executive performance, and that the magnitude of improvement would be greater than either training mode alone. Considering the role of cardiac autonomic control and cerebral oxygenation on aging-induced cognitive decline, the second purpose of this study was to assess the impact of simultaneous combined training on cerebral oxygenation and baroreflex sensitivity. We hypothesized that the improvement in executive functions was associated with the improvement of baroreflex sensitivity and / or an improvement of the brain oxygenation.

## **METHODS**

## Population

Forty-seven men (n=16) and women (n=19) aged between 55 and 76 years old were recruited to participate in this study. Their characteristics are presented in Table 1. None of them suffered from cognitive impairment (as defined as a MSSE score inferior or equal to 24), or from cardiovascular, metabolic, neurological, and psychiatric diseases. Participants who were prescribed with a pharmacological treatment that could modify cardiovascular or neuromuscular functions were excluded. The protocol was reviewed and approved by a national ethics committee for non-interventional research (CERSTAPS # 2017-11-17) and was conducted by recognized ethical standards and national/international laws. All participants signed a written statement of informed consent.

## <u>Study design</u>

Participants completed two experimental sessions within a 2-week period, before and after the completion of a training program involving two 30-min sessions per week for 12 weeks. The first experimental session was dedicated to the explanation of the study, the written informed consent, and the first part of the neuropsychological assessment (MOCA, BDI, Stroop Test, TMT, Rey words test, Digit Symbol Substitution Test). During the second experimental session, participants completed the second part of the neuropsychological assessment (N-back test), as well as the BRS test and the submaximal intensity exercise test. After the baseline evaluation, participants were randomly assigned to one of three experimental groups: (1) exercise training (ET); (2) cognitive training (CT); (3) simultaneously combined exercise and cognitive training (ST).

## Test and measures

## **Physiological assessment**

**Submaximal intensity exercise test**. This test was performed on a stationary bicycle (Monark LC6, Monark Exercise AB, Vansbro, Sweden). Theoretical maximal heart rate was determined by the formula proposed by Gellish and colleagues [43]

## Eq. 1 $tHR_{max} = 207 - 0.7 * age$

The test began at 30W and increased by 20 W every two minutes until the participant reached 60% of  $tHR_{max}$ . Participants had to pedal at 60 rotations per minute (rpm). Oxygen uptake (VO<sub>2</sub> ml.kg<sup>-1</sup>.min<sup>-1</sup>) was determined continuously on a 30-s basis using a portable cardiopulmonary exercise testing system (Metalyser Cortex 3B, CORTEX Biophysik GmbH, Germany). Gas analyzers were calibrated before each test using ambient air and a gas mixture of known concentrations (15 % O<sub>2</sub> and 5 % CO<sub>2</sub>). The turbine was calibrated before each test using a 3-1

syringe at several flow rates. The highest VO<sub>2</sub> over a 30-s period during the last stage was considered as the oxygen uptake at 60% of  $tHR_{max}$  ( $\dot{VO}_{2peak(60\%)}$ , ml.kg<sup>-1</sup>.min<sup>-1</sup>). Heart rate was measured continuously using a heart rate monitor (Polar RS800 cx, Polar Electro, Kempele, Finland).

**Baroreflex sensitivity**. Baroreflex sensitivity testing (BRS) is an established tool for assessing cardiac autonomic control and has proven to be a valuable source of information in clinical management, particularly in the assessment of prognosis in a variety of cardiac diseases (LaroVere et al, 2008). In this study, baroreflex sensitivity data were obtained by a non-invasive continuous blood pressure monitor (Finapres NOVA, FMS Company, the Netherlands). Arterial blood pressure was obtained continuously and noninvasively from a finger cuff by using the system with the volume-clamp technique maintaining the diameter of the artery under an inflated finger cuff at a set point, thereby determining arterial pressure with time changes. Diodes were located in the finger cuff, on the side of the finger, to detect changes in artery diameter and changes in the inflation of the cuff to maintain the diameter at the set point. The cuff was inflated or deflated via an air bladder connected to an air hose and pump. The software, using a mathematical model, generates an aortic pulse waveform from the finger arterial pressure wave. This computation considers changes in the pulse pressure and waveform shape as the pressure pulse is transmitted down the brachial arteries to the finger arteries. The arterial blood pressure measurement obtained from a finger was validated by Guelen et al, (2008). Also, an electrocardiogram (ECG) was obtained using 3 leads. The first electrode was positioned within the frame of rib cage, right under the clavicle near shoulder, the second electrode was placed below left clavicle, at the same level of the previous electrode and the last one was positioned above the left anterior superior iliac spine. This procedure was recommended by the manufacturer. The ECG and blood pressure signals were measured continuously by a Finapres (Finapres NOVA, FMS Company, the Netherlands). The blood pressure following each R peak was recorded. Spontaneous baroreflex sequences corresponding to the activation of the baroreflex are defined as a variation in the same direction, over at least three consecutive heartbeats, of the systolic blood pressure and the RR interval. For each sequence, a linear regression was used to estimate the slope of the relationship. The average baroreflex slope of all sequences was then calculated. This method, which allows non-invasive and non-traumatic measurement of baroreflex without pharmacological stimulation, was described previously (Bertinieri G et al, 1985, Yamamoto et al, 1991, Westerhof et al, 2004). All this method is performed by the software. The signals were measured continuously during a first period of five minutes in a supine position (in spontaneous breathing rate), followed by 5 minutes in a supine position with a controlled breathing rate (12 cycles per minute), followed by five minutes in a supine position (spontaneous breathing rate) and then 5 minutes in a standing position (spontaneous breathing rate). The BRS (in ms/mmHg) was calculated for each period of 5 minutes. The algorithms used by the software to calculate BRS included parameters sur such as age, body height, body weight, and blood pressure. This short-term supine and standing resting recording during spontaneous and/or paced breathing is usually carried out for clinical pruposes (LaRovere et al, 2008).

**Prefrontal cortex oxygenation measurement.** Cerebral oxygenation and more precisely the concentration changes of  $[HbO_2]$  ( $\Delta[HbO_2]$ ) and [HHb] ( $\Delta[HHb]$ ) were recorded during the n-back test with the PortaLite fNIRS system (Artinis Medical Systems, Elst, Netherlands) and during Stroop and TMT test with the OxyMon fNIRS system (Artinis Medical Systems, Elst, Netherlands). These two systems utilize near-infrared light, which penetrates the skull and brain but is absorbed by hemoglobin (Hb) chromophores in capillary, arteriolar, and venular beds (M. Ferrari and V. Quaresima). The light was transmitted with two wavelengths, 764 and 857 nm,

and data were sampled with a frequency of 10 Hz. This procedure measures relative changes in [HbO<sub>2</sub>] and [HHb] using the modified Beer-Lambert law. This law takes into account the differential pathlength factor (DPF), which is determined using the following formula: DPF ( $\lambda$ = 807 nm, A) =  $4.99 + 0.067 \times (age 0.814)$ . The region of interest is the Fp1 and Fp2 of the prefrontal cortex (PFC) and were located using the 10/20 international system. Concerning the n-back task, the placement of optode was already described in previous studies (Goenarjo et al, 2019). The patch used in this study for Stroop and TMT task used eight optical channels, comprising four emitters and four receptors, covered the right and left DLPFC and ventrolateral PFC (VLPFC) (Brodmann areas, BAs 9/46 and 47/45/44),. The distance between each emitter and receptor was 3.5 cm. The placement of this patch was already used in previous studies (Agbangla et al, Pellegrini et al, 2022). The sensors were shielded from ambient light with a black cloth. Oxysoft version 3.0 (Artinis Medical Systems, Elst, Netherlands) was used for data collection. Initially resting PFC oxygenation was acquired in a seated position for 2 min. Because continuous-wave technology does not allow quantifying absolute concentration due to the incapacity of measuring optical path lengths, the mean of [HbO<sub>2</sub>] and [HHb] during the total duration of each block of cognitive task, were compared to the minute preceding each block. During all the procedure, our participants were asked to always face forward during the test, avoid making a sudden head movement, clenching their jaw, frowning, and other facial expressions. This pre-processing analysis was already used in previous studies (Pellegrini et al, 2020). To analyze the fNIRS results, we first performed a channel-by-channel analysis. In the absence of interaction between the channels, we averaged the channels together from the right and left hemispheres and also averaged the hemispheres together.

## **Neuropsychological Assessment**

**Global cognitive test**. The Montreal Cognitive Assessment test (MOCA) was used to evaluate global intellectual efficiency. Briefly, this test consists of a 30-points test divided into eight parts and fourteen subtests. It's a very complete test because it targets most cognitive functions: visuospatial skills, executives' functions, attention, working memory, short-term memory delayed recall, language. A score higher than 26 is a normal score, below this one, people have a mild cognitive impairment [47]. The dependant variable was the total score.

**Executives' functions**. The Trail Making Test (TMT) and the Modified Stroop Color test were used to assess executives' function and attentional abilities. Part A of TMT evaluates the information processing speed, while part B assesses cognitive flexibility or switching. In part A, the task consisted in connecting numbers (from 1 to 25) with straight lines as fast as possible. In part B, participants had to alternate between letters in alphabetical order and numbers in ascending order (1-A-2-B-3C, etc.) as fast as possible. The dependent variable was accuracy and time to complete each part.

The Computerized Modified Stroop task was used to assess executive functions. The test used in this study is based on the Modified Stroop Color Test and included three experimental conditions: naming, inhibiting, and switching. This task was already used in several articles [45], [48]–[50] with older. Each block lasted between 2–4 minutes and was interspersed with 60-s resting blocks. Overall, there were three experimental task blocks (1 naming, 1 inhibition, and 1 switching) and 2 resting blocks, for a total length between 8–14 min. In total, there were 60 Naming trials (Block 1), 60 Inhibition trials (Block 2), and 60 Switching trials (Block 3). All trials began with a fixation cross (or square for switching condition) for 1.5 s, and all visual stimuli appeared in the center of the computer screen for 2.5 s. Participants responded with two fingers (index and major finger) from each of their hands on an AZERTY keyboard. In the Naming block, participants were presented with a visual stimulus of the name of colors (RED/BLUE/GREEN/YELLOW) in French presented in the color that is congruent with the word (i.e., RED presented in red ink). Participants were asked to identify the color of the ink with a button press. In the Inhibition block, each stimulus consisted of a color word (RED/BLUE/GREEN/YELLOW) printed in the incongruent ink color (e.g., the word RED was presented in blue ink). Participants were asked to identify the color of the ink (e.g., blue). In the Switching block, in 25% of the trials, a square replaced the fixation cross. When this occurred, participants were instructed to read the word instead of identifying the color of the ink (e.g., RED). As such, within the Switching block, there were both inhibition trials in which the participant had to inhibit their reading of the word and correctly identify the color of the ink, and there were switch trials in which the participant had to switch their response mode to read the word instead of identifying the color of the ink when a square appeared before the word presented. Visual feedback on performance was presented after each trial. A practice session was completed before the acquisition run to ensure the participants understood the task. The practice consisted of a shorter version of the task. Dependent variables were reaction times (ms) and the number of errors committed (%). We also calculated two scores, inhibition score which is the result of the inhibition bloc's score minus naming blocs' performances on number or error and corresponds to pure inhibition capacities. The second, flexibility cost is equal to results of flexibility bloc's score minus inhibition blocs' performances and reflect pure flexibility capacities.

**Memory.** Verbal memory and learning abilities were assessed with the Rey Words test [51]. Participants had to learn fifteen words list in five tries. The total number of words recalled was recorded. Memory span was also assessed; the forward memory span evaluates the capacity of short-term memory and the backward memory span assesses working memory corresponding to the maximum information kept in mind and the ability to manipulate it. The dependant variable was the number of elements in the last trial.

Working memory. The 2-back was used to assess working memory. A series of numbers (0-9) were presented auditorily, and the participants were asked to respond to the stimuli that they hear 2 items back. The cognitive load was manipulated by using two different conditions: single- and dual-task, and not by manipulating. In the single-task condition, participants were instructed to perform either a self-paced walk (motor task) or the 2-back task (cognitive task). In the dual-task condition, participants performed both self-paced walking and 2-back tasks simultaneously. The numbers were pseudo-randomly ordered to ensure that there were no repeats (1-1) and no sequenced series (1-2-3). The numbers were presented through wireless headphones (RS 165, Sennheiser Electronic, Wedemark, Germany). A participant might hear the series..."0 - 4 - 1 - 5.....", once they hear..."1", they would have to say "0", and once they hear "5" they would have to say "4", always keeping in mind the digit heard two positions back. The experimenter noted the participant's responses and their accuracy. All participants were given instructions before each task condition (simple- or dual-task). The practice trial consisted of two trials for each task condition. Then, the experimental test consisted of 10 blocks organized as follows: single-task 2-back (2 blocks), single-task (self-selected pace) walking (1 block), dual-task 2- back (4 blocks), single-task (self-selected pace) walking (1 block), and single-task 2-back (2 blocks). Each block of single and dual-task 2-back consisted of 12 items. Each item was presented in 1.5 s, and there was a 1 s response interval (total trial time 2.5 s) before the next item was presented. Each block lasted 40 s with a 60-second rest period between blocks. The dependent variable was accuracy. This procedure was already used by two other teams [52], [53].

**Psychomotor speed** was assessed by the Digit Symbol Substitution Test (DSST). The participant had to associate symbols to numbers (1-9), in a table of numbers by referring to a response key. People had 120 s to draw as many symbols as possible. The dependant variable was the number of complete cells.

## Training protocol

**Exercise training (ET).** Participants completed 30 minutes of stationary bicycle (Velo-Cognitif, REV'LIM, Limoges, France) at 60 % of their theoretical maximal heart rate ( $tHR_{max}$ ). The corresponding power was estimated during the submaximal intensity exercise test. Heart rate was recorded continuously to adjust power output during the session. The choice of an exercise intensity of 60% of  $tHR_{max}$  is based on several observations suggesting that cognitive performance was improved in optimal way at this intensity [54] and fulfills the recommendations by Lauenroth et al [35].

**Cognitive training (CT).** Participants completed 30 minutes of cognitive games in PRESCO (HappyNeuron, Grenade Sur Garonne, France). Each session consisted of 4 different games (among 32 possible games), which were the same for all participants, but with a level of difficulty adapted to their cognitive capacities. An example of one session was presented elsewhere (Pellegrini-Laplagne et al, 2022) The 32 games were used during the 12-week program in order to target all cognitive domains, with the same programming for all participants (i.e. 4 different games by sessions during 8 sessions, this sequence being repeated thrice). A detail of each session was presented in the Table 1.

**Simultaneously combined exercise and cognitive training (ST).** The participants performed both programmes simultaneously, with the same exercise intensities, durations and planning of the cognitive games explained above, using a specific device (Velo-Cognitif, REV'LIM, Limoges, France). This device has been previously described by Pellegrini-Laplagne et al, (2022).

## Statistical analysis

Standard statistical methods were used for the calculation of means and standard deviations. Normal Gaussian distribution of the data was verified by the Shapiro–Wilks's test and homoscedasticity by a modified Levene Test. Repeated ANOVA measures were used to test the interaction between cognitive performances x training group (simultaneous, cognitive, or physical training). If a significant main or interaction were found, a Bonferroni post-hoc test was realized. Correlation analysis was also performed using spearman correlation. All statistical analyses were made with SPSS 17.0 and all statistical analyses with a p-value < 0,05 were considered significant. Effects sizes (ES) were also calculated with Hedges' g formula, previously described by Dupuy et al (2015), and interpreted with Cohen's scale, where  $EF \le 0.2$  (trivial), > 0.2 (small), > 0.5 (moderate), > 0.8 (large).

## **RESULTS**

## Participants

The study included 74 healthy participants aged between 56 and 73 years old. We conducted the study between January 2019 and May 2021. The most important campaign began in January 2020 and was interrupted because of the first isolation period due to the COVID-19 pandemic. This attrition explains differences in sample sizes between experimental groups. Two participants decided to withdraw from the study, because of illness or difficulties to respect the frequency of the sessions (Figure 2). The final sample of this study was 35 participants. Characteristics of the participants are presented in Table 2. There were no significant differences between groups regarding age, education level, physiological characteristics, psychological health, and cognitive status.

## Physiological assessment

We observed no effect of time on the power output at 60% of  $tHR_{max}$ , on the  $\dot{V}O_2$  at this power output and on arterial blood pressure. As shown in Table 3, we found a trend toward an interaction between groups and time [F<sub>(2,25)</sub>=2,72; p=0.08] regarding the power output at 60% of  $tHR_{max}$ , with a moderate increase in ET (ES=0.65), a small increase in ST (ES=0.30 and no difference in CT (ES=-0.14). We found an interaction between groups and time on the BRS measured during controlled breathing [F<sub>(2,25)</sub>=4,28; p = 0,025]. In this condition, BRS increased in ST (p<0.01, ES= 0.74), while it did not change in CT (ES=-0.16) or in ET (ES=-0.40). Detailed results are presented in Figure 3. We found no differences in the three other condition of the BRS test.

PFC oxygenation changes during the computerized modified Stroop test and the TMT are presented in Figures 4 and 5. We found an interaction between groups and time for  $\Delta$ [HBO<sub>2</sub>] during the flexibility task [F<sub>(228)</sub>=3,58; p=0,04]. Post hoc analysis revealed a decrease in  $\Delta$ [HBO<sub>2</sub>] for CT (p=0.03, ES= -1,38), while there were no changes in ET and ST. We also found an overall decrease in  $\Delta$ [HBO<sub>2</sub>] after training during the TMT (flexibility) test, in left, right and total PFC [F<sub>(128)</sub>=8,72, p<0,001, ES= -0,58; F<sub>(128)</sub>=6,16, p=0,01, ES= -0,71; F<sub>(128)</sub>=8,93; p<0,005, ES= -0,80]. As presented in figure 5,  $\Delta$ [HHb] remained stable both during the computerized modified Stroop test and the TMT. PFC oxygenation changes during the 2-back test are presented in figure 6. We found a main effect of training on  $\Delta$ [HBO<sub>2</sub>] in the left PFC following in dual task condition [F<sub>(129)</sub>=5,09; p=0,03, ES=0,53], but not following single task condition [F<sub>(129)</sub>=1,72; p=0,20, ES=0,45]. In right PFC,  $\Delta$ [HBO<sub>2</sub>] remains stable in both single (ES=-0,03) and dual (ES=0,08) conditions.  $\Delta$ [HHb] is also unchanged during all 2-back test.

## Cognitive assessment

As shown in Table 4, we found a main effect of time on five scores: reaction time  $[F_{(1,29)}=18,19; p=0,001; ES=-0.5]$  and errors' number  $[F_{(1,29)}=12,91; p<0,00; ES=-0.6]$  during TMT B; accuracy of flexibility  $[F_{(1,29)}=9,05; p=0,005; ES=-0.76]$  and of flexibility cost  $[F_{(1,29)}=8,49; p=0,006; ES=0.71]$  during the computerized modified Stroop test; accuracy during the single task of the 2-back test  $[F_{(1,29)}=4,78; p=0,03; ES=0.3]$ . Taking together, these results show that all groups improved both their flexibility scores and their working memory in the single task condition.

The ANOVA analysis also revealed an interaction between groups and time regarding the two scores of cognitive function. The first one concerns the forward span measure, that reflects episodic memory  $[F_{(229)}=4,22; p=0,02; ES=0,17]$ . This score did not change for CT (ES=0,25) and ST (ES=0,17) and was reduced for ET (p=0,02; ES= -1,29). We also found an interaction between groups and time for the cost of inhibition  $[F_{(228)}=6,44; p=0,004; ES= -0,02)]$ . This cost

did not change for CT (ES=0,5) and ET (ES=0,6) groups, whereas it decreased in ST group (p=0,02; ES=-0,9). Detailed results are presented in Table 4.

## Association between physiological and cognitive measures

The magnitude of change in the cost of inhibition was associated to the magnitude of change in BRS during the controlled breathing condition (r=0,40; p=0,03). As shown in Figure 7, this association indicates that participants who increased the more their BRS were those who displayed the greatest improvement in the cost of inhibition. No other correlations were found between physiological measures and other cognitive performances.

## **DISCUSSION**

The study investigated the impact of a simultaneous combination of exercise and cognitive training on several cognitive domains and some of their physiological determinants in healthy older adults, in comparison with exercise and cognitive training alone. Based on Kraft theory [27], we hypothesized that simultaneous combined training would be effective in improving executive performance, and that the magnitude of improvement would be greater than either training mode alone. Considering the role of cardiac autonomic control and cerebral oxygenation on aging-induced cognitive decline, the second purpose of this study was to assess the impact of simultaneous combined training on cerebral oxygenation and baroreflex sensitivity. We hypothesized that the improvement in executive functions was associated with the improvement of baroreflex sensitivity and / or an improvement of the brain oxygenation. Our main findings were : (1) an improvement in flexibility abilities during the computerized modified Stroop test and the TMT B, as well as an improvement in working memory during the single-task of the 2-back test, whatever the group; (2) an improvement in inhibition performance and BRS during controlled breathing that was specific to ST, the magnitude of improvement of each variable being positively associated; (3), a decrease in total  $\Delta$ [HBO<sub>2</sub>] in the CT group during the flexibility block of the computerized modified Stroop test, and for all groups during TMT B, both in the left and right PFC.

Our study confirms previous observations which showed that combined exercise and cognitive training was an effective intervention to improve cognitive performance or limit cognitive impairment in older people [9], [55]. It is worth noting that CT and ET also improved executive functions like flexibility and working memory in our study. This specific effect of each mode of training implemented separately has already been reported for ET, both in young or healthy older adults [50], [56], but also for CT in older adults [57]–[59]. The novelty of this study was the better efficiency of ST to improve executive functions when compared with either modality alone. These results are in agreement with five other studies reporting a greater executive functions enhancement after a combined exercise and cognitive training when compared with exercise training alone [9], [60]–[63]. Among these studies, only two of them were interested in aging and confirmed this result in healthy older adults [9], [55]. The protocol used by Raichlens et al. [9] was very close to the protocol used in the present study, since it consisted in three 30-min sessions per week for twelve weeks, using multi-domain cognitive games and moderate intensity continuous exercises. They reported a greater improvement following ST in a serial subtraction test during a dual-task condition (walking), thus suggesting a greater impact on divided attention, executive control, and working memory [9]. In the same way, Theill and colleagues [55] reported a greater improvement in a test involving visual memory and learning after ST when compared with either training modality alone. Similarly to the present study and the one of Raichlen et al. [9] they proposed two 40-minutes sessions per week during ten weeks of simultaneous working memory training and moderate intensity brisk walk training. So, according to our findings, they observed dual task training superiority on cognitive training alone after a combined training of independents cognitive and physical tasks. They confirmed observations made with the first experimental condition, thus suggesting that this greater improvement in executive performance could be reached with different modalities of ST.

The specific effect of ST on inhibition is in accordance with the conclusions of the recent metaanalysis of Dhir et al [64]. Indeed, the authors reported that a program combining moderate intensity continuous exercises and multi-domains cognitive tasks improved inhibitory control, particularly when proposed to healthy older adults experiencing the normal age-related cognitive decline [64]. This result may have important clinical implications since the ability to exert an inhibitory control over behavioral urges or unhelpful thoughts plays a key role in mental health and wellbeing [65]. Similarly, it is important to keep in mind that inhibition plays a key role in executive functions. According to Miyake [66], executive functions are numerous, but they are interrelated and share some underlying commonalities. In this context, inhibitory control is a key component of an efficient executive control. In fact, several studies suggested that the alteration in inhibitory processes could play a major role in the age-related decline in several cognitive domains [67]. Also, as mentioned by Boucard et al. [68], inhibition performance appears to be particularly sensitive to daily physical and cognitive activity, because of its elementary and ubiquitous properties in the construction of executive functions. This hypothesis has been validated in interventional study by Albinet et al (2016), who observed in elderly people, after a 5-month swimming training programme, a specific improvement of the inhibition process among other tested executive functions. Our results abound in the sense of this hypothesis, since inhibition was the single executive function that was sensitive to simultaneous training.

To the best of our knowledge, this is the first study that assessed BRS changes after CT, ET or ST in healthy older adults. We observed an improvement in BRS during controlled breathing that was specific to ST; the magnitude of this improvement was positively associated to the gains in inhibition. The fact that this observation was limited to controlled breathing is probably due to the fact that breathing frequency influences significantly cardiac autonomic control ( Li et al, 2018) and its reproductibility (Driscoll et al, 2000, Melo et al, 2018, Gisselman et al, 2020). Therefore, controlling breathing frequencey reduces physiological noise, improves the signal-to-noise ratio and in fine increases statistical power, which is of great importance in our study if we consider the relatively small sample size. Furthermore, although we don't have any comparison, this result is not surprising considering the positive impact of exercise training on BRS [69]. Indeed, in their narrative review [69], La Rovere et al, indicated that exercise training limited the effect of aging on BRS, as well as its consequences on blood pressure and its underlying determinants. In addition, a recent study reported that a cognitive training, involving tasks that target processing speed and attention, improved these cognitive domains as well as cardiac autonomic control (as measured by heart rate variability) in older adults suffering from amnesia and mild cognitive impairment [70]. Surprisingly, we found no changes in BRS following CT and ET. A possible explanation may be the training dose [71]. In fact, the protocol by Lin et al. [70] consisted in four 1-hour sessions per week for six weeks, which represented a 4-time higher dose than in our study. The issue of a possible dose-response relationship should be explored more specifically.

As previously mentioned, we observed a positive association between BRS and inhibition changes after ST, participants who improved the most BRS being those who improved the most the cost of inhibition. This result is in agreement with the predictions of the neuro-visceral integration model by Thayer and Lane [72], which suggests that resting cardiac autonomic

control relates to PFC activity and predicts executive performance. In fact, these authors suggest the existence of a common reciprocal inhibitory cortical-subcortical neural circuit that serves as a structural link between psychological processes such as emotion and cognition, and health-related physiological processes, and that this circuit can be indexed with cardiac autonomic control [72]. Several studies have already shown that resting BRS was associated to better cognitive performance [73], [74], and that exercise training induced improvement in executive performance was mediated by a higher cardiac autonomic control [75], [76]. As mentioned earlier, Albinet et al. (2016) observed a specific improvement in inhibition after 5 months of swimming training in elderly people. This improvement seemed to be explained and mediated by an improvement in the cardiac parasympathetic control, as measured by heart rate variability. Our results are in accordance with these observations since cardiac autonomic control is a key component of BRS.

As for BRS, this is the first study that assessed PFC oxygenation changes after CT, ET or ST in healthy older adults. Our main result was a decrease in  $\Delta$ [HBO<sub>2</sub>] in all groups during TMT B, both in the left and right PFC, together with an improvement in performance during this test. This result is in accordance with a recent study which reported that older subjects displayed a lower cerebral oxygenation ( $\Delta$ [HBO<sub>2</sub>]) during a Stroop task after aerobic training [77]. It should be kept in mind that the functional activity of the brain is related to the blood supply [78] and that neuronal activation requires an increase in cerebral blood flow and metabolism [79]. This mechanism has been illustrated by Mehagnoul-Schipper et al. [80] who reported simultaneous increases in cerebral blood volume and cerebral oxygenation during cognitive activation in older adults. It also appears that the magnitude of this adaptation is proportional to the complexity of the cognitive task [46]. Our results, and more specifically the decrease in  $\Delta$ [HBO<sub>2</sub>] after the interventions, therefore suggest that one of the benefits of CT, ET and ST is to decrease the level of cortical activation required for a specific cognitive task. This is in accordance with observations made by Voelcker-Rehage et al. [81], who reported that 12 months of aerobic and coordination training resulted in a decreased task-related activation in the PFC during a Flanker task (a measure of cognitive inhibition). Interestingly, we observed similar results in CT. This adaptation was expected, since it has been shown that cognitive training improved brain plasticity, which plays an important role in neuronal activity [82]. Nevertheless, this hypothesis of lower cortical activation after a training programme, demonstrating some form of neuronal economy, needs to be confirmed.

Many physiological mechanisms at the molecular level can explain the cognitive improvements observed in the elderly people after a physical training. A number of molecules that are considered as growth factors are released during exercise, thus promoting cerebral plasticity. For example, it is well established that molecules such as BDNF, IGF or VEGF will induce synaptogenesis, neurogenesis or angiogenesis (Stimpson et al, 2018; Cotman et Berchtold, 2002; Ploughman, 2008; Dupuy et al, 2019). All these molecular factors potentially induce structural changes in the cerebral vasculature, and contribute to an increase of the volumes of the grey nuclei, the hippocampus, and the grey and white matters. More specifically to simultaneous training, Eduard Kraft (2012) makes the following synergetic hypothesis: 1) physical activity facilitates synaptic plasticity and neurogenesis via neurotrophic factors such as BDNF or IGF-1; 2) cognitive stimulation would guide synaptic plasticity in the brain areas activated by the cognitive task, through the selection of the most active synapses 3) The combination of the two would make it possible to include new neurons and create new synapses (thanks to increased neurogenesis and synaptogenesis) in brain networks selected by cognitive activity. In other words, physical stimulation would catalyze the effect of cognitive stimulation and therefore the restructuring of the neural network. These mechanisms are probably involved in the structural changes in the neural circuits that control both baroreflex regulation and the cognitive processes involved in inhibition. Future neuroimaging studies may allow us to validate these hypotheses.

Though this study had several strengths, it was not without limitations, and the interpretation of our results requires some caution. One of the main limitations is the small sample size, which limits statistical power and in fine the generalization of our results As mentioned in Figure 2, many participants were unable to complete this study due to lockdown of the COVID. The size of this study, which involved people aged 55 and over, does not allow us to distinguish a possible specific effect in middle-aged people (55-65 years) and in the older adults (>65 years). The same comment applies to the men and women involved in this study. Future studies with a larger sample size would allow us to address these limitations. Nevertheless, the calculation of effect size by Hedge's g allows us to appreciate our results from a clinical point of view. Although the size of this study may be considered a limitation, it must kept in mind that this is the first study to compare simultaneous cognitive and physical training to physical and cognitive training alone on cognitive function in healthy older adults. This is also the first study in healthy older people to propose physiological mechanisms that allow a better understanding of cognitive outcomes. Finally, it is not excluded that the superiority of ST training when compared with ET and CT training, is dependent on the dose of training. However, in the context of two training sessions per week, DT training may improve some cognitive functions and cardiovascular indices, but not CT or ET. It would be interesting to see whether this superiotity of training is still observed when the intensity and/or frequency of training is increased in future comprehensive studies.

## **Conclusion:**

The aim of this study was to investigate the effect of a simultaneous combination of exercise and cognitive training on several cognitive domains and some of their physiological determinants in healthy older adults, in comparison with exercise and cognitive training alone. Considering the role of cardiac autonomic control and cerebral oxygenation on aging-induced cognitive decline, a second purpose of this study was to assess the impact of simultaneous combined training on cerebral oxygenation and baroreflex sensitivity. Our main findings were (1) an improvement in flexibility abilities during the computerized modified Stroop test and the TMT B, as well as an improvement in working memory during the single-task of the 2-back test, whatever the group; (2) an improvement in inhibition performance and BRS during controlled breathing that was specific to ST, the magnitude of improvement of each variable being positively associated; (3), a decrease in total HBO<sub>2</sub> in the CT group during the flexibility block of the computerized modified Stroop test, and for all groups during TMT B, both in the left and right PFC. This study brings additional evidence that simultaneous exercise and cognitive training is more efficient in improving executive performance than either training modality alone. It also provides new information regarding the mechanisms that subtend this effect and underscores the interest of the neurovisceral integrative model by Thayer and Lane (2000) to provide a theoretical framework. Future studies should investigate the impact of simultaneous cognitive and physical training on larger cohorts with more intense sessions (longer or more session per week).

#### **References:**

- .] R. S. Wilson *et al.*, "The Influence of Cognitive Decline on Well-Being in Old Age," *Psychology and Aging*, vol. 28, no. 2, pp. 304–313, 2013, doi: 10.1037/a0031196.
- Y. S. Borson, "Cognition, aging, and disabilities: Conceptual issues," *Physical Medicine and Rehabilitation Clinics of North America*, vol. 21, no. 2. pp. 375–382, May 2010. doi: 10.1016/j.pmr.2010.01.001.
- [4] A. M. Fjell and K. B. Walhovd, "Structural Brain Changes in Aging: Courses, Causes and Cognitive Consequences," 2010.
- S. Colcombe and A. F. Kramer, "Fitness effects on the cognitive function of older adults: A metaanalytic study," *Psychological Science*, vol. 14, no. 2, pp. 125–130, Mar. 2003, doi: 10.1111/1467-9280.t01-1-01430.
- <sup>5</sup>] K. D. Monahan, "NEW INVESTIGATOR AWARD OF THE NEURAL CONTROL AND AUTONOMIC REGULATION SECTION Effect of aging on baroreflex function in humans," *Am J Physiol Regul Integr Comp Physiol*, vol. 293, pp. 3–12, 2007, doi: 10.1152/ajpregu.00031.2007.-Arterial.
- 6] P. Nicolini *et al.*, "Autonomic dysfunction in mild cognitive impairment: Evidence from power spectral analysis of heart rate variability in a cross-sectional case-control study," *PLoS ONE*, vol. 9, no. 5, May 2014, doi: 10.1371/journal.pone.0096656.
- '] S. A. Schapkin and G. Freude, "Cardiovascular Costs of Working Memory Performance: Effects of Age and Performance Feedback."
- A. Zeki Al Hazzouri, M. N. Haan, Y. Deng, J. Neuhaus, and K. Yaffe, "Reduced heart rate variability is associated with worse cognitive performance in elderly Mexican Americans," *Hypertension*, vol. 63, no. 1, pp. 181–187, Jan. 2014, doi: 10.1161/HYPERTENSIONAHA.113.01888.
- <sup>1</sup>] D. A. Raichlen *et al.*, "Effects of simultaneous cognitive and aerobic exercise training on dual-task walking performance in healthy older adults: Results from a pilot randomized controlled trial," *BMC Geriatrics*, vol. 20, no. 1, Mar. 2020, doi: 10.1186/s12877-020-1484-5.
- 0] P. W. Ku, Y. te Liu, M. K. Lo, L. J. Chen, and B. Stubbs, "Higher levels of objectively measured sedentary behavior is associated with worse cognitive ability: Two-year follow-up study in community-dwelling older adults," *Experimental Gerontology*, vol. 99, pp. 110–114, Dec. 2017, doi: 10.1016/j.exger.2017.09.014.
- N. Owen, P. B. Sparling, G. N. Healy, D. W. Dunstan, and C. E. Matthews, "Sedentary behavior: Emerging evidence for a new health risk," *Mayo Clinic Proceedings*, vol. 85, no. 12. Elsevier Ltd, pp. 1138–1141, 2010. doi: 10.4065/mcp.2010.0444.
- 2] M. Prince, R. Bryce, E. Albanese, A. Wimo, W. Ribeiro, and C. P. Ferri, "The global prevalence of dementia: A systematic review and metaanalysis," *Alzheimer's and Dementia*, vol. 9, no. 1. Elsevier Inc., pp. 63-75.e2, 2013. doi: 10.1016/j.jalz.2012.11.007.
- 3] L. S. Colzato, "Editorial," *Journal of Cognitive Enhancement*, vol. 1, no. 1, pp. 1–2, Mar. 2017, doi: 10.1007/s41465-016-0001-4.
- 4] D. E. Barnes *et al.*, "The mental activity and exercise (MAX) trial: A randomized controlled trial to enhance cognitive function in older adults," *JAMA Internal Medicine*, vol. 173, no. 9, pp. 797–804, May 2013, doi: 10.1001/jamainternmed.2013.189.

- 5] A. la Rue, "Healthy Brain Aging: Role of Cognitive Reserve, Cognitive Stimulation, and Cognitive Exercises," *Clinics in Geriatric Medicine*, vol. 26, no. 1. pp. 99–111, Feb. 2010. doi: 10.1016/j.cger.2009.11.003.
- 6] S. M. Landau *et al.*, "Association of lifetime cognitive engagement and low β-amyloid deposition," *Archives of Neurology*, vol. 69, no. 5, pp. 623–629, May 2012, doi: 10.1001/archneurol.2011.2748.
- 7] M. J. Valenzuela, P. Sachdev, W. Wen, X. Chen, and H. Brodaty, "Lifespan mental activity predicts diminished rate of hippocampal atrophy," *PLoS ONE*, vol. 3, no. 7, Jul. 2008, doi: 10.1371/journal.pone.0002598.
- 8] J. Reijnders, C. van Heugten, and M. van Boxtel, "Cognitive interventions in healthy older adults and people with mild cognitive impairment: A systematic review," *Ageing Research Reviews*, vol. 12, no. 1. pp. 263–275, Jan. 2013. doi: 10.1016/j.arr.2012.07.003.
- 9] L. M. J. Sanders, T. Hortobágyi, S. la B. van Gemert, E. A. van der Zee, and M. J. G. van Heuvelen, "Dose-response relationship between exercise and cognitive function in older adults with and without cognitive impairment: A systematic review and meta-analysis," *PLoS ONE*, vol. 14, no. 1. Public Library of Science, Jan. 01, 2019. doi: 10.1371/journal.pone.0210036.
- [0] F. T. Chen, J. L. Etnier, K. H. Chan, P. K. Chiu, T. M. Hung, and Y. K. Chang, "Effects of Exercise Training Interventions on Executive Function in Older Adults: A Systematic Review and Meta-Analysis," *Sports Medicine*, vol. 50, no. 8. Springer, pp. 1451–1467, Aug. 01, 2020. doi: 10.1007/s40279-020-01292-x.
- [1] S. Ludyga, M. Gerber, S. Brand, E. Holsboer-Trachsler, and U. Pühse, "Acute effects of moderate aerobic exercise on specific aspects of executive function in different age and fitness groups: A metaanalysis," *Psychophysiology*, vol. 53, no. 11, pp. 1611–1626, Nov. 2016, doi: 10.1111/psyp.12736.
- [2] J. M. Northey, N. Cherbuin, K. L. Pumpa, D. J. Smee, and B. Rattray, "Exercise interventions for cognitive function in adults older than 50: A systematic review with meta-Analysis," *British Journal* of Sports Medicine, vol. 52, no. 3. BMJ Publishing Group, pp. 154–160, Feb. 01, 2018. doi: 10.1136/bjsports-2016-096587.
- [3] A. S. Buchman, P. A. Boyle, L. Yu, R. C. Shah, R. S. Wilson, and D. A. Bennett, "Total daily physical activity and the risk of AD and cognitive decline in older adults," *Neurology*, vol. 78, no. 17, pp. 1323– 1329, Apr. 2012, doi: 10.1212/WNL.0b013e3182535d35.
- [4] C. W. Cotman, N. C. Berchtold, and L. A. Christie, "Exercise builds brain health: key roles of growth factor cascades and inflammation," *Trends in Neurosciences*, vol. 30, no. 9. Elsevier Ltd, pp. 464– 472, 2007. doi: 10.1016/j.tins.2007.06.011.
- [5] C. M. Stillman and K. I. Erickson, "Physical activity as a model for health neuroscience," Annals of the New York Academy of Sciences, 2018, doi: 10.1111/nyas.13669.
- C. M. Stillman, I. Esteban-Cornejo, B. Brown, C. M. Bender, and K. I. Erickson, "Effects of Exercise on Brain and Cognition Across Age Groups and Health States," *Trends in Neurosciences*, vol. 43, no. 7. Elsevier Ltd, pp. 533–543, Jul. 01, 2020. doi: 10.1016/j.tins.2020.04.010.
- E. Kraft, "Cognitive function, physical activity, and aging: Possible biological links and implications for multimodal interventions," *Aging, Neuropsychology, and Cognition*, vol. 19, no. 1–2, pp. 248–263, Jan. 2012, doi: 10.1080/13825585.2011.645010.
- [8] P. D. Bamidis *et al.*, "A review of physical and cognitive interventions in aging," *Neuroscience and Biobehavioral Reviews*, vol. 44. Elsevier Ltd, pp. 206–220, 2014. doi: 10.1016/j.neubiorev.2014.03.019.

- [9] K. Fabel, S. A. Wolf, D. Ehninger, H. Babu, P. Leal-Galicia, and G. Kempermann, "Additive effects of physical exercise and environmental enrichment on adult hippocampal neurogenesis in mice," *Frontiers in Neuroscience*, vol. 3, no. NOV, 2009, doi: 10.3389/neuro.22.002.2009.
- [0] A. M. Smith, K. M. Spiegler, B. Sauce, C. D. Wass, T. Sturzoiu, and L. D. Matzel, "Voluntary aerobic exercise increases the cognitive enhancing effects of working memory training," *Behavioural Brain Research*, vol. 256, pp. 626–635, Nov. 2013, doi: 10.1016/j.bbr.2013.09.012.
- <sup>51</sup> J. L. Tait, R. L. Duckham, C. M. Milte, L. C. Main, and R. M. Daly, "Influence of sequential vs. simultaneous dual-task exercise training on cognitive function in older adults," *Frontiers in Aging Neuroscience*, vol. 9, no. NOV. Frontiers Media S.A., Nov. 07, 2017. doi: 10.3389/fnagi.2017.00368.
- <sup>12</sup> F. Herold, D. Hamacher, L. Schega, and N. G. Müller, "Thinking while moving or moving while thinking concepts of motor-cognitive training for cognitive performance enhancement," *Frontiers in Aging Neuroscience*, vol. 10, no. AUG, Aug. 2018, doi: 10.3389/fnagi.2018.00228.
- [3] P. Fissler, O. Küster, W. Schlee, and I. T. Kolassa, "Novelty interventions to enhance broad cognitive abilities and prevent dementia: Synergistic approaches for the facilitation of positive plastic change," in *Progress in Brain Research*, vol. 207, Elsevier B.V., 2013, pp. 403–434. doi: 10.1016/B978-0-444-63327-9.00017-5.
- [4] L. L. F. Law, F. Barnett, M. K. Yau, and M. A. Gray, "Effects of combined cognitive and exercise interventions on cognition in older adults with and without cognitive impairment: A systematic review," *Ageing Research Reviews*, vol. 15, no. 1. Elsevier Ireland Ltd, pp. 61–75, 2014. doi: 10.1016/j.arr.2014.02.008.
- <sup>5</sup>] A. Lauenroth, A. E. Ioannidis, and B. Teichmann, "Influence of combined physical and cognitive training on cognition: A systematic review," *BMC Geriatrics*, vol. 16, no. 1. BioMed Central Ltd., Jul. 18, 2016. doi: 10.1186/s12877-016-0315-1.
- [6] C. Joubert and H. Chainay, "Aging brain: The effect of combined cognitive and physical training on cognition as compared to cognitive and physical training alone – A systematic review," *Clinical Interventions in Aging*, vol. 13. Dove Medical Press Ltd., pp. 1267–1301, 2018. doi: 10.2147/CIA.S165399.
- [7] X. Zhu, S. Yin, M. Lang, R. He, and J. Li, "The more the better? A meta-analysis on effects of combined cognitive and physical intervention on cognition in healthy older adults," *Ageing Research Reviews*, vol. 31. Elsevier Ireland Ltd, pp. 67–79, Nov. 01, 2016. doi: 10.1016/j.arr.2016.07.003.
- [8] H. M. Gavelin *et al.*, "Combined physical and cognitive training for older adults with and without cognitive impairment: A systematic review and network meta-analysis of randomized controlled trials," *Ageing Research Reviews*, vol. 66. Elsevier Ireland Ltd, Mar. 01, 2021. doi: 10.1016/j.arr.2020.101232.
- [9] W. Guo, M. Zang, S. Klich, A. Kawczyński, M. Smoter, and B. Wang, "Effect of combined physical and cognitive interventions on executive functions in older adults: A meta-analysis of outcomes," *International Journal of Environmental Research and Public Health*, vol. 17, no. 17, pp. 1–19, Sep. 2020, doi: 10.3390/ijerph17176166.
- [0] B. Wollesen, A. Wildbredt, K. S. van Schooten, M. L. Lim, and K. Delbaere, "The effects of cognitivemotor training interventions on executive functions in older people: A systematic review and metaanalysis," *European Review of Aging and Physical Activity*, vol. 17, no. 1. BioMed Central Ltd, Jul. 02, 2020. doi: 10.1186/s11556-020-00240-y.

- [1] F. Gheysen *et al.*, "Physical activity to improve cognition in older adults: Can physical activity programs enriched with cognitive challenges enhance the effects? A systematic review and meta-analysis," *International Journal of Behavioral Nutrition and Physical Activity*, vol. 15, no. 1. BioMed Central Ltd., Jul. 04, 2018. doi: 10.1186/s12966-018-0697-x.
- [42] M. A. Fiatarone Singh *et al.*, "The Study of Mental and Resistance Training (SMART) Study-Resistance Training and/or Cognitive Training in Mild Cognitive Impairment: A Randomized, Double-Blind, Double-Sham Controlled Trial," *Journal of the American Medical Directors Association*, vol. 15, no. 12, pp. 873–880, Dec. 2014, doi: 10.1016/j.jamda.2014.09.010.
- [3] R. L. Gellish, B. R. Goslin, R. E. Olson, A. McDonald, G. D. Russi, and V. K. Moudgil, "Longitudinal modeling of the relationship between age and maximal heart rate," *Medicine and Science in Sports and Exercise*, vol. 39, no. 5, pp. 822–829, May 2007, doi: 10.1097/mss.0b013e31803349c6.
- "Assessment of cardiovascular autonomic nervous system during vigorous tests," *Medecine des Maladies Metaboliques*, vol. 2, no. 1, pp. 68–73, 2008, doi: 10.1016/S1957-2557(08)70023-3.
- [5] R. Goenarjo *et al.*, "Cerebral oxygenation reserve: The relationship between physical activity level and the cognitive load during a stroop task in healthy young males," *International Journal of Environmental Research and Public Health*, vol. 17, no. 4, Feb. 2020, doi: 10.3390/ijerph17041406.
- [6] N. F. Agbangla, M. Audiffren, J. Pylouster, and C. T. Albinet, "Working memory, cognitive load and cardiorespiratory fitness: Testing the CRUNCH model with near-infrared spectroscopy," *Brain Sciences*, vol. 9, no. 2, 2019, doi: 10.3390/brainsci9020038.
- [47] Z. S. Nasreddine *et al.*, "The Montreal Cognitive Assessment, MoCA: A Brief Screening Tool For Mild Cognitive Impairment." [Online]. Available: www.mocatest.
- [8] O. Dupuy *et al.*, "Higher levels of cardiovascular fitness are associated with better executive function and prefrontal oxygenation in younger and older women," *Frontiers in Human Neuroscience*, vol. 9, no. FEB, Feb. 2015, doi: 10.3389/fnhum.2015.00066.
- [9] R. Goenarjo, O. Dupuy, S. Fraser, N. Berryman, A. Perrochon, and L. Bosquet, "Cardiorespiratory fitness and prefrontal cortex oxygenation during Stroop task in older males," *Physiology & Behavior*, vol. 242, p. 113621, Dec. 2021, doi: 10.1016/j.physbeh.2021.113621.
- [0] S. Mekari *et al.*, "Effect of high intensity interval training compared to continuous training on cognitive performance in young healthy adults: A pilot study," *Brain Sciences*, vol. 10, no. 2, Feb. 2020, doi: 10.3390/brainsci10020081.
- [1] A. Rey, "L'examin clinique en psychologie," Presses Universitaires de France, 1958.
- <sup>52</sup> R. Goenarjo, O. Dupuy, S. Fraser, A. Perrochon, N. Berryman, and L. Bosquet, "Cardiorespiratory fitness, blood pressure, and cerebral oxygenation during a dual-task in healthy young males," *Behavioural Brain Research*, vol. 380, Feb. 2020, doi: 10.1016/j.bbr.2019.112422.
- [3] S. A. Fraser, O. Dupuy, P. Pouliot, F. Lesage, and L. Bherer, "Comparable cerebral oxygenation patterns in younger and older adults during dual-task walking with increasing load," *Frontiers in Aging Neuroscience*, vol. 8, no. OCT, Oct. 2016, doi: 10.3389/fnagi.2016.00240.
- [4] K. Lambourne and P. Tomporowski, "The effect of exercise-induced arousal on cognitive task performance: A meta-regression analysis," *Brain Research*, vol. 1341. pp. 12–24, Jun. 23, 2010. doi: 10.1016/j.brainres.2010.03.091.

- <sup>55</sup>] N. Theill, V. Schumacher, R. Adelsberger, M. Martin, and L. Jäncke, "Effects of simultaneously performed cognitive and physical training in older adults," *BMC Neuroscience*, vol. 14, Sep. 2013, doi: 10.1186/1471-2202-14-103.
- <sup>6</sup>] D. Predovan, S. A. Fraser, M. Renaud, and L. Bherer, "The effect of three months of aerobic training on stroop performance in older adults," *Journal of Aging Research*, vol. 2012, 2012, doi: 10.1155/2012/269815.
- [7] F. D. Wolinsky, F. W. Unverzagt, D. M. Smith, R. Jones, A. Stoddard, and S. L. Tennstedt, "The ACTIVE Cognitive Training Trial and Health-Related Quality of Life: Protection That Lasts for 5 Years Downloaded from," 2006. [Online]. Available: http://biomedgerontology.oxfordjournals.org/
- [8] A. Corbett *et al.*, "The Effect of an Online Cognitive Training Package in Healthy Older Adults: An Online Randomized Controlled Trial," *Journal of the American Medical Directors Association*, vol. 16, no. 11, pp. 990–997, Nov. 2015, doi: 10.1016/j.jamda.2015.06.014.
- <sup>59</sup> C. Basak, S. Qin, and M. A. O'Connell, "Differential effects of cognitive training modules in healthy aging and mild cognitive impairment: A comprehensive meta-analysis of randomized controlled trials.," *Psychology and Aging*, vol. 35, no. 2, pp. 220–249, 2020, doi: 10.1037/pag0000442.
- 10] H. Yokoyama *et al.*, "The effect of cognitive-motor dual-task training on cognitive function and plasma amyloid β peptide 42/40 ratio in healthy elderly persons: A randomized controlled trial," *BMC Geriatrics*, vol. 15, no. 1, May 2015, doi: 10.1186/s12877-015-0058-4.
- M. Hiyamizu, S. Morioka, K. Shomoto, and T. Shimada, "Effects of dual task balance training on dual task performance in elderly people: A randomized controlled trial," *Clinical Rehabilitation*, vol. 26, no. 1, pp. 58–67, Jan. 2012, doi: 10.1177/0269215510394222.
- [52] S. Falbo, G. Condello, L. Capranica, R. Forte, and C. Pesce, "Effects of Physical-Cognitive Dual Task Training on Executive Function and Gait Performance in Older Adults: A Randomized Controlled Trial," *BioMed Research International*, vol. 2016, 2016, doi: 10.1155/2016/5812092.
- [3] J. León, A. Ureña, M. J. Bolaños, A. Bilbao, and A. Oña, "A combination of physical and cognitive exercise improves reaction time in persons 61-84 years old," *Journal of Aging and Physical Activity*, vol. 23, no. 1, pp. 72–77, Jan. 2015, doi: 10.1123/JAPA.2012-0313.
- [4] S. Dhir, W. P. Teo, S. R. Chamberlain, K. Tyler, M. Yücel, and R. A. Segrave, "The Effects of Combined Physical and Cognitive Training on Inhibitory Control: A Systematic Review and Meta-Analysis," *Neuroscience and Biobehavioral Reviews*, vol. 128. Elsevier Ltd, pp. 735–748, Sep. 01, 2021. doi: 10.1016/j.neubiorev.2021.07.008.
- [55] A. Bari and T. W. Robbins, "Inhibition and impulsivity: Behavioral and neural basis of response control," *Progress in Neurobiology*, vol. 108. pp. 44–79, Sep. 2013. doi: 10.1016/j.pneurobio.2013.06.005.
- [6] A. Miyake, N. P. Friedman, M. J. Emerson, A. H. Witzki, A. Howerter, and T. D. Wager, "The Unity and Diversity of Executive Functions and Their Contributions to Complex 'Frontal Lobe' Tasks: A Latent Variable Analysis," *Cognitive Psychology*, vol. 41, no. 1, pp. 49–100, 2000, doi: 10.1006/cogp.1999.0734.
- [7] L. Hasher and R. T. Zacks, "WORKING MEMORY, COMPREHENSION, AND AGING: A REVIEW AND A NEW VIEW."
- [6] G. K. Boucard, C. T. Albinet, C. A. Bouquet, D. Clarys, M. Audiffren, and A. Bugaiska, "Impact of Physical Activity on Executive Functions in Aging: A Selective Effect on Inhibition Among Old Adults," 2012.

- M. T. la Rovere and G. D. Pinna, "Beneficial effects of physical activity on baroreflex control in the elderly," *Annals of Noninvasive Electrocardiology*, vol. 19, no. 4. Blackwell Publishing Inc., pp. 303– 310, 2014. doi: 10.1111/anec.12170.
- <sup>'0]</sup> F. v. Lin *et al.*, "Processing speed and attention training modifies autonomic flexibility: A mechanistic intervention study," *NeuroImage*, vol. 213, Jun. 2020, doi: 10.1016/j.neuroimage.2020.116730.
- <sup>'1</sup>] P. D. Bamidis *et al.*, "Gains in cognition through combined cognitive and physical training: The role of training dosage and severity of neurocognitive disorder," *Frontiers in Aging Neuroscience*, vol. 7, no. JUL, 2015, doi: 10.3389/fnagi.2015.00152.
- <sup>'2</sup>] J. F. Thayer and R. D. Lane, "A model of neurovisceral integration in emotion regulation and dysregulation," 2000. [Online]. Available: www.elsevier.com/locate/jad
- '3] T. Tarumi *et al.*, "Central artery stiffness, baroreflex sensitivity, and brain white matter neuronal fiber integrity in older adults," *NeuroImage*, vol. 110, pp. 162–170, Apr. 2015, doi: 10.1016/j.neuroimage.2015.01.041.
- '4] M. saint Martin, E. Sforza, C. Thomas-Anterion, J. C. Barthélémy, and F. Roche, "Baroreflex sensitivity, vascular risk factors, and cognitive function in a healthy elderly population: The PROOF cohort," *Journal of the American Geriatrics Society*, vol. 61, no. 12, pp. 2096–2102, Dec. 2013, doi: 10.1111/jgs.12548.
- '5] A. Abou-Dest, C. T. Albinet, G. Boucard, and M. Audiffren, "Swimming as a positive moderator of cognitive aging: A cross-sectional study with a multitask approach," *Journal of Aging Research*, vol. 2012, 2012, doi: 10.1155/2012/273185.
- <sup>'6</sup>] C. T. Albinet, G. Boucard, C. A. Bouquet, and M. Audiffren, "Increased heart rate variability and executive performance after aerobic training in the elderly," *European Journal of Applied Physiology*, vol. 109, no. 4, pp. 617–624, Jul. 2010, doi: 10.1007/s00421-010-1393-y.
- <sup>'7</sup>] C. Coetsee and E. Terblanche, "Cerebral oxygenation during cortical activation: the differential influence of three exercise training modalities. A randomized controlled trial," *European Journal of Applied Physiology*, vol. 117, no. 8, pp. 1617–1627, Aug. 2017, doi: 10.1007/s00421-017-3651-8.
- C. Voelcker-Rehage and C. Niemann, "Structural and functional brain changes related to different types of physical activity across the life span," *Neuroscience and Biobehavioral Reviews*, vol. 37, no. 9. pp. 2268–2295, Nov. 2013. doi: 10.1016/j.neubiorev.2013.01.028.
- <sup>'9</sup>] M. H. Davenport *et al.*, "Cerebrovascular Reserve: The Link Between Fitness and Cognitive Function?," 2012. [Online]. Available: www.acsm-essr.org
- [0] D. J. Mehagnoul-Schipper *et al.*, "Simultaneous measurements of cerebral oxygenation changes during brain activation by near-infrared spectroscopy and functional magnetic resonance imaging in healthy young and elderly subjects," *Human Brain Mapping*, vol. 16, no. 1, pp. 14–23, 2002, doi: 10.1002/hbm.10026.
- [1] C. Voelcker-Rehage, B. Godde, and U. M. Staudinger, "Cardiovascular and coordination training differentially improve cognitive performance and neural processing in older adults," *Frontiers in Human Neuroscience*, no. MARCH, 2011, doi: 10.3389/fnhum.2011.00026.
- [2] S. B. Chapman *et al.*, "Neural mechanisms of brain plasticity with complex cognitive training in healthy seniors," *Cerebral Cortex*, vol. 25, no. 2, pp. 396–405, Feb. 2015, doi: 10.1093/cercor/bht234.

ew references added:

- Driscoll D, Dicicco G. The effects of metronome breathing on the variability of autonomic activity measurements. J Manipulative Physiol Ther. 2000 Nov-Dec;23(9):610-4. doi: 10.1067/mmt.2000.110944. PMID: 11145801.
- Li C, Chang Q, Zhang J, Chai W. Effects of slow breathing rate on heart rate variability and arterial baroreflex sensitivity in essential hypertension. Medicine (Baltimore). 2018 May;97(18):e0639. doi: 10.1097/MD.00000000010639. PMID: 29718876; PMCID: PMC6392805.
- Melo HM, Martins TC, Nascimento LM, Hoeller AA, Walz R, Takase E. Ultra-short heart rate variability recording reliability: The effect of controlled paced breathing. Ann Noninvasive Electrocardiol. 2018 Sep;23(5):e12565. doi: 10.1111/anec.12565. Epub 2018 Jun 4. PMID: 29863781; PMCID: PMC6931441.
- Gisselman AS, D'Amico M, Smoliga JM. Optimizing Intersession Reliability of Heart Rate Variability-The Effects of Artifact Correction and Breathing Type. J Strength Cond Res. 2020 Nov;34(11):3199-3207. doi: 10.1519/JSC.00000000002258. PMID: 33105371.
- Stimpson N. J., Davison G., Javadi A.-H. (2018). Joggin' the Noggin: Towards a Physiological Understanding of Exercise-Induced Cognitive Benefits. *Neurosci. Biobehav. Rev.* 88 177–186. 10.1016/j.neubiorev.2018.03.018
- Dupuy O., Goenarjo R., Fraser S. A., Bherer L., Bosquet L. (2019). Master Athletes and cognitive performance: what are the potential explanatory neurophysiological mechanisms? *Mov. Sport Sci.* 104 55–67. 10.1051/sm/2019023
- Cotman C. W., Berchtold N. C. (2002). Exercise: a behavioral intervention to enhance brain health and plasticity. *Trends Neurosci*. 25 295–301. 10.1016/s0166-2236(02)02143-4
- Ploughman M. (2008). Exercise is brain food: The effects of physical activity on cognitive function. *Dev. Neurorehabil.* 11 236–240. 10.1080/17518420801997007
- Sahathevan R, Brodtmann A, Donnan GA (2012) Dementia, stroke, and vascular risk factors; a review. Int J Stroke 7:61–73
- Ogoh S. Relationship between cognitive function and regulation of cerebral blood flow. J Physiol Sci. 2017 May;67(3):345-351. doi: 10.1007/s12576-017-0525-0. Epub 2017 Feb 3. PMID: 28155036.
- de Aquino Lemos V, Antunes HK, dos Santos RV, Lira FS, Tufik S, de Mello MT (2012) High altitude exposure impairs sleep patterns, mood, and cognitive functions. Psychophysiology 49:1298–1306
- LiuK,SunG,LiB,JiangQ,YangX,LiM,LiL,QianS,ZhaoL, Zhou Z, von Deneen KM, Liu Y (2013) The impact of passive hyperthermia on human attention networks: an fMRI study. Behav Brain Res 243:220–230
- SunG,YangX,JiangQ,LiuK,LiB,LiL,ZhaoL,LiM(2012), Hyperthermia impairs the executive function using the Attention Network Test. Int J Hyperthermia 28:621–626
- Virues-Ortega J, Buela-Casal G, Garrido E, Alcazar B (2004) Neuropsychological functioning associated with high-altitude exposure. Neuropsychol Rev 14:197–224
- Nakai M (1985) An increase in cerebral blood flow elicited by electrical stimulation of the solitary nucleus in rats with cervi- cal cordotomy and vagotomy. Jpn J Physiol 35(1):57–70
- Nakai M, Iadecola C, Reis DJ (1982) Global cerebral vasodi- lation by stimulation of rat fastigial cerebellar nucleus. Am J Physiol 243(2):H226–H235
- Iadecola C, Nakai M, Mraovitch S, Ruggiero DA, Tucker LW, Reis DJ (1983) Global increase in cerebral metabolism and blood flow produced by focal electrical stimulation of dorsal medullary reticular formation in rat. Brain Res 272(1):101–114
- Ogoh S, Tarumi T. Cerebral blood flow regulation and cognitive function: a role of arterial baroreflex function. J Physiol Sci. 2019 Nov;69(6):813-823. doi: 10.1007/s12576-019-00704-6. Epub 2019 Aug 23. PMID: 31444691.
- Ding, K.; Tarumi, T.; Wang, C.; Vernino, S.; Zhang, R.; Zhu, D.C. Central autonomic network functional connectivity: correlation with baroreflex function and cardiovascular variability in older adults. *BRAIN STRUCTURE & FUNCTION* **2020**, *225*, 1575-1585, doi:10.1007/s00429-020-02075-w.
- Kimmerly, D.S. A review of human neuroimaging investigations involved with central autonomic regulation of baroreflex-mediated cardiovascular control. *AUTONOMIC NEUROSCIENCE-BASIC & CLINICAL* 2017, 207, 10-21, doi:10.1016/j.autneu.2017.05.008.
- Bertinieri G, Di Rienzo M, Caval- lazzi A, Ferrari AU, Pedotti A, Mancia G : A new approach to analysis of the arterial baroreflex. J Hypertens 1985 ; 3 : S79-S81.
- Guelen, Iljaa; Westerhof, Berend Ea; van der Sar, Gertrude La; van Montfrans, Gert Ab; Kiemeneij, Ferdinandc; Wesseling, Karel Ha; Bos, Willem Jan WdValidation of brachial artery pressure

reconstruction from finger arterial pressure, Journal of Hypertension: July 2008 - Volume 26 - Issue 7 - p 1321-1327 doi: 10.1097/HJH.0b013e3282fe1d28

- Yamamoto Y, Hughson RL. Coarse- graining spectral analysis : new method for studying heart rate variability, J Appl Physiol 1991 ; 71 (3) : 1143- 50.
- Westerhof BE, Gisolf J, Stok WJ, Wesseling KH, Karemaker JM. Time-domain cross-correlation baroreflex sensitivity: performance on the EUROBAVAR data set. J Hypertens. 2004 Jul;22(7):1371-80. doi: 10.1097/01.hjh.0000125439.28861.ed. PMID: 15201554.
- La Rovere MT, Pinna GD, Raczak G. Baroreflex sensitivity: measurement and clinical implications. Ann Noninvasive Electrocardiol. 2008 Apr;13(2):191-207. doi: 10.1111/j.1542-474X.2008.00219.x. PMID: 18426445; PMCID: PMC6931942.
- Agbangla NF, Audiffren M, Pylouster J, Albinet CT. Working Memory, Cognitive Load andCardiorespiratory Fitness: Testing the CRUNCHModel with Near-Infrared Spectroscopy. Brain Sci. 2019 Feb 9;9(2):38. doi: 10.3390/brainsci9020038. PMID: 30744137; PMCID: PMC6406418.
- Pellegrini-Laplagne M, Dupuy O, Sosner P, Bosquet L. Acute Effect of a Simultaneous Exercise and Cognitive Task on Executive Functions and Prefrontal Cortex Oxygenation in Healthy Older Adults. Brain Sci. 2022 Mar 28;12(4):455. doi: 10.3390/brainsci12040455. PMID: 35447987; PMCID: PMC9027217.
- Goenarjo R, Dupuy O, Fraser S, Perrochon A, Berryman N, Bosquet L. Cardiorespiratory fitness, blood pressure, and cerebral oxygenation during a dual-task in healthy young males. Behav Brain Res. 2020 Feb 17;380:112422. doi: 10.1016/j.bbr.2019.112422. Epub 2019 Dec 11. PMID: 31837344.
- Saint Martin, M.; Roche, F.; Thomas-Anterion, C.; Barthelemy, J.C.; Sforza, E.; Cerebrovasc, P.R.C. Eight-Year Parallel Change in Baroreflex Sensitivity and Memory Function in a Sample of Healthy Older Adults. *JOURNAL OF THE AMERICAN GERIATRICS SOCIETY* **2015**, *63*, 270-275, doi:10.1111/jgs.13252
- Saint Martin, M.; Sforza, E.; Thomas-Anterion, C.; Barthelemy, J.C.; Roche, F.; Grp, P.S. Baroreflex Sensitivity, Vascular Risk Factors, and Cognitive Function in a Healthy Elderly Population: The PROOF Cohort. *JOURNAL OF THE AMERICAN GERIATRICS SOCIETY* **2013**, *61*, 2096-2102, doi:10.1111/jgs.12548.
- Dupuy, O.; Gauthier, C.J.; Fraser, S.A.; Desjardins-Crepeau, L.; Desjardins, M.; Mekary, S.; Lesage, F.; Hoge, R.D.; Pouliot, P.; Bherer, L. Higher levels of cardiovascular fitness are associated with better executive function and prefrontal oxygenation in younger and older women. *FRONTIERS IN HUMAN NEUROSCIENCE* **2015**, *9*, doi:10.3389/fnhum.2015.00066.
- Mekari S, Fraser S, Bosquet L, Bonnéry C, Labelle V, Pouliot P, Lesage F, Bherer L. The relationship between exercise intensity, cerebral oxygenation and cognitive performance in young adults. Eur J Appl Physiol. 2015 Oct;115(10):2189-97. doi: 10.1007/s00421-015-3199-4. Epub 2015 Jun 11. PMID: 26063061.
- Tarumi, T.; de Jong, D.L.K.; Zhu, D.C.; Tseng, B.Y.; Liu, J.; Hill, C.; Riley, J.; Womack, K.B.; Kerwin, D.R.; Lu, H., et al. Central artery stiffness, baroreflex sensitivity, and brain white matter neuronal fiber integrity in older adults. *NEUROIMAGE* **2015**, *110*, 162-170, doi:10.1016/j.neuroimage.2015.01.041.

**cknowledgment:** This study was supported in part by the European Union and the New Aquitaine region through the Habisan program (CPER- FEDER)

# **Figure Captions:**

Figure 1: Velo-cognitif® device.



#### Figure 2: Flow chart of the study.





BRS, Baroreflex sensitivity; ET, Exercise training; CT, Cognitive training; ST, Simultaneous training \* Significant interaction Time x Group, p < 0.05



Figure 4: Effect of simultaneous, physical and cognitive training on oxyheamoglobin during Stroop and Trail test.  $\Delta$  [HbO2], Oxyhemoglobin; ET, Exercise training; CT, Cognitive training; ST, Simultaneous training \*\* = p < 0.01



Figure 5: Effect of simultaneous, physical and cognitive training on deoxyheamoglobin during Stroop and Trail test.  $\Delta$ [HHb], Deoxyhemoglobin; ET, Exercise training; CT, Cognitive training; ST, Simultaneous training \* = p < 0.05; \*\* = p < 0.01



Figure 6: Effect of simultaneous, physical and cognitive training on cerebral oxygenation during 2-Back test in single and dual-task condition. For one group, first column corresponds to pre-test and second column to post-test.  $\Delta$ [HbO2], Oxyhemoglobin;  $\Delta$ [HHb], Deoxyhemoglobin; ET, Exercise training; CT, Cognitive training; ST, Simultaneous training; \* = p<0,05.



*Figure 7: Correlation between baroreflex sensitivity and inhibition performance changes after training. BRS, Baroreflex sensitivity* 



1 adie 1 : Pla	anning of cognitive tr	aining on 12 weeks (2 Game 1	z sessions per week) Game 2	Game 3	Game 4	
Session 1 / 13	Stimulated cognitive function	Language and vocabulary	Language and vocabulary	Language and vocabulary	Processing speed	
	Name of the Game‡	Mot coupés	Videz votre sac	Que d'accros dans cette histoire	Haute tension	
Session 2 / 14	Stimulated cognitive function	Auditory memory	Auditory memory	Verbal memory	Language and reasoning	
	Name of the Game	Vous avez un message	Mot, où êtes vous	Mémoire d'éléphant	Dechiffrement	
Session 3 /	Stimulated cognitive function	Language and reasoning	Visual-spatial skills	Visual-spatial skills	Language and vocabulary	
15	Name of the Game	Menez l'enquête	Tours de main	Attention ça tourne	Broderies	
Session 4 /	Stimulated cognitive function	Mémoire visuelle	Mémoire visuelle	Mémoire visuelle	Language and reasoning	
10	Name of the Game	Formes et couleurs	Jeux du blason	Images déplacées	écrire dans les étoiles	
Session 5 /	Stimulated cognitive function	Processing speed	Attention	Attention	Calcul (+attention)	
17	Name of the Game	Attrapez la coccinelle	Attention concentrez-vous	Cherchez l'intrus	mettez de l'ordre dans ces comptes	
Session 6 / 18	Stimulated cognitive function	Visual and verbal memory	Visual and verbal memory	Visual and verbal memory	Calcul	
	Name of the Game	Le tours du monde en 80 voyages	On se connait	Garçon svp	Au bout du compte	
Session 7 /	Stimulated cognitive function	Calcul	Executive function	Executive function	Auditory memory	
19	Name of the Game	A vous de compter	Tours de Hanoï	Vive l'alternance	chant d'oiseau	
Session 8 / 20	Stimulated cognitive function	Mémoire Spatiale	Mémoire Spatiale	Mémoire Spatiale	Executive function	
	Name of the Game	Retrouvez votre chemin	Objet où êtes vous	Mémorisez par sous ensemble	N back	
Session 9 /	Stimulated cognitive function	Language and vocabulary	Language and vocabulary	Language and vocabulary	Processing speed	
21	Name of the Game	Mot coupés	Videz votre sac	Que d'accros dans cette histoire	Haute tension	
Session 10 /	Stimulated cognitive function	Auditory memory	Verbal memory	Verbal memory	Language and reasoning	
22	Name of the Game	Vous avez un message	Mot, où êtes vous	Mémoire d'éléphant	Dechiffrement	
Session 11 /	Stimulated cognitive function	Language and reasoning	Visual-spatial skills	Visual-spatial skills	Language and vocabulary	
23	Name of the Game	Menez l'enquête	Tours de main	Attention ça tourne	Broderies	
Session 12 /	Stimulated cognitive function	Visual memory	Visual memory	Visual memory	Language and reasoning	
24	Name of the Game	Formes et couleurs	Jeux du blason	Images déplacées	écrire dans les étoiles	
‡: The name	es of the game are pre	sented in French lang	guage since are issue	from Software Happy	Neuron®	

		Overall (n=35)	ET (n=7)	CT (n=11)	ST (n=17)	p- value
6	Women	19	4	5	10	-
Sex	Men	16	3	6	7	-
Age (years)	)	$63.9\pm4.4$	$63.7\pm5.1$	$63.1\pm2.7$	$64.6\pm5.0$	0.47
Height (cm	)	$167.5\pm8.8$	$167.3\pm10.5$	$168.2\pm10.0$	$167.1\pm7.8$	0.84
Weight (Kg	g)	$66.9 \pm 11.2$	$65.8 \pm 11.8$	68.1 ± 11.3	$66.5 \pm 11.5$	0.88
Body fat (k	Kg)	$18.2\pm4.9$	$16.1 \pm 2.8$	$17.7\pm4.6$	$19.3\pm5.6$	0.58
Lean body	mass (Kg)	$48.8\pm9.9$	$49.6\pm10.4$	$50.2 \pm \! 10.7$	$47.4\pm9.7$	0.82
Education	level†	17.5±2.8	17.2±1.8	16.6±2.6	18.2±3.1	0.50
BDI		$0.96 \pm 1.20$	$0.83\pm0.75$	$0.82 \pm 1.25$	$1.13\pm1.35$	0.70
MOCA		$27.7\pm1.6$	$28.0\pm1.4$	$27.3\pm1.7$	$27.8\pm1.5$	0.74

Table 2 : Characteristics of the participants

BDI: Beck's Depression Inventory, MoCA: Montreal Cognitive Assessment, ET, Exercise training; CT, Cognitive training; ST, Simultaneous training; †: The level of education was calculated according to the number of years of schooling. (Kindergarten, primary, secondary, high school and university).

	Overall (n=35)			ET (n=7)			CT (n=11)				ST (n=17)				
	Pre		Post		Pre		Post		Pre		Post		Pre		Post
sMAP (W)	86.7 28.3	±	94.2 30.3	±	81.4 25.4	±	101.4 27.9	±	98.9 30.2	±	93.6 34.4	±	82.0 28.1	±	91.3 ± 29.7
$VO_{2peak(60\%)}(l.min^{-1})$	1.28 0.46	±	1.41 0.46	±	1.30 0.47	±	1.61 0.42	±	1.37 0.56	±	1.44 0.52	±	1.22 0.39	±	$\begin{array}{cc} 1.31 & \pm \\ 0.43 & \end{array}$
VO <sub>2peak(60%)</sub> (ml.min <sup>-1</sup> .kg <sup>-1</sup> )	18.7 4.8	±	20.8 5.3	±	20.4 4.9	±	24.4 4.4	±	20.2 6.0	±	21.4 6.2	±	17.0 3.5	±	$19.1\pm4.5$
SBP (mmHg)	122.2 12.1	±	122.9 12.4	±	118.5 27.9	±	121.8 12.1	±	123.8 16.3	±	126.1 14.7	±	123.0 8.5	±	121.5± 11.4
DBP (mmHg)	76.8 8.0	±	77.6 8.0	±	72.9 9.7	±	74.0 7.5	±	79.7 9.7	±	82.0 8.6	±	76.6 5.0	±	$76.4\pm7.1$

#### Table 3: Effect of training intervention on cardiovascular measures

sMAP: sub–Maximal Aerobic Power. VO<sub>2</sub> peak: peak oxygen uptake at 60% of maximal heart rate; SBP: systolic blood pressure; DBP: diastolic blood pressure; *ET, Exercise training; CT, Cognitive training; ST, Simultaneous training.* 

		Total (n=35)		ET (n=7)			CT (n=	11)	ST (n=17)		
		Pre	Post	Pre	Post		Pre	Post	Pre	Post	
BDI		$0.96 \pm 1.2$	0.84 0.97	$0.83\pm0.75$	0.83 0.40	±	0.81 ± 1.2	0.45 = 0.93	$1.1\pm1.3$	$1.1 \pm 1.1$	
Forward span		6.3 ± 1.2 <sup>1</sup>	$6.3\pm1.2$	$6.4 \pm 0.78$	5.3 0.75*	±	$5.8\pm1.1$	$6.1\pm0.94$	$6.5\pm1.3$	$6.8\pm1.3$	
Backward span		$4.9\pm1.2$	5.2±1.3	$4.6\pm0.97$	4.7 ± 1.1	1	$4.6\pm1.2$	$4.7\pm1.2$	$5.2\pm1.3$	$5.6\pm1.3$	
DDST		53.5 ± 9.2	$55.6\pm9.0$	$53.0\pm7.7$	53.0 7.6	±	54.9 ± 7.2	$56.9 \pm 6.9$	52.7 11.3	$55.9\pm10.8$	
Rey's Words	Immediate recall	57.0 ± 7.4	57.3 ± 6.2	$58.7\pm6.7$	57.7 4.8	±	56.4 ± 4.9	56.3 ± 6.3	$56.7\pm9.0$	$57.8\pm 6.9$	
тмт а	Errors (n)	$0.08\pm0.2$	0.02 0.16	$0.14\pm0.37$	0.14 0.37	±	0	0	0.11 0.33	0	
TMT A	RT (ms)	$35.8\pm8.1$	$34.2 \pm 9.9$	$32.8\pm7.4$	33.3 10.6	±	38.6 ± 9.2	35.2 = 10.7	35.2 ± 7.5	$34.1\pm9.7$	
BDI Forward span Backward span DDST Rey's Words TMT A TMT B	Errors (n)	$0.60\pm0.8$	0.20 0.53 ª	$1.14 \pm 1.06$	0.28 0.75	±	$\begin{array}{c} 0.81 \\ \pm 0.75 \end{array}$	0.18 = 0.60	0.23 0.56	$0.17\pm0.39$	
	RT (ms)	$81.1\pm25$	69.2 20.0 ª	$89.6 \pm 27.6$	69.6 28.1	±	85.0 ± 29.3	67.0 = 13.9	75.2 22.2	$70.3 \pm 20.6$	
	Naming (Errors. n)	$0.91 \pm 1.7$	0.51 0.65	$0.71\pm0.95$	0.42 0.53	±	$1.1 \pm 1.1$	0.27 = 0.46	$0.88 \pm 2.5$	$0.70\pm0.77$	
	Naming (RT. ms)	808.7 106.1	788.6 127.7	774.5 ± 57.3	725.8 135.4	±	$\begin{array}{rrr} 754.3 & \pm \\ 98.8 \end{array}$	734.5 = 118.6	857.9 107.2	$849.5\pm106.3$	
	Inhibition (Errors. n)	$1.1\pm1.7$	0.60 0.94	$0.28\pm0.48$	0.71 0.95	±	$1.0\pm1.4$	$0.90 \pm 1.2$	$1.5\pm2.1$	$0.35\pm0.70$	
	Inhibition (RT. ms)	974.1 128.1	955.8 127.5	908.7 ± 105.2	891.5 147.9	±	$\begin{array}{rrr} 933.5 & \pm \\ 114.8 \end{array}$	939.6 = 144.3	1029.0 127.5	$992.8\pm99.4$	
GTROOR	Inhibition cost <i>(Errors.</i> n)	$0.38\pm1.7$	$\substack{0.08 \pm 1.}{}_{\text{b}}$	-0.42 ± 0.78	-0.28 1.4	±	-0.09 ± 1.5	0.63 ± 1.5	1.06 ± 1.8	$-0.43 \pm 1.1$ *	
STROOP	Inhibition cost (RT. ms)	166.3 108.1	167.2 109.4	134.2 ± 102.8	165.7 131.6	±	179.2 ± 70.1	205.1 = 111.8	171.1 131.1	$143.3\pm97.6$	
	Flexibility (Errors. n)	$12.4\pm7.9$	$8.0\pm5.5$	$14.8\pm7.5$	12.6 6.5	±	$12.7 \pm 8.5$	$8.1\pm5.8$	$11.2 \pm 7.9$	$6.1\pm3.7$	
	Flexibility (RT. ms)	1351.8 157.8	1320.9 13.3	1244.4 ± 107.3	1272.8 195.3	±	$\begin{array}{r} 1372.1 \ \pm \\ 130.7 \end{array}$	1291.0 = 207.1	1382.8 177.7	1360.1±164.4	
	Flexibility cost <i>(Errors.</i> n)	$11.3 \pm 7.5$	$7.4\pm5.3$	$14.6\pm7.6$	11.8 6.0	±	11.7 ± 7.8	$7.2\pm5.9$	$9.7\pm7.3$	$5.7\pm3.4$	
	Flexibility cost ( <i>RT. ms</i> )	543.1 194.1	532.2 196.0	469.9 ± 116.7	547.0 201.2	±	614.7 ± 179.3	556.4 = 209.9	524.9 219.7	$510.6\pm194.6$	
	Single task (Accuracy)	$32.8\pm8.9$	35.1 ± 5.	$34.7\pm9.2$	35.1 7.5	±	29.7 ± 11.0	35.1 ± 5.3	$33.9\pm7.4$	$35.1\pm5.9$	
N-BACK	Dual Task (Accuracy)	$30.9\pm7.8$	$32.0\pm7.1$	$30.8\pm8.3$	30.4 8.2	±	29.5 ± 10.1	31.7 ± 8.:	$31.8 \pm 6.5$	$32.9\pm 6.2$	
	Single task (walk. m)	$42.2\pm7.6$	$41.6\pm8.2$	$42.9\pm7.2$	45.0 6.6	±	$\begin{array}{rrr} 42.5 & \pm \\ 8.5 \end{array}$	$39.6\pm7.0$	41.6 ± 7.1	$41.6\pm9.3$	
	Dual Task (walk. m)	$40.6\pm8.1$	40.1 ± 8	$42.6\pm7.2$	43.0 6.7	±	$\begin{array}{rrr} 40.4 & \pm \\ 8.8 \end{array}$	38.3 ± 7.3	$39.8\pm8.4$	$40.0\pm9.5$	

# Table 4: Effect of training intervention on cognitive measures

BDI: Beck's Depression Inventory; DSST: Digit Symbol Substitution Test; TMT: Trail Making Test; RT: Reaction Time; n: number; m: meter. **a**: main effect of time (p<0.05). **b**: interaction Time by Group; \* different from pré (p<0.05), *ET, Exercise training; CT, Cognitive training; ST, Simultaneous training*