



Exploring the IRE1 interactome: from canonical signaling functions to unexpected roles

Simon Le Goupil, Hadrien Laprade, Marc Aubry, Eric Chevet

► To cite this version:

Simon Le Goupil, Hadrien Laprade, Marc Aubry, Eric Chevet. Exploring the IRE1 interactome: from canonical signaling functions to unexpected roles. *Journal of Biological Chemistry*, 2024, 300 (4), pp.107169. 10.1016/j.jbc.2024.107169 . hal-04540346

HAL Id: hal-04540346

<https://hal.science/hal-04540346>

Submitted on 10 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Exploring the IRE1 interactome: From canonical signaling functions to unexpected roles

Received for publication, November 13, 2023, and in revised form, March 4, 2024. Published, Papers in Press, March 15, 2024,
<https://doi.org/10.1016/j.jbc.2024.107169>

Simon Le Gouplil^{1,2,*}, Hadrien Laprade^{1,2}, Marc Aubry^{1,2}, and Eric Chevet^{1,2}

From the ¹INSERM U1242, University of Rennes, Rennes, France; ²Centre de Lutte contre le cancer Eugène Marquis, Rennes, France

Reviewed by members of the JBC Editorial Board. Edited by Ronald Wek

The unfolded protein response is a mechanism aiming at restoring endoplasmic reticulum (ER) homeostasis and is likely involved in other adaptive pathways. The unfolded protein response is transduced by three proteins acting as sensors and triggering downstream signaling pathways. Among them, inositol-requiring enzyme 1 alpha (IRE1 α) (referred to as IRE1 hereafter), an endoplasmic reticulum–resident type I transmembrane protein, exerts its function through both kinase and endoribonuclease activities, resulting in both X-box binding protein 1 mRNA splicing and RNA degradation (regulated ire1 dependent decay). An increasing number of studies have reported protein–protein interactions as regulators of these signaling mechanisms, and additionally, driving other noncanonical functions. In this review, we deliver evolutive and structural insights on IRE1 and further describe how this protein interaction network (interactome) regulates IRE1 signaling abilities or mediates other cellular processes through catalytic-independent mechanisms. Moreover, we focus on newly discovered targets of IRE1 kinase activity and discuss potentially novel IRE1 functions based on the nature of the interactome, thereby identifying new fields to explore regarding this protein's biological roles.

either retaining improperly folded proteins in the ER to allow further folding cycles or to promote the degradation of terminally misfolded proteins through ER-associated degradation (ERAD) (10). This process is critical to ensure ER homeostasis, integrating protein synthesis rates, folding, and degradation/export through the secretory pathway (6). Under some circumstances, ER homeostasis can be disrupted when protein folding demand in the ER exceeds the ER capacity to handle productive protein folding. This leads to an accumulation of improperly folded proteins in this compartment, a situation known as ER stress (11). To cope with ER stress, organisms/cells have evolved a strategy termed the unfolded protein response (UPR). The UPR is a signaling pathway triggered upon ER stress mainly to restore ER homeostasis (adaptive UPR). However, if this process fails, the UPR will engage cells on a death path (terminal UPR) (12, 13). Over the years, many studies showed the involvement of UPR signaling in a wide range of diseases such as cancers, metabolic, or degenerative disorders (14, 15).

The UPR is transduced by three ER-resident transmembrane sensors named respectively the PKR-like endoplasmic reticulum kinase (PERK), the activating transcription factor 6 (ATF6), and the inositol-requiring enzyme 1 (IRE1). PERK is a type 1 transmembrane protein with a luminal sensing domain and a cytosolic kinase catalytic domain. Activated PERK enables the phosphorylation of the eukaryotic translation initiation factor 2A (eIF2 α), thereby decreasing overall mRNA translation and therefore the ER protein load (16). However, phosphorylated eIF2 α allows bypassing the reading of upstream ORF on translated mRNA by ribosomes to activate the selective synthesis of several proteins, including the ATF4 transcription factor (17). ATF4 targets the transcription of genes whose products are involved in redox homeostasis, amino acid metabolism, or autophagy (18, 19) (Fig. 1). The ATF4-dependent transcription network is also contributing to a negative feedback loop allowing translation to resume (20). Indeed, ATF4 controls the expression of the phosphatase subunit GADD34 which associates with the protein phosphatase 1 catalytic subunit to dephosphorylate eIF2 α hence restoring translation (21). Upon accumulation of improperly folded proteins in the ER lumen, the type II ER transmembrane protein ATF6 is translocated in the Golgi apparatus through coat protein complex II transport vesicles, where it is cleaved by the site 1/2 proteases (S1P

Endoplasmic reticulum stress and the unfolded protein response

The endoplasmic reticulum (ER) is one of the largest organelles in eukaryotic cells. It was observed for the first time at the end of the 19th century by Charles Garnier and named “ergastoplasm” (1) then defined as ER in 1945 by Porter, Claude, and Fullam (2). Beyond its functions in the regulation of calcium, cells' redox status and lipid homeostasis, the ER has a predominant role in protein synthesis, productive folding, and trafficking (3–6). Indeed, about one-third of the proteome passes through the ER to acquire proper folding before reaching its final destination (7). The ER lumen exhibits a calcium-rich, oxidative environment populated by ER-resident enzymes and foldases (e.g., protein disulfide isomerase (PDIs), chaperones, N-linked glycosylation machinery), which ensure proper protein folding (8, 9). In addition, a protein quality control system is in charge of

* For correspondence: Simon Le Gouplil, simon.legouplil@univ-rennes1.fr.






Figure 1. The unfolded protein response. Schematic representation of the IRE1, PERK, and ATF6 branches of the UPR. ATF6, activating transcription factor 6; IRE1, inositol-requiring enzyme 1; PERK, PKR-like endoplasmic reticulum kinase.

and S2P) leading to the release of ATF6 cytosolic domain (CD) (ATF6f) in the cytosol (22). ATF6f is a leucine zipper (bZIP) motif containing transcription factor that mostly controls the expression of genes coding for ER chaperones/foldases, thereby leading to an increase in productive protein folding in the ER (23, 24) (Fig. 1).

The most evolutionary conserved stress transducer of the UPR is IRE1, discovered in yeast and mammals in the 90s (25, 26). IRE1 is an ER-resident type 1 transmembrane protein that comprises both BiP-binding (binding-immuno-globulin protein) and misfolded protein binding domain in the ER lumen as well as kinase and endoribonuclease catalytic domains in the cytosol. Upon ER stress, IRE1 oligomerizes and transautophosphorylates, triggering two outcomes managed by the RNase domain (25, 27). First, the nonconventional splicing of X-box binding protein 1 (XBP1) mRNA, where IRE1 removes a 26-nt intron, with further ligation of the remaining 5' and 3' fragments by the tRNA ligase RtcB to form the spliced XBP1 (XBP1s) mRNA (28–30). This splicing event causes a frameshift in the XBP1s transcript, leading to translation of a protein containing a transcription activator domain in the C-terminal end. Hence XBP1s acts as a transcription factor regulating the expression of enzymes involved in lipid biogenesis, of ER

chaperones, and ERAD components (31–34). Second, the RNase activity termed as regulated IRE1-dependent decay (RIDD) is responsible for cleaving a pool of RNAs (mRNA, rRNA, miRNA), aiming at decreasing translation or at inducing novel cellular programs (35, 36) (Fig. 1). Importantly, RIDD specificity still remains unclear and might be cell specific. To date, a consensus sequence was identified for RIDD substrates (CUGCAG) within a stem-loop motif (37), but a recent study unveiled targeting of RNAs lacking those criteria and termed this activity as RIDD-lacking endomotif (38). In addition to their signaling, the three UPR sensors exert a crosstalk between each other as shown by the upregulation of XBP1 expression by both ATF6f and ATF4 (39, 40) or the competitive role of PERK and IRE1 regarding expression of the death receptor 5 (41). Some hypotheses propose a hierarchy among the UPR sensors in human, suggesting that low ER stress is solely resolved by the ATF6 branch, while mild or high ER stress additionally requires IRE1 and PERK branches. Moreover, upon persistent ER stress, it is thought that IRE1 signaling declines in the benefit of PERK, to mediate CHOP-induced apoptosis (42). Numerous studies also point toward scaffolding functions of IRE1 either to regulate its signaling or to mediate other cellular processes (translation, cytoskeleton ...). In this

review, we propose an overview of IRE1 biochemistry and describe the known IRE1 protein–protein interactions (PPis) (Table 1) and their biological outputs.

IRE1 across evolution

IRE1 α and IRE1 β paralogs

The evolution of the IRE1 gene might have encountered a duplication event, as, unlike in *Saccharomyces cerevisiae*, two IRE1 paralogs exist in mammals, namely IRE1 α and IRE1 β , respectively encoded by two genes: ERN1 and ERN2,

respectively localized on 17q23.3 and 16p12.2 chromosomal regions and displaying 39% sequence identity (43). Both proteins conserved their structural domains. Whereas IRE1 α is ubiquitously expressed, IRE1 β is preferentially found in mucosal epithelia (44, 45). Regarding their functions, both IRE1 α and IRE1 β exerts RIDD activity and IRE1 β was found to inhibit mRNA translation by degrading rRNA (43). However, while IRE1 α -mediated XBP1 mRNA splicing has been largely described, it remains unclear whether IRE1 β is capable of such activity in a physiological setting. Due to nonconserved amino acids in the kinase domain, IRE1 β has impaired

Table 1

List of described IRE1 interactors

Biological functions	Protein ID	Protein full name	Reference
IRE1 activation	Misfolded proteins AGR2* BIP ERDJ4 HSP47 PDIA1 PDIA6	Misfolded proteins Anterior gradient protein 2 Endoplasmic reticulum chaperone BiP DnaJ homolog subfamily B member 9 Serpine family H member 1 Protein disulfide isomerase A1 Protein disulfide isomerase A6	(85–87) (51, 95) (88) (88) (96) (99) (98)
IRE1 Oligomerization	c-ABL NMIIIB RAC Sec61 β Sec63	c-ABL Nonmuscle myosin heavy chain IIB RAC Protein transport protein Sec61 subunit beta Translocation protein Sec63 homolog	(104, 105) (106) (103) (101, 102) (101, 102)
IRE1 RNase activity	BAK BAX BI-1 BIM c-ABL NCK PUMA RNH1	Bcl-2 homologous antagonist/killer Apoptosis regulator BAX BAX inhibitor 1 Bcl-2-like protein 11 Tyrosine-protein kinase ABL1 Cytoplasmic protein NCK1 Bcl-2-binding component 3 Ribonuclease inhibitor	(109) (109) (112, 113) (110) (104) (114) (110) (111)
IRE1 Phospho regulation	AMPK Dcr2 ERK PKA PP2A PP2Ce Ptcp2 PTP1B RACK1 RPAP2 SCP3	AMP-activated protein kinase Phosphatase DCR2 (<i>S. cerevisiae</i>) Extracellular signal-regulated kinase Protein kinase A Protein phosphatase 2A Protein phosphatase 1L Serine/threonine phosphatase of type 2C (<i>S. cerevisiae</i>) Tyrosine-protein phosphatase nonreceptor type 1 Receptor for activated C-kinase 1 RNA polymerase II-associated protein 2 Small carboxy-terminal domain phosphatase 3	(118, 119) (116) (120) (122) (118) (117) (26) (105, 121) (118) (41) (120)
PTM and IRE1 degradation	BIM CHIP M50 MITOL OS9 RNF13 SEL1L SYN1V1	Bcl-2-like protein 11 E3 ubiquitin-protein ligase CHIP M50 viral protein E3 ubiquitin-protein ligase MARCHF5 OS9 endoplasmic reticulum lectin E3 ubiquitin-protein ligase RNF13 Protein sel-1 homolog 1 E3 ubiquitin-protein ligase synoviolin	(110) (129) (128) (131) (127) (130) (126, 127) (126, 127)
Apoptosis	AIP1 ASK1 JNK MITOL TRAF2	ASK1-interacting protein-1 Apoptotic signaling kinase 1 Mitogen-activated protein kinase 8 E3 ubiquitin-protein ligase MARCHF5 TNF receptor-associated factor 2	(135, 136) (135) (131) (135)
Cytoskeleton	CORO1 FLNA PKC α TUBA TUBB	Coronin Filamin A Protein kinase C alpha type Tubulin A Tubulin B	(111) (139) (139) (111) (111)
SRP and ribosome	RPLs SRPs	Ribosome protein large subunits Signal recognition particles	(143) (143)
Ca $^{2+}$ homeostasis	INSP3R STIM1	Inositol 1,4,5-trisphosphate receptor type 3 Stromal interaction molecule 1	(148) (149)
IRE1 Kinase target	AMPK BCL2 FMRP IKK PUM1 Pumilio SPL	AMP-activated protein kinase Apoptosis regulator Bcl-2 Fragile X messenger ribonucleoprotein 1 IkB α kinase Pumilio homolog 1 Pumilio (<i>Drosophila melanogaster</i>) Sphingosine 1-phosphate lyase	(119) (119) (150) (152) (151) (151) (153)

IRE1, inositol-requiring enzyme 1; PTM, posttranslational modification.

(*) indicates IRE1 β specific interactions.

phosphorylation and reduced oligomerization abilities (46–48). Even so, few studies were able to observe a slight XBP1 splicing activity for IRE1 β in *in vitro* splicing experiments and in overexpressing cells (49–51). In the latter case, the splicing activity is dependent on cell type and appeared to be reduced in mucosal cell line where IRE1 β is usually expressed (51). Concomitantly, IRE1 β was found to act as a negative regulator of IRE1 α , as its overexpression is able to decrease IRE1 α -dependent XBP1 splicing (47). Indeed, IRE1 β conserved a capacity to oligomerize with IRE1 α and by doing so, acts as a dominant negative by preventing IRE1 α functions (47). In this regard, IRE1 β is thought to attenuate ER stress upon inflammation induced diseases in mucosal epithelium (44). For the continuation of this review, we will

mainly consider IRE1 α (and refer to it as IRE1 unless specified).

IRE1 across evolution

IRE1 can be divided into three main domains: an N-terminal luminal domain (NLD), a transmembrane domain (TMD), and a CD, the latter comprising both kinase and RNase catalytic activities. These domains are all conserved in metazoan, fungi, and plant orthologs, as shown by the yeast *versus* human IRE1 comparison (Fig. 2A) (52). However, compared to human IRE1, the amino acid sequence differs across the reigns. Indeed, while the BLAST alignment of protein sequences reveals a high degree of conservation in mammals (>90% identity), a low IRE1 sequence conservation is found between human and the other




Figure 2. Evolutive insights on IRE1. *A*, schematic representation of yeast (above) and human (below) IRE1 sequence displaying the signal peptide (SP) with the three mains domains: luminal (NLD), a transmembrane domain, and the cytosolic domain exerting kinase and RNase activities. *B*, comparison of IRE1 protein sequence identity and similarity in different species using BLAST. *C*, GO Manhattan plot using g:Profiler of yeast (up) and human (bottom) IRE1 interactome from curated databases (APID, String and Biogrid). Black arrows highlight terms of interest, and values in brackets represent the number of enriched terms in each Gene Ontology. IRE1, inositol-requiring enzyme 1; NLD, N-terminal luminal domain.

species (approximatively 30% identity according to species) (Fig. 2B) (53). This trend is comparable with the overall similarity of the sequences (>90% in mammals while approximatively 50% in other eukaryotes and <40% in plants) (Fig. 2B). This suggests that both IRE1 kinase and RNase domains share common features and functions across species despite the changes in sequences and implies that the luminal domain diverges more and may show different characteristics which may be linked to (i) IRE1 activation mode in different species and/or (ii) the ability of IRE1 luminal domain to directly and selectively bind to specific motifs which may differ between species. However, IRE1 activity presents significant differences in yeast species. In *S. cerevisiae*, IRE1 can splice HAC1 mRNA (XBP1 functional homolog in yeast) together with the *S. cerevisiae* Rlg1p ligase (RtcB homolog in yeast) (27, 54) and to a much lesser extent RIDD (55). Whereas in *S. pombe*, IRE1 only degrades RNA through RIDD since further analysis found that *S. pombe* lacks HAC1 (or orthologs) (55, 56). Therefore, one can hypothesize that all eukaryotes may express an IRE1 capable of both splicing and RIDD activities (with different selectivity) and that the occurrence of splicing does not depend on the IRE1 RNase domain but on the expression of the target gene. This suggests an evolutive trait of the IRE1 pathway in managing ER stress, by modifying signaling characteristics and likely to expand its biological functions during evolution. This correlates also with the diversification of the UPR, with the appearance of the PERK and ATF6 arms (57) in higher order organisms (metazoans and chordates).

Comparison of IRE1 interactomes across evolution

To analyze the diversity of IRE1 PPis across evolution, we collected IRE1 interactome data from *S. cerevisiae* and *Homo sapiens*, the most documented species on public repositories (agile protein interactomes dataserver, String and Biogrid) (58–60). Interestingly, agile protein interactomes dataserver, the most complete database reports 300,000 interactions among the 5800 proteins studied in yeast while in human, 700,000 interactions are distributed among 14,500 proteins. Regarding IRE1, analysis of those repositories documents a total of 101 interactions for human against 121 for yeast. Gene Ontology analysis carried out through gProfiler revealed a different pattern of enrichment between human and yeast IRE1 human interactome. Indeed, every ontology (molecular function, biological process, and cellular component) displays much more terms in human than in yeast (Fig. 2C) (61). Regarding the content of these ontologies, while in yeast IRE1 interactors are mainly linked to stress-related processes such as protein folding or ERAD, human IRE1 interactors are connected with many more diverse processes in addition to the UPR, from signaling pathways (with the mitogen-activated protein kinase (MAPK) signaling), trafficking (coat protein complex II) to apoptosis or ubiquitination (Fig. 2C). We analyzed the number of terms enriched between the interactomes of a significant number of *H. sapiens* and *S. cerevisiae* orthologs and found that the richer and denser functional enrichment observed in the human IRE1 interactome is significantly different (p value = 0.0014, not shown) than the mean of interactome differences of other

orthologs. Altogether, it suggests that IRE1 is part of a specific subset of proteins, which gains new functions across evolution through their protein interaction network.

IRE1 structural insights

IRE1 luminal domain

A first crystal structure of the conserved core region of IRE1 NLD was obtained in *S. cerevisiae* in 2005 (62). It shows a dimer harboring a putative protein/protein interface (Fig. 3A). This structure describes that the IRE1 NLD is able to dimerize through both sides of the NLD and suggests that IRE1 could form high order oligomers necessary for its activation. This crystalized dimer also shows a “groove” structure between the two monomers similar to the major histocompatibility peptide-binding complexes (MHCs) (62). Interestingly, mutations in the NLD contact sites were shown to abolish IRE1 activation (63). In comparison, human IRE1 NLD dimer structure, obtained by crystallography, reveals a triangle shape of each monomer, composed of three β -sheets that appear different from the yeast NLD. This structural difference could be explained by the length of this domain in both species (Fig. 2A) and by the previously discussed poor conservation of the sequence as shown by the amino acid percentage identity between human and yeast (Fig. 2B). Upon dimerization, one face of a triangle interacts with another similar structure in an antiparallel manner, creating an MHC-like motif (Fig. 3B). This motif presents an interface able to form hydrophobic and hydrogen bonds, the last especially through the K121 amino acid (64). The following MHC-like structure is comparable to that of yeast, but the closed groove suggests a different peptide binding capacity in sensing unfolded protein (64). Of note, cancer-associated somatic mutations in the luminal domain of human IRE1 were reported to affect IRE1 signaling (65).

IRE1 transmembrane domain

The IRE1 TMD consists in a single amino acid stretch exhibiting a juxta-membrane amphipathic helix (AH) containing polar residues facing the lumen and a transmembrane helix (66) (Fig. 3, C and D). Interestingly, analysis of the TMD from different species revealed a conservation of these AH and transmembrane helix from mammals to plants (66). In yeast, a study identified a link between IRE1 activation and lipid sensing properties regulated by IRE1 AH (66). It was found that the AH could sense lipid composition of the ER membrane and that mutation in this domain led to impaired HAC1 mRNA splicing (66). They further discussed a model in which IRE1 AH could locally compress the membrane, allowing stable formation of oligomers (66). Another study showed that lipid unbalance induces UPR and that IRE1 counteracts membrane stress-induced cell death by reprogramming protein homeostasis and maintains membrane functions without affecting its composition (67). The same team recently reported that spliced HAC1 regulates the expression of different genes according to either lipid bilayer stress or unfolded protein stress (68). In human, IRE1 TMD has also been documented for lipid-sensing properties with IRE1-detecting lipid




Figure 3. Comparative structure of IRE1 in *Homo sapiens* and *Saccharomyces cerevisiae*. Dimer structures of luminal domain of (A) yeast (PDB: 2BE1) and (B) human (PDB: 2HZ6) IRE1. Each color represents a monomer. Helical wheel schematic representation of IRE1 yeast (C) and (D) human amphipathic helix from the transmembrane domain. The arrow represents the hydrophobic moment. E, crystal structure of dimeric IRE1 cytosolic domain of yeast (PDB: 2RIO). F, crystal structures of human dimeric IRE1 cytosolic domain with back-to-back (PDB: 6W3C, left) and face-to-face (PDB: 3P23, right) conformations. G, crystal structure of yeast IRE1 cytosolic oligomer (PDB: 3FBV). Each color represents a monomer, and the black arrow illustrates the progressive 52° rotation between each dimer. IRE1, inositol-requiring enzyme 1.

bilayer stress to trigger UPR activation (69, 70). Moreover, by using fluorescence complementation assays, Cho and colleagues showed in cells that lipid saturation of the ER membrane with palmitate induces IRE1 (lacking NLD) dimerization through the TMD, by a mechanism involving W457 and S450

residues (71). In addition, the IRE1 TMD has also been proposed to be involved in the interaction with the membrane-embedded translocon component Sec61β. Indeed, alteration of IRE1 TMD by replacing it with calnexin TMD abolished its interaction with the translocon (72).

IRE1 CD

The CD of IRE1 is the signaling region of IRE1. As such, it shows two main domains exerting, respectively a serine/threonine kinase activity and an RNase activity. The former leads to IRE1 transautophosphorylation upon oligomerization, thereby leading to the activation of the latter. IRE1 RNase induces the noncanonical splicing of HAC1 mRNA in yeast and of XBP1 mRNA in human, with further ligation by the Rlg1 (yeast) or RtcB (human) RNA ligases, respectively. This process allows for the synthesis of the spliced form of the XBP1s/Hac1^{1p} transcription factors, thereby promoting the expression of UPR target genes (28, 73, 74). IRE1 cytosolic regions presents two main fragments: a kinase domain (yeast 674–980; human 571–832), and a RNase domain (yeast 983–1115; human 835–963) (Fig. 2A). Structural evidence in yeast showed that the phosphorylated IRE1 kinase/RNase domain is able to dimerize in its active form (75). This dimer is arranged in a back-to-back orientation, meaning that the kinase domain of each monomer is not in contact with their target, thus suggesting that the phosphorylation occurs by contact with another dimer (Fig. 3E). This activating IRE1 transautophosphorylation is further supported by the ability of the CD to form higher order oligomers (76). The phosphorylation occurring on S840, S841, and T844 in yeast IRE1 leads to the formation of both intramolecular and intermolecular (with another IRE1) salt bridges promoting the self-activation of the RNase domain (76). In contrast, hyperphosphorylation of a 28 amino acid loop (residues 864–892) containing seven serine and two threonine residues may participate in the deactivation of IRE1 by promoting its disassembly, a regulatory mechanism to adapt to a sustained ER stress (77). In human, the crystal structure of the IRE1 CD was found to resemble that of the yeast IRE1, with a back-to-back orientation bringing the RNase domains in close proximity, compatible with the cleavage of known substrates (78, 79) (Fig. 3F, left). This would be relevant regarding the modest conservation of sequence (~60% similarity) (Fig. 2B). Although, another structure was obtained after IRE1 dephosphorylation, called the face-to-face conformation (Fig. 3F, right), where the RNase domains of two monomers have minimal interactions. This last orientation brings another possibility regarding the transautophosphorylation, where the kinase domain of a monomer is in close contact with the activation loop of a monomer to achieve the reciprocal phosphorylation at S724, S726, and S729 (80, 81). Despite the lack of observation of this crystal structure in yeast, the interface between the two kinase domains is conserved, suggesting that such conformation exists for all IRE1 homologs. These two states are part of a model in which IRE1 first acquires the face-to-face conformation and then proceeds to the transphosphorylation of the activation loop, leading to reorganization in a back-to-back conformation that brings two RNase domains in proximity, thus promoting IRE1 activity. Additionally, an oligomeric structure of yeast IRE1 CDs was obtained by crystallography, displaying fourteen monomers assembled in seven back-to-back dimers exerting a 52 degree rotation between each dimer, therefore confirming the ability of the CD to form high order oligomer (Fig. 3G)

(76). Nevertheless, this structure does not take into consideration the constraints imposed by the membrane insertion of IRE1. A foci structure containing five IRE1 dimers was further observed in yeast by microscopy. Altogether, it shows that IRE1 monomers gather into larger signaling platform at the ER (63). To allow such a complex, the IRE1 linker sequence positioned between the kinase/RNase and the TMDs might be of importance by providing flexibility to the protein. In agreement with this, an *in vitro* experiment evaluating the oligomerization capacity of the cytosolic portion of IRE1 showed that deletion of the linker region prevents formation of oligomers (76). To date, there is no available crystal structure of a full-length IRE1 in any species, thereby preventing the elucidation of the exact conformation of IRE1 active form, since its oligomerization and activation may also be governed (at least in part) by the luminal domain.

While IRE1 activation might be subjected to different modes that remain debated (see Activation in ER lumen below), it is possible that the changes in sequence and structure between yeast and human IRE1 (Figs. 2A and 3, A and B) lead to modulation of the signaling properties. Especially, as the luminal domain is the most divergent between species, it can confer a variability in binding misfolded proteins or other partners, leading to the complexification and fine-tuning of this signaling. This would be concomitant with the appearance of other UPR sensors in metazoans. Nonetheless, one must notice that despite the three UPR sensors are expressed in *Caenorhabditis elegans*, the IRE1–XBP1s axis was shown to represent the main path to cope with ER stress in this organism (82). Additionally, while the CD retains a similar structure between species, the changes in sequence and length of this domain (~50 amino acids more in the yeast IRE1 compared to human) may modify the oligomerization kinetic as yeast IRE1 was observed to form higher order oligomer, which were not observed in human so far. Those differences also emerge as a parameter regulating the binding and response to small molecules. Indeed, quercetin, a flavonoid, was first found to bind to yeast IRE1 in a small pocket formed by kinase and RNase domain residues (amino acids 838–844 and 1036–1040, called Q-site) (83). This compound induces a modification of yeast IRE1 nucleotide-binding site and further activates IRE1. Later, an O-glycoside quercetin derivative, the quercitrin, was characterized to bind to another binding site, located at the dimer interface region of the human RNase domain (amino acids 830–834). This molecule leads to an opposite effect, by inhibiting IRE1 (84). Altogether, it suggests that IRE1 from yeast and human respond differently to external stimuli.

Protein-protein interactions regulating IRE1 signaling: the IRE1 signalosome

Protein–protein interactions

Activation in ER lumen

IRE1 luminal domain has mainly been described as responsible for sensing misfolded proteins. Yet these sensing mechanisms are not fully understood and three models are currently proposed. On the one hand, the “direct association”

model where peptides exposed at the surface of improperly folded proteins bind directly to IRE1 NLD, promoting its oligomerization and further activation. This model was first thought to occur in yeast but not human IRE1 (85, 86), however, a recent study described this mechanism in human where the direct binding induces a conformational change and leads to IRE1 oligomerization (87). On the other hand, the “competition models” presents IRE1 as a signal transducer (instead of direct sensor), where the chaperone BiP interacts with IRE1 NLD to repress IRE1 signaling. Here, two mechanisms were proposed and are still a matter of debate in the field. One sustains that in the absence of improperly folded proteins, the ER chaperone ERDJ4 binds IRE1 NLD and recruits BiP substrate binding domain to IRE1 NLD, stabilizing its monomeric form and inhibiting its oligomerization (88, 89). The other mechanism involves direct binding of BiP nucleotide binding domain to IRE1 NLD, where misfolded proteins bind BiP substrate binding domain, triggering an allosteric conformational change, releasing BiP from IRE1 and enabling its oligomerization (90–92). In both cases, upon presence of misfolded proteins, BiP is titrated away from IRE1, thus allowing its oligomerization and further activation. Such mechanisms do not seem to exist in *S. cerevisiae* (with the BiP ortholog kar2) (93, 94). Interestingly, a similar mechanism seems to occur for IRE β with the ER chaperone AGR2, where the last binds to IRE β luminal domain to favorize the monomeric state (51, 95).

In addition to those models, other PPis in the ER lumen were shown to modulate IRE1 activation. Immunoprecipitation coupled to mass spectrometry experiments, followed by biochemical approaches confirmation, showed that the ER chaperone HSP47 interacts with IRE1 NLD with high affinity upon ER stress, competing with BiP binding and leading to IRE1 oligomerization or stabilization of the oligomeric state (96). At last, oxidation of C148 in IRE1 luminal domain allows the protein disulfide isomerase PDIA6 to form disulfide bonds with oligomeric IRE1 and converting it into monomeric entities therefore limiting its signaling (97, 98). ER-stress induced phosphorylation of another protein disulfide isomerase, PDIA1, was also reported to interact with IRE1 NLD, leading to an attenuation of excessive IRE1 activation (99).

Oligomerization

IRE1 oligomerization is necessary to ensure signaling. IRE1 can dimerize and possibly form high order oligomers composed of several dimers. However, it remains unclear how the degree of oligomerization defines IRE1 signaling outputs. While IRE1 tetramers are thought to preferentially catalyze XBP1 splicing, IRE1 dimers tend to be more prone to mediate RIDD (55, 76). Using single molecule tracking and HaloTag, IRE1 was shown to mainly exist in an inactive back-to-back dimer (100). Upon ER stress, IRE1 dimers form transient higher order oligomers, allowing the reciprocal dimers to be positioned in a “face-to-face like” configuration. Therefore, it is interesting to focus on interactions able to fine tune IRE1

oligomerization state. In this regard, studies reported that the translocon subunit Sec61 β interacts with IRE1 TMD, forming a complex with another translocon subunit Sec63, preventing IRE1 oligomerization and further activation by recruiting BiP (72, 101, 102). Additionally, it was shown that the ribosome-associated complex (RAC) chaperone is involved in the regulation of IRE1 oligomerization state. Observation of IRE1 cluster foci using microscopy revealed that RAC promotes IRE1 high order oligomerization to enhance XBP1 mRNA splicing but not RIDD (103). However, the physical interaction between RAC and IRE1 needs further confirmation. The Abelson tyrosine-protein kinase 1 (c-ABL) was found to interact with IRE1 CD upon ER stress. Independently of its kinase activity, c-ABL acts as a scaffold protein and promotes IRE1 oligomerization and further hyperactivation, leading to RIDD and terminal UPR (104). Moreover, c-ABL modulates the IRE1 signaling cascade by phosphorylating the tRNA ligase RtcB at least on three tyrosine residues (306, 316, 475), with Y306 phosphorylation reducing RtcB interaction with IRE1 and subsequently XBP1 mRNA splicing (105). One might also hypothesize that dissociation of the IRE1–RtcB complex by c-ABL might leads to the formation of IRE1 clusters able to perform RIDD. Altogether, it confers a role of switch to c-ABL, able to fine-tune IRE1 signaling through different mechanisms. At last, a link between IRE1 signaling and cytoskeleton was unveiled. The nonmuscle myosin IIB (NMIIB) interacts with IRE1 during ER stress. While it seems that NMIIB is not directly involved in IRE1 phosphorylation and dimerization, NMIIB drives IRE1 foci formation, promoting splicing of XBP1 mRNA (106). Interestingly, this mechanism was found to require the ATPase activity of NMIIB as well as a functional actin network, suggesting that the actomyosin contractility of NMIIB is required for IRE1 α signaling (106).

RNase activity

Regarding interactions directly regulating the RNase activity, the previously described relationship between IRE1 and c-ABL was also found to act as a genotoxic stress sensor, by promoting RIDD activation and further cleavage of PPP2R1A and RuvB-like AAA ATPase 1 mRNA, two proteins involved in dephosphorylation of DNA damage proteins (107). The apoptosis regulator BAX (BAX) and Bcl-2 Homologous Antagonist/Killer (BAK), two proapoptotic proteins (108), were shown by coimmunoprecipitation to bind IRE1 (preferentially in its active form) upon high ER stress and were found to promote XBP1 mRNA splicing (109). Similarly, other PPis were found to enhance IRE1 signaling. This is illustrated by the association of IRE1 with the Bcl-2 family members Bcl-2-like protein 11 (BIM) and PUMA, leading to sustained XBP1 mRNA splicing upon ER stress (110).

Conversely, some PPis were shown to impair IRE1 RNase activity. An IRE1 immunoprecipitation coupled to mass spectrometry experiment identified RNH1, a ribonuclease inhibitor, to be present in complex with IRE1. This result was confirmed using both immunoprecipitation and proximity ligation. Surprisingly, ER stress increases this interaction

between IRE1 and RNH1, thereby reducing XBP1 mRNA splicing (111). This counterintuitive outcome might indicate that IRE1–RNH1 interaction acts as a regulatory mechanism, where RNH1 decreases IRE1 activity upon prolonged ER stress. The BAX inhibitor-1 (BI-1) was found to compete with BAX and BAK for IRE1. BI-1 interacts with the IRE1 C-terminal region and reduces IRE1 RNase activity (112, 113). Under basal conditions, the interaction of the SH2/SH3-containing adaptor protein 1 with IRE1 carboxy-terminal region was shown to repress IRE1, a mechanism proposed to avoid a signaling leakage (114).

Protein modification

Phosphorylation-dependent regulations of IRE1 signaling

Evidences unveiled a role of IRE1 phosphorylation in determining its signaling outcomes (115). As such, IRE1 faces phosphorylation-dephosphorylation cycles relying on PPis. These regulating events were studied and revealed interesting mechanisms. Regarding the downregulation of IRE1 signaling through dephosphorylation, two yeast phosphatases—Ptc2p serine/threonine phosphatase of type 2C and Dcr2—were found to interact with phosphorylated IRE1 and further dephosphorylate it to downregulate the splicing of HAC1 mRNA (26, 116). Interestingly, it was found that Dcr2 specifically interacts with IRE1 when phosphorylated at S840 and S841 residues (116). This finding was extended to human thanks to another study describing IRE1 interaction with the ER membrane phosphatase PP2Ce, the human homolog of the *S. cerevisiae* Ptcp2 (117). Here, PP2Ce dephosphorylates IRE1 S724 upon ER stress, acting as a negative regulator of IRE1 signaling. Of note, both IRE1 α and IRE1 β were shown to partner with PP2Ce, unveiling a probable common regulatory mechanism (117). Coimmunoprecipitation assays found that the scaffold protein RACK1 interacts with IRE1 linker domain (between the kinase and the TMD) (118, 119). In basal condition, RACK1 recruits the phosphatase PP2A, which dephosphorylate IRE1 (120). A recent article uncovered through screening different serine-threonine phosphatases that the small carboxy-terminal domain phosphatase 3 was able to interact directly with IRE1 and further dephosphorylate the protein (120). We previously mentioned the crosstalk between the UPR sensors. In this regard, a study investigated the role of the PERK pathway in PPi-dependent attenuation of IRE1 signaling. Upon acute ER stress, PERK-prolonged activation leads to the phosphatase RPAP2 to interact and dephosphorylate IRE1, thereby decreasing its signaling (41). Another mechanism involving the protein tyrosine phosphatase PTP1B was shown to regulate both catalytic and scaffolding functions of IRE1 (105, 121). Here, PTP1B deficiency or catalytically inactive impairs XBP1 splicing and c-Jun-N-terminal-kinase (JNK) activation (121), while further investigations revealed that PTP1B dephosphorylates RtcB, thus enhancing its association with IRE1 for XBP1 mRNA splicing (105). However, to date, direct interaction between IRE1 and PTP1B requires confirmation.

Conversely, other proteins were found to phosphorylate IRE1. Interestingly, the previously mentioned IRE1–RACK1 complex was found to play another role. Upon increased ER stress, RACK1 recruits the AMP-activated protein kinase, which phosphorylates IRE1 (118, 119). Hence, RACK1 acts as a platform regulating IRE1 signaling according to ER stress through the phosphorylation status. Glucagon-stimulated PKA was reported to phosphorylate IRE1, indicating a role in glucose metabolism and suggesting an ER-lumen-independent activation (122). At last, a mechanism by which KRas mutated tumors elaborate tumor resistance to KRas inhibitors was shown to involve ERK1/2-mediated phosphorylation of IRE1, thus preventing its degradation by the ERAD machinery (120). This led us to further document the role of post-translational modifications (PTMs) of IRE1 and its degradation.

PTMs and IRE1 degradation

N-glycosylation is a PTM occurring during synthesis and maturation of ER proteins, participating in their folding and functionality (8, 123). IRE1 is N-glycosylated at N176 (124, 125) but there is no clear evidence mentioning a role for this PTM in regulating IRE1 activity. Despite that, few studies reported that mutations preventing this PTM do not impair IRE1 folding nor oligomerization properties (124, 125). Also, N-glycosylation disruption causes ER stress, and some might hypothesize that it could be a sensor-initiating IRE1 cascade.

Alternatively, ERAD was shown to control IRE1 expression and activity, as shown by the SYVN1/SEL1L-dependent ubiquitination of IRE1, promoting its degradation (126, 127). In this case, IRE1 is sent to ERAD in basal conditions, where both BiP and the lectin protein OS9 interacts with IRE1 in the lumen, recruiting the SYVN1/SEL1L complex and subsequently allowing IRE1 cytosolic ubiquitination by the E3 ligase SYNV1 (127). OS9 being a lectin, we suggest that it recognizes IRE1 through its N-glycan chain, therefore conferring the N-glycan a role in IRE1 stability. Cytomegalovirus M50 was found to take advantage of this ERAD-dependent IRE1 degradation, where the M50 viral protein interacts and tethers with both IRE1 and SEL1L to promote IRE1 degradation (128). IRE1 was also found to be a substrate of the E3 ubiquitin-protein ligase CHIP (also involved in ERAD). Authors proposed a dual role for CHIP-dependent IRE1 ubiquitination during ER stress: first, ubiquitination of IRE1 K545, enhancing IRE1 phosphorylation and second, of IRE1 K828, facilitating recruitment of TNF receptor-associated factor 2 (TRAF2) for JNK pathway activation (129). This is supported by another study, where the E3 ligase RNF13 interacts with IRE1, exerting enhanced XBP1 splicing and JNK activation (130). On another note, a recent study identified IRE1 to be a substrate of the mitochondria ubiquitin ligase MARCHF5 (MITOL), which ubiquitinylate IRE1 K481 with a K63 ubiquitin chain at mitochondria-ER contact sites (131). This ubiquitination was proposed to disrupt BIM stabilizing interaction with IRE1 oligomers, thus inhibiting IRE1 signaling (110, 131).

IRE1 PPis associated with other functions/outcomes**Protein–protein interactions***Apoptosis*

IRE1 signaling has been connected to the activation of the apoptosis cascade. Indeed, upon prolonged ER stress, IRE1 RIDD activity cleaves nonselectively several prosurvival miRNAs. Among them, miR-17, a miRNA known to repress TXNIP expression, subsequently triggers caspase 1 and 2/interleukin-1 β -dependent apoptosis (36, 132). Of note another report showed that ER stress did not lead to the activation of caspase 2 to initiate apoptosis (133). Several other studies have also connected IRE1 PPis with cell death, occurring during terminal UPR (134). This is linked to the activation of the proapoptotic JNK pathway, which first involves the interaction of IRE1 with TRAF2 and the further recruitment ASK1, also known as apoptotic signaling kinase 1 (135, 136). ASK1 may then phosphorylate JNK, thereby initiating the JNK pathway and leading to apoptosis. A new actor in the TRAF2/ASK1/JNK axis was also identified as the ASK1-interacting-protein 1 (AIP1), which, upon ER stress, is recruited on IRE1 prior to TRAF2 and further promotes JNK pathway activation (137). Additionally, this study presented evidence that AIP1 could enhance IRE1 dimerization (137). We previously mentioned IRE1 interactions with the Bcl-2 family member BIM and PUMA. These interactions occur through the BH3 domain of these proteins, which is necessary for their canonical proapoptotic signaling (109). It was shown that knock out of BIM and PUMA reduces both IRE1 RIDD and JNK activation (110). Therefore, BIM and PUMA might promote apoptosis by sustaining cleavage of prosurvival miRNAs by RIDD and activation of the JNK pathway. Accordingly, the other members of the Bcl2-2 family BAX and BAK, also containing BH3 domains, may act in a similar manner by interacting with IRE1 (109). Therefore, it confers a noncanonical role of these BH3 domain proteins in inducing cell death. Conversely, we also hypothesize on this model that the previously described protein BI-1 can inhibit cell death by competing with BAX and BAK interactions with IRE1. More recently, another IRE1 PPi was reported to exert antiapoptotic functions, involving the previously described interaction between IRE1 and MITOL, shown to inhibit JNK phosphorylation (131).

Cytoskeleton

In the past years, few studies reported a crosstalk between UPR and cytoskeleton as mentioned with NMIIB or through the interaction of the actin-binding protein FLNA with PERK (106, 138). Interestingly, a mechanism involving both IRE1 and FLNA showed that IRE1 promotes cell migration through FLNA phosphorylation. In this study, IRE1 associates with FLNA and acts as a recruiting platform for the kinase PKC α , enabling the phosphorylation of FLNA, which in turns drives actin cytoskeleton remodeling and cell migration (139). This interaction was shown to participate in migration of cells infected by toxoplasma, where the parasite engages the

IRE1–FLNA axis (140). Additionally, several high-throughput interactome screenings identified other cytoskeletal proteins such as CORO1 proteins or tubulin α and β , however, they require further validation (111, 139). IRE1 interacting with cytoskeleton proteins may also participate in membrane contact sites between ER and plasma membrane (ER-PM contact sites). This complex is known to regulate Ca^{2+} and lipid homeostasis, two processes associated with IRE1 signaling (141, 142).

Translation machinery: signal recognition particle and ribosome

High translational load can result in nonefficient protein folding in the ER by overwhelming the folding capacity of this compartment, hence, triggering ER stress. Given the proximity of IRE1 with the protein synthesis machinery, a crosstalk has been discovered between IRE1 signaling and mRNA translation. Cross-linking experiments revealed interactions between IRE1 and translating ribosomes at the cytosolic ER surface, through the ribosomal subunits RPL9, RPL10A, RPS14, and RPS24 (143). In the same study, immunoprecipitation analyses showed IRE1 interaction with proteins from the signal recognition particle (SRP) (143). These results correlate with studies describing that XBP1 mRNA begins translation before being brought to the ER membrane by the SRP pathway to undergo IRE1-dependent splicing (144–146). Moreover, we can hypothesize a role for IRE1 in monitoring the newly synthesized polypeptides entering the ER during the cotranslational translocation, as a way to avoid protein accumulation into the ER. Of note, those interactions with SRP and ribosomal proteins did not modify the catalytic functions of IRE1, thus suggesting a catalytic-independent regulation of the UPR which remains to be further investigated.

Calcium homeostasis

The ER is an intracellular organelle known to be involved in the maintenance of Ca^{2+} homeostasis. It was suggested that IRE1 could play a role in regulating Ca^{2+} metabolism independently of its enzymatic activities (147, 148). Indeed, IRE1 CD was shown to interact with the inositol-1,4,5-trisphosphate receptors. This PPi allows mitochondria to localize close to the ER in the mitochondria-associated ER membrane, another membrane contact site, and further promote calcium transfer to support mitochondrial respiration (148). This is supported by recent findings describing the interaction between IRE1 and the stromal interaction molecule 1 (STIM1) in the ER lumen (149). This interaction connects the UPR and Ca^{2+} metabolism, where STIM1 promotes XBP1 mRNA splicing upon ER Ca^{2+} store depletion and IRE1 promotes STIM1-dependent ER-PM contact sites, triggering the store operated Ca^{2+} entry pathway.

Direct IRE1 kinase-dependent phosphorylation

For many years, the only known substrate of the IRE1 kinase domain was IRE1 itself. However, few recent publications reported novel targets for IRE1 kinase activity. The previously described IRE1–RACK1 axis was shown to participate in BCL-2 phosphorylation by IRE1 at S70, a phosphorylation site

known to exert antiapoptotic functions (119). Liu and colleagues also showed that IRE1 interacts and promotes AMP-activated protein kinase phosphorylation exerting prosurvival signaling, nevertheless the direct/indirect modification remains unclear (119). The fragile X mental retardation protein (FMRP) was also identified as a new IRE1 kinase substrate, possibly regulating cholesterol transport and efferocytosis (150). In addition, pumilio, a RNA-binding protein in *drosophila*, was shown to be phosphorylated by IRE1, allowing its binding to XBP1 mRNA and protecting it against IRE1 RIDD activity (151). PUM1, the human homolog of pumilio, has also been shown to be phosphorylated by IRE1 (151). Another study has reported an interaction between IRE1 and

the I κ B α kinase complex (IKK). The use of a kinase defective IRE1 mutant (K599A) revealed a reduced phosphorylation of I κ B α kinase, rescued by transfection with a WT IRE1 construct (152). This links IRE1 with the NF- κ B pathway, involved in many cellular processes such as inflammation or apoptosis. At last, the sphingosine 1-phosphate lyase (SPL) was shown to be phosphorylated by IRE1, inhibiting its enzymatic activity and resulting in activation of a mitochondrial-dependent UPR (153). Even though the experiments performed were mostly *in vitro*, these discoveries open a new field in the exploration of IRE1 kinase-dependent signaling and raise questions about potential implication of IRE1 in processes not limited to the UPR.




Figure 4. Summary of the current IRE1 interactome. Clusters of IRE1 interactors regulating IRE1 signaling (A) or exerting noncanonical functions (B). C, integrated view of IRE1 interactions mediating its signaling and other functions. (*) indicates IRE1 β -specific interactions. AIP1, ASK1-interacting protein 1; FMRP, fragile X mental retardation protein; IKK, I κ B α kinase complex; IRE1, inositol-requiring enzyme 1; MITOL, mitochondria ubiquitin ligase MARCHF5; SPL, sphingosine 1-phosphate lyase

Concluding remarks and outstanding questions

IRE1 first appeared as a transducer of the UPR, in order to restore a balance in ER homeostasis and help cells adapt to stress. The evolutive interactome unveils possible new roles of IRE1 in mammals. IRE1 is connected with immunity, inflammation, and metabolism and is therefore involved in disease progression (154, 155). While most of the studies focus on IRE1 RNase activity, understanding the dynamic of IRE1 interactions may also provide useful insights on disease development and potential therapies. This review summarizes the known IRE1 interactors and classifies them according to their role in regulating IRE1 signaling (Fig. 4A) or exerting noncanonical (RNase independent) functions (Fig. 4B). While a majority of the interactors were characterized to regulate IRE1 canonical signaling, an increasing number of evidences tend to attribute to IRE1 a wide range of diverse functions (Fig. 4C).

IRE1 signaling can therefore be regulated by interactors at every step of its activation. We term this protein interaction network regulating IRE1 signaling functions as the IRE1 signalosome. The latter can be modulated by specific signals; hence, deciphering how the regulation of the IRE1 cascade exerts different cellular outcomes might represent an appealing avenue to better understand IRE1 signaling. We found recent studies describing new targets of the IRE1 kinase activity. Protein phosphorylation is a crucial mechanism in transducing signals and regulating cellular functions (156). Considering IRE1 as a regular kinase opens a new field of investigations. Studying the sequence and modeling the complex of the known IRE1-dependent phosphorylation could provide some insights on potential other targets. Lastly, IRE1 is known for its role in regulating the translational load through RIDD in the cytosol. However, some studies reported a possible localization of IRE1 in the nucleus (143, 157–159) and confirming this would require further investigations. Considering the continuity between ER membrane and nuclear envelope, we can hypothesize a nonconventional localization of IRE1, where it would unveil a new landscape of regulators and regulate gene expression independently from RIDD.

Acknowledgments—We thank Dr Xavier Guillory for his assistance regarding IRE1 structural representations.

Author contributions—S. L. G., H. L., M. A., and E. C. investigation; S. L. G., H. L., M. A., and E. C. conceptualization; S. L. G., H. L., M. A., and E. C. writing-original draft; S. L. G., H. L., M. A., and E. C. writing-review and editing.

Funding and additional information—This work was funded by grants from Institut National du Cancer (INCa, PLBio) and Fondation pour la Recherche Médicale (FRM; EQU202403018041), INSERM (International Research Project – TUPRIC) to E. C. S. L. G. was funded by a PhD scholarship from the French Ministry of research and H. L. was funded by a postdoctoral fellowship from Région Bretagne.

Conflict of interest—E. C. is the founder of Thabor Therapeutics (<https://www.thabor-tx.com/>). The other authors declare that they have no conflicts of interest with the contents of this article.

Abbreviations—The abbreviations used are: AH, amphipathic helix; ASK1, apoptotic signaling kinase 1; ATF, activating transcription factor; BI-1, BAX inhibitor-1; CD, cytosolic domain; ER, endoplasmic reticulum; ERAD, ER-associated degradation; IRE1, inositol-requiring enzyme 1; JNK, c-Jun-N-terminal-kinase; MHC, major histocompatibility peptide-binding complex; NLD, N-terminal luminal domain; NM IIB, nonmuscle myosin IIB; PDI, protein disulfide isomerase; PERK, PKR-like endoplasmic reticulum kinase; PTM, posttranslational modification; RAC, ribosome-associated complex; RIDD, regulated IRE1-dependent decay; RNH, ribonuclease inhibitor; SRP, signal recognition particle; STIM1, stromal interaction molecule 1; UPR, unfolded protein response; XBP1, X-box binding protein 1.

References

- Garnier, C. (1897) *Les filaments basaux des cellules glandulaires*. Note préliminaire, Bibliographie Anatomique [Paris]: [Berger-Levrault]
- Porter, K. R., Claude, A., and Fullam, E. F. (1945) A study of tissue culture cells by electron microscopy. *J. Exp. Med.* **81**, 233–246
- Ashby, M. C., and Tepikin, A. V. (2001) ER calcium and the functions of intracellular organelles. *Semin. Cell Dev. Biol.* **12**, 11–17
- Fagone, P., and Jackowski, S. (2009) Membrane phospholipid synthesis and endoplasmic reticulum function. *J. Lipid Res.* **50**, S311–S316
- Braakman, I., and Hebert, D. N. (2013) Protein folding in the endoplasmic reticulum. *Cold Spring Harb. Perspect. Biol.* **5**, a013201
- Almanza, A., Carlesso, A., Chintha, C., Creedican, S., Doultinos, D., Leuzzi, B., et al. (2019) Endoplasmic reticulum stress signalling - from basic mechanisms to clinical applications. *FEBS J.* **286**, 241–278
- Ghaemmaghami, S., Huh, W. K., Bower, K., Howson, R. W., Belle, A., Dephoure, N., et al. (2003) Global analysis of protein expression in yeast. *Nature* **425**, 737–741
- Hanson, S. R., Culyba, E. K., Hsu, T. L., Wong, C. H., Kelly, J. W., and Powers, E. T. (2009) The core trisaccharide of an N-linked glycoprotein intrinsically accelerates folding and enhances stability. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 3131–3136
- Ellgaard, L., and Helenius, A. (2003) Quality control in the endoplasmic reticulum. *Nat. Rev. Mol. Cell Biol.* **4**, 181–191
- Olzmann, J. A., Kopito, R. R., and Christianson, J. C. (2013) The mammalian endoplasmic reticulum-associated degradation system. *Cold Spring Harb. Perspect. Biol.* **5**, a013185
- Schröder, M., and Kaufman, R. J. (2005) The mammalian unfolded protein response. *Annu. Rev. Biochem.* **74**, 739–789
- Shore, G. C., Papa, F. R., and Oakes, S. A. (2011) Signaling cell death from the endoplasmic reticulum stress response. *Curr. Opin. Cell Biol.* **23**, 143–149
- McGrath, E. P., Centonze, F. G., Chevet, E., Avril, T., and Lafont, E. (2021) Death sentence: the tale of a fallen endoplasmic reticulum. *Biochim. Biophys. Acta Mol. Cell Res.* **1868**, 119001
- Hughes, D., and Mallucci, G. R. (2019) The unfolded protein response in neurodegenerative disorders – therapeutic modulation of the PERK pathway. *FEBS J.* **286**, 342–355
- Madden, E., Logue, S. E., Healy, S. J., Manie, S., and Samali, A. (2019) The role of the unfolded protein response in cancer progression: from oncogenesis to chemoresistance. *Biol. Cell* **111**, 1–17
- Hetz, C., Zhang, K., and Kaufman, R. J. (2020) Mechanisms, regulation and functions of the unfolded protein response. *Nat. Rev. Mol. Cell Biol.* **21**, 421–438
- Vattem, K. M., and Wek, R. C. (2004) Reinitiation involving upstream ORFs regulates ATF4 mRNA translation in mammalian cells. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 11269–11274
- B'chir, W., Maurin, A. C., Carraro, V., Averous, J., Jousse, C., Muranishi, Y., et al. (2013) The eIF2α/ATF4 pathway is essential for stress-induced autophagy gene expression. *Nucleic Acids Res.* **41**, 7683–7699

19. Harding, H. P., Zhang, Y., Zeng, H., Novoa, I., Lu, P. D., Calfon, M., et al. (2003) An integrated stress response regulates amino acid metabolism and resistance to oxidative stress. *Mol. Cell* **11**, 619–633
20. Novoa, I., Zhang, Y., Zeng, H., Jungreis, R., Harding, H. P., and Ron, D. (2003) Stress-induced gene expression requires programmed recovery from translational repression. *EMBO J.* **22**, 1180–1187
21. Novoa, I., Zeng, H., Harding, H. P., and Ron, D. (2001) Feedback inhibition of the unfolded protein response by GADD34-mediated dephosphorylation of eIF2 α . *J. Cell Biol.* **153**, 1011–1022
22. Haze, K., Yoshida, H., Yanagi, H., Yura, T., and Mori, K. (1999) Mammalian transcription factor ATF6 is synthesized as a transmembrane protein and activated by proteolysis in response to endoplasmic reticulum stress. *Mol. Biol. Cell* **10**, 3787–3799
23. Wu, J., Rutkowski, D. T., Dubois, M., Swathirajan, J., Saunders, T., Wang, J., et al. (2007) ATF6 α Optimizes long-term endoplasmic reticulum function to protect cells from chronic stress. *Dev. Cell* **13**, 351–364
24. Shoulders, M. D., Ryno, L. M., Genereux, J. C., Moresco, J. J., Tu, P. G., Wu, C., et al. (2013) Stress-independent activation of XBP1s and/or ATF6 reveals three functionally diverse ER proteostasis environments. *Cell Rep.* **3**, 1279–1292
25. Mori, K., Ma, W., Gething, M. J., and Sambrook, J. (1993) A transmembrane protein with a cdc2+/CDC28-related kinase activity is required for signaling from the ER to the nucleus. *Cell* **74**, 743–756
26. Welihinda, A. A., Tirasophon, W., Green, S. R., and Kaufman, R. J. (1998) Protein serine/threonine phosphatase Ptc2p negatively regulates the unfolded-protein response by dephosphorylating Ire1p kinase. *Mol. Cell. Biol.* **18**, 1967–1977
27. Cox, J. S., Shamu, C. E., and Walter, P. (1993) Transcriptional induction of genes encoding endoplasmic reticulum resident proteins requires a transmembrane protein kinase. *Cell* **73**, 1197–1206
28. Jurkin, J., Henkel, T., Nielsen, A. F., Minnich, M., Popow, J., Kaufmann, T., et al. (2014) The mammalian tRNA ligase complex mediates splicing of XBP1 mRNA and controls antibody secretion in plasma cells. *EMBO J.* **33**, 2922–2936
29. Kosmaczewski, S. G., Edwards, T. J., Han, S. M., Eckwahl, M. J., Meyer, B. I., Peach, S., et al. (2014) The RtcB RNA ligase is an essential component of the metazoan unfolded protein response. *EMBO Rep.* **15**, 1278–1285
30. Lu, Y., Liang, F. X., and Wang, X. (2014) A synthetic biology approach identifies the mammalian UPR RNA ligase RtcB. *Mol. Cell* **55**, 758–770
31. Sidrauski, C., and Walter, P. (1997) The transmembrane kinase Ire1p is a site-specific Endonuclease that initiates mRNA splicing in the unfolded protein response. *Cell* **90**, 1031–1039
32. Lee, A. H., Iwakoshi, N. N., and Glimcher, L. H. (2003) XBP-1 regulates a subset of endoplasmic reticulum resident chaperone genes in the unfolded protein response. *Mol. Cell. Biol.* **23**, 7448–7459
33. Lee, A. H., Scapa, E. F., Cohen, D. E., and Glimcher, L. H. (2008) Regulation of hepatic lipogenesis by the transcription factor XBP1. *Science* **320**, 1492–1496
34. Yamamoto, K., Suzuki, N., Wada, T., Okada, T., Yoshida, H., Kaufman, R. J., et al. (2008) Human HRD1 promoter Carries a functional unfolded protein response element to which XBP1 but not ATF6 directly binds. *J. Biochem.* **144**, 477–486
35. Hollien, J., and Weissman, J. S. (2006) Decay of endoplasmic reticulum-localized mRNAs during the unfolded protein response. *Science* **313**, 104–107
36. Upton, J. P., Wang, L., Han, D., Wang, E. S., Huskey, N. E., Lim, L., et al. (2012) IRE1 α cleaves select microRNAs during ER stress to Derepress translation of Proapoptotic caspase-2. *Science* **338**, 818–822
37. Oikawa, D., Tokuda, M., Hosoda, A., and Iwawaki, T. (2010) Identification of a consensus element recognized and cleaved by IRE1 α . *Nucleic Acids Res.* **38**, 6265–6273
38. Le Thomas, A., Ferri, E., Marsters, S., Harnoss, J. M., Lawrence, D. A., Zuazo-Gaztelu, I., et al. (2021) Decoding non-canonical mRNA decay by the endoplasmic-reticulum stress sensor IRE1 α . *Nat Commun* **12**, 7310
39. Yoshida, H., Matsui, T., Yamamoto, A., Okada, T., and Mori, K. (2001) XBP1 mRNA is induced by ATF6 and spliced by IRE1 in response to ER stress to Produce a highly active transcription factor. *Cell* **107**, 881–891
40. Tsuru, A., Imai, Y., Saito, M., and Kohno, K. (2016) Novel mechanism of enhancing IRE1 α -XBP1 signalling via the PERK-ATF4 pathway. *Sci. Rep.* **6**, 24217
41. Chang, T. K., Lawrence, D. A., Lu, M., Tan, J., Harnoss, J. M., Marsters, S. A., et al. (2018) Coordination between two branches of the unfolded protein response determines apoptotic cell fate. *Mol. Cell* **71**, 629–636.e5
42. Lin, J. H., Li, H., Yasumura, D., Cohen, H. R., Zhang, C., Panning, B., et al. (2007) IRE1 signaling affects cell fate during the unfolded protein response. *Science* **318**, 944–949
43. Iwawaki, T., Hosoda, A., Okuda, T., Kamigori, Y., Nomura-Furuwatari, C., Kimata, Y., et al. (2001) Translational control by the ER transmembrane kinase/ribonuclease IRE1 under ER stress. *Nat. Cell Biol.* **3**, 158–164
44. Bertolotti, A., Wang, X., Novoa, I., Jungreis, R., Schlessinger, K., Cho, J. H., et al. (2001) Increased sensitivity to dextran sodium sulfate colitis in IRE1 β -deficient mice. *J. Clin. Invest.* **107**, 585–593
45. Martino, M. B., Jones, L., Brighton, B., Ehre, C., Abdulah, L., Davis, C. W., et al. (2013) The ER stress transducer IRE1 β is required for airway epithelial mucin production. *Mucosal Immunol.* **6**, 639–654
46. Imagawa, Y., Hosoda, A., Sasaka, S., Tsuru, A., and Kohno, K. (2008) RNase domains determine the functional difference between IRE1 α and IRE1 β . *FEBS Lett.* **582**, 656–660
47. Grey, M. J., Cloots, E., Simpson, M. S., LeDuc, N., Serebrenik, Y. V., De Luca, H., et al. (2020) IRE1 β negatively regulates IRE1 α signaling in response to endoplasmic reticulum stress. *J. Cell Biol.* **219**, e201904048
48. Cloots, E., Simpson, M. S., De Nolf, C., Lencer, W. I., Janssens, S., and Grey, M. J. (2021) Evolution and function of the epithelial cell-specific ER stress sensor IRE1 β . *Mucosal Immunol.* **14**, 1235–1246
49. Calfon, M., Zeng, H., Urano, F., Till, J. H., Hubbard, S. R., Harding, H. P., et al. (2002) IRE1 couples endoplasmic reticulum load to secretory capacity by processing the XBP-1 mRNA. *Nature* **415**, 92–96
50. Feldman, H. C., Vidadal, V. N., Potter, Z. E., Papa, F. R., Backes, B. J., and Maly, D. J. (2019) Development of a chemical Toolset for studying the Paralog-specific function of IRE1. *ACS Chem. Biol.* **14**, 2595–2605
51. Cloots, E., Guillet, P., Provost, M., Neidhardt, L., Van de Velde, E., Fayazpour, F., et al. (2023) Activation of goblet-cell stress sensor IRE1 β is controlled by the mucin chaperone AGR2. *EMBO J.* **43**, 1–24
52. Zhang, L., Zhang, C., and Wang, A. (2016) Divergence and conservation of the major UPR branch IRE1-bZIP signaling pathway across eukaryotes. *Sci. Rep.* **6**, 27362
53. Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. (1990) Basic local alignment search tool. *J. Mol. Biol.* **215**, 403–410
54. Cox, J. S., and Walter, P. (1996) A novel mechanism for regulating activity of a transcription factor that controls the unfolded protein response. *Cell* **87**, 391–404
55. Tam, A. B., Koong, A. C., and Niwa, M. (2014) Ire1 has Distinct catalytic mechanisms for XBP1/HAC1 splicing and RIDD. *Cell Rep.* **9**, 850–858
56. Kimmig, P., Diaz, M., Zheng, J., Williams, C. C., Lang, A., Aragón, T., et al. (2012) The unfolded protein response in fission yeast modulates stability of select mRNAs to maintain protein homeostasis. *Elife* **1**, e00048
57. Hollien, J. (2013) Evolution of the unfolded protein response. *Biochim. Biophys. Acta* **1833**, 2458–2463
58. Alonso-López, D., Campos-Laborie, F. J., Gutiérrez, M. A., Lambourne, L., Calderwood, M. A., Vidal, M., et al. (2019) APID database: redefining protein–protein interaction experimental evidences and binary interactomes. *Database (Oxford)* **2019**, baz005
59. Szklarczyk, D., Gable, A. L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., et al. (2019) STRING v11: protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res.* **47**, D607–D613
60. Oughtred, R., Rust, J., Chang, C., Breitkreutz, B. J., Stark, C., Willem, A., et al. (2021) The BioGRID database: a comprehensive biomedical resource of curated protein, genetic, and chemical interactions. *Protein Sci.* **30**, 187–200
61. Kolberg, L., Raudvere, U., Kuzmin, I., Adler, P., Vilo, J., and Peterson, H. (2023) g:Profilor—interoperable web service for functional enrichment

- analysis and gene identifier mapping (2023 update). *Nucleic Acids Res.* **51**, W207–W212
62. Credle, J. J., Finer-Moore, J. S., Papa, F. R., Stroud, R. M., and Walter, P. (2005) On the mechanism of sensing unfolded protein in the endoplasmic reticulum. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 18773–18784
 63. Aragón, T., van Anken, E., Pincus, D., Serafimova, I. M., Korenykh, A. V., Rubio, C. A., et al. (2009) Messenger RNA targeting to endoplasmic reticulum stress signalling sites. *Nature* **457**, 736–740
 64. Zhou, J., Liu, C. Y., Back, S. H., Clark, R. L., Peisach, D., Xu, Z., et al. (2006) The crystal structure of human IRE1 luminal domain reveals a conserved dimerization interface required for activation of the unfolded protein response. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 14343–14348
 65. Lhomond, S., Avril, T., Dejeans, N., Voutetakis, K., Doultsinos, D., McMahon, M., et al. (2018) Dual IRE1 RNase functions dictate glioblastoma development. *EMBO Mol. Med.* **10**, e7929
 66. Halbleib, K., Pesek, K., Covino, R., Hofbauer, H. F., Wunnicke, D., Hänelt, I., et al. (2017) Activation of the unfolded protein response by lipid bilayer stress. *Mol. Cell* **67**, 673–684.e8
 67. Thibault, G., Shui, G., Kim, W., McAlister, G. C., Ismail, N., Gygi, S. P., et al. (2012) The membrane stress response buffers lethal effects of lipid disequilibrium by reprogramming the protein homeostasis network. *Mol. Cell* **48**, 16–27
 68. Ho, N., Yap, W. S., Xu, J., Wu, H., Koh, J. H., Goh, W. W. B., et al. (2020) Stress sensor Ire1 deploys a divergent transcriptional program in response to lipid bilayer stress. *J. Cell Biol.* **219**, e201909165
 69. Volmer, R., van der Ploeg, K., and Ron, D. (2013) Membrane lipid saturation activates endoplasmic reticulum unfolded protein response transducers through their transmembrane domains. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 4628–4633
 70. Kono, N., Amin-Wetzel, N., and Ron, D. (2017) Generic membrane-spanning features endow IRE1α with responsiveness to membrane aberrancy. *Mol. Biol. Cell* **28**, 2318–2332
 71. Cho, H., Stanzione, F., Oak, A., Kim, G. H., Yerneni, S., Qi, L., et al. (2019) Intrinsic structural features of the human IRE1α transmembrane domain sense membrane lipid saturation. *Cell Rep.* **27**, 307–320.e5
 72. Li, X., Sun, S., Appathurai, S., Sundaram, A., Plumb, R., and Mariappan, M. (2020) A molecular mechanism for turning off IRE1α signaling during endoplasmic reticulum stress. *Cell Rep.* **33**, 108563
 73. Sidrauski, C., Cox, J. S., and Walter, P. (1996) tRNA ligase is required for regulated mRNA splicing in the unfolded protein response. *Cell* **87**, 405–413
 74. Peschek, J., Acosta-Alvear, D., Mendez, A. S., and Walter, P. (2015) A conformational RNA zipper promotes intron ejection during non-conventional XBP1 mRNA splicing. *EMBO Rep.* **16**, 1688–1698
 75. Lee, K. P. K., Dey, M., Neculai, D., Cao, C., Dever, T. E., and Sicheri, F. (2008) Structure of the dual enzyme Ire1 reveals the basis for catalysis and regulation in nonconventional RNA splicing. *Cell* **132**, 89–100
 76. Korenykh, A. V., Egea, P. F., Korostelev, A. A., Finer-Moore, J., Zhang, C., Shokat, K. M., et al. (2009) The unfolded protein response signals through high-order assembly of Ire1. *Nature* **457**, 687–693
 77. Rubio, C., Pincus, D., Korenykh, A., Schuck, S., El-Samad, H., and Walter, P. (2011) Homeostatic adaptation to endoplasmic reticulum stress depends on Ire1 kinase activity. *J. Cell Biol.* **193**, 171–184
 78. Concha, N. O., Smallwood, A., Bonnette, W., Totoritis, R., Zhang, G., Federowicz, K., et al. (2015) Long-range inhibitor-induced conformational regulation of human IRE1α endoribonuclease activity. *Mol. Pharmacol.* **88**, 1011–1023
 79. Joshi, A., Newbatt, Y., McAndrew, P. C., Stubbs, M., Burke, R., Richards, M. W., et al. (2015) Molecular mechanisms of human IRE1 activation through dimerization and ligand binding. *Oncotarget* **6**, 13019–13035
 80. Ali, M. M. U., Bagratuni, T., Davenport, E. L., Nowak, P. R., Silva-Santisteban, M. C., Hardcastle, A., et al. (2011) Structure of the Ire1 autophosphorylation complex and implications for the unfolded protein response: structure of the Ire1 autophosphorylation complex and implications for the UPR. *EMBO J.* **30**, 894–905
 81. Prischl, F., Nowak, P. R., Carrara, M., and Ali, M. M. U. (2014) Phosphoregulation of Ire1 RNase splicing activity. *Nat. Commun.* **5**, 3554
 82. Shen, X., Ellis, R. E., Sakaki, K., and Kaufman, R. J. (2005) Genetic interactions due to constitutive and inducible gene regulation mediated by the unfolded protein response in *C. elegans*. *PLOS Genet.* **1**, e37
 83. Wiseman, R. L., Zhang, Y., Lee, K. P. K., Harding, H. P., Haynes, C. M., Price, J., et al. (2010) Flavonol activation defines an unanticipated ligand-binding site in the kinase-RNase domain of IRE1. *Mol. Cell* **38**, 291–304
 84. Amarasinghe, K. N., Pelizzari-Raymundo, D., Carlesso, A., Chevet, E., Eriksson, L. A., and Jalil Mahdizadeh, S. (2022) Sensor dimer disruption as a new mode of action to block the IRE1-mediated unfolded protein response. *Comput. Struct. Biotechnol. J.* **20**, 1584–1592
 85. Gardner, B. M., and Walter, P. (2011) Unfolded proteins are Ire1-activating Ligands that directly induce the unfolded protein response. *Science* **333**, 1891–1894
 86. Kimata, Y., and Kohno, K. (2011) Endoplasmic reticulum stress-sensing mechanisms in yeast and mammalian cells. *Curr. Opin. Cell Biol.* **23**, 135–142
 87. Karagöz, G. E., Acosta-Alvear, D., Nguyen, H. T., Lee, C. P., Chu, F., and Walter, P. (2017) An unfolded protein-induced conformational switch activates mammalian IRE1. *Elife* **6**, e30700
 88. Amin-Wetzel, N., Saunders, R. A., Kamphuis, M. J., Rato, C., Preissler, S., Harding, H. P., et al. (2017) A J-protein Co-chaperone recruits BiP to monomerize IRE1 and repress the unfolded protein response. *Cell* **171**, 1625–1637.e13
 89. Amin-Wetzel, N., Neidhardt, L., Yan, Y., Mayer, M. P., and Ron, D. (2019) Unstructured regions in IRE1α specify BiP-mediated destabilization of the luminal domain dimer and repression of the UPR. *Elife* **8**, e50793
 90. Carrara, M., Prischl, F., Nowak, P. R., Kopp, M. C., and Ali, M. M. (2015) Noncanonical binding of BiP ATPase domain to Ire1 and Perk is dissociated by unfolded protein CH1 to initiate ER stress signaling. *Elife* **4**, e03522
 91. Kopp, M. C., Nowak, P. R., Larburu, N., Adams, C. J., and Ali, M. M. (2018) *In vitro* FRET analysis of IRE1 and BiP association and dissociation upon endoplasmic reticulum stress. *Elife* **7**, e30257
 92. Kopp, M. C., Larburu, N., Durairaj, V., Adams, C. J., and Ali, M. M. U. (2019) UPR proteins IRE1 and PERK switch BiP from chaperone to ER stress sensor. *Nat. Struct. Mol. Biol.* **26**, 1053–1062
 93. Kimata, Y., Oikawa, D., Shimizu, Y., Ishiwata-Kimata, Y., and Kohno, K. (2004) A role for BiP as an adjustor for the endoplasmic reticulum stress-sensing protein Ire1. *J. Cell Biol.* **167**, 445–456
 94. Pincus, D., Chevalier, M. W., Aragón, T., van Anken, E., Vidal, S. E., El-Samad, H., et al. (2010) BiP binding to the ER-stress sensor Ire1 tunes the Homeostatic Behavior of the unfolded protein response. *PLoS Biol.* **8**, e1000415
 95. Neidhardt, L., Cloots, E., Friemel, N., Weiss, C. A. M., Harding, H. P., McLaughlin, S. H., et al. (2023) The IRE1β-mediated unfolded protein response is repressed by the chaperone AGR2 in mucin producing cells. *EMBO J.* **43**, 1–35
 96. Sepulveda, D., Rojas-Rivera, D., Rodríguez, D. A., Groenendyk, J., Köhler, A., Lebeaupin, C., et al. (2018) Interactome screening identifies the ER luminal chaperone Hsp47 as a regulator of the unfolded protein response transducer IRE1α. *Mol. Cell* **69**, 238–252.e7
 97. Liu, C. Y., Xu, Z., and Kaufman, R. J. (2003) Structure and intermolecular interactions of the luminal dimerization domain of human IRE1α. *J. Biol. Chem.* **278**, 17680–17687
 98. Eletto, D., Eletto, D., Dersh, D., Gidalevitz, T., and Argon, Y. (2014) Protein disulfide isomerase A6 controls the decay of IRE1α signaling via disulfide-dependent association. *Mol. Cell* **53**, 562–576
 99. Yu, J., Li, T., Liu, Y., Wang, X., Zhang, J., Wang, X., et al. (2020) Phosphorylation switches protein disulfide isomerase activity to maintain proteostasis and attenuate ER stress. *EMBO J.* **39**, e103841
 100. Belyi, V., Zuazo-Gaztelu, I., Alamban, A., Ashkenazi, A., and Walter, P. (2022) Endoplasmic reticulum stress activates human IRE1α through reversible assembly of inactive dimers into small oligomers. *Elife* **11**, e74342
 101. Plumb, R., Zhang, Z. R., Appathurai, S., and Mariappan, M. (2015) A functional link between the co-translational protein translocation pathway and the UPR. *Elife* **4**, e07426

102. Sundaram, A., Plumb, R., Appathurai, S., and Mariappan, M. (2017) The Sec61 translocon limits IRE1 α signaling during the unfolded protein response. *Elife*. **6**, e27187
103. Wu, I. H., Yoon, J. S., Yang, Q., Liu, Y., Skach, W., and Thomas, P. (2021) A role for the ribosome-associated complex in activation of the IRE1 branch of UPR. *Cell Rep.* **35**, 109217
104. Morita, S., Villalta, S. A., Feldman, H. C., Register, A. C., Rosenthal, W., Hoffmann-Petersen, I. T., et al. (2017) Targeting ABL-IRE1 α signaling spares ER-Stressed Pancreatic β cells to reverse autoimmune diabetes. *Cell Metab.* **25**, 1207
105. Papaioannou, A., Centonze, F., Metais, A., Maurel, M., Negroni, L., Gonzalez-Quiroz, M., et al. (2022) Stress-induced tyrosine phosphorylation of RtcB modulates IRE1 activity and signaling outputs. *Life Sci. Alliance* **5**, e202201379
106. He, Y., Beatty, A., Han, X., Ji, Y., Ma, X., Adelstein, R. S., et al. (2012) Nonmuscle Myosin II B links cytoskeleton to IRE1 α signaling during ER stress. *Dev. Cell* **23**, 1141–1152
107. Dufey, E., Bravo-San Pedro, J. M., Eggers, C., González-Quiroz, M., Urra, H., Sagredo, A. I., et al. (2020) Genotoxic stress triggers the activation of IRE1 α -dependent RNA decay to modulate the DNA damage response. *Nat. Commun.* **11**, 2401
108. Vandenabeele, P., Bultynck, G., and Savvides, S. N. (2022) Pore-forming proteins as drivers of membrane permeabilization in cell death pathways. *Nat. Rev. Mol. Cell Biol.* **24**, 312–333
109. Hetz, C., Bernasconi, P., Fisher, J., Lee, A. H., Bassik, M. C., Antonsson, B., et al. (2006) Proapoptotic BAX and BAK modulate the unfolded protein response by a direct interaction with IRE1 α . *Science* **312**, 572–576
110. Rodriguez, D. A., Zamorano, S., Lisbona, F., Rojas-Rivera, D., Urra, H., Cubillos-Ruiz, J. R., et al. (2012) BH3-only proteins are part of a regulatory network that control the sustained signalling of the unfolded protein response sensor IRE1 α : BH3-only controls the sustained signalling of IRE1 α . *EMBO J.* **31**, 2322–2335
111. Tavernier, Q., Bennana, E., Poindessous, V., Schaeffer, C., Rampoldi, L., Pietrancosta, N., et al. (2018) Regulation of IRE1 RNase activity by the Ribonuclease inhibitor 1 (RNH1). *Cell Cycle* **17**, 1901–1916
112. Lisbona, F., Rojas-Rivera, D., Thielen, P., Zamorano, S., Todd, D., Martinon, F., et al. (2009) BAX inhibitor-1 is a negative regulator of the ER stress sensor IRE1 α . *Mol. Cell* **33**, 679–691
113. Baily-Maitre, B., Belgardt, B. F., Jordan, S. D., Coornaert, B., von Freyend, M. J., Kleinridders, A., et al. (2010) Hepatic Bax inhibitor-1 inhibits IRE1 α and protects from obesity-associated insulin resistance and glucose intolerance. *J. Biol. Chem.* **285**, 6198–6207
114. Nguyén, D. T., Kebache, S., Fazel, A., Wong, H. N., Jenna, S., Emadali, A., et al. (2004) Nck-dependent activation of Extracellular signal-regulated kinase-1 and regulation of cell survival during endoplasmic reticulum stress. *Mol. Biol. Cell* **15**, 4248–4260
115. Han, D., Lerner, A. G., Vande Walle, L., Upton, J. P., Xu, W., Hagen, A., et al. (2009) IRE1 α kinase activation modes control alternate endoribonuclease outputs to determine divergent cell fates. *Cell* **138**, 562–575
116. Guo, J., and Polymenis, M. (2006) Dcr2 targets Ire1 and downregulates the unfolded protein response in *Saccharomyces cerevisiae*. *EMBO Rep.* **7**, 1124–1127
117. Lu, G., Ota, A., Ren, S., Franklin, S., Rau, C. D., Ping, P., et al. (2013) PPM1l encodes an inositol requiring-protein 1 (IRE1) specific phosphatase that regulates the functional outcome of the ER stress response. *Mol. Metab.* **2**, 405–416
118. Qiu, Y., Mao, T., Zhang, Y., Shao, M., You, J., Ding, Q., et al. (2010) A crucial role for RACK1 in the regulation of glucose-stimulated IRE1 α activation in Pancreatic β cells. *Sci. Signal.* **3**, ra7
119. Liu, D., Liu, X., Zhou, T., Yao, W., Zhao, J., Zheng, Z., et al. (2016) IRE1–RACK1 axis orchestrates ER stress preconditioning-elicited cytoprotection from ischemia/reperfusion injury in liver. *J. Mol. Cell Biol.* **8**, 144–156
120. Lv, X., Lu, X., Cao, J., Luo, Q., Ding, Y., Peng, F., et al. (2023) Modulation of the proteostasis network promotes tumor resistance to oncogenic KRAS inhibitors. *Science* **381**, eabn4180
121. Gu, F., Nguyen, D. T., Stuible, M., Dubé, N., Tremblay, M. L., and Chevet, E. (2004) Protein-tyrosine phosphatase 1B potentiates IRE1 signaling during endoplasmic reticulum stress. *J. Biol. Chem.* **279**, 49689–49693
122. Mao, T., Shao, M., Qiu, Y., Huang, J., Zhang, Y., Song, B., et al. (2011) PKA phosphorylation couples hepatic inositol-requiring enzyme 1 α to glucagon signaling in glucose metabolism. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 15852–15857
123. Shental-Bechor, D., and Levy, Y. (2008) Effect of glycosylation on protein folding: a close look at thermodynamic stabilization. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 8256–8261
124. Liu, C. Y., Schröder, M., and Kaufman, R. J. (2000) Ligand-independent dimerization activates the stress response kinases IRE1 and PERK in the lumen of the endoplasmic reticulum. *J. Biol. Chem.* **275**, 24881–24885
125. Liu, C. Y., Wong, H. N., Schauerte, J. A., and Kaufman, R. J. (2002) The protein kinase/endoribonuclease IRE1 α that signals the unfolded protein response has a luminal N-terminal ligand-independent dimerization domain. *J. Biol. Chem.* **277**, 18346–18356
126. Gao, B., Lee, S. M., Chen, A., Zhang, J., Zhang, D. D., Kannan, K., et al. (2008) Synoviolin promotes IRE1 ubiquitination and degradation in synovial fibroblasts from mice with collagen-induced arthritis. *EMBO Rep.* **9**, 480–485
127. Sun, S., Shi, G., Sha, H., Ji, Y., Han, X., Shu, X., et al. (2015) IRE1 α is an endogenous substrate of endoplasmic-reticulum-associated degradation. *Nat. Cell Biol.* **17**, 1546–1555
128. Hinte, F., Müller, J., and Brune, W. (2021) Viral-mediated tethering to SEL1L facilitates endoplasmic reticulum-associated degradation of IRE1. *J. Virol.* **95**, e01990–e02020
129. Zhu, X., Zhang, J., Sun, H., Jiang, C., Dong, Y., Shan, Q., et al. (2014) Ubiquitination of inositol-requiring enzyme 1 (IRE1) by the E3 ligase CHIP mediates the IRE1/TRAFF/JNK pathway. *J. Biol. Chem.* **289**, 30567–30577
130. Arshad, M., Ye, Z., Gu, X., Wong, C. K., Liu, Y., Li, D., et al. (2013) RNF13, a RING Finger protein, mediates endoplasmic reticulum stress-induced apoptosis through the inositol-requiring enzyme (IRE1 α)/c-Jun NH2-terminal kinase pathway. *J. Biol. Chem.* **288**, 8726–8736
131. Takeda, K., Nagashima, S., Shiiba, I., Uda, A., Tokuyama, T., Ito, N., et al. (2019) MITOL prevents ER stress-induced apoptosis by IRE1 α ubiquitylation at ER–mitochondria contact sites. *EMBO J.* **38**, e100999
132. Lerner, A. G., Upton, J. P., Praveen, P. V. K., Ghosh, R., Nakagawa, Y., Igbaria, A., et al. (2012) IRE1 α induces thioredoxin-interacting protein to activate the NLRP3 inflammasome and promote programmed cell death during endoplasmic reticulum stress. *Cell Metab.* **16**, 250–264
133. Sandow, J. J., Dorstyn, L., O'Reilly, L. A., Tailler, M., Kumar, S., Strasser, A., et al. (2014) ER stress does not cause upregulation and activation of caspase-2 to initiate apoptosis. *Cell Death Differ.* **21**, 475–480
134. Merksamer, P. I., and Papa, F. R. (2010) The UPR and cell fate at a glance. *J. Cell Sci.* **123**, 1003–1006
135. Urano, F., Wang, X., Bertolotti, A., Zhang, Y., Chung, P., Harding, H. P., et al. (2000) Coupling of stress in the ER to activation of JNK protein kinases by transmembrane protein kinase IRE1. *Science* **287**, 664–666
136. Nishitoh, H., Matsuzawa, A., Tobiume, K., Saegusa, K., Takeda, K., Inoue, K., et al. (2002) ASK1 is essential for endoplasmic reticulum stress-induced neuronal cell death triggered by expanded polyglutamine repeats. *Genes Dev.* **16**, 1345–1355
137. Luo, D., He, Y., Zhang, H., Yu, L., Chen, H., Xu, Z., et al. (2008) AIP1 is critical in transducing IRE1-mediated endoplasmic reticulum stress response. *J. Biol. Chem.* **283**, 11905–11912
138. van Vliet, A. R., Giordano, F., Gerlo, S., Segura, I., Van Eygen, S., Molenberghs, G., et al. (2017) The ER stress sensor PERK Coordinates ER–plasma membrane contact site formation through interaction with filamin-A and F-actin remodeling. *Mol. Cell* **65**, 885–899.e6
139. Urra, H., Henriquez, D. R., Cánovas, J., Villarroel-Campos, D., Carreras-Sureda, A., Pulgar, E., et al. (2018) IRE1 α governs cytoskeleton remodelling and cell migration through a direct interaction with filamin A. *Nat. Cell Biol.* **20**, 942–953
140. Augusto, L., Martynowicz, J., Amin, P. H., Alakhras, N. S., Kaplan, M. H., Wek, R. C., et al. (2020) Toxoplasma gondii Co-opts the unfolded

- protein response to enhance migration and Dissemination of infected Host cells. *mBio* **11**, e00915–e00920
141. Zaman, M. F., Nenadic, A., Radojičić, A., Rosado, A., and Beh, C. T. (2020) Sticking with it: ER-PM membrane contact sites as a Coordinating Nexus for regulating lipids and proteins at the cell Cortex. *Front. Cell Dev. Biol.* **8**, 675
 142. Li, C., Qian, T., He, R., Wan, C., Liu, Y., and Yu, H. (2021) Endoplasmic reticulum–plasma membrane contact sites: regulators, mechanisms, and physiological functions. *Front. Cell Dev. Biol.* **9**, 627700
 143. Acosta-Alvear, D., Karagöz, G. E., Fröhlich, F., Li, H., Walther, T. C., and Walter, P. (2018) The unfolded protein response and endoplasmic reticulum protein targeting machineries converge on the stress sensor IRE1. *Elife* **7**, e43036
 144. Yanagitani, K., Imagawa, Y., Iwawaki, T., Hosoda, A., Saito, M., Kimata, Y., et al. (2009) Cotranslational targeting of XBP1 protein to the membrane promotes Cytoplasmic splicing of its Own mRNA. *Mol. Cell* **34**, 191–200
 145. Yanagitani, K., Kimata, Y., Kadokura, H., and Kohno, K. (2011) Translational pausing ensures membrane targeting and Cytoplasmic splicing of XBP1u mRNA. *Science* **331**, 586–589
 146. Kanda, S., Yanagitani, K., Yokota, Y., Esaki, Y., and Kohno, K. (2016) Autonomous translational pausing is required for XBP1u mRNA recruitment to the ER via the SRP pathway. *Proc. Natl. Acad. Sci. U. S. A.* **113**, E5886–E5895
 147. Son, S. M., Byun, J., Roh, S. E., Kim, S. J., and Mook-Jung, I. (2014) Reduced IRE1α mediates apoptotic cell death by disrupting calcium homeostasis via the InsP3 receptor. *Cell Death Dis.* **5**, e1188
 148. Carreras-Sureda, A., Jaña, F., Urra, H., Durand, S., Mortenson, D. E., Sagredo, A., et al. (2019) Non-canonical function of IRE1α determines mitochondria-associated endoplasmic reticulum composition to control calcium transfer and bioenergetics. *Nat. Cell Biol.* **21**, 755–767
 149. Carreras-Sureda, A., Zhang, X., Laubry, L., Brunetti, J., Koenig, S., Wang, X., et al. (2023) The ER stress sensor IRE1 interacts with STIM1 to promote store-operated calcium entry, T cell activation, and muscular differentiation. *Cell Rep.* **42**, 113540
 150. Yildirim, Z., Baboo, S., Hamid, S. M., Dogan, A. E., Tufanlı, O., Robichaud, S., et al. (2022) Intercepting IRE1 kinase-FMRP signaling prevents atherosclerosis progression. *EMBO Mol. Med.* **14**, e15344
 151. Cairrão, F., Santos, C. C., Le Thomas, A., Marsters, S., Ashkenazi, A., and Domingos, P. M. (2022) Pumilio protects Xbp1 mRNA from regulated Ire1-dependent decay. *Nat. Commun.* **13**, 1587
 152. Tam, A. B., Mercado, E. L., Hoffmann, A., and Niwa, M. (2012) ER stress activates NF-κB by integrating functions of basal IKK activity, IRE1 and PERK. *PLoS One* **7**, e45078
 153. Yildirim, A. D., Citir, M., Dogan, A. E., Veli, Z., Yildirim, Z., Tufanlı, O., et al. (2022) ER stress-induced sphingosine-1-phosphate lyase phosphorylation potentiates the mitochondrial unfolded protein response. *J. Lipid Res.* **63**, 100279
 154. Huang, S., Xing, Y., and Liu, Y. (2019) Emerging roles for the ER stress sensor IRE1α in metabolic regulation and disease. *J. Biol. Chem.* **294**, 18726–18741
 155. Dong, H., Adams, N. M., Xu, Y., Cao, J., Allan, D. S. J., Carlyle, J. R., et al. (2019) The IRE1 endoplasmic reticulum stress sensor activates natural killer cell immunity in part by regulating c-Myc. *Nat. Immunol.* **20**, 865–878
 156. Arditò, F., Giuliani, M., Perrone, D., Troiano, G., and Lo Muzio, L. (2017) The crucial role of protein phosphorylation in cell signaling and its use as targeted therapy (Review). *Int. J. Mol. Med.* **40**, 271–280
 157. Niwa, M., Sidrauski, C., Kaufman, R. J., and Walter, P. (1999) A role for Presenilin-1 in nuclear accumulation of Ire1 fragments and induction of the mammalian unfolded protein response. *Cell* **99**, 691–702
 158. Lee, K., Tirasophon, W., Shen, X., Michalak, M., Prywes, R., Okada, T., et al. (2002) IRE1-mediated unconventional mRNA splicing and S2P-mediated ATF6 cleavage merge to regulate XBP1 in signaling the unfolded protein response. *Genes Dev.* **16**, 452–466
 159. Goffin, L., Vodala, S., Fraser, C., Ryan, J., Timms, M., Meusburger, S., et al. (2006) The unfolded protein response transducer Ire1p Contains a nuclear localization sequence recognized by Multiple β Importins. *Mol. Biol. Cell* **17**, 5309–5323