

EGU24-419, updated on 12 Apr 2024 https://doi.org/10.5194/egusphere-egu24-419 EGU General Assembly 2024 © Author(s) 2024. This work is distributed under the Creative Commons Attribution 4.0 License.



Exploring the landscape heterogeneity and the hydrological diversity in three contrasted observatories of the French critical zone research infrastructure OZCAR

Julien Ackerer¹, Sylvain Kuppel², Isabelle Braud³, Sylvain Pasquet⁴, Ophélie Fovet⁵, Anne Probst⁶, Marie Claire Pierret⁷, Laurent Ruiz⁵, Tiphaine Tallec⁸, Nolwen Lesparre⁷, Sylvain Weill⁷, Christophe Flechard⁵, Jean Luc Probst⁶, Jean Marçais³, Agnes Riviere⁹, Florence Habets¹⁰, Sandrine Anquetin¹, and Jerome Gaillardet¹¹

¹Institut des Géosciences de l'Environnement (IGE), IRD, UGA, Saint Martin D'heres, France (julien.ackerer@orange.fr) ²Géosciences Environnement Toulouse, CNRS-IRD-UPS-CNES, 14 Avenue Edouard Belin, 31400, Toulouse, France

³INRAE, RiverLy, 5 Rue de la Doua, 69625, Villeurbanne, France

⁴Observatoire des Sciences de l'Univers, ECCE TERRA, UAR 3455, CNRS, Sorbonne Université, Paris, France & UMR 7619 METIS, Sorbonne Université, CNRS, EPHE, 75005 Paris, France.

⁵INRAE, 65 rue de St Brieuc, 35042, Rennes, France.

⁶Laboratoire Ecologie Fonctionnelle et Environnement, ENSAT, Avenue de l'Agrobiopole, 31326, Castanet-Tolosan, France ⁷Institut Terre et Environnement de Strasbourg, ITES, 5 rue Descartes, 67084, Strasbourg, France.

⁸Centre d'Etudes Spatiales de la Biosphère, CESBIO, 13 Avenue du Colonel Roche, 31400, Toulouse, France

⁹Centre de Géosciences Mines Paris, PSL, 60 Bd St Michel, 75006, Paris, France.

¹⁰Laboratoire de Géologie, CNRS UMR 8538, École Normale Supérieure, PSL University, IPSL, Paris, France.

¹¹Institut de Physique du Globe, IPGP, 1 rue Jussieu, 75005, Paris, France.

The French OZCAR critical zone network offers the opportunity to conduct multi-site studies and to explore the critical zone functioning under contrasted climate, geology, vegetation and land use. In this study, an integrated modeling of the water cycle is performed with the ecohydrological model EcH₂O-iso in three long-term observatories: (1) the Naizin watershed characterized by an oceanic climate, a metamorphic bedrock and an intensive agriculture (north-west of France, AgrHyS observatory); (2) the Aurade watershed, a watershed with a warmer semi-continental oceanic climate, a sedimentary geological substratum and a crop cover with a wheat-sunflower rotation (south-west of France, Aurade observatory) and; (3) the Strengbach watershed characterized by a mountain climate, a granitic bedrock, and a beech-spruce forest cover (north-east of France, OHGE observatory).

Modeling robustness is evaluated by taking advantage of the large database for critical zone sciences including stream flow, water level in piezometers, and evapotranspiration fluxes measured from climatological stations and flux-towers located in the watersheds. Our comparative study brings these general outcomes: (1) the long term CZ evolution controlling the regolith thickness strongly impacts the total water storage in watersheds; (2) the Quaternary geomorphological evolution influences the current hydrological partitioning and the separation of

hydrologically active and inactive water storage; (3) Both internal watershed characteristics and external forcings, such as current atmospheric forcing and recent land use need to be considered to infer stream persistence and to understand hydrological diversity; and (4) the observed hydrological diversity cannot be fully understood without considering a continuum of time scales in CZ evolution.

Overall, this work illustrates the strength of critical zone networks, allowing a new level of multisite and comparative studies that are crossing several observatories and encompassing a wide diversity of geology and climate.