



**HAL**  
open science

# SETS OF CROSS NUMBERS OF SEQUENCES OVER FINITE ABELIAN GROUPS

Aqsa Bashir, Wolfgang Schmid

► **To cite this version:**

Aqsa Bashir, Wolfgang Schmid. SETS OF CROSS NUMBERS OF SEQUENCES OVER FINITE ABELIAN GROUPS. 2024. hal-04546837v2

**HAL Id: hal-04546837**

**<https://hal.science/hal-04546837v2>**

Preprint submitted on 20 Jan 2025

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# SETS OF CROSS NUMBERS OF SEQUENCES OVER FINITE ABELIAN GROUPS

AQSA BASHIR AND WOLFGANG A. SCHMID

ABSTRACT. Let  $G$  be a finite abelian group with  $\exp(G)$  the exponent of  $G$ . Then  $W(G)$  denotes the set of cross numbers of minimal zero-sum sequences over  $G$  and  $w(G)$  denotes the set of all cross numbers of non-trivial zero-sum free sequences over  $G$ . It is clear that  $W(G)$  and  $w(G)$  are bounded subsets of  $\frac{1}{\exp(G)}\mathbb{N}$  with maximum  $K(G)$  and  $k(G)$ , respectively (here  $K(G)$  and  $k(G)$  denote the large and the small cross number of  $G$ , respectively). We give results on the structure of  $W(G)$  and  $w(G)$ . We first show that both sets contain long arithmetic progressions and that only close to the maximum there might be some gaps. Then, we provide groups for which  $W(G)$  and  $w(G)$  actually are arithmetic progressions, and argue that this is rather a rare phenomenon. Finally, we provide some results in case there are gaps.

## 1. INTRODUCTION

Let  $(G, +)$  be a finite abelian group. For a sequence  $S = g_1 \dots g_\ell$  over  $G$ , that is a collection of elements  $g_i$  of  $G$  where repetitions are allowed (for a formal definition and other undefined terminology see below), the cross number of  $S$  is defined as

$$k(S) = \frac{1}{\text{ord}(g_1)} + \dots + \frac{1}{\text{ord}(g_\ell)}.$$

This can be seen as a weighted version of the length  $\ell$  of the sequence. The term cross number was introduced by Krause in 1984 [23, 24]. The cross number is an interesting zero-sum constant and in particular plays an important role in the factorization theory of Krull monoids, see for example [7, 8, 11, 12, 13, 5, 16, 17, 9, 18, 25, 21, 22, 27, 3] for various contributions on the cross number.

In analogy with the Davenport constant of a finite abelian group, one defines the large cross number of  $G$ , denoted  $K(G)$  as the maximal cross number of a minimal zero-sum sequence, and the small cross number of  $G$ , denoted  $k(G)$ , as the maximal cross number of a zero-sum free sequence. Recall that the large Davenport constant of a finite abelian group  $G$ , denoted  $D(G)$ , is defined as the maximal length of a minimal zero-sum sequence of  $G$  while the small Davenport constant, denoted  $d(G)$ , is the maximal length of a zero-sum free sequence.

For the small Davenport constant it is obvious that zero-sum free sequences of each length up to the maximum exist as every subsequence of a zero-sum free sequence is zero-sum free. Also, for the large Davenport constant one can see without difficulty that for each length smaller than the large Davenport constant there exists a minimal zero-sum sequence of that length as well; it suffices to replace two elements in a minimal zero-sum sequence by their sum to obtain a minimal zero-sum sequence whose length is diminished by 1.

By contrast, for the cross number the analogous problem is more subtle. Of course, we cannot obtain all the rational numbers up to the maximum as cross numbers, as the cross number of each sequence over  $G$  is easily seen to be an integral multiple of  $\frac{1}{\exp(G)}$ . The actual question is if each integral multiple of  $\frac{1}{\exp(G)}$  up to the maximum occurs as a cross number. The basic reasoning that we just recalled for the Davenport constant does not translate to this situation, and indeed in earlier work on the subject it was

---

2020 *Mathematics Subject Classification.* 11B13, 11B30, 20M13.

*Key words and phrases.* finite abelian groups, zero-sum sequences, cross numbers.

This work was supported by the Austrian Science Fund FWF (Project Number W1230) and by the ANR (project ANR-21-CE39-0009 - BARRACUDA).

proved that, depending on the group, there are certain restrictions on the values of the cross numbers of minimal zero-sum sequences, see [4, 1]. In recent literature, the sets of values attained by arithmetic invariants (including the sets of elasticities, of catenary degrees, and more) found wide attention, because knowing all the values of the invariants allows a finer understanding of the problem than just knowing its maximum, see for example [14, 2, 6, 15, 28].

In the present paper we continue the investigations on the cross numbers of minimal zero-sum sequences and in addition study the values of cross numbers of zero-sum free sequences. The latter problem did not yet get much attention. It turns out that, depending on the group, the two sets can be essentially identical, with one being a shift of the other by  $\frac{1}{\exp(G)}$ , yet they can also be quite different.

## 2. PRELIMINARIES

Let  $\mathbb{N}$  denote the set of positive integers and let  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ . For integers  $a, b \in \mathbb{Z}$ , we set  $[a, b] = \{z \in \mathbb{Z} \mid a \leq z \leq b\}$  the interval of integers. Given subsets  $A, B \subseteq \mathbb{Z}$  and  $\lambda \in \mathbb{R}$ , we set  $A + B = \{a + b \mid a \in A, b \in B\}$  the sumset of  $A$  and  $B$ , and we set  $\lambda A = \{\lambda a \mid a \in A\}$  the dilation of  $A$  by  $\lambda$ ; since we hardly use it, we do not introduce a notation for the  $s$ -fold sumset of  $A$ , that is the sumset of  $s$  copies of  $A$ .

For  $n \in \mathbb{N}$  let  $C_n$  be a cyclic group of order  $n$ ; we use additive notation. Let  $(G, +, 0)$  be an additive finite abelian group. There exist unique integers  $1 < n_1 \mid \cdots \mid n_r$  such that  $G \cong C_{n_1} \oplus \cdots \oplus C_{n_r}$ . Moreover, there exist unique prime-powers  $1 < q_1 \leq \cdots \leq q_s$  such that  $G \cong C_{q_1} \oplus \cdots \oplus C_{q_s}$ . The integer  $n_r$  is called the exponent of the group, and the order  $\text{ord}(g)$  of each element  $g \in G$  divides  $\exp(G)$ . One calls  $r$  the rank of  $G$  and  $s$  the total rank of  $G$ . In case  $\exp(G)$  is a prime-power, the two coincide and one calls  $G$  a  $p$ -group. A  $p$ -group is called an elementary  $p$ -group if the exponent is prime. For a given prime  $p$  one calls the number of prime powers  $q_i$  that are a  $p$ -power the  $p$ -rank of  $G$ , denoted by  $r_p(G)$ . It is non-zero if and only if the prime  $p$  divides the exponent.

Note that if  $|G| = 1$ , then the exponent is 1, while all the ranks are 0.

As indicated in the introduction, the focus of the paper is on cross numbers of sequences over a finite abelian group  $G$ . Informally, a sequence over a finite abelian group  $G$  is a collection of elements of  $G$  where repetitions of elements are allowed yet the ordering of the terms is typically disregarded. Formally, a sequence over  $G$  is an element of the free abelian monoid  $\mathcal{F}(G)$  over  $G$ . That is, a sequence  $S$  over  $G$  can be written uniquely as  $S = \prod_{g \in G} g^{v_g}$  with  $v_g \in \mathbb{N}_0$  for each  $g \in G$ ; moreover it can be written as  $S = g_1 \dots g_\ell$  where  $g_i \in G$  for each  $i \in [1, \ell]$  and these elements are unique up to ordering. The neutral element of  $\mathcal{F}(G)$  is denoted by 1, unless there is a risk of confusion, and is called the trivial sequence. Sometimes we want to consider only sequences that contain elements in a subset  $G_0 \subseteq G$ ; in that case we use the notation  $\mathcal{F}(G_0)$ .

One calls

- $\sigma(S) = \sum_{g \in G} v_g g = \sum_{i=1}^{\ell} g_i$  the sum of  $S$ ,
- $|S| = \sum_{g \in G} v_g = \ell$  the length of  $S$ ,
- $\mathbf{k}(S) = \sum_{g \in G} \frac{v_g}{\text{ord}(g)} = \sum_{i=1}^{\ell} \frac{1}{\text{ord}(g_i)}$  the cross number of  $S$ .

For a sequence  $S$ , a subsequence  $T$  of  $S$  is a divisor of  $S$  in  $\mathcal{F}(G)$ . We denote by  $\Sigma(S) = \{\sigma(T) : 1 \neq T \mid S\}$  the set of sums of non-trivial subsequences of  $S$ . Moreover,  $\text{supp}(S) = \{g_1, \dots, g_\ell\}$ , called the support of  $S$ , denotes the set of elements that occur in  $S$ . A sequence  $S$  is called a zero-sum sequence if  $\sigma(S) = 0$  and it is called zero-sum free if  $0 \notin \Sigma(S)$ . A non-trivial zero-sum sequence is called a minimal zero-sum sequence (or an atom) if it has no proper non-trivial subsequence that is a zero-sum sequence. Equivalently,  $S = g_1 \dots g_\ell$  is a zero-sum sequence if  $\sum_{i=1}^{\ell} g_i = 0$ , while it is zero-sum free if  $\sum_{i \in I} g_i \neq 0$  for each  $\emptyset \neq I \subseteq [1, \ell]$  (note that  $I = [1, \ell]$  is possible). Furthermore, it is a minimal zero-sum sequence if it is non-trivial with  $\sum_{i=1}^{\ell} g_i = 0$ , yet  $\sum_{i \in I} g_i \neq 0$  for each proper subset  $\emptyset \neq I \subsetneq [1, \ell]$ .

We denote by  $\mathcal{B}(G)$  the set of all zero-sum sequences over  $G$ , by  $\mathcal{A}(G)$  the set of all minimal zero-sum sequences over  $G$  and by  $\mathcal{A}^*(G)$  the set of all non-trivial zero-sum free sequences over  $G$  (note that the trivial sequence is also zero-sum free according to the definition given above).

While it is not central for our current investigation, we mention in passing that  $\mathcal{B}(G)$  is a submonoid of  $\mathcal{F}(G)$ ; this submonoid is atomic and its irreducible elements are precisely the elements of  $\mathcal{A}(G)$ .

The central objects for the current paper are the following two sets. The set of cross numbers of all minimal zero-sum sequences over  $G$ , denoted

$$W(G) = \{k(S) \mid S \in \mathcal{A}(G)\}$$

has been investigated, e.g., in [1, 4, 13]. We also study the related set

$$w(G) = \{k(S) \mid S \in \mathcal{A}^*(G)\}$$

the set of cross numbers of all non-trivial zero-sum free sequences over  $G$ ; sometimes it is technically advantageous to consider  $w^\circ(G) = w(G) \cup \{0\}$ , which would correspond to considering the trivial sequence as well.

As mentioned in the introduction the large cross number of  $G$  is defined by

$$K(G) = \max\{k(S) \mid S \in \mathcal{A}(G)\}$$

and the small cross number of  $G$ , for  $|G| \neq 1$ , by

$$k(G) = \max\{k(S) \mid S \in \mathcal{A}^*(G)\}.$$

In other words,  $K(G) = \max W(G)$  and  $k(G) = \max w(G)$ . If  $|G| = 1$ , then  $K(G) = 1$  and we set  $k(G) = 0$ .

Let  $G = C_{q_1} \oplus \cdots \oplus C_{q_s}$  be a direct sum decomposition of  $G$  into cyclic groups of prime power order. Set

$$k^*(G) = \sum_{i=1}^s \frac{q_i - 1}{q_i} \quad \text{and} \quad K^*(G) = \frac{1}{\exp(G)} + k^*(G).$$

Suppose  $\{e_1, \dots, e_s\}$  is an independent generating set of  $G$  with  $\text{ord}(e_i) = q_i$  for each  $i \in [1, s]$ . Then  $T = e_1^{q_1-1} \cdots e_s^{q_s-1}$  is zero-sum free and  $S = T(e_1 + \cdots + e_s)$  is a minimal zero-sum sequence. This shows that  $K^*(G) \leq K(G)$  and  $k^*(G) \leq k(G)$ . In principle one could use any set of independent elements, but using a set with elements of prime power order yields the best bound.

Given such a lower bound, the question arises if equality holds. Equality holds in particular for  $p$ -groups (see [10, Proposition 5.1.8 and Theorem 5.5.9]) and for some other special cases (see e.g. [11, 21, 22]); we invoke some results in later sections. No example is known where equality does not hold. Krause and Zahlten conjectured in [24, Page 688] that the equality  $K(G) = K^*(G)$  holds for all cyclic groups  $G$ , but even this remains open.

The two constants  $k(G)$  and  $K(G)$  are closely related. It is easy to see that  $\frac{1}{\exp(G)} + k(G) \leq K(G)$ , yet it is not known if equality always holds. However, in case  $K(G) = K^*(G)$ , we have

$$\frac{1}{\exp(G)} + k^*(G) \leq \frac{1}{\exp(G)} + k(G) \leq K(G) = K^*(G) = \frac{1}{\exp(G)} + k^*(G)$$

and thus  $k(G) = k^*(G)$  and also  $\frac{1}{\exp(G)} + k(G) = K(G)$ .

Since we use them sometimes in the current paper, we recall a few results for the Davenport constant. The equality  $D(G) = |G|$  for finite cyclic groups is known and easy to see. Moreover for  $p$ -groups  $G = C_{q_1} \oplus \cdots \oplus C_{q_s}$  one has  $D(G) = 1 + \sum_{i=1}^s (q_i - 1)$ . See for instance [10, Chapter 5] for these and further results.

We end this section with some results on  $W(G)$  and  $w(G)$  that we use frequently.

For every finite abelian group, with  $|G| \neq 1$ , one has  $W(G) \subseteq \frac{1}{\exp(G)}[2, \exp(G)K(G)]$  and  $w(G) \subseteq \frac{1}{\exp(G)}[1, \exp(G)k(G)]$ . If  $|G|$  is even and  $\exp(G) = 2^k m$  with an odd  $m$  and  $G$  does not contain a subgroup of the form  $C_{2^k}^2$ , then  $W(G) \subseteq \frac{2}{\exp(G)}[1, \frac{\exp(G)K(G)}{2}]$ , see [4, Lemma 1].

In the other direction for every finite abelian group with  $|G|$  odd and  $|G| \neq 1$ , one has  $\frac{1}{\exp(G)}[2, \exp(G)] \subseteq \mathbf{W}(G)$ . If  $|G|$  is even, then in general one only has  $\frac{2}{\exp(G)}[1, \frac{\exp(G)}{2}] \subseteq \mathbf{W}(G)$ ; however, if  $\exp(G) = 2^k m$  with an odd  $m$  and  $G$  contains a subgroup of the form  $C_{2^k}^2$ , then we have  $\frac{1}{\exp(G)}[2, \exp(G)] \subseteq \mathbf{W}(G)$  as in the case of groups of odd order, see [4, Theorem 2].

### 3. RESULTS ON $\mathbf{w}(G)$

In the current section, we obtain some results on the structure of the set  $\mathbf{w}(G)$ . More specifically, we show in two different ways that the set is in some sense close to an arithmetic progression, and that deviations can only occur for values close to the maximum  $\mathbf{k}(G)$ . In later sections, we discuss that while for  $G$  a  $p$ -group and in a few other cases, the set is indeed an arithmetic progression, this is not always the case.

First, we establish and recall a few simple lemmas. It is easy to see that  $\mathbf{w}(G)$  contains all small elements; the result for  $\mathbf{W}(G)$  is recalled at the end of the preliminaries.

**Lemma 3.1.** *Let  $G$  be a finite abelian group. Then*

$$\frac{1}{\exp(G)}[1, \exp(G) - 1] \subseteq \mathbf{w}(G).$$

*Proof.* We need to show that for each  $j \in [1, \exp(G) - 1]$ , there is some  $S \in \mathcal{A}^*(G)$  with  $\mathbf{k}(S) = \frac{j}{\exp(G)}$ . Assume  $g \in G$  with  $\text{ord}(g) = \exp(G)$ . Then  $S_j = g^j$  is a zero-sum free sequence for each  $j \in [1, \exp(G) - 1]$  and  $\mathbf{k}(S_j) = \frac{j}{\exp(G)}$ .  $\square$

While this result is basic, it is indeed sharp in some cases, namely when  $G$  is cyclic group of prime power order as in this case, and in this case only, we have  $\mathbf{k}(G) = \frac{\exp(G)-1}{\exp(G)}$ . The following lemma shows that this is indeed the only case. Later, we establish results that yield the existence of larger arithmetic progressions in  $\mathbf{w}(G)$  in case the group has a large rank.

**Lemma 3.2.** *Let  $G$  be a finite abelian group and let  $H \subseteq G$  be a subgroup. Then  $\mathbf{w}(H) \subseteq \mathbf{w}(G)$ , and equality holds if and only if  $H = G$ .*

*Proof.* It is immediate from the definition that  $\mathcal{A}^*(H) \subseteq \mathcal{A}^*(G)$  and thus  $\mathbf{w}(H) \subseteq \mathbf{w}(G)$  by definition. Clearly,  $H = G$  yields  $\mathbf{w}(H) = \mathbf{w}(G)$  and it remains to show the converse. Note by [10, Proposition 5.1.11] that

$$\mathbf{k}(G) \geq \mathbf{k}(H) + \frac{\mathbf{k}(G/H)}{\exp(H)}.$$

Since  $\mathbf{w}(H) = \mathbf{w}(G)$  implies  $\mathbf{k}(H) = \mathbf{k}(G)$ , it means  $\mathbf{k}(G/H) = 0$ , and hence  $|G/H| = 1$  (which is same as  $H = G$ ).  $\square$

The following lemma is a key tool for the current section.

**Lemma 3.3.** *Let  $G_1$  and  $G_2$  be non-trivial finite abelian groups. Then*

$$\mathbf{w}^\circ(G_1) + \mathbf{w}^\circ(G_2) \subseteq \mathbf{w}^\circ(G_1 \oplus G_2) \text{ and } \mathbf{w}(G_1) + \mathbf{w}^\circ(G_2) \subseteq \mathbf{w}(G_1 \oplus G_2).$$

*Proof.* For  $i \in [1, 2]$ , let  $q_i \in \mathbf{w}^\circ(G_i)$ . Since  $\mathbf{w}^\circ(G_i) \subseteq \mathbf{w}^\circ(G_1 \oplus G_2)$  by Lemma 3.2, we may assume that  $q_1 > 0$  and  $q_2 > 0$ . Then there are zero-sum free sequences  $S_i \in \mathcal{F}(G_i)$  with  $\mathbf{k}(S_i) = q_i$ . Since  $S_1 S_2 \in \mathcal{F}(G_1 \oplus G_2)$  is zero-sum free, we obtain that

$$q_1 + q_2 = \mathbf{k}(S_1) + \mathbf{k}(S_2) = \mathbf{k}(S_1 S_2) \in \mathbf{w}^\circ(G_1 \oplus G_2).$$

The second claim is obvious from the first, since 0 is not contained in the left-hand set.  $\square$

We now combine these results with a result from additive combinatorics to show that in certain cases  $\mathbf{w}(G)$  is close to an arithmetic progression.

**Theorem 3.4.** *Let  $G$  be a finite abelian group with  $\exp(G) = n \geq 2$ .*

1. *If  $G = G_1 \oplus \dots \oplus G_s$ , where  $s \in \mathbb{N}$ ,  $\exp(G_i) = n$  for all  $i \in [1, s]$ , then*

$$\frac{1}{n}[1, (n-1)s] \subseteq \mathfrak{w}(G).$$

2. *There exist constants  $c, s^* \in \mathbb{N}$  such that, for all  $s \geq s^*$ ,*

$$\frac{1}{\exp(G)}[1, s \exp(G)k(G) - c] \subseteq \mathfrak{w}(G^s).$$

To see the relevance of the second part of this result, we point out that, for every finite abelian group  $G$  and for every  $s \in \mathbb{N}$ , we have

$$k^*(G^s) = sk^*(G).$$

Thus, since there is no example known of a finite abelian group  $G'$  where  $k(G') \neq k^*(G')$ , there is no example known of a group  $G$  and an integer  $s \in \mathbb{N}$ , for which

$$k(G^s) \neq sk(G).$$

However, if for some finite abelian group  $G$ , we have  $k(G^s) = sk(G)$  for all  $s \in \mathbb{N}$ , then the second part of the theorem shows that for all  $s \in \mathbb{N}$  the set  $\mathfrak{w}(G^s)$  is an arithmetic progression, apart from a globally bounded upper part; to see this it suffices to recall that  $\mathfrak{w}(G^s) \subseteq \frac{1}{\exp(G^s)}[1, \exp(G^s)k(G^s)]$  and that  $\exp(G^s) = \exp(G)$ .

*Proof.* 1. Since  $\frac{1}{n}[1, n-1] \subseteq \mathfrak{w}(G_i)$  for all  $i \in [1, s]$ , we obtain that

$$\begin{aligned} \frac{1}{n}[1, (n-1)s] &\subseteq \underbrace{\frac{1}{n}[0, n-1] + \dots + \frac{1}{n}[0, n-1]}_s \subseteq \mathfrak{w}^\circ(G_1) + \dots + \mathfrak{w}^\circ(G_s) \\ &\subseteq \mathfrak{w}^\circ(G), \end{aligned}$$

2. We set  $A = \exp(G)\mathfrak{w}^\circ(G) \subseteq \mathbb{N}_0$ . Since  $\gcd(A) = 1$  by 1, it follows from [26, Theorem 1.1] that there exists integer  $c \in \mathbb{N}$  such that the  $s$ -fold sumset of  $A$  has the form

$$A' \uplus [c, s \max A - c] \uplus A'' \subseteq \exp(G)\mathfrak{w}^\circ(G^s),$$

with  $A' \subseteq [0, c-2]$  and  $A'' \subseteq s \max A - c + [2, c]$ , for all  $s \geq \max\{1, (|A| - 2)(\max A - 1) \max A\}$ ; we set  $s^*$  equal to this value.

Now suppose in addition that  $(n-1)s^* \geq c$ . Then the first assertion implies that  $[1, c] \subseteq \exp(G)\mathfrak{w}^\circ(G^s)$ . Thus, we obtain that

$$[0, s \max A - c] \uplus A'' \subseteq \exp(G)\mathfrak{w}^\circ(G^s),$$

and hence the assertion follows.  $\square$

There are better estimates for the constant  $c$  than the one given in [26]; see for example [20, 19], however we do not pursue this route for improvement. Instead, we present our second approach to the problem of determining that  $\mathfrak{w}(G)$  contains all small elements. The approach is similar, in that we use again Lemma 3.3. However, we do not impose anymore that all the groups have the same exponent. This has the advantage of being able to apply the result to any group. The drawback is that the application of Lemma 3.3 is less direct. To overcome this issue we need a result on set-addition, which we give in Lemma 3.7.

The main result we obtain is the following theorem. As a corollary we obtain a complete description of  $\mathfrak{w}(G)$  for  $G$  a  $p$ -group.

**Theorem 3.5.** *Let  $G = H \oplus C_n$  with  $\exp(G) = n$ . Then*

$$\frac{1}{n}[1, n-1 + nk^*(H)] \subseteq \mathfrak{w}(G).$$

**Corollary 3.6.** *Let  $G$  be a  $p$ -group. Then*

$$w(G) = \frac{1}{\exp(G)}[1, \exp(G)k(G)].$$

The proof of the following lemma is basic and chances are the result is somewhere in the literature, we include it for lack of a suitable reference. We recall that  $\Delta(A)$  denotes the set of successive distance of  $A$ , that is for  $A = \{a_1, \dots, a_k\}$  with  $a_i < a_{i+1}$  the set is given by  $a_{i+1} - a_i$  for  $i \in [1, k-1]$ .

**Lemma 3.7.** *Let  $A \subseteq \mathbb{Z}$  be a finite set with  $|A| \geq 2$  and  $\max \Delta(A) \leq l$ . Then  $[0, l-1] + A = [\min A, \max A + l - 1]$ .*

*Proof.* Let  $n \in [\min A, \max A + l - 1]$ . We show that  $n \in [0, l-1] + A$ . Let  $a \in A$  be maximal with  $a \leq n$ , which exists since  $\min A \leq n$ . We note that  $n - a \leq l - 1$ . This is due to the fact that if  $a \neq \max A$ , then  $[a+1, a+l]$  contains an element of  $A$  by the condition on  $\Delta(A)$ , while if  $a = \max A$ , then the assertion follows from the fact that  $0 \leq n - a \leq l - 1$ .  $\square$

In the following result we allow rather arbitrary direct sum decomposition of  $G$  into cyclic groups, note though that the exponent of  $G$  is indeed  $n$ . Usually, starting from a given group, the best way to decompose is to impose that each  $q_i$  is a prime power.

**Proposition 3.8.** *Let  $G = \bigoplus_{i=1}^t C_{q_i} \oplus C_n$  where each  $q_i$  divides  $n$ . Then*

$$\sum_{i=1}^t \frac{1}{q_i} [0, q_i - 1] + w^\circ(C_n) \subseteq w^\circ(G)$$

and

$$\sum_{i=1}^t \frac{1}{q_i} [0, q_i - 1] + w(C_n) \subseteq w(G)$$

*Proof.* By repeated application of Lemma 3.3 we have

$$\sum_{i=1}^t w^\circ(C_{q_i}) + w^\circ(C_n) \subseteq w^\circ(G)$$

and

$$\sum_{i=1}^t w(C_{q_i}) + w(C_n) \subseteq w(G).$$

Now, by Lemma 3.1 we have  $\frac{1}{q_i} [0, q_i - 1] \subseteq w^\circ(C_{q_i})$  and the claims follow.  $\square$

Combining the result with earlier lemmas we get a main technical result of this section.

**Proposition 3.9.** *Let  $G = \bigoplus_{i=1}^t C_{q_i} \oplus C_n$  where each  $q_i$  divides  $n$ . Then*

$$\frac{1}{n} [0, n - 1 + n \sum_{i=1}^t \frac{q_i - 1}{q_i}] \subseteq w^\circ(G)$$

and

$$\frac{1}{n} [1, n - 1 + n \sum_{i=1}^t \frac{q_i - 1}{q_i}] \subseteq w(G)$$

*Proof.* We apply the Proposition 3.8 to obtain  $\sum_{i=1}^t \frac{1}{q_i} [0, q_i - 1] + w^\circ(C_n) \subseteq w^\circ(G)$ . Now by Lemma 3.1, we have  $\frac{1}{n} [0, n - 1] \subseteq w^\circ(C_n)$ . By Lemma 3.7, we have  $\frac{n}{q_i} [0, q_i - 1] + [0, n - 1] = [0, n - 1 + \frac{n(q_i - 1)}{q_i}]$ . Then the claim follows by a simple induction.  $\square$

We now conclude this section with the proof of the main results.

*Proof of Theorem 3.5 and Corollary 3.6.* The theorem is a direct consequence of Proposition 3.9; it suffices to impose that  $q_i$  is a prime power for each  $q_i$  and to recall the definition of  $k^*(H)$ . To get the corollary it suffices to note that if  $G$  is a  $p$ -group and we write  $G = H \oplus C_n$  with  $\exp(G) = n$ , then  $k(G) = k^*(G) = \frac{n-1}{n} + k^*(H)$ .  $\square$

#### 4. RESULTS ON $W(G)$

The purpose of this section is to obtain results along the lines of the ones for  $w(G)$  presented in the preceding section. We recall the following result due to Chapman and Geroldinger [4, Theorem 4] that gives a complete description of  $W(G)$  for  $p$ -groups, like we established it in Corollary 3.6 for  $w(G)$ . It is interesting to observe the difference for 2-groups.

**Theorem 4.1.** *Let  $G = C_{n_1} \oplus \cdots \oplus C_{n_r}$  be a finite abelian  $p$ -group with  $1 = n_0 < n_1 \mid \cdots \mid n_r$ .*

1. *Suppose that  $p$  is either odd or that  $p = 2$  with  $n_{r-1} = n_r$ . Then*

$$W(G) = \frac{1}{\exp(G)} [2, \exp(G)K(G)].$$

2. *Suppose  $p = 2$  and  $n_{r-1} < n_r$ . Then*

$$W(G) = \frac{2}{\exp(G)} [1, \frac{\exp(G)}{2}K(G)].$$

The goal of the remainder of the section is to obtain a result like Proposition 3.9 for  $W(G)$  instead of  $w(G)$ . The overall strategy of the proof is the same but the problem is more subtle as we do not have a result like Lemma 3.3. Moreover, we impose right away that each  $q_i$  is a prime power, which anyway is the most relevant case.

**Proposition 4.2.** *Let  $G = \bigoplus_{i=1}^t C_{q_i} \oplus C_n$  where each  $q_i$  is a prime power that divides  $n$ . Then*

$$\sum_{i=1}^t \frac{1}{q_i} [0, q_i - 1] + W(C_n) \subseteq W(G)$$

The proof is similar to that of Proposition 3.8, but a problem is that we cannot simply take the concatenation of zero-sum free sequences. Instead we need to maintain the property that the sequence is a minimal zero-sum sequence while at the same time controlling the cross number. Recall that if  $U_i$  is a minimal zero-sum sequence over  $G_i$  and  $g_i \mid U_i$  for  $i \in [1, 2]$ , then  $(g_1^{-1}U_1)(g_2^{-1}U_2)(g_1 + g_2)$  is a minimal zero-sum sequence over  $G_1 \oplus G_2$  of length  $|U_1| + |U_2| - 1$ . However, to control the cross number we need some information on the orders of the elements involved.

To this end we establish the following lemma. Let  $p$  be a prime, we denote by  $v_p(n)$  the  $p$ -adic valuation, or  $p$ -valuation for short, of a positive natural number  $n$ .

**Lemma 4.3.** *Let  $n \geq 2$ . For each  $w \in W(C_n)$ , there is an  $A \in \mathcal{A}(C_n)$  with  $k(A) = w$  and for each prime divisor  $p$  of  $n$  there exists a  $g \in \text{supp}(A)$  such that  $v_p(\text{ord}(g)) = v_p(n)$ .*

*Proof.* Let  $w \in W(C_n)$ . Then there exists a  $T \in \mathcal{A}(C_n)$  with  $k(T) = w$ . Let  $p_0 \mid n$  be a prime and let  $g \in \text{supp}(T)$  be an element whose order has maximal  $p_0$ -valuation among the orders of elements in  $\text{supp}(T)$ . If  $v_{p_0}(\text{ord}(g)) = v_{p_0}(n)$ , there is nothing to do. Thus, assume that  $v_{p_0}(\text{ord}(g)) < v_{p_0}(n)$ . There exists some  $g_0 \in C_n$  such that  $p_0 g_0 = g$ . We have  $\text{ord}(g_0) = p_0 \text{ord}(g)$  and this means  $v_{p_0}(\text{ord}(g_0)) = v_{p_0}(\text{ord}(g)) + 1$ , while  $v_p(\text{ord}(g_0)) = v_p(\text{ord}(g))$  for every other prime  $p$ . Now, the sequence  $S = (Tg^{-1})(g_0)^{p_0}$  is a minimal zero-sum sequence with the same cross number  $k(S) = k(T) - \frac{1}{\text{ord}(g)} + \frac{p_0}{\text{ord}(g_0)} = k(T) = w$ .

Now if  $v_{p_0}(\text{ord}(g_0)) = v_{p_0}(n)$ , we are done otherwise we repeat the same steps starting with  $S$  instead of  $T$  until we obtain a sequence that contains an element whose order has  $p_0$ -valuation equal to that of  $n$ .

Repeating the argument for each prime  $p_0$  dividing  $n$ , the assertion follows. Note that the above mentioned process only affects the valuations for the prime  $p_0$ , whence there is no interference between the different steps.  $\square$



With this lemma at hand we can prove the proposition.

*Proof of Proposition 4.2.* Assume  $C_{q_i} = \langle e_i \rangle$  where  $\text{ord}(e_i) = q_i$  for  $i \in [1, t]$ . Let  $[1, t] = \uplus_{j=1}^s I_j$  where the order of  $q_i$  for  $i \in I_j$  is a  $p_j$ -power for some prime  $p_j$ ; in addition, we assume that all the  $I_j$  are non-trivial and the  $p_j$  are pairwise distinct. In other words,  $\bigoplus_{j=1}^s (\bigoplus_{i \in I_j} \langle e_i \rangle)$  is a decomposition into  $p$ -groups of  $\bigoplus_{i=1}^t C_{q_i}$ , and  $|I_j|$  is the  $p_j$ -rank of the group.

We have

$$G = C_n \oplus \bigoplus_{j=1}^s (\bigoplus_{i \in I_j} \langle e_i \rangle).$$

Let  $w_G = w_c + \sum_{i=1}^t \frac{j_i}{q_i} \in \mathbb{W}(C_n) + \sum_{i=1}^t \frac{1}{q_i} [0, q_i - 1]$  where  $w_c \in \mathbb{W}(C_n)$  and  $j_i \in [0, q_i - 1]$ , which we can also write as

$$w_G = w_c + \sum_{j=1}^s \sum_{i \in I_j} \frac{j_i}{q_i}.$$

Then by Lemma 4.3, there exists some  $S_c \in \mathcal{A}(C_n)$  such that  $\mathbf{k}(S_c) = w_c$  and for each  $i \in [1, s]$  the sequence  $S_c$  contains an element  $g_i$  whose  $p_i$ -valuation is maximal, that is, its  $p_i$ -valuation is  $\mathbf{v}_{p_i}(n)$ . Note that the  $g_i$  are not necessarily distinct, which causes slight complication.

We now construct, recursively, a sequence with the desired cross number. For clarity, we present the first step in detail and then briefly line out the general step. Let  $g_1$  be an element in  $S_c$  such that the  $p_1$ -valuation of  $g_1$  is maximal. Let

$$g_1^* = g_1 - \sum_{i \in I_1} j_i e_i,$$

then  $\text{ord}(g_1^*) = \text{ord}(g_1)$ . The sequence

$$S_1 = g_1^{-1} g_1^* S_c \prod_{i \in I_1} e_i^{j_i}$$

is a minimal zero-sum sequence over  $C_n \oplus (\bigoplus_{i \in I_1} \langle e_i \rangle)$  and  $\mathbf{k}(S_1) = \mathbf{k}(S_c) - \frac{1}{\text{ord}(g_1)} + \frac{1}{\text{ord}(g_1^*)} + \sum_{i \in I_1} \frac{j_i}{q_i} = w_c + \sum_{i \in I_1} \frac{j_i}{q_i}$ . Moreover  $S_1$  contains an element whose order has  $p_i$ -valuation equal to  $\mathbf{v}_{p_i}(n)$  for each  $i \in [1, s]$ .

The result now follows by repeating this process. Assume that for  $k \in [1, s - 1]$  we have a sequence  $S_k$  over  $C_n \oplus \bigoplus_{j=1}^k (\bigoplus_{i \in I_j} \langle e_i \rangle)$  with cross number  $w_c + \sum_{j=1}^k \sum_{i \in I_j} \frac{j_i}{q_i}$  that contains an element of whose  $p_i$ -valuation is maximal, that is, its  $p_i$ -valuation is  $\mathbf{v}_{p_i}(n)$  for each  $i \in [1, s]$ .

Now let  $g_{k+1}$  be in  $S_k$  such that the  $p_{k+1}$ -valuation of  $g_{k+1}$  is maximal. Let

$$g_{k+1}^* = g_{k+1} - \sum_{i \in I_{k+1}} j_i e_i,$$

then  $\text{ord}(g_{k+1}^*) = \text{ord}(g_{k+1})$ . The sequence

$$S_{k+1} = g_{k+1}^{-1} g_{k+1}^* S_k \prod_{i \in I_{k+1}} e_i^{j_i}$$

is a minimal zero-sum sequence over  $C_n \oplus \bigoplus_{j=1}^{k+1} (\bigoplus_{i \in I_j} \langle e_i \rangle)$  and  $\mathbf{k}(S_{k+1}) = \mathbf{k}(S_k) - \frac{1}{\text{ord}(g_{k+1})} + \frac{1}{\text{ord}(g_{k+1}^*)} + \sum_{i \in I_{k+1}} \frac{j_i}{q_i} = w_c + \sum_{j=1}^{k+1} \sum_{i \in I_j} \frac{j_i}{q_i}$ . Moreover  $S_{k+1}$  contains an element whose order has  $p_i$ -valuation equal to  $\mathbf{v}_{p_i}(n)$  for each  $i \in [1, s]$ .

The proof is complete by observing that  $S_s$  has the required property.  $\square$

**Theorem 4.4.** *Let  $G = H \oplus C_n$  be a finite abelian group with  $\exp(G) = n$ .*

1. *If  $n$  is odd, then*

$$\frac{1}{n} [2, n + nk^*(H)] \subseteq \mathbb{W}(G).$$

2. If  $n$  is even, then

$$\frac{2}{n} \left[ 1, \frac{1}{2}(n + n(k^*(H) - \delta)) \right] \subseteq W(G)$$

where  $\delta = 0$  if  $nk^*(H)$  is even and  $\delta = 2^{-v_2(\exp(H))}$  otherwise.

We note that in case  $v_2(\exp(H)) < v_2(n)$ , we have  $W(G) \subseteq \frac{2}{n} [1, \frac{n}{2} K(G)]$  as recalled at the end of Section 2 and moreover  $nk^*(H)$  is even. Thus the second part of the result is quite tight, too, in this case. However, in case  $v_2(\exp(H)) = v_2(n)$ , there is a considerable discrepancy. In Proposition 5.4 we address this in a special case.

*Proof.* We want to apply Proposition 4.2. Let  $q_1, \dots, q_t$  be prime powers such that  $H \cong \bigoplus_{i=1}^t C_{q_i}$ , moreover we assume that  $q_1 \leq \dots \leq q_s$  are powers of 2 while the other  $q_i$  are odd. Let us denote  $H = H_2 \oplus H_o$  where  $|H_o|$  is odd and  $|H_2|$  is a power of 2, then  $H_2 \cong \bigoplus_{i=1}^s C_{q_i}$ .

Now,  $G \cong \bigoplus_{i=1}^t C_{q_i} \oplus C_n$  and each  $q_i$  divides  $n$ , thus Proposition 4.2 yields

$$\sum_{i=1}^t \frac{1}{q_i} [0, q_i - 1] + W(C_n) \subseteq W(G).$$

As observed at the end of Section 2 we know that if  $n$  is odd, then  $\frac{1}{n} [2, n] \subseteq W(C_n)$  and if  $n$  is even, then  $\frac{2}{n} [1, \frac{n}{2}] \subseteq W(C_n)$ . For  $n$  odd, using Lemma 3.7 as in the proof of Proposition 3.9, we get  $\sum_{i=1}^t \frac{1}{q_i} [0, q_i - 1] + \frac{1}{n} [2, n] = \frac{1}{n} [2, n + n \sum_{i=1}^t \frac{q_i - 1}{q_i}]$ , which is precisely  $\frac{1}{n} [2, n + nk^*(H)]$ .

Likewise, if  $n$  is even and  $q_i$  is odd, then  $q_i \mid \frac{n}{2}$ , and we get again by Lemma 3.7 that  $\sum_{i=s+1}^t \frac{1}{q_i} [0, q_i - 1] + \frac{2}{n} [1, \frac{n}{2}] = \frac{2}{n} [1, \frac{1}{2}(n + n \sum_{i=s+1}^t \frac{q_i - 1}{q_i})]$ , which is precisely  $\frac{2}{n} [1, \frac{1}{2}(n + nk^*(H_o))]$ .

It remains to treat the contribution of the even  $q_i$ . Using Lemma 3.7, we get  $\sum_{i=1}^s \frac{1}{q_i} [0, q_i - 1] = \frac{1}{q_s} [0, q_s k^*(H_2)]$ . Now, if  $q_s \mid \frac{n}{2}$ , we have again using Lemma 3.7 that  $\frac{1}{q_s} [0, q_s k^*(H_2)] + \frac{2}{n} [1, \frac{1}{2}(n + nk^*(H_o))] = \frac{2}{n} [1, \frac{1}{2}(n + nk^*(H))]$ . If  $q_s \nmid \frac{n}{2}$ , then we note that

$$\frac{2}{q_s} \left[ 0, \left\lfloor \frac{q_s}{2} k^*(H_2) \right\rfloor \right] \subseteq \frac{1}{q_s} [0, q_s k^*(H_2)],$$

and thus  $\frac{2}{q_s} [0, \lfloor \frac{q_s}{2} k^*(H_2) \rfloor] + \frac{2}{n} [1, \frac{1}{2}(n + nk^*(H_o))] \subseteq W(G)$ . Since  $\frac{q_s}{2} \mid \frac{n}{2}$ , we can apply Lemma 3.7 to get that

$$\frac{2}{q_s} \left[ 0, \left\lfloor \frac{q_s}{2} k^*(H_2) \right\rfloor \right] + \frac{2}{n} \left[ 1, \frac{1}{2}(n + nk^*(H_o)) \right]$$

equals

$$\frac{2}{n} \left[ 1, \frac{1}{2}(n + nk^*(H_o)) + \frac{n}{q_s} \left\lfloor \frac{q_s}{2} k^*(H_2) \right\rfloor \right].$$

Now, if  $q_s k^*(H_2)$  is even, this is again  $\frac{2}{n} [1, \frac{1}{2}(n + nk^*(H))]$ , while if  $q_s k^*(H_2)$  is odd, it equals  $\frac{2}{n} [1, \frac{1}{2}(n + n(k^*(H) - \frac{1}{q_s}))]$ . To conclude the proof it remains to observe that the parity of  $nk^*(H)$  and  $q_s k^*(H_2)$  coincide.  $\square$

## 5. WHEN ARE $w(G)$ AND $W(G)$ ARITHMETIC PROGRESSIONS?

In earlier sections we showed that for a wide variety of groups the initial parts of the sets  $w(G)$  and  $W(G)$  are arithmetic progressions with difference  $\frac{1}{\exp(G)}$  or sometimes  $\frac{2}{\exp(G)}$ . In the current section we study when the full sets actually are arithmetic progressions. We first recall that this is the case for  $p$ -groups, more precisely for  $G = C_{n_1} \oplus \dots \oplus C_{n_r}$  a finite abelian  $p$ -group with  $1 = n_0 < n_1 \mid \dots \mid n_r$  we have that  $W(G)$  and  $w(G)$  are arithmetic progressions with difference  $\frac{1}{\exp(G)}$ , unless  $p = 2$  and  $n_{r-1} < n_r$  in which case  $W(G)$  is an arithmetic progression with difference  $\frac{2}{\exp(G)}$ , see Theorem 4.1.

We now consider groups that are the direct sum of an elementary 2-group and a  $p$ -group and establish in some cases that sets of cross numbers are arithmetic progressions, too. While this is a quite special

class of group, there are reasons to assume that the phenomenon is quite rare; we discuss this at the end of this paper.

We start with a result for cyclic groups; the assertion on  $W(G)$  is known by a result of Baginski et al. [1, Theorem 2.1].

**Theorem 5.1.** *Let  $G = C_{2p^k}$  with  $k \in \mathbb{N}$  and  $p$  be a prime. Then*

$$W(G) = \frac{2}{\exp(G)}[1, \frac{\exp(G)}{2}K^*(G)] \quad \text{and} \quad w(G) = \frac{1}{\exp(G)}[1, \exp(G)k^*(G)].$$

*Proof.* The statement on  $W(G)$  was proved by Baginski et al. in [1, Theorem 2.1]. We need to prove the result for  $w(G)$ . Based on the outcome concerning  $W(G)$ , note that we have  $k(G) = k^*(G) = \frac{3p^k-2}{2p^k}$ . Now for some  $g \in G$  with  $\text{ord}(g) = 2p^k$ , take  $S = g^i \in \mathcal{A}^*(G)$  for  $i \in [1, 2p^k - 1]$  whence

$$\frac{1}{2p^k}[1, 2p^k - 1] \subseteq w(G) \subseteq \frac{1}{2p^k}[1, 3p^k - 2]$$

It remains to obtain the missing elements as cross numbers of a zero-sum free sequence. Write  $G = C_2 \oplus C_{p^k}$  and suppose  $(e, f)$  is a generating set of  $G$  with  $\text{ord}(e) = 2$  and  $\text{ord}(f) = p^k$ . For  $1 \leq l \leq \frac{p^k-1}{2}$ , set

$$B_l = e(e+f)f^{p^k-1-l} \in \mathcal{A}^*(G) \quad \text{and} \quad B'_l = B_l(e+f)^{-1} \in \mathcal{A}^*(G)$$

then

$$k(B_l) = \frac{3p^k - 1 - 2l}{2p^k} \quad \text{and} \quad k(B'_l) = k(B_l) - \frac{1}{2p^k}.$$

Altogether,  $B_l$  and  $B'_l$  give us  $[2p^k, 3p^k - 3] \subseteq 2p^k w(G)$ . Now if we put  $l = 0$  in  $B_l$  then  $B_0 \in \mathcal{A}(G)$  but  $B'_0 = B_0(e+f)^{-1} \in \mathcal{A}^*(G)$  and  $k(B'_0) = \frac{3p^k-2}{2p^k}$  and the proof is complete.  $\square$

Combining this result with results from the preceding section we obtain more general results. We start by a result for  $w(G)$ , which are again easier to obtain.

**Proposition 5.2.** *Let  $G_p$  be a  $p$ -group for an odd prime  $p$  and let  $G = C_2^r \oplus G_p$  for some  $r \in \mathbb{N}$ . Then*

$$\frac{1}{\exp(G)}[1, \exp(G)k^*(G)] \subseteq w(G) \subseteq \frac{1}{\exp(G)}[1, \exp(G)k(G)]$$

*In particular, if  $k(G) = k^*(G)$ , then equality holds and  $w(G)$  is an arithmetic progression.*

*Proof.* The inclusion  $w(G) \subseteq \frac{1}{\exp(G)}[1, \exp(G)k(G)]$  is obvious. We show  $\frac{1}{\exp(G)}[1, \exp(G)k^*(G)] \subseteq w(G)$ . Let  $G = \sum_{i=1}^t C_{q_i} \oplus C_n$  where  $n = \exp(G)$  and the  $q_i$  are prime-powers, that is in the current case either a  $p$ -power or 2. By Proposition 3.8 we know that  $\sum_{i=1}^t \frac{1}{q_i}[0, q_i - 1] + w(C_n) \subseteq w(G)$ . By Theorem 5.1, we know that  $w(C_n) = \frac{1}{n}[1, nk^*(C_n)]$ . Using Lemma 3.7, in the same way as in the proof of Proposition 3.9, we see that  $\sum_{i=1}^t \frac{1}{q_i}[0, q_i - 1] + \frac{1}{n}[1, nk^*(C_n)] = \frac{1}{n}[1, nk^*(G)]$ , and the claim is established.  $\square$

We now want a similar result for  $W(G)$ . We need a technical lemma.

**Lemma 5.3.** *Let  $G$  be a cyclic  $p$ -group with  $G = \langle f \rangle$  where  $\text{ord}(f) = p^k$  for some odd prime  $p$  and some  $k \in \mathbb{N}$ .*

1. *For each  $j \in [1, p^k - 1]$ , there exists some  $S_j \in \mathcal{A}^*(G)$  such that  $\sigma(S_j) = -f$  and  $k(S_j) = \frac{j}{p^k}$ .*
2. *For each  $j \in [1, p^k - 2]$ , there exists some  $T_j \in \mathcal{A}^*(G)$  such that  $\sigma(T_j) = -2f$ ,  $-f \notin \Sigma(T_j)$  and  $k(T_j) = \frac{j}{p^k}$ .*

*Proof.* 1. Take  $S_j = f^{j-1}((p^k - j)f)$  for some  $j \in [1, p^k - 1]$ . Then  $S_j$  is a zero-sum free sequence with sum  $-f$ . We have  $k(S_j) = \frac{j}{p^k}$  unless  $p^k - j$  is divisible by  $p$ . Suppose  $p^k - j$  is a multiple of  $p$ , that means  $j$  is a multiple of  $p$ , but then  $j - 1$  is not a multiple of  $p$ . Therefore,  $S'_j = f^{j-2}(2f)((p^k - j - 1)f)$  is a zero-sum free sequence that fulfills the requirements.

2. Take  $T_j = f^{j-1}((p^k - j - 1)f)$  for some  $j \in [1, p^k - 2]$ , then  $T_j$  is zero-sum free with sum  $-2f$  and  $-f \notin \Sigma(T_j)$ . The cross number is  $\frac{j}{p^k}$  unless  $p^k - j - 1$  is divisible by  $p$ . If  $p^k - j - 1$  is a multiple of  $p$ , then  $p^k - j - 2$  is not and therefore  $T'_j = f^{j-2}(2f)((p^k - j - 2)f)$  has the required properties.  $\square$

**Proposition 5.4.** *Let  $G_p$  be a  $p$ -group for an odd prime  $p$  and let  $G = C_2^r \oplus G_p$  for some  $r \in \mathbb{N}$ .*

1. *If  $r = 1$ , then*

$$\frac{2}{\exp(G)} \left[ 1, \frac{\exp(G)}{2} \mathbf{K}^*(G) \right] \subseteq \mathbf{W}(G) \subseteq \frac{2}{\exp(G)} \left[ 1, \frac{\exp(G)}{2} \mathbf{K}(G) \right]$$

*In particular, if  $\mathbf{K}(G) = \mathbf{K}^*(G)$  then equality holds and  $\mathbf{W}(G)$  is an arithmetic progression with difference  $\frac{2}{\exp(G)}$ .*

2. *If  $r \geq 2$ , then*

$$\frac{1}{\exp(G)} \left[ 2, \exp(G) \mathbf{K}^*(G) \right] \subseteq \mathbf{W}(G) \subseteq \frac{1}{\exp(G)} \left[ 2, \exp(G) \mathbf{K}(G) \right]$$

*In particular, if  $\mathbf{K}(G) = \mathbf{K}^*(G)$  then equality holds and thus  $\mathbf{W}(G)$  is an arithmetic progression with difference  $\frac{1}{\exp(G)}$ .*

*Proof.* The inclusions for  $\mathbf{W}(G)$  are the well-known ones, see the end of Section 2. It remains to show that  $\mathbf{W}(G)$  contains the claimed elements. Assume  $G_p = C_{p^{k_1}} \oplus \cdots \oplus C_{p^{k_l}}$  for some  $k_i \in \mathbb{N}$ ,  $i \in [1, l]$  and  $k_1 \leq \cdots \leq k_l$ . By Proposition 4.2, it suffices to show the results for  $G = C_2^r \oplus C_{p^{k_l}}$  only. Therefore, from now on assume  $G = C_2^r \oplus C_{p^k}$  for some  $k \in \mathbb{N}$ .

The first assertion, then is precisely Theorem 5.1. We turn to the second assertion. We show the result for  $r = 2$  only, again the rest follows from Proposition 4.2. Write

$$G = C_2 \oplus C_2 \oplus C_{p^k} = \langle e_1 \rangle \oplus \langle e_2 \rangle \oplus \langle f \rangle$$

where  $\text{ord}(f) = p^k$  and  $\text{ord}(e_i) = 2$  for  $i \in [1, 2]$ .

Then  $\mathbf{K}^*(G) = 2\frac{1}{2} + \frac{p^k-1}{p^k} + \frac{1}{2p^k} = 2 - \frac{1}{2p^k}$ . Now,  $\frac{1}{\exp(G)} [2, \exp(G)] \subseteq \mathbf{W}(G)$  by [4, Theorem 2]. For  $j \in [1, p^k - 1]$ , let  $A_j = e_1 e_2 (e_1 + e_2 + f) S_j \in \mathcal{A}(G)$  where  $S_j \in \mathcal{A}^*(G)$  such that  $\sigma(S_j) = -f$  and  $k(S_j) = \frac{j}{p^k}$ , which exists by Lemma 5.3. We have  $k(A_j) = 1 + \frac{1+2j}{2p^k}$  for  $j \in [1, p^k - 1]$ .

For  $j \in [1, p^k - 2]$ , let  $A'_j = e_1 e_2 (e_1 + f)(e_2 + f) T_j \in \mathcal{A}(G)$  where  $T_j \in \mathcal{A}^*(G)$  such that  $\sigma(T_j) = -2f$ ,  $-f \notin \Sigma(T_j)$ , and  $k(T_j) = \frac{j}{p^k}$ , which exists by Lemma 5.3. We have  $k(A'_j) = 1 + \frac{2+2j}{2p^k}$  for  $j \in [1, p^k - 2]$ . Whence  $1 + \frac{1}{2p^k}$  and  $1 + \frac{2}{2p^k}$  are the only cross numbers which are not yet realized by some minimal zero-sum sequence in  $G$  but  $S = e_1 e_2 (e_1 + f)(e_2 - f) \in \mathcal{A}(G)$  with  $k(S) = 1 + \frac{1}{2p^k} + \frac{1}{2p^k}$  and  $S = (e_1 + e_2 - f)(e_1 + f)(e_2 + f) f^{p^k-1} \in \mathcal{A}(G)$  with  $k(S) = 1 + \frac{1}{2p^k}$ .

Thus we infer that  $\frac{1}{2p^k} [2, 2p^k \mathbf{K}^*(G)] \subseteq \mathbf{W}(G)$ . The equality, under the assumption of  $\mathbf{K}^*(G) = \mathbf{K}(G)$ , is now obvious.  $\square$

In order to have unconditional results of the above mentioned type we study  $\mathbf{K}(G)$  for these types of groups, which also yields the result for  $k(G)$ . We stress that for  $k(G)$  more general results are obtained in [22, Theorem 7], however we could not see how to obtain the result for  $\mathbf{K}(G)$  from those results and thus present proofs even if they are quite similar.

**Proposition 5.5.** *Let  $H$  be a finite abelian group of odd order. If  $\mathbf{K}(H) = \mathbf{K}^*(H)$  and  $\sum_{d|\exp(H)} \frac{1}{d} \leq 2$ , then  $\mathbf{K}(C_2 \oplus H) = \mathbf{K}^*(C_2 \oplus H)$ .*

*Proof.* Let  $C_2 = \langle e \rangle$  with  $\text{ord}(e) = 2$  and let  $H$  be a finite abelian group of odd order with  $\sum_{d|\exp(H)} \frac{1}{d} < 2$  and  $\mathsf{K}^*(H) = \mathsf{K}(H)$ , which implies in particular that  $\mathsf{k}^*(H) = \mathsf{k}(H)$ . Let  $S \in \mathcal{A}(C_2 \oplus H)$  and assume for a contradiction that  $\mathsf{k}(S) > \mathsf{K}^*(C_2 \oplus H)$ . The condition on the cross number gives

$$\mathsf{k}(S) \geq \mathsf{K}^*(C_2 \oplus H) + \frac{1}{2\exp(H)} = \mathsf{k}(H) + \frac{1}{2} + \frac{1}{\exp(H)}.$$

We distinguish two cases.

**Case I:** There is an element, say  $e$ , in  $\text{supp}(S)$  with order 2.

In this case we have  $\mathsf{k}(Se^{-1}) \geq \frac{1}{\exp(H)} + \mathsf{k}(H) = \mathsf{K}(H)$ . Consider  $\pi : C_2 \oplus H \rightarrow (C_2 \oplus H)/\langle e \rangle \cong H$ . We note that  $\pi(Se^{-1})$  is a zero-sum sequence. Additionally, since the sum of  $Se^{-1}$  is  $e$ , it necessarily contains an element of even order.

This implies that

$$\mathsf{k}(\pi(Se^{-1})) > \mathsf{k}(Se^{-1}) = \mathsf{K}(H),$$

which means  $\pi(Se^{-1})$  is not a minimal zero-sum sequence. Yet this yields a contradiction to  $S$  being a minimal zero-sum sequence, as  $Se^{-1}$  has a proper subsequence with sum  $e$  or 0, which together with  $e$  yields a proper zero-sum sequence of  $S$ .

**Case II:** There is no element of order 2 in  $\text{supp}(S)$ . Let us write  $S = S_H S_{e+H}$  where  $S_H \in \mathcal{F}(H)$  and  $S_{e+H} \in \mathcal{F}(e+H)$ . Now for some  $g_1 g_2 \mid S_{e+H}$  with  $\text{ord}(g_1) = \text{ord}(g_2)$ , consider  $g_1 + g_2$ . Then  $\text{ord}(g_1 + g_2) \mid \frac{\text{ord}(g_1)}{2}$ , in particular  $\frac{1}{\text{ord}(g_1)} + \frac{1}{\text{ord}(g_2)} \leq \frac{1}{\text{ord}(g_1 + g_2)}$ . Thus without loss, we can assume that  $S_{e+H}$  contains at most 1 element of each order as if there are two elements of the same order in  $S_{e+H}$ , we can replace the two by their sum, which maintains the property that the sequence is a minimal zero-sum sequence and does not decrease the cross number. Thus we assume  $S = S_H R$  where  $R$  is a sequence in  $e+H$  which contains at most one element of each order. Then

$$\mathsf{k}(R) \leq \sum_{1 \neq d|\exp(H)} \frac{1}{2d} = \frac{1}{2} \sum_{1 \neq d|\exp(H)} \frac{1}{d}$$

and

$$\begin{aligned} \mathsf{k}(S_H) &= \mathsf{k}(S) - \mathsf{k}(R) \\ &\geq \mathsf{k}(H) + \frac{1}{\exp(H)} + \frac{1}{2} - \mathsf{k}(R). \end{aligned}$$

Now  $\sum_{1 \neq d|\exp(H)} \frac{1}{d} \leq 1$  implies that

$$\mathsf{k}(S_H) \geq \frac{1}{\exp(H)} + \mathsf{k}(H)$$

which implies that  $S_H$  is not a zero-sum free sequence in  $H$ . The only way how this does not contradict the fact that  $S$  is a minimal zero-sum sequence is that  $S = S_H$  that is  $R$  is trivial. Yet, in this case  $\mathsf{k}(S_H) \geq \mathsf{k}(H) + \frac{1}{\exp(H)} + \frac{1}{2} > \mathsf{k}(H) + \frac{1}{\exp(H)}$ , and thus  $S$  is not a minimal zero-sum sequence in  $C_2 \oplus H$ , which again shows that it is not a minimal zero-sum sequence.  $\square$

For any  $p$ -group  $H = G_p$ , Proposition 5.5 implies that  $\mathsf{K}^*(C_2 \oplus G_p) = \mathsf{K}(C_2 \oplus G_p)$ .

**Theorem 5.6.** *Let  $G = C_2^2 \oplus G_p$  where  $G_p$  is a  $p$ -group for some odd prime  $p$ . Then  $\mathsf{K}^*(G) = \mathsf{K}(G)$ .*

*Proof.* Assume  $\exp(G_p) = p^k$  for some  $k \in \mathbb{N}$  so  $\exp(G) = 2p^k$  and  $\mathsf{K}^*(G) = 1 + \mathsf{k}^*(G_p) + \frac{1}{2p^k}$ . We know  $\mathsf{K}^*(G) \leq \mathsf{K}(G)$ . Assume  $S \in \mathcal{A}(G)$  with  $\mathsf{k}(S) > \mathsf{K}^*(G)$ . We must show that no such  $S$  exists. We have  $\mathsf{k}(S) \geq \mathsf{K}^*(G) + \frac{1}{2p^k} = \mathsf{k}(G_p) + \frac{1}{p^k} + 1$ . We distinguish three cases.

**Case I:**  $\text{supp}(S)$  contains two elements of order 2, say  $e_1$  and  $e_2$ . Let  $C_2^2 = \langle e_1, e_2 \rangle$ . Then

$$\mathsf{k}(S(e_1 e_2)^{-1}) \geq \mathsf{k}^*(G_p) + \frac{1}{p^k} = \mathsf{K}^*(G_p) = \mathsf{K}(G_p)$$

Now let  $\pi : G \rightarrow G/\langle e_1, e_2 \rangle \cong G_p$  be the canonical epimorphism. Then  $\pi(S(e_1e_2)^{-1}) \in \mathcal{A}(G/\langle e_1, e_2 \rangle)$  but

$$k(\pi(S(e_1e_2)^{-1})) > k(S(e_1e_2)^{-1}) \geq K(G_p)$$

because there is at least one element in  $S(e_1e_2)^{-1}$  of even order, therefore  $\pi(S(e_1e_2)^{-1}) > K(G_p)$  contradicting our assumption.

**Case II:**  $\text{supp}(S)$  contains exactly one element of order 2, say  $e$ . Let  $\pi_1 : G \rightarrow G/\langle e \rangle \cong G_p \oplus C_2$  be the canonical epimorphism. Then  $\pi_1(Se^{-1}) \in \mathcal{A}(G/\langle e \rangle)$  but

$$\begin{aligned} k(\pi_1(Se^{-1})) &> k(Se^{-1}) \\ &> K^*(G) - \frac{1}{2} \\ &= K^*(G/\langle e \rangle) \\ &= K(G/\langle e \rangle) \end{aligned}$$

the first inequality is strict because there is at least one element in  $Se^{-1}$  of even order and the last equality is due to Proposition 5.5, a contradiction.

**Case III:**  $\text{supp}(S)$  does not contain an element of order 2.

Now set  $S = \prod_{i=1}^k S_{p^i} \prod_{j=1}^k S_{2p^j}$  where  $S_{p^i}$  consists of elements of order  $p^i$  for  $i \in [1, k]$  and  $S_{2p^j}$  consists of elements of order  $2p^j$  for  $j \in [1, k]$ . Note that without loss we can assume  $|S_{2p^j}| \leq 3$  for all  $j \in [1, k]$  as we can replace two elements of even order whose sum is not of even order by their sum (compare this to the argument in Case II of the proof of Proposition 5.5). Then

$$\begin{aligned} k\left(\prod_{i=1}^k S_{p^i}\right) &= k(S) - k\left(\prod_{j=1}^k S_{2p^j}\right) \\ &\geq k(S) - \frac{3}{2} \sum_{j=1}^k p^{-j} \\ &> k(S) - \frac{3}{2} \frac{1}{p-1} \\ &> k(S) - 1 \\ &\geq k(G_p) + \frac{1}{p^k} \\ &= K(G_p) \end{aligned}$$

a contradiction to  $S \in \mathcal{A}(G)$  and hence the assertion follows. □

**Corollary 5.7.** *Let  $G_p$  be a  $p$ -group for an odd prime  $p$  and let  $G = C_2^r \oplus G_p$  for some  $r \in \mathbb{N}$ .*

1. *If  $r = 1$ , then*

$$W(G) = \frac{2}{\exp(G)} \left[ 1, \frac{\exp(G)}{2} K^*(G) \right].$$

2. *If  $r = 2$ , then*

$$W(G) = \frac{1}{\exp(G)} \left[ 2, \exp(G) K^*(G) \right].$$

*Proof.* This follows directly from Propositions 5.4, 5.5 and Theorem 5.6. □

The second point yields a class of groups, other than  $p$ -groups of odd order for which  $W(G)$  is an arithmetic progressions with difference  $\frac{1}{\exp(G)}$ . This problem was raised in [4, Remark 5].

**Corollary 5.8.** *Let  $G_p$  be a  $p$ -group for an odd prime  $p$  and let  $G = C_2^r \oplus G_p$  for some  $r \in \mathbb{N}$ . If  $r \leq 2$ , then*

$$w(G) = \frac{1}{\exp(G)} [1, \exp(G)k^*(G)].$$

*Proof.* This follows directly from Propositions 5.2, 5.5 and Theorem 5.6.  $\square$

We end this section with a reflection of the problem to which extent the sets  $W(G)$  and  $w(G)$  determine the structure of  $G$ . Note that if  $|G|, |G'| \in \{1, 2\}$ , then  $W(G) = W(G') = \{1\}$ .

**Proposition 5.9.** *Let  $G$  and  $G'$  be finite abelian groups.*

1. *Let  $|G|, |G'| \geq 3$ . If  $W(G) = W(G')$  then  $\exp(G) = \exp(G')$  and  $K(G) = K(G')$ .*
2. *Let  $G$  be a  $p$ -group for some odd prime  $p$ . We have  $W(G) = W(G')$  if and only if  $\exp(G) = \exp(G')$  and  $K(G) = K(G')$ .*
3. *Let  $G$  be a  $p$ -group for some prime  $p$ . We have  $w(G) = w(G')$  if and only if  $\exp(G) = \exp(G')$  and  $k(G) = k(G')$ .*

*Proof.* 1. Let  $W(G) = W(G')$ . Since  $\min W(G) = \frac{2}{\exp(G)}$  and  $\min W(G') = \frac{2}{\exp(G')}$ , we get from  $W(G) = W(G')$  that  $\exp(G) = \exp(G')$ . Similarly, since  $\max W(G) = K(G)$  and  $\max W(G') = K(G')$ , we get  $K(G) = K(G')$ .

2. Assume first that  $\exp(G) = p^k$  for some  $k \in \mathbb{N}$ . We only need to show the reverse implication of 1. and then the assertion is complete. If  $\exp(G) = \exp(G')$ , then  $\exp(G') = p^k$  and  $G'$  is also a  $p$ -group. Now let  $K(G) = K(G')$ . Then by Theorem 4.1, we have  $W(G) = \frac{1}{\exp(G)} [2, \exp(G)K(G)] = \frac{1}{\exp(G')} [2, \exp(G')K(G')] = W(G')$ .

3. Suppose  $w(G) = w(G')$ . Then  $\frac{1}{\exp(G)} = \min w(G) = \min w(G') = \frac{1}{\exp(G')}$  whence  $\exp(G) = \exp(G')$ . Moreover, we also get  $k(G) = \max w(G) = \max w(G') = k(G')$ . Conversely, suppose  $\exp(G) = p^k$  for some  $k \geq 1$  and let  $p^k = \exp(G) = \exp(G')$ . Then  $G'$  is also a  $p$  group having the same exponent as  $G$ . Now if  $k(G) = k(G')$  too, then Corollary 3.6 implies that  $w(G) = w(G')$ .  $\square$

The following remark highlights some examples that complement the preceding result.

**Remark 5.10.**

1. Let  $G = C_{2^{k_1}} \oplus \dots \oplus C_{2^{k_r}}$  with  $1 \leq k_1 \leq \dots \leq k_{r-1} = k_r$  and  $G' = C_{2^{l_1}} \oplus \dots \oplus C_{2^{l_s}}$  with  $1 \leq l_1 \leq \dots \leq l_{s-1} < l_s$ . Assume that  $k_r = l_s$ , that is,  $\exp(G) = \exp(G')$ . For suitable choices of  $r, s$  and the  $k_i, l_i$  it is possible to have  $K(G) = K(G')$ , too. Indeed, for instance take  $G = C_4^3$  and  $G' = C_2^3 \oplus C_4$  then  $\exp(G) = \exp(G') = 4$  and  $K(G) = K(G') = \frac{10}{4}$ . But in any case, Theorem 4.1 tells that  $W(G) \neq W(G')$ . Thus the Proposition 5.9.2 only works for odd primes.
2. For some  $p$ -groups  $G, G'$ , the group rank can still differ even if  $\exp(G) = \exp(G')$ ,  $K(G) = K(G')$  and  $W(G) = W(G')$ . For instance, if  $G = C_3^4 \oplus C_9$  and  $G' = C_9^4$  then  $W(G) = W(G')$  by Theorem 4.1 but  $G$  and  $G'$  have different rank. Therefore,  $\exp(G) = \exp(G')$ ,  $K(G) = K(G')$  and  $W(G) = W(G')$  does not determine the structure of the group.

## 6. GAP STRUCTURE IN $W(C_p^r \oplus C_q^s)$

The goal of this section is to highlight the fact that there are gaps in the set of cross numbers for the group  $G = C_p^r \oplus C_q^s$  with  $p > q$  odd primes and  $r, s \in \mathbb{N}$  at least if  $p$  is large relative to  $q$ . Note that  $r = s = 1$  is studied in detail in [1].

Since it is needed in this section we recall that for a finite abelian group  $G$ , the constant  $\eta(G)$  is defined to be the smallest integer  $t$  such that every sequence  $S$  over  $G$  with  $|S| \geq t$  has a zero-sum subsequence  $T$  with length  $|T| \in [1, \exp(G)]$ , such a subsequence is called short. It is easy to see that this constant is finite, for example  $\exp(G)|G|$  is a trivial upper bound. In general, the exact value is unknown. By definition, we have  $D(G) \leq \eta(G)$ . The exact value of  $\eta(G)$  is known for groups of rank at most two and it is known that  $\eta(G) \leq |G|$  always holds, we refer to [10, Chapter 5] for these and other results.



The main results are the two corollaries at the end. We start with a few technical results. The following lemma is useful to establish a link between  $w(G)$  and  $W(G)$ .

**Lemma 6.1.** *Let  $G = C_p^r \oplus C_q^s$  for  $r, s \in \mathbb{N}$  and  $p > q$  be primes. Let  $S \in \mathcal{A}(G)$  such that*

$$k(S) > \max\left\{\frac{1+r(p-1)}{p}, \frac{1+s(q-1)}{q}\right\}.$$

*Then  $S$  contains an element of order  $pq$ .*

*Proof.* Assume to the contrary that  $S$  does not contain an element of order  $pq$ . Then  $S$  contains either only elements of order  $p$  or element of order  $q$ . This contradicts the assumption on the cross number.  $\square$

The following result allows to focus on  $w(G)$  when trying to show the existence of gaps.

**Proposition 6.2.** *Let  $G = C_p^r \oplus C_q^s$  for  $r, s \in \mathbb{N}$  and  $p > q$  be primes. Then  $W(G) \subseteq \frac{1}{pq} + w(G)$ .*

*Proof.* Let  $S \in \mathcal{A}(G)$ . We need to show that  $k(S) \in \frac{1}{pq} + w(G)$ . If  $k(S) > \max\left\{\frac{1+r(p-1)}{p}, \frac{1+s(q-1)}{q}\right\}$ , then by Lemma 6.1  $S$  contains an element  $g$  of order  $pq$  and thus  $g^{-1}S$  is zero-sum free with cross number  $k(S) - \frac{1}{pq}$ , which proves the claim. If  $k(S) \leq \max\left\{\frac{1+r(p-1)}{p}, \frac{1+s(q-1)}{q}\right\}$ , then  $k(S)$  is in  $\frac{1}{pq} + w(G)$ , simply as by Theorem 3.5 the set contains every integral multiple of  $\frac{1}{pq}$  from  $\frac{2}{pq}$  up to  $\frac{pq+(r-1)(p-1)+(s-1)(q-1)}{pq}$ .  $\square$

The following result gives some insight into the structure of zero-sum free sequences of large cross number for groups of exponent  $pq$  for two odd primes  $p, q$  under the assumption that  $p$  is larger than  $q$ ; this assumption is implicit in the condition on  $t$ , if  $p$  is not large enough no such  $t$  exists and the result is void. In other words the result is only relevant for  $p > \eta(C_q^s)$ .

**Proposition 6.3.** *Let  $G = C_p^r \oplus C_q^s$  for  $r, s \in \mathbb{N}$  and  $p > q$  primes. Let  $S \in \mathcal{A}^*(G)$  such that*

$$k(S) \geq k^*(G) - \frac{t}{pq}$$

*where  $t \in \mathbb{N}$  such that  $t \leq p - \eta(C_q^s)$ . Then  $S$  contains  $s(q-1)$  elements of order  $q$ .*

We first establish a lemma for sequences that do not contain an element of order at most  $q$ .

**Lemma 6.4.** *Let  $G = C_p^r \oplus C_q^s$  for  $r, s \in \mathbb{N}$  and  $p > q$  be primes. Let  $S \in \mathcal{F}(G \setminus C_q^s)$  such that*

$$k(S) \geq \frac{r(p-1)}{p} + \frac{\eta(C_q^s)}{pq}.$$

*Then  $S$  is not zero-sum free.*

*Proof.* We write  $S = S_p S_{pq}$  where the elements in  $S_p$  and  $S_{pq}$  have order  $p$  and  $pq$ , respectively.

Let us write  $S_{pq} = Q_1 \dots Q_l R$  such that  $\sigma(Q_i) \in C_p^r$  and  $|Q_i| \leq q$  and  $l$  is maximal. In other words, the images of sequences  $Q_i$  are short zero-sum sequences over  $G/C_p^r \cong C_q^s$ . Thus, we have that  $|R| \leq \eta(C_q^s) - 1$ . Note that

$$pqk(S_p S_{pq}) = q|S_p| + |S_{pq}| \leq q|S_p| + ql + \eta(C_q^s) - 1$$

and  $pqk(S_p S_{pq}) \geq qr(p-1) + \eta(C_q^s)$  by our assumption that  $k(S) \geq \frac{r(p-1)}{p} + \frac{\eta(C_q^s)}{pq}$ . Therefore  $|S_p| + l > r(p-1)$ . Thus,  $S_p \sigma(Q_1) \dots \sigma(Q_l)$  is a sequence over  $C_p^r$  of length greater than  $r(p-1)$  and hence cannot be zero-sum free. It follows that  $S_p Q_1 \dots Q_l$  is not zero-sum free either, which shows the claim.  $\square$

*Proof of Proposition 6.3.* We write  $S = S_p S_q S_{pq}$  where the order of elements in  $S_j$  is  $j$  for each  $j \in \{p, q, pq\}$ . Since  $S$  is zero-sum free it is clear that  $|S_q|$  is at most  $s(q-1)$ . Assume  $|S_q| < s(q-1)$ . Then  $k(S_p S_{pq}) \geq \frac{r(p-1)}{p} + \frac{1}{q} - \frac{t}{pq}$ . Since  $p-t \geq \eta(C_q^s)$  by assumption, it follows that  $k(S_p S_{pq}) \geq \frac{r(p-1)}{p} + \frac{\eta(C_q^s)}{pq}$ . Now, by Lemma 6.4 the sequence  $S_p S_{pq}$  is not zero-sum free and thus  $S$  is not zero-sum free. This contradiction establishes the claim.  $\square$



We now determine the large values in the set of cross numbers.

**Corollary 6.5.** *Let  $G = C_p^r \oplus C_q^s$  for  $r, s \in \mathbb{N}$  and  $p > q$  odd primes such that  $p \geq \eta(C_q^s) + 2q$ . Then*

$$w(G) \subseteq \frac{1}{pq}[1, pqk^*(G)] \setminus \left\{ \frac{1}{pq}[pqk^*(G) - (2q - 3), pqk^*(G) - (q + 1)] \cup \frac{1}{pq}[pqk^*(G) - (q - 2), pqk^*(G) - 1] \right\}$$

while  $\frac{1}{pq}\{pqk^*(G) - (2q - 2), pqk^*(G) - q, pqk^*(G) - (q - 1), pqk^*(G)\} \subseteq w(G)$  and moreover  $k(G) = k^*(G)$ .

*Proof.* To prove the first inclusion and that  $k(G) = k^*(G)$ , let  $S \in \mathcal{A}^*(G)$  such that  $k(S) \geq k^*(G) - \frac{2q-1}{pq}$ , say  $k(S) = k^*(G) - \frac{t}{pq}$  with  $t \leq 2q - 1$ . We need to show that  $t$  is not in  $[1, q - 2] \cup [q + 1, 2q - 3]$  and that  $t \geq 0$ . Then, at the end of the proof we establish examples of sequences that yield the values  $\frac{1}{pq}\{pqk^*(G) - (2q - 2), pqk^*(G) - q, pqk^*(G) - (q - 1), pqk^*(G)\}$ , corresponding to  $t$  equal to  $2q - 2, q, q - 1$ , and 0.

We decompose  $S$  into subsequences according to the order of the elements, namely, let  $S_q, S_p$  and  $S_{pq}$  be the subsequences of  $S$  of elements of order  $q, p$  and  $pq$ , respectively.

First, we recall a result on  $S_q$  and then we investigate  $S_p S_{pq}$ . By Proposition 6.3 we know that  $|S_q| = s(q - 1)$ . Thus  $|S_q| = D(C_q^s) - 1$  and  $S_q$  is a zero-sum free sequence of maximal length over  $C_q^s$ . Therefore  $\Sigma(S_q) = C_q^s \setminus \{0\}$  (see [10, Proposition 5.1.4]). Assume  $\pi : G \rightarrow C_p^r$  to be the canonical epimorphism. It follows that  $\pi(S_p S_{pq})$  is zero-sum free in  $C_p^r$ . We get that  $|\pi(S_p S_{pq})| \leq r(p - 1)$ .

This fairly strong condition on  $|\pi(S_p S_{pq})|$  and thus on  $|S_p S_{pq}|$  has consequences for the values the cross number of  $S_p S_{pq}$  can attain. Since we know that  $k(S_q) = \frac{s(q-1)}{q}$ , this directly yields results on  $k(S)$ .

We analyze these restrictions in detail. We observe that  $k(S_p S_{pq}) = \frac{|S_p|}{p} + \frac{|S_{pq}|}{pq}$ . Moreover, we know that  $|S_p| + |S_{pq}| \leq r(p - 1)$ . In particular,  $k(S_p S_{pq})$  is at most  $\frac{|S_p| + |S_{pq}|}{p}$ , and from  $|S_p S_{pq}| \leq r(p - 1)$  it follows that  $k(S_p S_{pq}) \leq \frac{r(p-1)}{p}$ . Since  $k(S_q) = \frac{s(q-1)}{q}$ , we get that  $k(S) \leq \frac{s(q-1)}{q} + \frac{r(p-1)}{p} = k^*(G)$ . Thus there is no zero-sum free sequence of cross-number strictly greater than  $k^*(G)$  and thus  $k^*(G) = k(G)$ . By the same argument, if  $|S_p S_{pq}| \leq r(p - 1) - 2$ , then  $k(S) \leq k^*(G) - \frac{2q}{pq}$ , a contradiction to our assumption on  $k(S)$ .

Therefore,  $|S_p S_{pq}| \in \{r(p - 1) - 1, r(p - 1)\}$ . Let us observe possible values for  $k(S_p S_{pq})$ . For ease of notation we set  $M = \frac{r(p-1)}{p}$ , so that  $k^*(G) = \frac{s(q-1)}{q} + M$ . We include all the relevant values in the table. Note that for  $|S_{pq}|$  larger than the values in the table, the value of the cross number of  $k(S)$  is below bound we assumed initially.

$ S_p  +  S_{pq} $	$ S_{pq} $	$k(S_p S_{pq})$
$r(p - 1)$	0	$\frac{r(p-1)}{p} = M$
	1	$\frac{r(p-1)}{p} - \frac{1}{p} + \frac{1}{pq} = M - \frac{q-1}{pq}$
	2	$\frac{r(p-1)}{p} - \frac{2}{p} + \frac{2}{pq} = M - \frac{2(q-1)}{pq}$
$r(p - 1) - 1$	0	$\frac{r(p-1)}{p} - \frac{1}{p} = M - \frac{q}{pq}$
	1	$\frac{r(p-1)}{p} - \frac{2}{p} + \frac{1}{pq} = M - \frac{2q-1}{pq}$
	2	$\frac{r(p-1)}{p} - \frac{3}{p} + \frac{2}{pq} = M - \frac{3q-2}{pq}$

Thus it can be seen that  $\{k^*(G) - \frac{q-2}{pq}, \dots, k^*(G) - \frac{1}{pq}\} \cap w(G) = \emptyset$  and  $\{k^*(G) - \frac{2q-3}{pq}, \dots, k^*(G) - \frac{q+1}{pq}\} \cap w(G) = \emptyset$ , as those values do not appear in the table. This establishes our claim.

It remains to show that  $\frac{1}{pq}\{pqk^*(G) - (2q - 2), pqk^*(G) - q, pqk^*(G) - (q - 1), pqk^*(G)\} \subseteq w(G)$ . We note that there are choices of the parameters in the table that yield these values. It remains to show that zero-sum free sequences with the parameters as indicated in the table actually exist. We outline the argument below.

Of course a zero-sum free sequence  $S_q$  over  $C_q^s$  of length  $s(q-1)$  exists, and we also have zero-sum free sequence  $S'_p$  and  $S''_p$  over  $C_p^r$  of length  $r(p-1)$  and length  $r(p-1)-1$ , respectively. The sequences  $S_q S'_p$  and  $S_q S''_p$  are zero-sum free and yield the values in the table corresponding to  $|S_{pq}| = 0$ .

Moreover for  $h \in C_q^s \setminus \{0\}$  and  $g \mid S'_p$ , the sequences  $g^{-1}(g+h)S'_p$  and  $g^{-1}(g+h)S'_p S_q$  are still zero-sum free (the projection of the former to  $C_p^r$  remains unchanged and is thus zero-sum free) and contain exactly one element of order  $pq$ . Likewise for  $g_1 g_2 \mid S'_p$  the sequences  $g_1^{-1}(g_1+h)g_2^{-1}(g_2+h)S'_p$  and  $g_1^{-1}(g_1+h)g_2^{-1}(g_2+h)S'_p$  are still zero-sum free and contain exactly two elements of order  $pq$ . These yield sequences for the values corresponding to  $|S_p| + |S_{pq}| = r(p-1)$  and  $|S_{pq}|$  equal to 1 and 2. The same argument for  $S''_p$ , yields the values corresponding to  $|S_p| + |S_{pq}| = r(p-1) - 1$  and  $|S_{pq}|$  equal to 1 and 2.  $\square$

The proof does not really make use of the fact that  $q$  is odd. But the claim is void for  $q = 2$ . The situation is somewhat different for the proof below.

**Corollary 6.6.** *Let  $G = C_p^r \oplus C_q^s$  for  $r, s \in \mathbb{N}$  and  $p > q$  odd primes such that  $p \geq \eta(C_q^s) + 2q$ . Then*

$$W(G) \subseteq \frac{1}{pq}[2, pqK^*(G)] \setminus \left\{ \frac{1}{pq}[pqK^*(G) - (2q-3), pqK^*(G) - (q+1)] \cup \frac{1}{pq}[pqK^*(G) - (q-2), pqK^*(G) - 1] \right\}$$

while  $\frac{1}{pq}\{pqK^*(G) - (2q-2), pqK^*(G) - q, pqK^*(G) - (q-1), pqK^*(G)\} \subseteq W(G)$  and moreover  $K(G) = K^*(G)$ .

*Proof.* We know by Lemma 6.4 that  $W(G) \subseteq \frac{1}{pq} + w(G)$ . This proves the claim by invoking Corollary 6.5, except for the part regarding the existence of the values. However, to see this it suffices to note that we can construct the zero-sum free sequences in the proof of Corollary 6.5 in such a way that the sum has order  $pq$ . It is clear that the projection of the sum on  $C_p^r$  is non-zero. If the projection of the sum on  $C_q^s$  is 0, then we could just use a different element  $h$  for our construction; indeed every other element of order  $q$  would do in that case, and there is more than one, as  $q \neq 2$ .  $\square$

It would be possible to obtain more precise results on the sets of cross numbers in case  $p$  is much larger than  $q$ . Broadly speaking the gap structure in  $W(C_p^r \oplus C_q^s)$  follows the same pattern as that in the cyclic case discussed in [1].

We conclude by pointing out that the phenomenon we observed in this section suggest that sets of cross numbers are rarely arithmetic progressions. The point is that zero-sum free sequences whose cross number is maximal should contain only elements whose order is not the exponent. Starting from such a sequence, minor changes will usually not result in a sequence whose cross number is merely  $\frac{1}{\exp(G)}$  less than the original sequence.

#### ACKNOWLEDGMENTS

The authors are grateful to A. Geroldinger for encouragement and guidance during the work on this paper. In particular, they thank him for his contribution of Theorem 3.4. They thank the referee for a very careful reading and numerous important remarks.

#### REFERENCES

- [1] P. Baginski, S.T. Chapman, K. McDonald, and L. Pudwell, *On cross numbers of minimal zero sequences in certain cyclic groups*, *Ars Comb.* **70** (2004), 47 – 60.
- [2] T. Barron, C. O'Neill, and R. Pelayo, *On the set of elasticities in numerical monoids*, *Semigroup Forum* **94** (2017), 37 – 50.
- [3] N. Bushaw and G. Hurlbert, *Thresholds for zero-sums with small cross numbers in abelian groups*, *Integers* **24** (2024), Paper A93, 12pp.
- [4] S.T. Chapman and A. Geroldinger, *On cross numbers of minimal zero sequences*, *Australas. J. Comb.* **14** (1996), 85 – 92.

- [5] S. Elledge and G.H. Hurlbert, *An application of graph pebbling to zero-sum sequences in abelian groups*, Integers **5(1)** (2005), Paper A17, 10pp.
- [6] Y. Fan and A. Geroldinger, *Minimal relations and catenary degrees in Krull monoids*, J. Commut. Algebra **11** (2019), 29 – 47.
- [7] A. Geroldinger, *On a conjecture of Kleitman and Lemke*, J. Number Theory **44** (1993), 60 – 65.
- [8] ———, *The cross number of finite abelian groups*, J. Number Theory **48** (1994), 219 – 223.
- [9] A. Geroldinger and D.J. Grynkiewicz, *On the structure of minimal zero-sum sequences with maximal cross number*, J. Combinatorics and Number Theory **1 (2)** (2009), 9 – 26.
- [10] A. Geroldinger and F. Halter-Koch, *Non-Unique Factorizations. Algebraic, Combinatorial and Analytic Theory*, Pure and Applied Mathematics, vol. 278, Chapman & Hall/CRC, 2006.
- [11] A. Geroldinger and R. Schneider, *The cross number of finite abelian groups II*, European J. Combin. **15** (1994), 399 – 405.
- [12] ———, *The cross number of finite abelian groups III*, Discrete Math. **150** (1996), 123 – 130.
- [13] ———, *On minimal zero sequences with large cross number*, Ars Comb. **46** (1997), 297 – 303.
- [14] A. Geroldinger and Q. Zhong, *The set of minimal distances in Krull monoids*, Acta Arith. **173** (2016), 97 – 120.
- [15] ———, *Sets of arithmetical invariants in transfer Krull monoids*, J. Pure Appl. Algebra **223** (2019), 3889 – 3918.
- [16] B. Girard, *Inverse zero-sum problems and algebraic invariants*, Acta Arith. **135** (2008), 231 – 246.
- [17] ———, *A new upper bound for the cross number of finite abelian groups*, Israel J. Math. **172** (2009), 253 – 278.
- [18] ———, *On a combinatorial problem of Erdős, Kleitman and Lemke*, Adv. Math. **231** (2012), 1843 – 1857.
- [19] A. Granville, G. Shakan, and A. Walker, *Effective results on the size and structure of sumsets*, Combinatorica **43** (2023), 1139 – 1178.
- [20] A. Granville and A. Walker, *A tight structure theorem for sumsets*, Proc. Amer. Math. Soc. **149** (2021), no. 10, 4073 – 4082.
- [21] X. He, *Cross number invariants of finite abelian groups*, J. Number Theory **136** (2014), 100 – 117.
- [22] B. Kim, *The cross number of minimal zero-sum sequences over finite abelian groups*, J. Number Theory **157** (2015), 99 – 122.
- [23] U. Krause, *A characterization of algebraic number fields with cyclic class group of prime power order*, Math. Z. **186** (1984), 143 – 148.
- [24] U. Krause and C. Zahlten, *Arithmetic in Krull monoids and the cross number of divisor class groups*, Mitt. Math. Ges. Hamb. **12** (1991), 681 – 696.
- [25] D. Kriz, *On a conjecture concerning the maximal cross number of unique factorization indexed sequences*, J. Number Theory **133** (2013), 3033 – 3056.
- [26] M.B. Nathanson, *Additive Number Theory: Inverse Problems and the Geometry of Sumsets*, Springer, 1996.
- [27] L. Wang, *Tiny zero-sum sequences over some special groups*, Open Math. **18** (2020), 820 – 828.
- [28] Q. Zhong, *On elasticities of locally finitely generated monoids*, J. Algebra **534** (2019), 145 – 167.

UNIVERSITY OF GRAZ, NAWI GRAZ, DEPARTMENT OF MATHEMATICS AND SCIENTIFIC COMPUTING, HEINRICHSTRASSE 36, 8010 GRAZ, AUSTRIA

*Email address:* [aqsa.bashir@uni-graz.at](mailto:aqsa.bashir@uni-graz.at)

LABORATOIRE ANALYSE, GÉOMÉTRIE ET APPLICATIONS, LAGA, UNIVERSITÉ SORBONNE PARIS NORD, CNRS, UMR 7539, F-93430, VILLETANEUSE, FRANCE, AND, LABORATOIRE ANALYSE, GÉOMÉTRIE ET APPLICATIONS (LAGA, UMR 7539), CO-MUE UNIVERSITÉ PARIS LUMIÈRES, UNIVERSITÉ PARIS 8, CNRS, 93526 SAINT-DENIS CEDEX, FRANCE

*Email address:* [schmid@math.univ-paris13.fr](mailto:schmid@math.univ-paris13.fr), [wolfgang.schmid@univ-paris8.fr](mailto:wolfgang.schmid@univ-paris8.fr)