



HAL
open science

Group Dispersal Modelling revisited

Mario Ayala, Jérôme Coville, Samuel Soubeyrand

► **To cite this version:**

Mario Ayala, Jérôme Coville, Samuel Soubeyrand. Group Dispersal Modelling revisited. 2024. hal-04561082

HAL Id: hal-04561082

<https://hal.science/hal-04561082>

Preprint submitted on 10 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

GROUP DISPERSAL MODELLING REVISITED

MARIO AYALA^{1,2}, JEROME COVILLE^{2,3}, SAMUEL SOUBEYRAND²

¹ **Technische Universität München,
School of Computation Information and Technology,
Boltzmannstr. 3, 85748 Garching, Germany**

² **INRAE, BioSP,
228 route de l'aérodrome, 84914 Avignon, France.**

³ **Institut Camille Jordan, UMR 5208 - Université of Lyon 1**

ABSTRACT. In this paper we revisit the notion of grouped dispersal that have been introduced by Soubeyrand and co-authors [14] to model the simultaneous (and hence dependent) dispersal of several propagules from a single source in a homogeneous environment. We built a time continuous measure valued process that takes into account the main feature of a grouped dispersal and derive its infinitesimal generator. To cope with the multigeneration aspect associated to the demography we introduce two types of propagules in the description of the population which is one of the main innovations here. We also provide a rigorous description of the process and its generator. We derive as well, some large population asymptotics of the process unveiling the degenerate ultra parabolic system of PDE satisfied by the density of population. Finally, we also show that such a PDE system has a non-trivial solution which is unique in a certain functional space.

1. INTRODUCTION

The notion of grouped dispersal has been introduced to analyse situations where several entities move or are moved together from one location to another one. This dispersal mechanism has been studied mainly for animals, e.g. juvenile tarantulas [11] and female viscachas [2], and for seeds eaten and transported by large mammals [7, 15] or birds [10]. It also appears in epidemiological contexts to understand the creation of multiple secondary foci of infection of airborne pathogens [14]. In particular, [14] proposes a mechanical description of this process assuming a hierarchical structure of dependence on the “propagules” transport.

Namely, the Grouped Dispersal Model, GDM for short, introduced in [14] relies on the following assumptions:

- (1) The numbers of propagules in each new group are independently generated from a counting distribution.
- (2) The barycenters of groups are independently transported according to a dispersal kernel.
- (3) Within each group, propagules follow independent Brownian motions, which are centered around the respective group barycenter.
- (4) Each Brownian motion, representing the trajectory of a propagule, is stopped at a stopping time proportional to the distance between the source and the deposit location of the group's barycenter.

Starting with an initial source at 0, the above assumptions lead to consider a time discrete process where the numbers of groups and propagules by group are generated randomly with some prescribed law and that the position of the deposited propagule satisfies

$$X_{jn} = X_j + B_{jn}(\nu \|X_j\|),$$

E-mail address: mario.ayala@tum.de, jerome.coville@inrae.fr, samuel.soubeyrand@inrae.fr.

Part of this work was done while M. Ayala was at INRAE supported by the project ARCHIV of the Agence Nationale de la Recherche (ANR-18-CE32-0004).

where X_j is the location of the barycenter of the group j , X_{jn} is the location of the n -th propagule of the group j and B_{jn} is a Brownian motion followed by the propagule. The stopping time is proportional to the distance of the barycenter $\|X_j\|$ by a uniform constant factor ν .

This formulation allows the formation of patchy patterns with the existence of secondary foci compatible with observed data. However, the resulting multigeneration process induces some additional constraints: in particular, all the propagules become sources at once, which prevents a natural identification of a continuous time version of the GDM process, and as well, makes difficult the derivation of large population asymptotics.

In this article, we revisit the notion of grouped dispersal, and propose new modelling tools that in a sense generalise the approach and stochastic process defined in [14]. In particular, our aim is to set up a flexible approach describing continuous time group dispersal of particles for which large population asymptotics can be easily derived, and that can also be extended to the description of the movement of a broader variety of geometric objects, e.g. disks, cylinders [13]. To this end, we use the methodology introduced in [5], and in subsequent works ([4, 3, 8]), to build a measure-valued stochastic model representing the dynamics of the particles. Then we establish some large population asymptotics of the constructed measure value GDM stochastic process and give a proper analysis of the ultra parabolic PDE system obtained.

Grouped dispersal Model. Unlike the GDM of [14], which assumes indistinguishable particles that are sources of other particles and also can travel, in order to make sense to a continuous-time measure-valued version of the GDM model we introduce two populations denoted by $\nu_s(t)$ and $\nu_p(t)$. These two populations represent respectively a population of seeds that travels through space, and a population of plants that become new sources of seeds. We assume that these two populations live in the spatial domain $\bar{\mathcal{X}}$ which is the closure of an open connected subset \mathcal{X} of \mathbb{R}^d for some $d \geq 1$. Working as in [5, 4, 3, 8], let us introduce for a Polish space X the notation $\mathcal{M}_{\text{pt}}(X)$ denoting the space of finite point measures on X endowed with the topology of weak convergence, and let us represent then the populations $\nu_s(t)$ and $\nu_p(t)$ as finite point measures on the adequate Polish spaces as follows.

First, let us consider the population of plants $\nu_p(t)$ which we simply represent by means of a finite point measure on $\bar{\mathcal{X}}$. That is for $t \geq 0$, the measure is given by

$$\nu_p(t) = \sum_{i=1}^{N_p(t)} \delta_{x_i(t)},$$

where

$$N_p(t) := \int_{\bar{\mathcal{X}}} 1 \cdot \nu_p(t)(dx)$$

is the total number of plants, and

$$(x_1(t), \dots, x_{N_p(t)}(t)) \in \bar{\mathcal{X}}^{N_p(t)}$$

corresponds to an arbitrary ordering of the $N_p(t)$ positions (the positions of plants are fixed).

Similarly, we represent the population of seeds $\nu_s(t)$ by means of a finite point measure, but this time on the product space $\bar{\mathcal{X}} \times \bar{\mathcal{X}}$:

$$\nu_s(t) = \sum_{i=1}^{N_s(t)} \delta_{y_i(t), z_i(t)},$$

where $N_s(t)$ denotes the total number of seeds at time t , and for a seed i the first coordinate, $y_i(t)$ represents the location of its parent plant (which is fixed in time) and the second coordinate, $z_i(t)$ denotes the location of the seed at time t .

In addition to this point measure description, we also require that the evolution of these measures results from the following rules:

- i): The groups of seeds generated by a plant are independently generated at exponentially distributed random times, and the number of seeds in a new group is generated according to a counting distribution $p: \mathbb{N} \cup \{0\} \rightarrow [0, 1]$.

- ii): The barycenters of the groups at time zero after release are set according to a dispersal kernel $D: \bar{\mathcal{X}} \times \bar{\mathcal{X}} \rightarrow \mathbb{R}$. Barycenters work as the initial position in space of a newly generated group of seeds. Here we can think that for a given position $x \in \bar{\mathcal{X}}$, the deposit location of the barycenter is drawn from a probability distribution with density $D(x, y)$ with respect to some measure $\bar{D}(dy)$ on $\bar{\mathcal{X}}$.
- iii): Within each group, seeds follow independent Itô's diffusions with infinitesimal generator L given by:

$$(1.1) \quad L\phi(y) = a(y) \cdot \nabla\phi(y) + \frac{\sigma(y)^2}{2} \Delta\phi(y),$$

where $\phi: \bar{\mathcal{X}} \rightarrow \mathbb{R}$ is an element of one of the following domains:

$$(1.2) \quad D_r(L) = \left\{ \phi \in \mathcal{C}^2(\mathcal{X}) : \frac{\partial}{\partial n} \phi(y) = 0, \quad \forall y \in \partial\mathcal{X} \right\},$$

or

$$(1.3) \quad D_k(L) = \{ \phi \in \mathcal{C}^2(\mathcal{X}) : \phi(y) = 0, \quad \forall y \in \partial\mathcal{X} \}.$$

Notice that the first domain D_r encodes diffusion of seeds with normal reflection in the boundary of \mathcal{X} , while the second represents *killing* of seeds when touching the boundary. Moreover, we also consider the possibility of \mathcal{X} being the whole \mathbb{R}^d , in which case the boundary condition on the domain is omitted.

- iv): Each diffusion, representing the trajectory of a seed currently alive, is independently of each other and stopped at an exponentially distributed random time of a given parameter function λ (for example with mean proportional to the distance between the source and the deposit location of the group's barycenter). At the stopping time, the seed is assumed to mature and become a new plant with fixed position being equal to the position of the *mother seed* at the time of maturity.

Infinitesimal generator. Let us consider the $\mathcal{M}_{\text{pt}}(\mathcal{X}) \times \mathcal{M}_{\text{pt}}(\mathcal{X}^2)$ -valued process

$$\{\nu_t : t \geq 0\} = \{(\nu_p(t), \nu_s(t)) : t \geq 0\}.$$

The evolution of this process can be described in terms of the infinitesimal generator:

$$(1.4) \quad \begin{aligned} \mathcal{L}F_\phi(\nu) &= \mathcal{L}F_\phi(\nu_p, \nu_s), \\ &= \int_{\bar{\mathcal{X}}^2} \lambda(x, y) [F_\phi(\nu_p + \delta_y, \nu_s - \delta_{x,y}) - F_\phi(\nu_p, \nu_s)] \nu_s(dx, dy) \\ &+ \sum_{\kappa \in \mathbb{N}} q(\kappa) \int_{\bar{\mathcal{X}}^2} D(x, y) [F_\phi(\nu_p, \nu_s + \kappa\delta_{x,y}) - F_\phi(\nu_p, \nu_s)] dy \nu_p(dx) \\ &+ \left(\int_{\bar{\mathcal{X}}^2} L_z \phi_s(y, z) \nu_s(dy, dz) \right) \partial_z F(\langle \phi, \nu \rangle) + \left(\int_{\bar{\mathcal{X}}^3} \frac{(\sigma(z))^2}{2} |\nabla_z \phi_s(y, z)|^2 \nu_s(dy, dz) \right) \partial_z^2 F(\langle \phi, \nu \rangle), \end{aligned}$$

where:

$$(1.5) \quad F_\phi(\nu) := F_\phi(\nu_p, \nu_s) := F(\langle \phi_p, \nu_p \rangle, \langle \phi_s, \nu_s \rangle),$$

for $F \in \mathcal{C}^2(\mathbb{R}^2; \mathbb{R})$, and a pair of functions $\phi := \{\phi_p, \phi_s\}$ being such that:

- ϕ_α is measurable for $\alpha \in \{p, s\}$.
- For every $y \in \bar{\mathcal{X}}$, $\phi_s(y, \cdot) : \bar{\mathcal{X}} \rightarrow \mathbb{R}$ is in the domain $D_\iota(L)$ for $\iota \in \{r, k\}$.

Remark 1. Thereafter, whenever we need to refer to a particular choice of cylindrical function we will just need to specify the choice of F and the pair $\phi = \{\phi_p, \phi_s\}$. Moreover, for such ϕ we introduce the additional notation

$$(1.6) \quad \langle \phi, \nu \rangle := \langle \phi_p, \nu_p \rangle_{\bar{\mathcal{X}}} + \langle \phi_s, \nu_s \rangle_{\bar{\mathcal{X}}^2},$$

where $\nu = (\nu_p, \nu_s)$.

Each of the terms on the RHS of (1.4) corresponds to a type of event in the dynamics:

- The first term corresponds to maturation of seeds to become plants. This happens at rate $\lambda(x, y)$, where $x \in \bar{\mathcal{X}}$ is the location of the seed that produced this seed, and $y \in \bar{\mathcal{X}}$ is the current position of the seed undergoing maturation.

- The second terms corresponds to the release of a group of κ new seeds, which are displaced from their mother plant's position $x \in \mathcal{X}$ to a new position $y \in \mathcal{X}$ according to the displacement kernel D .
- The last two terms correspond to the diffusion of seeds. They come from an application of Itô's formula to integrals of the finite point measure ν_s . See for example [4].

Describing the dynamics in terms of the infinitesimal generator (1.4) is non-rigorous and requires some extra effort to verify that the process it represents is indeed well-defined. In particular we need the following assumption:

Assumption 1. The model parameters are as follows:

- (1) The counting distribution $q: \mathbb{N} \rightarrow \mathbb{R}$, has finite first and second moments, i.e.:

$$(1.7) \quad \mu_1 := \sum_{\kappa \in \mathbb{N}} q(\kappa) \kappa < \infty,$$

and

$$(1.8) \quad \sqrt{\mu_2} := \sum_{\kappa \in \mathbb{N}} q(\kappa) \kappa^2 < \infty.$$

- (2) The dispersal kernel $D(x, y)$ is of compact support in its second argument. Moreover, it is a probability density with respect to some measure \bar{D} on $\bar{\mathcal{X}}$.
- (3) There exists $\bar{\lambda} > 0$ such that the diffusion stopping rate $\lambda: \bar{\mathcal{X}} \times \bar{\mathcal{X}} \rightarrow \mathbb{R}$ is uniformly bounded:

$$\sup_{(x, y) \in \bar{\mathcal{X}}^2} \lambda(x, y) =: \bar{\lambda} < \infty.$$

We refer to the Appendix for a rigorous description of the processes in terms of Poisson point measures.

Main Results. Before stating our main result, the following theorem verifies that indeed we have a process which is well-defined.

Theorem 1.1. Let $\nu_0 = (\nu_p(0), \nu_s(0))$ be such that we have:

$$(1.9) \quad \mathbb{E}(\langle \mathbf{1}, \nu_0 \rangle) < \infty,$$

where

$$(1.10) \quad \langle \mathbf{1}, \nu_0 \rangle = \int_{\mathcal{X}} 1 \nu_p(0)(dx) + \int_{\mathcal{X}^2} 1 \nu_s(0)(dx, dy).$$

Then, under Assumption 1, part 1), the process $(\nu_t)_{t \geq 0} = (\nu_p(t), \nu_s(t) : t \geq 0)$ satisfies:

$$(1.11) \quad \mathbb{E} \left(\sup_{t \in [0, T]} \langle \mathbf{1}, \nu_t \rangle \right) < \infty.$$

In particular, we also have that the process ν_t is well-defined.

For a proof we refer to the Appendix, where we proved (1.11) for higher-order moments, and where we also showed that the dynamics of the process $(\nu_t)_{t \geq 0}$ indeed corresponds to the one described by the infinitesimal generator \mathcal{L} .

Our first main result concerns the derivation of scaling limits for the joint populations of plants and seeds. In order to derive this result, we introduce a rescaling parameter K which will tend to infinity. This parameter represents the order of the size of the populations and it is used to rescale our populations as follows:

$$(1.12) \quad \nu_p^K(t) = \frac{1}{K} \sum_{i=1}^{N_p(t)} \delta_{x_i(t)}, \quad \text{and} \quad \nu_s^K(t) = \frac{1}{K} \sum_{i=1}^{N_s(t)} \delta_{x_i(t), y_i(t)},$$

where we consider the processes ν_p^K and ν_s^K as taking values on the spaces $\frac{1}{K} \mathcal{M}_{\text{pt}}(\bar{\mathcal{X}})$ and $\frac{1}{K} \mathcal{M}_{\text{pt}}(\bar{\mathcal{X}}^2)$ respectively. Additionally, we assume that the model parameters also depend on the scaling parameter K as follows:

$$(1.13) \quad \lambda_K(x, y) = \frac{1}{K} \lambda(x, y) \quad \text{and} \quad D_K(x, y) = \frac{1}{K} D(x, y),$$

where λ and D satisfy Assumption 1.

The rescaling (1.12) can be interpreted as saying that the initial populations of plants and seeds are roughly of the same order, namely the order K . To make this idea more precise we assume that at time zero a law of large numbers is satisfied:

Assumption 2. Assume that the sequence of measures $\{\nu_0^{(K)}\}_{K \geq 1} = \{(\nu_p^{(K)}(0), \nu_s^{(K)}(0))\}_{K \geq 1}$ converges weakly to a deterministic measure $\xi(0) = (\xi_p(0), \xi_s(0))$. Moreover, assume that the measures $\xi_p(0)$, and $\xi_s(0)$, have densities $f_0 : \mathcal{X} \rightarrow \mathbb{R}_+$, and $g_0 : \mathcal{X}^2 \rightarrow \mathbb{R}_+$, with respect to the Lebesgue measures on \mathcal{X} , and \mathcal{X}^2 , respectively.

Under Assumption 1 and Assumption 2 we have the following result:

Theorem 1.2. *Let the sequence of initial measures $\{\nu_0^{(K)}\}_{K \geq 1} = \{(\nu_p^{(K)}(0), \nu_s^{(K)}(0))\}_{K \geq 1}$ be such that:*

$$(1.14) \quad \sup_{K \in \mathbb{N}} \mathbb{E} \left[\langle 1, \nu_p(0)^{(K)} \rangle^3 + \langle 1, \nu_s(0)^{(K)} \rangle^3 \right] < \infty.$$

Then, for all $T > 0$, the sequence $\{\nu(t)^{(K)} : t \in [0, T]\}_{K \geq 1} = \{(\nu_p^{(K)}(t), \nu_s^{(K)}(t)) : t \in [0, T]\}_{K \geq 1}$ converges in law in $\mathbb{D}([0, T], \mathcal{M}_F(\mathcal{X}) \times \mathcal{M}_F(\mathcal{X}^2))$ to a deterministic $\xi_t = (f(t, x) dx, g(t, x, y) dx dy)$, element of $\mathbb{C}([0, T], \mathcal{M}_F(\mathcal{X}) \times \mathcal{M}_F(\mathcal{X}^2))$, where f and g are the unique solution of the following system:

$$(1.15) \quad \partial_t f(t, x) = \int_{\bar{\mathcal{X}}} \lambda(z, x) g(t, z, x) dz,$$

and

$$(1.16) \quad \partial_t g(t, x, y) = L^* g(t, x, y) - \lambda(x, y) g(t, x, y) + \mu_1 D(x, y) f(t, x),$$

with initial conditions

$$(1.17) \quad f(0, x) = f_0(x), \quad g(0, x, y) = g_0(x, y), \quad \forall (x, y) \in \mathcal{X}^2,$$

and for all $t \geq 0, x \in \mathcal{X}$ the functions $g(t, x, \cdot)$ satisfy for all $y \in \partial \mathcal{X}$:

$$(1.18) \quad \nabla_y g(t, x, y) \cdot \vec{n} = 0,$$

with \vec{n} the normal of $\partial \mathcal{X}$ for the reflecting case, and

$$(1.19) \quad g(t, x, y) = 0,$$

for the case in which seeds are killed in the boundary.

Of interest by itself, the degenerate PDE system obtained falls into the class of ultra-parabolic PDE system. However, it turns out that classical theory do not apply here and an existence theory has to be established. In the next results, we provide a first existence theorem in the particular situation of a standard diffusion with reflecting boundary case in a bounded domain \mathcal{X} , leading to consider the following coupled PDE system:

$$(1.20) \quad \partial_t f(t, y) = \int_{\mathcal{X}} \lambda(z, y) g(t, z, y) dz \quad \text{for all } t > 0, y \in \mathcal{X},$$

$$(1.21) \quad \partial_t g(t, x, y) = \nabla_y (A \cdot \nabla_y g(t, x, y)) - \lambda(x, y) g(t, x, y) + \mu_1 D(x, y) f(t, x) \quad \text{for all } t > 0, (x, y) \in \mathcal{X}^2,$$

$$(1.22) \quad \nabla_y g(t, x, y) \cdot \vec{n} = 0 \quad \text{for all } t > 0, x \in \mathcal{X}, y \in \partial \mathcal{X},$$

$$(1.23) \quad f(0, x) = f_0(x) \quad \text{for all } x \in \mathcal{X},$$

$$(1.24) \quad g(0, x, y) = 0 \quad \text{for all } (x, y) \in \mathcal{X}^2,$$

with A an elliptic matrix and \mathcal{X} a bounded domain of \mathbb{R}^d .

Theorem 1.3. *Let \mathcal{X} be a smooth bounded domain, with at least a $C^{1,\alpha}$ boundary. Assume further that $\lambda, D \in W^{1,\infty}(\mathcal{X}^2)$, $f_0 \in H^1(\mathcal{X})$ and A is an elliptic matrix. Then there exists a (f, g) such that $f \in C^1((0, T), H^1(\mathcal{X}))$, $g \in C^1((0, T), H^1(\mathcal{X}) \times H^1(\mathcal{X}))$ solution to (1.20)–(1.24). In addition, if D, λ are non negative functions, this solution is unique in this class of function.*

A careful inspection of the proofs shows that our arguments can readily be transposed to Dirichlet system and so we have

Theorem 1.4. *Let \mathcal{X} be a smooth bounded domain, with at least a $C^{1,\alpha}$ boundary. Assume further that $\lambda, D \in W^{1,\infty}(\mathcal{X}^2)$, $f_0 \in H_0^1(\mathcal{X})$, A an elliptic matrix. Then there exists a (f, g) such that $f \in C^1((0, T), H_0^1(\mathcal{X}))$, $g \in C^1((0, T), H_0^1(\mathcal{X}) \times H_0^1(\mathcal{X}))$ solution to (1.20)–(1.24). In addition, if D, λ are positive functions, this solution is unique in this class of function.*

Organization of the paper. The rest of our paper is organized as follows. In Section 2 we provide a specific example in order to make transparent the connections of this setting with the one introduced earlier in [14]. Moreover, in the same section by means of computer simulations we show that our modeling framework can indeed produce patchy patterns for various common choices of dispersal kernels. In the same section we also present how our modelling framework can be used to obtain analytical expressions for observables such as the total number of plants at a given time. In Section 3 we prove our main theorems using the standard compactness-uniqueness approach. There, we also include the derivation of some standard martingale properties that are used in our proofs. In Section 4, we provide a preliminary analysis of the degenerate PDE ultra parabolic system resulting of the large population asymptotic of the GDM process and prove the existence results. Finally, in the Appendix we include the rigorous definition of the processes on path-space and the proof of some of the necessary propositions in order to rigorously construct a solution to the PDE system.

2. A MEASURE VALUED ADAPTATION OF THE GDM

As an example of the class of processes we have just introduced, we consider the setting given in Section 4.1 in [14] adapted to our framework. This means:

- The spatial domain is

$$\bar{\mathcal{X}} = [-100, 100] \times [-100, 100].$$

- The counting distribution of the number of particles is given by a negative binomial distribution with mean μ_1 and size parameter $s = \mu_1^2 / (\mu_2 - \mu_1)$, i.e.,

$$(2.1) \quad q(\kappa) = \frac{\Gamma(\kappa + s)}{\Gamma(s)\Gamma(\kappa + 1)} \left(\frac{\mu_1}{\mu_2}\right)^s \left(1 - \frac{\mu_1}{\mu_2}\right)^\kappa.$$

- The dispersal kernel $D : \bar{\mathcal{X}} \times \bar{\mathcal{X}} \rightarrow \mathbb{R}$ is of exponential form, i.e., given by

$$(2.2) \quad D(x, y) = \frac{1}{2\pi\beta^2} e^{-\frac{\|x-y\|}{\beta}}.$$

- At time zero the initial populations of plants and seeds are given by

$$(2.3) \quad \nu_p(0) = \delta_0, \quad \text{and} \quad \nu_s(0) = 0.$$

Moreover, we will assume the seed's maturity rate (i.e. diffusion stopping time's rate) to be

$$(2.4) \quad \lambda(x, y) = \lambda_0 \|y - x\|,$$

for some $\lambda_0 \geq 0$.

2.1. Simulations. In this section we show some simulations of our model. Let us first introduce some notation to present an algorithm to simulate this process. The notation $\text{Exp}(u)$ denotes an exponentially distributed random variable of parameter u . For the measures $\nu_p(t)$, the notation $\vec{v}_p(t)$ denotes an ordered array with the positions of each plant. Analogously for the measure $\nu_s(t)$. The command `RandomChoice(A, Weights= B)` selects a random element from A according to the weights given by B . For both the sake of generality and shortness of exposition, the command `Update diffusion(ν_s, dt)` does as its names suggests, updates the position of the population of seeds according to the relevant Itô diffusion with the specified time step.

Algorithm 1: Simulating the measure valued group dispersal model

```

 $T_0 \leftarrow 0;$ 
 $\nu_p(T_0) \leftarrow \delta_0;$ 
 $N_p(T_0) \leftarrow 1;$ 
 $\nu_s(T_0) \leftarrow \emptyset;$ 
 $N_s(T_0) \leftarrow 0;$ 
 $k \leftarrow 0;$ 
while  $T_k < T_{max}$  do
  for  $i \leftarrow 0$  to  $N_s(T_k)$  do
     $R_{i,k} \leftarrow \int \lambda(x, y) \nu_s(T_k)(dx, dy)$ 
  end for
   $R_k \leftarrow N_p(T_k) + \sum_i R_{i,k};$ 
   $\epsilon_k \leftarrow \text{Simulate}(\text{Exp}(R_k));$ 
   $T_{k+1} \leftarrow T_k + \epsilon_k;$ 
   $\theta_k \leftarrow \text{Simulate}(\text{Unif}([0, 1]));$ 
  if  $0 \leq \theta_k \leq \frac{N_p(T_k)}{R_k}$  then
     $p \leftarrow \text{RandomChoice}(\nu_p(T_k), \text{Weights} = [1, \dots, 1]);$ 
     $n \leftarrow \text{Simulate}(\text{Progeny});$ 
     $y \leftarrow \text{Simulate}(D(\vec{\nu}_p(T_k))(p));$ 
     $\nu_s(T_k) \leftarrow \nu_s(T_k) + n \cdot \delta_{\nu_p(T_k)(p), y};$ 
     $N_s(T_{k+1}) \leftarrow N_s(T_k) + n;$ 
  else
     $s \leftarrow \text{RandomChoice}(\nu_s(T_k), \text{Weights} = [R_{i,k}]);$ 
     $(x, y) \leftarrow \vec{\nu}_s(T_k)(s);$ 
     $\nu_s(T_k) \leftarrow \nu_s(T_k) - \delta_{x, y};$ 
     $\nu_p(T_k) \leftarrow \nu_p(T_k) + \delta_y;$ 
     $N_s(T_{k+1}) \leftarrow N_s(T_k) - 1;$ 
     $N_p(T_{k+1}) \leftarrow N_p(T_k) + 1;$ 
  end if
   $\nu_p(T_{k+1}) \leftarrow \nu_p(T_k);$ 
   $\nu_s(T_{k+1}) \leftarrow \text{Update diffusion}(\nu_s(T_k), \epsilon_k);$ 
   $k \leftarrow k + 1;$ 
end while

```

The simulations that we present in this section differ in the choice of the dispersal kernel. Here we focus on the two dimensional case, and use the following dispersal kernels: Gaussian, exponential, and inverse power law. Simulations are run with parameters given by Table 1 below and an Itô diffusion without drift and $\sigma^2 = 5$. The parameters for the dispersal kernels are pick such that an average displacement of 10 units is enforced.

Parameter	Value
\mathcal{X}	$[-100, 100] \times [-100, 100]$
$q(\kappa)$	$\frac{\Gamma(\kappa+s)}{\Gamma(s)\Gamma(\kappa+1)} \left(\frac{\mu_1}{\mu_2}\right)^s \left(1 - \frac{\mu_1}{\mu_2}\right)^\kappa$
μ_i	$\mu_1 = 1$ and $\mu_2 = 25$
$\lambda(x, y)$	$0.05 \cdot \ x\ $
$\nu_p(0)$	δ_0
$\nu_s(0)$	0

TABLE 1. Parameters used in simulations. The dispersal kernel is different in each simulation case.

For the three cases of dispersal kernel mentioned above, we show a run of the simulations on which patchy patterns emerge. All simulations are stopped until a total population of 2000 plants is reached. After this, a smoothing is performed using the Gaussian kde method from the `scipy` library of Python3.

Exponential kernel. In this case the kernel is given by:

$$(2.5) \quad D(x, y) = \frac{1}{2\pi\beta^2} e^{-\frac{\|x-y\|}{\beta}}$$

for some $\beta > 0$.




FIGURE 1. Patchy pattern observed in a simulation using an exponential dispersal kernel with mean displacement equal to 10, and where seeds follow diffusion with killing in the boundary. Left: pattern of particles. Right: estimated point intensity obtained by kernel smoothing with a Gaussian kernel.




FIGURE 2. Patchy pattern observed in a simulation using an exponential dispersal kernel with mean displacement equal to 10, and where seeds follow diffusion with reflection in the boundary. Left: pattern of particles. Right: estimated point intensity obtained by kernel smoothing with a Gaussian kernel.

Gaussian kernel. The Gaussian kernel case is given by:

$$(2.6) \quad D(x, y) = \frac{1}{2\pi\beta^2} e^{-\frac{\|x-y\|^2}{\beta^2}}$$

for $\beta > 0$.




FIGURE 3. Patchy pattern observed in a simulation using a Gaussian dispersal kernel with mean displacement 10, and where seeds follow diffusion with killing in the boundary. Left: pattern of particles. Right: estimated point intensity obtained by kernel smoothing with a Gaussian kernel.




FIGURE 4. Patchy pattern observed in a simulation using a Gaussian dispersal kernel with mean displacement 10, and where seeds follow diffusion with reflection in the boundary. Left: pattern of particles. Right: estimated point intensity obtained by kernel smoothing with a Gaussian kernel.

Power-law kernel. For $\beta > 0$ and $a > 2$, the power-law kernel takes the form.

$$(2.7) \quad D(x, y) = \frac{(a-2)(a-1)}{2\pi\beta^2} \left(1 + \frac{\|x-y\|}{\beta}\right)^{-a}$$




FIGURE 5. Patchy pattern observed in a simulation using a power-law dispersal kernel of parameters $\beta = 5$, and $a = 4$ (i.e. with mean displacement 10), and where seeds follow diffusion with killing in the boundary. Left: pattern of particles. Right: estimated point intensity obtained by kernel smoothing with a Gaussian kernel.




FIGURE 6. Patchy pattern observed in a simulation using a power-law dispersal kernel of parameters $\beta = 5$, and $a = 4$ (i.e. with mean displacement 10), and where seeds follow diffusion with reflection in the boundary. Left: pattern of particles. Right: estimated point intensity obtained by kernel smoothing with a Gaussian kernel.

2.2. Population level descriptors as observables. Here, we focus on the simple situation in which seeds mature at constant rate, i.e., $\lambda(x, y) = 1$ for all $x, y \in \mathcal{X}$. Let us denote by $N_p(t)$ and $N_s(t)$ the random number of plants and seeds at time t , respectively. These two important quantities can be obtained by integration of the measures $\nu_p(t)$ and $\nu_s(t)$ against the constant function one:

$$(2.8) \quad N_p(t) = \langle 1, \nu_p(t) \rangle, \text{ and } N_s(t) = \langle 1, \nu_s(t) \rangle,$$

where we have abused notation by representing with same symbol, $\langle \cdot, \cdot \rangle$, integration over the spaces \mathcal{X} and \mathcal{X}^2 .

Using the algorithm of Section 2.1, we obtain the following quantitative scenario for the expected number of plants:




FIGURE 7. Expected number of plants with a time horizon $T = 20$. Simulations were run with an exponential dispersal kernel (truncated to $[-100, 100] \times [-100, 100]$), and a constant maturity rate $\lambda = 1$.

Thanks to the knowledge of the generator \mathcal{L} have the following result concerning the expectation of $N_p(t)$ and $N_s(t)$ when starting from a single plant at the origin.

Proposition 2.1. *Let $\nu(0) = (\delta_0, 0)$, then we have:*

$$(2.9) \quad \mathbb{E}N_p(t) = \frac{\left((\sqrt{4\mu+1}-1)e^{\frac{1}{2}(\sqrt{4\mu+1}-1)t} + (\sqrt{4\mu+1}+1)e^{-\frac{1}{2}(\sqrt{4\mu+1}+1)t} \right)}{2\sqrt{4\mu+1}},$$

$$(2.10) \quad \mathbb{E}N_s(t) = \frac{\mu \left(e^{\frac{1}{2}(\sqrt{4\mu+1}-1)t} - e^{-\frac{1}{2}(\sqrt{4\mu+1}+1)t} \right)}{\sqrt{4\mu+1}}.$$

where μ is given by:

$$(2.11) \quad \mu = \sum_{\kappa} q(\kappa)\kappa.$$

3. PROOFS

We will not show Theorem 1.2 directly, instead we are going to provide a formulation of it in IDE form, i.e. in the form of Theorem 3.1 below.

3.1. Scaling limits: general statement. We first relax Assumption 2 by drooping the required existence of densities at time zero.

Assumption 3. Assume that the sequence of measures $\{\nu_0^{(K)}\}_{K \geq 1} = \{(\nu_p^{(K)}(0), \nu_s^{(K)}(0))\}_{K \geq 1}$ converges weakly to a deterministic measure $\xi(0) = (\xi_p(0), \xi_s(0))$.

Under Assumption 1 and Assumption 3 we have the following result:

Theorem 3.1. *Let the sequence of initial measures $\{\nu_0^{(K)}\}_{K \geq 1} = \{(\nu_p^{(K)}(0), \nu_s^{(K)}(0))\}_{K \geq 1}$ be such that for all $K \in \mathbb{N}$ we have:*

$$(3.1) \quad \mathbb{E} \left[\langle 1, \nu_p(0)^{(K)} \rangle^3 + \langle 1, \nu_s(0)^{(K)} \rangle^3 \right] < \infty.$$

Then, for all $T > 0$, the sequence $\{\nu(t)^{(K)} : t \in [0, T]\}_{K \geq 1} = \{(\nu_p^{(K)}(t), \nu_s^{(K)}(t)) : t \in [0, T]\}_{K \geq 1}$ converges in law in $\mathbb{D}([0, T], \mathcal{M}_F(\mathcal{X}) \times \mathcal{M}_F(\mathcal{X}^2))$ to a deterministic continuous function $\xi_t = (\xi_p(t), \xi_s(t))$ being the unique solution of the following system:

$$(3.2) \quad \langle \xi_p(t), \phi_p \rangle_{\bar{\mathcal{X}}} - \langle \xi_p(0), \phi_p \rangle_{\bar{\mathcal{X}}} = \int_0^t \int_{\bar{\mathcal{X}}^2} \lambda(x, y) \phi_p(y) \xi_s(s) (dx, dy) ds$$

$$(3.3) \quad \langle \xi_s(t), \phi_s \rangle_{\bar{\mathcal{X}}^2} - \langle \xi_s(0), \phi_s \rangle_{\bar{\mathcal{X}}^2} = \int_0^t \langle \xi_s(s), \mathcal{L}_y \phi_s(x, y) \rangle_{\bar{\mathcal{X}}^2} ds - \int_0^t \int_{\bar{\mathcal{X}}^2} \lambda(x, y) \phi_s(x, y) \xi_s(s) (dx, dy) ds$$

$$+ \mu_1 \int_0^t \int_{\bar{\mathcal{X}}^2} D(x, y) \phi_s(x, y) \xi_p(s) (dx, dy) ds$$

with initial conditions

$$(3.4) \quad \xi_p(0) = \xi \text{ and } \xi_s(0) = \xi,$$

and for all x with $\phi_s(x, \cdot)$ being in the domain of the diffusive generator L .

3.2. Martingale characterization. In this section we discuss the proofs of our main results. Let us start with some well-known martingales associated to Markov processes. We start with a simple application of Dynkin's theorem:

Theorem 3.2. *Let $\nu_0 = (\nu_p(0), \nu_s(0))$ be such that for some $r \geq 2$ we have:*

$$(3.5) \quad \mathbb{E}(\langle \mathbf{1}, \nu_p(0) \rangle^r + \langle \mathbf{1}, \nu_s(0) \rangle^r) < \infty,$$

and

$$(3.6) \quad \mu_r := \sum_{\kappa} q(\kappa) \kappa^r < \infty.$$

Let also F and ϕ , be such that there exists a C , possibly dependant on F , and ϕ , such that for all $\nu \in \mathcal{M}_p$:

$$(3.7) \quad |F_\phi(\nu)| + |\mathcal{L}F_\phi(\nu)| \leq C (1 + \langle \mathbf{1}, \nu \rangle^r).$$

Then, under Assumption 1, we have that the process

$$(3.8) \quad M_t(F_\phi) = F_\phi(\nu(t)) - F_\phi(\nu(0)) - \int_0^t \mathcal{L}F_\phi(\nu(s)) ds$$

is a càdlàg martingale.

Proof. From Proposition A.3 and Dynkin's theorem we know that $M_t(F_\phi)$ is a local martingale. Hence, it is enough to show that the R.H.S. of (3.8) is integrable. This is a consequence of assumption (3.7) and Proposition A.4. \square

Remark 2. Simple but tedious computations show that for, $1 \leq m \leq r - 1$, the functions

$$(3.9) \quad F_p^m(\nu) = (\langle \phi_p, \nu_p \rangle_{\bar{\mathcal{X}}})^m, \text{ and } F_s^m(\nu) = (\langle \phi_s, \nu_s \rangle_{\bar{\mathcal{X}}})^m$$

satisfy condition (3.7).

The following result is a consequence of Proposition 3.2 and Remark 2

Proposition 3.3. *Let ϕ_p and ϕ_s be as in Remark 2, then under Assumption 1, we have the following càdlàg martingales:*

(1) *For the population of plants:*

$$(3.10) \quad M_t^{\phi_p} = \langle \nu_p(t), \phi_p \rangle_{\bar{\mathcal{X}}} - \langle \nu_p(0), \phi_p \rangle_{\bar{\mathcal{X}}} - \int_0^t \int_{\bar{\mathcal{X}}^2} \lambda(x, y) \phi_p(y) \nu_s(s) (dx, dy) ds$$

is a càdlàg martingale with predictable quadratic variation

$$(3.11) \quad \langle M^{\phi_p} \rangle_t = \int_0^t \int_{\bar{\mathcal{X}}^2} \lambda(x, y) \phi_p(y)^2 \nu_s(s) (dx, dz) ds$$

(2) For the population of seeds:

$$\begin{aligned}
(3.12) \quad M_t^{\phi_s} &= \langle \nu_s(t), \phi_s \rangle_{\bar{\mathcal{X}}^2} - \langle \nu_s(0), \phi_s \rangle_{\bar{\mathcal{X}}^2} - \int_0^t \langle \nu_s(s), \mathcal{L}_y \phi_s(x, y) \rangle_{\bar{\mathcal{X}}^2} ds \\
&+ \int_0^t \int_{\bar{\mathcal{X}}^2} \lambda(x, y) \phi_s(x, y) \nu_s(s) (dx, dy) ds \\
&- \sum_{\kappa} q(\kappa) \kappa \int_0^t \int_{\bar{\mathcal{X}}^2} D(x, y) \phi_s(x, y) \nu_p(s) (dx, dy) ds
\end{aligned}$$

is a càdlàg martingale with predictable quadratic variation

$$\begin{aligned}
(3.13) \quad \langle M^{\phi_s} \rangle_t &= \int_0^t \int_{\bar{\mathcal{X}}^2} \sigma^S(y)^2 |\nabla_y \phi_s(x, y)|^2 \nu_s(s) (dx, dy) ds + \int_0^t \int_{\bar{\mathcal{X}}^2} \lambda(x, y) \phi_s(x, y)^2 \nu_s(s) (dx, dz) ds \\
&+ \sum_{\kappa} q(\kappa) \kappa^2 \int_0^t \int_{\bar{\mathcal{X}}^2} D(x, y) \phi_s(x, y)^2 \nu_p(s) (dx, dy) ds
\end{aligned}$$

Proof. Assume $r \geq 2$. By Proposition 3.2 and Remark 2 with $m = 1$, we have that $M_t^{\phi_p}$ and $M_t^{\phi_s}$ are càdlàg martingales. The predictable quadratic variation is also obtained by standard arguments. However, we show the simple case $\langle M^{\phi_p} \rangle_t$ for completeness of exposition.

Using Remark 2, with $q = 2$, we obtain that the following is a càdlàg martingale:

$$(3.14) \quad \langle \phi_p, \nu_p(t) \rangle^2 - \langle \phi_p, \nu_p(0) \rangle^2 - \int_0^t \int_{\bar{\mathcal{X}}^2} \lambda(x, y) \left[(\langle \phi_p, \nu_p(s) \rangle + \phi_p(y))^2 - \langle \phi_p, \nu_p(s) \rangle^2 \right] \nu_s(s) (dx, dy) ds,$$

On the other hand, using Itô's formula on (3.29) gives the additional martingale:

$$(3.15) \quad \langle \phi_p, \nu_p(t) \rangle^2 - \langle \phi_p, \nu_p(0) \rangle^2 - 2 \int_0^t \langle \phi_p, \nu_p(s) \rangle \int_{\bar{\mathcal{X}}^2} \lambda(x, y) \phi_p(y) \nu_s(s) (dx, dy) ds - \langle M^{\phi_p} \rangle_t.$$

We conclude using the uniqueness of the Doob-Meyer's decomposition. \square

We now present some ways on how to use Proposition and Proposition 3.3 to deduce some properties of our process. Let us start with the latter.

3.3. Proof: expected number of plants. Thanks to the Martingale characterization, the proof of Proposition 2.1 becomes easily accessible.

Proof of Proposition 2.1. We can use the martingale characterization given in Proposition 3.3, with $\phi_p(\cdot) = 1 = \phi_s(\cdot, \cdot)$, to obtain the system

$$(3.16) \quad \mathbb{E}N_p(t) = 1 + \int_0^t \mathbb{E}N_s(s) ds,$$

$$(3.17) \quad \mathbb{E}N_s(t) = \mu \cdot \int_0^t \mathbb{E}N_p(s) ds - \int_0^t \mathbb{E}N_s(s) ds,$$

with initial conditions $N_p(0) = 1$ and $N_s(0) = 0$. Solving this first order linear system finishes the proof. \square

3.4. Proof of Theorem 1.2. Here we show how Theorem 1.2 is an application of Theorem 3.1 under the additional assumption of having densities at time zero. In this section we additionally assume the following:

- The spatial diffusion to be given by standard reflected Brownian motion. This means zero drift, and $\sigma = 1$.
- The existence of densities at time zero, and the propagation of absolute continuity for later times.

Let us first start with the equations concerning the population of plants. Under the above assumptions we can rewrite (3.2) as follows:

$$(3.18) \quad \int_{\bar{\mathcal{X}}} \phi_p(x) f(t, x) dx - \int_{\bar{\mathcal{X}}} \phi_p(x) f(0, x) dx = \int_0^t \int_{\bar{\mathcal{X}}^2} \lambda(z, x) \phi_p(x) g(s, z, x) dz dx ds,$$

where $f(t, x)$ denotes the density of plants at position $x \in \bar{\mathcal{X}}$, at time t .

Differentiating with respect to time we obtain

$$(3.19) \quad \int_{\bar{\mathcal{X}}} \phi_p(x) \partial_t f(t, x) dx = \int_{\bar{\mathcal{X}}^2} \lambda(z, x) \phi_p(x) g(t, z, x) dz dx.$$

By DuBois-Reymond lemma, we have the strong form

$$(3.20) \quad \partial_t f(t, x) = \int_{\bar{\mathcal{X}}} \lambda(z, x) g(t, z, x) dz.$$

Analogously for (3.3) we have the strong formulation

$$(3.21) \quad \partial_t g(t, x, y) = \frac{1}{2} \Delta_y g(t, x, y) - \lambda(x, y) g(t, x, y) + \mu_1 D(x, y) f(t, x).$$

3.5. Proof of Theorem 3.1. For the proof of Theorem 3.1 we are going to use the standard compactness-uniqueness approach:

Step 1: show uniqueness of solutions. Let us first assume that $(\xi_p(t), \xi_s(t))_{t \geq 0}$, and $(\bar{\xi}_p(t), \bar{\xi}_s(t))_{t \geq 0}$ are solutions of (3.2)-(3.3). We want to show that for $\alpha \in \{p, s\}$ we have:

$$(3.22) \quad \|\xi_\alpha - \bar{\xi}_\alpha\|_\alpha = 0,$$

where for ν_α^1 and ν_α^2 their variation norm is given by

$$(3.23) \quad \|\nu_\alpha^1 - \nu_\alpha^2\| = \sup_{\substack{\phi_\alpha \in L^\infty \\ \|\phi_\alpha\|_\infty \leq 1}} |\langle \nu_\alpha^1 - \nu_\alpha^2, \phi_\alpha \rangle|,$$

where the inner products are taken in the appropriate spaces.

Let us first deal with the case $\alpha = p$. Let ϕ_p be such that $\|\phi_p\|_\infty \leq 1$, by (3.2) and Assumption 1, we have

$$(3.24) \quad |\langle \xi_p(t) - \bar{\xi}_p(t), \phi_p \rangle| \leq \bar{\lambda} \int_0^t \sup_{\phi \leq 1} |\langle \xi_s(t) - \bar{\xi}_s(t), \phi \rangle| ds.$$

For the case $\alpha = s$, we consider the semi-groups $P_s(t)$ corresponding to the diffusion process with generator L . Let us fix a function $\phi_s \in L^\infty(\mathcal{X}^2)$ with $\|\phi_s\| \leq 1$, take $t \in [0, T]$, and define the following function:

$$(3.25) \quad \phi_s(s, x, y) = P_s(t - s) \phi_s(x, y) \quad \forall (x, y) \in \mathcal{X}^2,$$

where the semigroup acts on the second variable of the original ϕ_s .

By construction $\phi_s(s, x, y)$ is a solution of the boundary problem:

$$(3.26) \quad \begin{aligned} \partial_s \phi_s(s, x, y) + L \phi_s(s, x, y) &= 0 \quad \text{on } [0, T] \times \mathcal{X}^2, \\ \nabla_y \phi_s(s, x, y) \cdot \bar{n} &= 0 \quad \text{on } [0, T] \times \mathcal{X} \times \partial \mathcal{X}, \\ \lim_{s \rightarrow t} \phi_s(s, x, y) &= \phi_s(x, y) \quad \text{on } \mathcal{X}^2. \end{aligned}$$

The weak time-space formulation of (3.3), is given by

$$(3.27) \quad \begin{aligned} \langle \xi_s(t), \phi_s(t, \cdot) \rangle - \langle \xi_s(0), \phi_s(0, \cdot) \rangle &= \int_0^t \int_{\mathcal{X}^2} (\mathcal{L}_y + \partial_s) \phi_s(s, x, y) \nu_s(t)(dx, dy) ds \\ &- \int_0^t \int_{\bar{\mathcal{X}}^2} \lambda(x, y) \phi_s(s, x, y) \nu_s(s)(dx, dy) ds + \mu_1 \int_0^t \int_{\bar{\mathcal{X}}^2} D(x, y) \phi_s(s, x, y) \nu_p(s)(dx) dy ds, \end{aligned}$$

and hence

$$(3.28) \quad |\langle \xi_s(t) - \bar{\xi}_s(t), \phi_s(x, y) \rangle| \leq \bar{\lambda} \int_0^t \sup_{\phi \leq 1} |\langle \xi_s(t) - \bar{\xi}_s(t), \phi \rangle| ds + \mu_1 \int_0^t \sup_{\phi \leq 1} |\langle \xi_p(t) - \bar{\xi}_p(t), \phi \rangle| ds,$$

where we used

$$\sup_{\phi \leq 1} |\langle \xi_p(t) - \bar{\xi}_p(t), P_s(t-s)\phi \rangle| \leq \sup_{\phi \leq 1} |\langle \xi_p(t) - \bar{\xi}_p(t), \phi \rangle|,$$

which is a consequence of the maximal principle applied to (3.26).

Taking first the sup on the LHS of each case, and then summation gives the inequality:

$$\sum_{\alpha \in \{p,s\}} \sup_{\phi \leq 1} |\langle \xi_\alpha(t) - \bar{\xi}_\alpha(t), \phi_\alpha \rangle| \leq C \left(\int_0^t \sum_{\alpha \in \{p,s\}} \sup_{\phi \leq 1} |\langle \xi_\alpha(s) - \bar{\xi}_\alpha(s), \phi_\alpha \rangle| ds \right),$$

and by Gronwall's lemma we conclude uniqueness.

Step 2: uniform estimates. Under the rescaling given by (1.12), a consequence of Proposition 3.3 is the following:

Proposition 3.4. *We have the following cádlág martingales:*

(1) *For the population of plants,*

$$(3.29) \quad M_t^{K, \phi_p} = \langle \nu_p^{(K)}(t), \phi_p \rangle_{\bar{\mathcal{X}}} - \langle \nu_p^{(K)}(0), \phi_p \rangle_{\bar{\mathcal{X}}} - \int_0^t \int_{\bar{\mathcal{X}}^2} \lambda(x, y) \phi_p(y) \nu_s^{(K)}(s)(dx, dy) ds$$

is a cádlág martingale with predictable quadratic variation

$$(3.30) \quad \langle M^{K, \phi_p} \rangle_t = \frac{1}{K} \int_0^t \int_{\bar{\mathcal{X}}^2} \lambda(x, y) \phi_p(y)^2 \nu_s^{(K)}(s)(dx, dz) ds.$$

(2) *For the population of seeds,*

$$(3.31) \quad \begin{aligned} M_t^{K, \phi_s} &= \langle \nu_s^{(K)}(t), \phi_s \rangle_{\bar{\mathcal{X}}^2} - \langle \nu_s^{(K)}(0), \phi_s \rangle_{\bar{\mathcal{X}}^2} - \int_0^t \langle \nu_s^{(K)}(t), \mathcal{L}_y \phi_s(x, y) \rangle_{\bar{\mathcal{X}}^2} ds \\ &\quad + \int_0^t \int_{\bar{\mathcal{X}}^2} \lambda(x, y) \phi_s(x, y) \nu_s^{(K)}(s)(dx, dy) ds \\ &\quad - \mu_1 \int_0^t \int_{\bar{\mathcal{X}}^2} D(x, y) \phi_s(x, y) \nu_p^{(K)}(s)(dx, dy) ds \end{aligned}$$

is a cádlág martingale with predictable quadratic variation

$$(3.32) \quad \begin{aligned} \langle M^{K, \phi_s} \rangle_t &= \frac{1}{K} \int_0^t \int_{\bar{\mathcal{X}}^2} \sigma^S(y)^2 |\nabla_y \phi_s(x, y)|^2 \nu_s^{(K)}(s)(dx, dy) ds + \frac{1}{K} \int_0^t \int_{\bar{\mathcal{X}}^2} \lambda(x, y) \phi_s(x, y)^2 \nu_s(s)(dx, dz) ds \\ &\quad + \frac{\mu_2}{K} \int_0^t \int_{\bar{\mathcal{X}}^2} D(x, y) \phi_s(x, y)^2 \nu_p^{(K)}(s)(dx, dy) ds. \end{aligned}$$

Notice that by (1.14), for $T > 0$, we have

$$(3.33) \quad \sup_{K \in \mathbb{N}} \mathbb{E} \left(\sup_{t \in [0, T]} \langle \nu_\alpha^{(K)}(t), 1 \rangle^3 \right) < \infty$$

for all $\alpha \in \{p, s\}$. This in turn implies the uniform estimate:

$$(3.34) \quad \sup_{K \in \mathbb{N}} \mathbb{E} \left(\sup_{t \in [0, T]} \langle \nu_\alpha^{(K)}(t), \phi_\alpha \rangle^3 \right) < \infty,$$

for ϕ_α bounded and measurable, and in particular $\phi_s \in D(L)$.

Step 3: tightness. Let us consider the spaces of finite measures $\mathcal{M}_F^p := \mathcal{M}_F(\mathcal{X})$ and $\mathcal{M}_F^s := \mathcal{M}_F(\mathcal{X}^2)$, both equipped with the vague topology. Here we show that the sequences of laws Q_α^K of the processes $\nu_\alpha^{(K)}$ are uniformly tight in $\mathbb{P}(\mathcal{D}([0, T], \mathcal{M}_F^\alpha))$, for $\alpha \in \{p, s\}$. We then use Theorem 2.1 from [12]. This means that we need to find dense sets D_α , dense in the space of continuous functions over \mathcal{X} , and \mathcal{X}^2 , respectively, such that the real-valued sequences $\langle \nu_\alpha^{(K)}, f \rangle$ are tight in $\mathbb{P}(\mathcal{D}([0, T], \mathbb{R}))$ for all f_α in D_α . For the sets D_α , we set them to be $C_b(\mathcal{X})$, and $D(\mathcal{L})$, respectively for $\alpha = p$ and $\alpha = s$, respectively. By Aldous and Rebolledo's criteria, it is enough to show:

$$(3.35) \quad \sup_{K \in \mathbb{N}} \mathbb{E} \left(\sup_{t \in [0, T]} |\langle \nu_\alpha(t)^{(K)}, f_\alpha \rangle| \right) < \infty,$$

and the tightness of the quadratic variation and the drift part of the martingales given in Proposition 3.4 above.

Notice that (3.35) is already given in (3.34). The tightness of quadratic variations and the drift part are done the same way. We then only show tightness for the quadratic variation part. Let $\delta > 0$, and consider stopping times τ, τ' such that

$$0 \leq \tau \leq \tau' \leq \tau + \delta \leq T.$$

By Doob's inequality and the estimate (3.33) we have

$$(3.36) \quad \mathbb{E} (\langle M_K^\alpha \rangle_\tau - \langle M_K^\alpha \rangle_{\tau'}) \leq C\delta.$$

Since similar estimates can be obtain for the drift part of the martingales, we conclude the proof of uniform tightness of the sequence Q_α^K .

Step 4: characterization of limit points. Now that we have shown tightness, we can assume convergence of a sub-sequence of the laws Q_α^K . Let us denote by Q_α the limiting law of this sub-sequence, that with an abuse of notation we denote again by Q_α^K . Let us denote by Λ_α a process with law Q_α . We need to verify that it is almost surely strongly continuous. We follow step 5, on page 54 of [8], with a small modification due to the branching mechanism. For $j \in \mathbb{N}$, let us introduce

$$(3.37) \quad \kappa_j = \inf \left\{ \kappa \in \mathbb{N} : \sum_{i=1}^{\kappa_j} p(i) \geq 1 - \frac{1}{2^j} \right\}.$$

Notice that by construction of our processes we have

$$(3.38) \quad \mathbb{P} \left(\sup_{t \in [0, T]} \sup_{f \leq 1} \left| \langle \nu_\alpha^{(K)}(t), f \rangle - \langle \nu_\alpha^{(K)}(t-), f \rangle \right| \leq \frac{\kappa_j}{K} \right) \geq 1 - \frac{1}{2^j}.$$

Similar arguments to those in [8], together with the continuity of the mapping $\nu \mapsto \sup_{t \in [0, T]} |\langle \nu(t), f \rangle - \langle \nu(t-), f \rangle|$ on path space, verify that indeed Λ_α is a.s. strongly continuous. The rest of the characterization is a standard use of the martingales given in (3.4), and follows the lines of [5, 8, 4].

4. PROPERTIES OF THE PDE SYSTEM

Let us now look a bit more at the properties of the PDE system satisfied by the population of seeds and plants, respectively g and f . To have a tractable analysis, we restrict our analysis to bounded domain \mathcal{X} and we will present our argument with $A(x) = Id$. All the arguments we use can be applied to a general elliptic matrix at the price of more complex notations. In this simplified situation, f and g satisfy

$$(4.1) \quad \partial_t f(t, y) = \int_{\mathcal{X}} \lambda(z, y) g(t, z, y) dz \quad \text{for all } t > 0, y \in \mathcal{X}$$

$$(4.2) \quad \partial_t g(t, x, y) = \Delta_y g(t, x, y) - \lambda(x, y) g(t, x, y) + \mu_1 D(x, y) f(t, x) \quad \text{for all } t > 0, (x, y) \in \mathcal{X}^2$$

$$(4.3) \quad \nabla_y g(t, x, y) \cdot \vec{n} = 0, \quad \text{for all } t > 0, x \in \mathcal{X}, y \in \partial \mathcal{X}$$

$$(4.4) \quad f(0, x) = f_0(x) \quad \text{for all } x \in \mathcal{X}$$

$$(4.5) \quad g(0, x, y) = 0 \quad \text{for all } (x, y) \in \mathcal{X}^2$$

In the first two subsections we prove that such problem has a non-trivial solution which is unique in a certain functional space. Then, we derive a reduced version of the model that may have some practical interest to exhibit some relevant parameters that explain exponential explosion of the solution.

4.1. Existence of a non trivial solution. From the equation satisfied by mature plants f (4.1) -(4.4), we have for $x \in \mathcal{X}$:

$$f(t, x) = f_0(x) + \int_0^t \int_{\mathcal{X}} \lambda(z, x) g(s, z, x) dz ds,$$

and we then have a single equation for the moving seed population for $(x, y) \in \mathcal{X}^2$, namely

$$\partial_t g(t, x, y) = \Delta_y g(t, x, y) - \lambda(x, y) g(t, x, y) + \mu_1 D(x, y) \left(f_0(x) + \int_0^t \int_{\mathcal{X}} \lambda(z, x) g(s, z, x) dz ds \right).$$

Observe that the above equation has some degeneracy in the variable x that makes its resolution non standard. Without the nonlocal term, such type of equation are usually called ultra parabolic equation in the literature [16].

A possible way to construct a solution, is to use a viscosity solution approach. Namely, let $\varepsilon > 0$, and introduce the approximation operator

$$\mathcal{L}_\varepsilon := \varepsilon \Delta_x + \Delta_y,$$

and the associated approximation problem:

$$(4.6) \quad \partial_t f_\varepsilon(t, x) = \int_{\mathcal{X}} \lambda(z, x) g_\varepsilon(t, z, x) dz \quad \text{for all } t > 0, x \in \mathcal{X},$$

$$(4.7) \quad \partial_t g_\varepsilon(t, x, y) = \mathcal{L}_\varepsilon [g_\varepsilon](t, x, y) - \lambda(x, y) g_\varepsilon(t, x, y) + \mu_1 D(x, y) f_\varepsilon(t, x) \quad \text{for all } t > 0, (x, y) \in \mathcal{X}^2,$$

$$(4.8) \quad \nabla_{x, y} g_\varepsilon(t, x, y) \cdot \vec{n} = 0, \quad \text{for all } t > 0, (x, y) \in \partial \mathcal{X}^2,$$

$$(4.9) \quad f(0, x) = f_0(x) \quad \text{for all } x \in \mathcal{X},$$

$$(4.10) \quad g(0, x, y) = 0 \quad \text{for all } (x, y) \in \mathcal{X}^2,$$

where \vec{n} is a normal vector of $\partial(\mathcal{X}^2)$, i.e. $\vec{n} = (\vec{n}', \vec{n})$ with \vec{n} a normal vector of $\partial \mathcal{X}$.

For any $\varepsilon > 0$, provide λ and D are smooth functions, we will show that the above system has indeed a solution.

Proposition 4.1. *Let \mathcal{X} be a smooth bounded domain, with at least a C^1 boundary. Assume further that $f_0 \in H^1(\mathcal{X})$, $f_0 \geq 0$, $\lambda, D \in W^{1, \infty}(\mathcal{X}^2)$ non negative functions. Then for all $T > 0$, there exists a couple of function $(f_\varepsilon, g_\varepsilon)$ satisfying $f_\varepsilon \in C^1((0, T), H^1(\mathcal{X}))$, $g_\varepsilon \in C^1((0, T), H^1(\mathcal{X}) \times H^1(\mathcal{X}))$ such that f_ε and g_ε are positive and $(f_\varepsilon, g_\varepsilon)$ is a solution to*

$$\begin{aligned} \partial_t f_\varepsilon(t, x) &= \int_{\mathcal{X}} \lambda(z, x) g_\varepsilon(t, z, x) dz && \text{for all } 0 < t < T, x \in \mathcal{X}, \\ \partial_t g_\varepsilon(t, x, y) &= \mathcal{L}_\varepsilon [g_\varepsilon](t, x, y) - \lambda(x, y) g_\varepsilon(t, x, y) + \mu_1 D(x, y) f_\varepsilon(t, x) && \text{for all } 0 < t < T, (x, y) \in \mathcal{X}^2, \\ \nabla_{x, y} g_\varepsilon(t, x, y) \cdot \vec{n} &= 0, && \text{for all } 0 < t < T, (x, y) \in \partial \mathcal{X}^2, \\ f_\varepsilon(0, x) &= f_0(x) && \text{for all } x \in \mathcal{X}, \\ g_\varepsilon(0, x, y) &= 0 && \text{for all } (x, y) \in \mathcal{X}^2. \end{aligned}$$

Having the existence of a positive solution of the regularised problem at hand, the trick is now to see whether we can extract a solution of the sequence $f_\varepsilon, g_\varepsilon$ as $\varepsilon \rightarrow 0$. To do so, we first establish a uniform estimate independent of ε .

Proposition 4.2. *Let \mathcal{X} be a smooth bounded domain, with at least a C^1 boundary. Assume further that $f_0 \in H^1(\mathcal{X})$, $f_0 \geq 0$, $\lambda, D \in W^{1, \infty}(\mathcal{X}^2)$ are non negatives functions. Let $f_\varepsilon, g_\varepsilon$ be positive functions satisfying $f_\varepsilon \in C^1((0, T), H^1(\mathcal{X}))$, $g_\varepsilon \in C^1((0, T), H^1(\mathcal{X}) \times H^1(\mathcal{X}))$ and such that $(f_\varepsilon, g_\varepsilon)$ is a solution to (4.6)-(4.10). Then f_ε and g_ε are increasing in time, and there exists C_0, C_1 independent of ε such that*

$$\|f_\varepsilon\|_2 \leq \|f_0\|_2 e^{C_0 t^2} \quad \text{and} \quad \|g_\varepsilon\|_2 \leq C_1 \|f_0\|_2 \int_0^t e^{C_0 \tau^2} d\tau.$$

Proof. For convenience, in the rest of the proof we drop the subscript ε of $f_\varepsilon, g_\varepsilon$ and consider f, g . Since f, g are positive, due to (4.6) the monotone increasing character of f is trivial. The monotone character of g is a consequence of the monotone character of f and the parabolic comparison principle. Indeed, for any real $h > 0$, let us define $w_h(t, x, y) := g(t + h, x, y) - g(t, x, y)$. By straightforward computation, since f is monotone increasing in time, we can check that for any $h > 0$, w_h satisfies for all $t > 0, (x, y) \in \mathcal{X}^2$:

$$\begin{aligned} \partial_t w_h(t, x, y) &= \mathcal{L}_\varepsilon[w_h](t, x, y) - \lambda(x, y)w_h(t, x, y) + \mu_1 D(x, y)(f(t + h, x) - f(t, x)) \geq \mathcal{L}_\varepsilon[w_h](t, x, y) - \lambda(x, y)w_h(t, x, y) \\ \nabla_{x, y} w_h(t, x, y) \cdot \vec{\mathbf{n}} &= 0, \\ w_h(0, x, y) &= g(h, x, y) \geq 0. \end{aligned}$$

As a consequence, by the Parabolic maximum principle, we get $w_h > 0$ for all t , meaning that $g(t + h, x, y) \geq g(t, x, y)$. The argument being true for any $h > 0$, this means that g is monotonic increasing in time.

Let us now obtain the estimate. First, let us multiply (4.7) by g and integrate over \mathcal{X}^2 . We then obtain,

$$\frac{1}{2} \frac{d}{dt} (\|g\|_2^2) = \int_{\mathcal{X}^2} g \Delta_y g + \varepsilon \int_{\mathcal{X}^2} g \Delta_x g - \int_{\mathcal{X}^2} \lambda(x, y) g^2 + \mu_1 \int_{\mathcal{X}^2} D(x, y) f(t, x) g(t, x, y)$$

Since g satisfies (4.8) and $\partial \mathcal{X}^2 = \partial \mathcal{X} \times \mathcal{X} \cup \mathcal{X} \times \partial \mathcal{X} \cup \partial \mathcal{X} \times \partial \mathcal{X}$, integrating by part the two first integral of the right hand side, we then get

$$\frac{1}{2} \frac{d}{dt} (\|g\|_2^2) = - \int_{\mathcal{X}^2} |\nabla_y g|^2 - \varepsilon \int_{\mathcal{X}^2} |\nabla_x g|^2 - \int_{\mathcal{X}^2} \lambda(x, y) g^2 + \mu_1 \int_{\mathcal{X}^2} D(x, y) f(t, x) g(t, x, y)$$

and thus, using Cauchy-Schwarz inequality,

$$\frac{1}{2} \frac{d}{dt} (\|g\|_2^2) \leq \mu_1 \int_{\mathcal{X}^2} D(x, y) f(t, x) g(t, x, y) \leq \mu_1 \|g\|_2 \left(\int_{\mathcal{X}} f^2(t, x) \left(\int_{\mathcal{X}} D^2(x, y) dy \right) dx \right)^{\frac{1}{2}} \leq \mu_1 \sqrt{\|\bar{D}\|_\infty} \|g\|_2 \|f\|_2,$$

By simplifying by $\|g\|_2$, and setting $C_1 := \mu_1 \sqrt{\|\bar{D}\|_\infty}$ we get,

$$\frac{d}{dt} (\|g\|_2) \leq C_1 \|f\|_2,$$

and therefore

$$(4.11) \quad \|g\|_2(t) \leq C_1 \int_0^t \|f\|_2(\tau) d\tau.$$

On the other hand, by multiplying by f the equation satisfied by f and integrating it over \mathcal{X} , we then get, using Cauchy-Schwarz and Jensen's inequality,

$$\frac{1}{2} \frac{d}{dt} \int_{\mathcal{X}} f^2(t, x) dx \leq \bar{\lambda} \|f\|_2 \|g\|_2.$$

Since $f > 0$ for all t , dividing the above inequality by $\|f\|_2^2$ and using the estimate (4.11) yield, after simplification,

$$\frac{d}{dt} (\ln(\|f\|_2)) \leq \bar{\lambda} \mu_1 \sqrt{\|\bar{D}\|_\infty} \int_0^t \frac{\|f\|_2(\tau)}{\|f\|_2(t)} d\tau.$$

Since f is monotone increasing and positive for all $t > \tau > 0$, $\frac{\|f\|_2(\tau)}{\|f\|_2(t)} \leq 1$ and we get

$$\frac{d}{dt} (\ln(\|f\|_2)) \leq \bar{\lambda} \mu_1 \sqrt{\|\bar{D}\|_\infty} t,$$

which then implies that

$$(4.12) \quad \|f\|_2 \leq \|f_0\|_2 e^{C_0 t^2},$$

with $C_0 := \frac{1}{2}\bar{\lambda}C_1$. The estimates on g is a straightforward consequence of (4.11) that is

$$(4.13) \quad \|g\|_2 \leq C_1 \|f_0\|_2 \int_0^t e^{C_0\tau^2} d\tau.$$

□

For any sequence $\varepsilon_n \rightarrow 0$, the estimate is unfortunately insufficient to make sequences $(f_{\varepsilon_n}, g_{\varepsilon_n})_{n \in \mathbb{N}}$ pre-compact in a reasonable functional space. To gain some compactness, provided some extra assumptions on λ and D , we obtain H^1 bounds independent of ε . Namely, we have

Proposition 4.3. *Let \mathcal{X} be a smooth bounded domain, with at least a C^1 boundary. Assume further that $f_0 \in H^1(\mathcal{X})$, $f_0 \geq 0$, $\lambda, D \in W^{1,\infty}(\mathcal{X}^2)$ are non negative functions. Let $f_\varepsilon, g_\varepsilon$ be positive functions satisfying $f_\varepsilon \in C^1((0, T), H^1(\mathcal{X}))$, $g_\varepsilon \in C^1((0, T), H^1(\mathcal{X}) \times H^1(\mathcal{X}))$ and such that $(f_\varepsilon, g_\varepsilon)$ is a solution to (4.6)–(4.10).*

Then there exists positive constants $C_0, C_1, C_2, C_3, C_4, C_5$ independent of ε such that

$$\|f_\varepsilon\|_{H^1} \leq \|\nabla f_0\|_2 + \|f_0\|_2 \left(e^{C_0 t^2} + C_0 \int_0^t \int_0^\tau e^{C_0 s^2} ds d\tau + C_1 \int_0^t \int_0^\tau \int_0^s e^{C_0 \sigma^2} d\sigma ds d\tau \right).$$

and

$$\begin{aligned} \|g_\varepsilon\|_{H^1} \leq C_1 \|\nabla f_0\|_2 + \|f_0\|_2 & \left(C_2 \int_0^t e^{C_0 \tau^2} d\tau + C_3 \int_0^t \int_0^\tau e^{C_0 s^2} ds d\tau \right. \\ & \left. + C_4 \int_0^t \int_0^\tau \int_0^s e^{C_0 \sigma^2} d\sigma ds d\tau + C_5 \int_0^t \int_0^\tau \int_0^s \int_0^\sigma e^{C_0 \omega^2} d\omega d\sigma ds d\tau \right). \end{aligned}$$

Moreover, for all $T > 0$, along any sequence $\varepsilon_n \rightarrow 0$ there exists a subsequence $(\varepsilon_{n_k})_{k \in \mathbb{N}}$ such that

$$f_{\varepsilon_{n_k}} \geq f_0 \quad \text{and} \quad \|g_{\varepsilon_{n_k}}\|_2(t) \geq \frac{1}{2} \|g_1\|_2(t) \quad \text{for} \quad 0 < t < T,$$

where g_1 is the unique solution of the ultra-parabolic equation

$$\begin{aligned} \partial_t g_1(t, x, y) &= \Delta_y g_1(t, x, y) - \lambda(x, y) g_1(t, x, y) + \mu_1 D(x, y) f_0(x) && \text{for all } t > 0, (x, y) \in \mathcal{X}^2, \\ \nabla_y g_1(t, x, y) \cdot \vec{n} &= 0, && \text{for all } t > 0, x \in \mathcal{X}, y \in \partial\mathcal{X}, \\ g_1(0, x, y) &= 0 && \text{for all } (x, y) \in \mathcal{X}^2. \end{aligned}$$

The proof of this proposition is rather standard knowing the estimate of Proposition 4.2 and can be founded in the appendix.

From these H^1 estimates, for all $T > 0$, we can check that g_ε is uniformly bounded in $H^1((0, T), H^1(\mathcal{X}^2))$ independently of ε . As well, we get that f_ε is uniformly bounded in $H^1((0, T), H^1(\mathcal{X}))$ independently of ε . Therefore, along any sequence $\varepsilon_n \rightarrow 0$, by a diagonal extraction process, we can extract a convergent subsequence $(f_n, g_n)_{n \in \mathbb{N}}$ such that $f_n, g_n \rightarrow f^*, g^*$ in $C((0, T), L^2(\mathcal{X}))$, $C((0, T), L^2(\mathcal{X}^2))$ and such that (f^*, g^*) satisfies (4.1) – (4.5) in the sense of distribution. Up to extraction of a subsequences, we also have

$$f^* \geq f_0, \quad \text{and} \quad \|g^*\|_2(t) \geq \frac{1}{2} \|g_1\|_2(t), \quad \text{for } t > 0.$$

By using the partial regularity of the singular system (4.1) – (4.5), using Sobolev embedding and bootstrap arguments, we conclude that $f^* \in C^1((0, T), H^1(\mathcal{X}))$, $g^* \in C^1((0, T), H^1(\mathcal{X}^2))$.

To complete the construction, we need to prove the singular system (4.1) – (4.5) have some uniqueness of the solution.

Uniqueness of solutions. Here we show that the solution obtain is unique in H^1 . More precisely, we show

Proposition 4.4. *Let \mathcal{X} be a smooth bounded domain, with at least a C^1 boundary. Assume further that $\lambda, D \in W^{1,\infty}(\mathcal{X}^2)$ and λ, D are non negative functions. Then the system (4.1)–(4.5) has a unique solution f, g such that $f \in C^1((0, T), H^1(\mathcal{X}))$, $g \in C^1((0, T), H^1(\mathcal{X}) \times H^1(\mathcal{X}))$.*

Proof. Let us assume that there exists (f_1, g_1) and (f_2, g_2) belonging to $C^1(\mathbb{R}^+, H^1(X)) \times C^1(\mathbb{R}^+, H^1(\mathcal{X}^2))$ that are two solution of (4.1) – (4.5). Thanks to the regularity of λ and D , the functions f_i, g_i are continuous in $x \in \mathcal{X}, (x, y) \in \mathcal{X}^2$ for all $t > 0$. Let us define $w := g_1 - g_2$, then thanks to (4.2), we can check that w satisfies

$$\begin{aligned}\partial_t w(t, x, y) &= \Delta_y w(t, x, y) - \lambda(x, y)w(t, x, y) + \mu_1 D(x, y) \int_0^t \int_{\mathcal{X}} \lambda(z, x)w(s, z, x) dz \\ \nabla_y w(t, x, y) \cdot \vec{n} &= 0, \\ w(0, x, y) &= 0.\end{aligned}$$

For any $\gamma \in \mathbb{R}$, let us consider $\tilde{w} := e^{\gamma t} w$. A trivial computation shows that \tilde{w} satisfies

$$\begin{aligned}\partial_t \tilde{w}(t, x, y) &= \Delta_y \tilde{w}(t, x, y) - (\lambda(x, y) - \gamma)\tilde{w}(t, x, y) + \mu_1 D(x, y) \int_0^t e^{\gamma(t-s)} \int_{\mathcal{X}} \lambda(z, x)\tilde{w}(s, z, x) dz \\ \nabla_y \tilde{w}(t, x, y) \cdot \vec{n} &= 0, \\ \tilde{w}(0, x, y) &= 0.\end{aligned}$$

Take now $\varepsilon > 0$ and consider the function $h_\varepsilon(t, x, y) := \tilde{w}(t, x, y) + \varepsilon$. At $t = 0$, $h_\varepsilon(0, x, y) = \varepsilon > 0$ and by continuity, h_ε is positive for later times, says for $t \leq t_\varepsilon$. We claim that

Claim 4.5. $h_\varepsilon(t, x, y) > 0$ for all $t > 0, (x, y) \in \mathcal{X}^2$.

Assume for the moment that the claim holds true, we are then done. Indeed, from the above claim we deduce that $\tilde{w}(t, x, y) \geq -\varepsilon$ for all $t > 0, (x, y) \in \mathcal{X}^2$ and since ε is arbitrary chosen this implies $\tilde{w}(t, x, y) \geq 0$ for all $t > 0, (x, y) \in \mathcal{X}^2$. By linearity this argument holds true for $-\tilde{w}$ and thus $\tilde{w} \equiv 0$ for all $t > 0, (x, y) \in \mathcal{X}^2$ showing thus the uniqueness of the solution. \square

Let us now prove the claim.

Proof. The result will follow from a form of maximum principle. First, a quick computation shows that h_ε satisfies

$$\begin{aligned}\partial_t h_\varepsilon(t, x, y) &= \Delta_y h_\varepsilon(t, x, y) - (\lambda(x, y) - \gamma)h_\varepsilon(t, x, y) + \mu_1 D(x, y) \int_0^t e^{\gamma(t-s)} \int_{\mathcal{X}} \lambda(z, x)h_\varepsilon(s, z, x) dz \\ &\quad + \varepsilon(\lambda(x, y) - \gamma) - \varepsilon D(x, y)\bar{\lambda}(x) \int_0^t e^{\gamma(t-s)} ds.\end{aligned}$$

By taking $\gamma \ll -1$ so that $\gamma^2 \geq \|D\|_\infty \|\bar{\lambda}\|_\infty$ we get,

$$\begin{aligned}\partial_t h_\varepsilon(t, x, y) &\geq \Delta_y h_\varepsilon(t, x, y) - (\lambda(x, y) - \gamma)h_\varepsilon(t, x, y) + \mu_1 D(x, y) \int_0^t e^{\gamma(t-s)} \int_{\mathcal{X}} \lambda(z, x)h_\varepsilon(s, z, x) dz \\ \nabla_y h_\varepsilon(t, x, y) \cdot \vec{n} &= 0, \\ h_\varepsilon(0, x, y) &= \varepsilon.\end{aligned}$$

Recall that h_ε is a smooth function in t and continuous in x, y and that $h_\varepsilon(0, x, y) = \varepsilon > 0$. So by continuity there exists $t_\varepsilon > 0$ such that $h_\varepsilon > 0$ for all $t \in (0, t_\varepsilon)$ and $(x, y) \in \mathcal{X}^2$. Let us define

$$t^* := \sup\{t > 0, |h(s, x, y)| > 0, \forall s \leq t, (x, y) \in \mathcal{X}^2\}.$$

Now, if $t^* = +\infty$ we are done. So assume by contradiction that $t^* < +\infty$. By definition of t^* , since μ_1, D, λ and h_ε are non negative quantities we have for all $t \leq t^*$,

$$\begin{aligned}\partial_t h_\varepsilon(t, x, y) &\geq \Delta_y h_\varepsilon(t, x, y) - (\lambda(x, y) - \gamma)h_\varepsilon(t, x, y) \geq \Delta_y h_\varepsilon(t, x, y) - \sigma h_\varepsilon(t, x, y) \\ \nabla_y h_\varepsilon(t, x, y) \cdot \vec{n} &= 0, \\ h_\varepsilon(0, x, y) &= \varepsilon.\end{aligned}$$

where $\sigma := |\lambda|_\infty + |\gamma|$. To obtain a contradiction, our aim is to show that $h_\varepsilon(t^*, x, y) > 0$ for all $(x, y) \in \bar{\mathcal{X}}^2$. To do so, let us denote by $\omega(t, x, y) := \varepsilon e^{-\sigma t} - h_\varepsilon(t, x, y)$ then a quick computation shows that

$$\begin{aligned}
\partial_t \omega(t, x, y) &\leq \Delta_y \omega(t, x, y) - \sigma \omega(t, x, y) && \text{for all } t \in (0, t^*), x \in \mathcal{X}, y \in \mathcal{X} \\
\nabla_y \omega(t, x, y) \cdot \vec{n} &= 0 && \text{for all } t \in (0, t^*), x \in \mathcal{X}, y \in \partial \mathcal{X} \\
\omega(0, x, y) &= 0
\end{aligned}$$

Let us now multiply by ω^+ the above inequality and integrate over \mathcal{X}^2 . After a straightforward computation we get,

$$\frac{1}{2} \frac{d}{dt} \left(\int_{\mathcal{X}^2} (\omega^+)^2(t, x, y) dx dy \right) \leq - \int_{\mathcal{X}^2} |\nabla \omega^+|^2(t, x, y) dy dx - \sigma \int_{\mathcal{X}^2} (\omega^+)^2(t, x, y) \leq 0.$$

Hence, for all $t \leq t^*$, we have

$$0 \leq \int_{\mathcal{X}^2} (\omega^+)^2(t, x, y) dx dy \leq \int_{\mathcal{X}^2} (\omega^+)^2(0, x, y) dx dy = 0.$$

Therefore we achieve for all $t \leq t^*$, $\omega^+(t, x, y) = 0$ for almost every $(x, y) \in \mathcal{X}^2$ meaning in particular that

$$h_\varepsilon(t^*, x, y) \geq \varepsilon e^{-\sigma t^*} > 0 \quad \text{almost everywhere.}$$

From the continuity of h_ε , it then follows that the inequality holds true for all $(x, y) \in \mathcal{X}^2$ and thus

$$h_\varepsilon(t^*, x, y) \geq \varepsilon e^{-\sigma t^*} > 0 \quad \text{for all } (x, y) \in \mathcal{X}^2.$$

From the later by using the continuity of h_ε with respect to time and space, we can then find $t' > t^*$ such that for all $t \leq t'$, $h_\varepsilon(t, x, y) > 0$ contradicting the definition of t^* . □

4.2. A reduced model. From a population dynamics point of view, a pertinent quantity to look at, is

$$\bar{g}(t, y) := \int_{\mathcal{X}} g(t, x, y) dx$$

representing the population of all seeds at the position y . A quick computation shows that

$$\partial_t f(t, x) = \int_{\mathcal{X}} \lambda(z, x) g(t, z, x) dz \quad \text{for all } t > 0, x \in \mathcal{X},$$

$$\partial_t \bar{g}(t, y) = \Delta_y \bar{g}(t, y) - \int_{\mathcal{X}} \lambda(x, y) g(t, x, y) dx + \mu_1 \int_{\mathcal{X}} D(x, y) f(t, x) dx \quad \text{for all } t > 0, y \in \mathcal{X},$$

or equivalently

$$\partial_t \bar{g}(t, y) = \Delta_y \bar{g}(t, y) - \int_{\mathcal{X}} \lambda(x, y) g(t, x, y) dx + \mu_1 \int_{\mathcal{X}} D(x, y) \left(f_0(x) + \int_0^t \int_{\mathcal{X}} \lambda(z, x) g(s, z, x) dz ds \right) dx,$$

which for seed's maturity rates $\lambda(x, y)$ only dependent of the deposit position, i.e. $\lambda(x, y) = \lambda(y)$, reduces to

$$\partial_t f(t, x) = \lambda(x) \bar{g}(t, x) \quad \text{for all } t > 0, x \in \mathcal{X}$$

$$\partial_t \bar{g}(t, y) = \Delta_y \bar{g}(t, y) - \lambda(y) \bar{g}(t, y) + \mu_1 \int_{\mathcal{X}} D(x, y) f(t, x) dx \quad \text{for all } t > 0, y \in \mathcal{X}.$$

This can also be rewritten as

$$\partial_t \bar{g}(t, y) = \Delta_y \bar{g}(t, y) - \lambda(y) \bar{g}(t, y) + \mu_1 \int_0^t \int_{\mathcal{X}} D(x, y) \lambda(x) \bar{g}(s, x) dx ds + F_0(y) \quad \text{for all } t > 0, y \in \mathcal{X},$$

where

$$F_0(y) := \mu_1 \int_{\mathcal{X}} D(x, y) f_0(x) dx.$$

By using that D is a probability density, and since x and y are interchangeable variable we may write the above equation as

$$\partial_t \bar{g}(t, y) = \Delta_y \bar{g}(t, y) + \lambda(y) \left(\mu_1 \int_0^t \bar{g}(s, y) ds - \bar{g}(t, y) \right) + \mu_1 \int_0^t \int_{\mathcal{X}} D(x, y) [\lambda(x) \bar{g}(s, x) - \lambda(y) \bar{g}(s, y)] dx ds + F_0(y).$$

Observe that a first natural time t_0 appears above which the population of moving seed will increase exponentially fast, t_0 being the times for which the "cumulative production rate of seed" by plant $\int_0^t \frac{\bar{g}(s, y)}{\bar{g}(t, y)} ds$ exceeds $\frac{1}{\mu_1}$.

APPENDIX A. RIGOROUS DEFINITION OF THE MODEL

A.1. Path-wise construction. We want to rigorously define a process $\{\nu_t : t \geq 0\}$ with generator \mathcal{L} given by (1.4). In order to do so, we consider the process $\{\nu_t : t \geq 0\}$ as an element of the path-space of right-continuous process $\mathcal{D}([0, \infty), \mathcal{M}_F)$, taking values in $M_F := \mathcal{M}_F(\bar{\mathcal{X}}) \times \mathcal{M}_F(\bar{\mathcal{X}}^2)$. In the vein of [5], we will construct this process in terms of Poisson measures. Hence, we introduce the following probabilistic objects:

Definition A.1. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a sufficiently large probability space. On this probability space we consider the following independent random elements:

- i):** An \mathcal{M}_F -valued random variable ν_0 of the form $(\nu_p(0), \nu_s(0))$, i.e., the initial distribution of plants and seeds.
- ii):** A Poisson random measures $Q_{dep}(ds, di, d\theta)$ on $[0, \infty) \times \mathbb{N} \times \mathbb{R}_+$, with intensity measure:

$$l(ds) \otimes \left(\sum_{j \geq 1} \delta_j(di) \right) \otimes l(d\theta),$$

where l denotes the Lebesgue measure on \mathbb{R}_+ .

- iii):** A Poisson random measure $Q_{dis}(ds, dk, dy, di, d\theta)$ on $[0, \infty) \times \mathbb{N} \times \bar{\mathcal{X}} \times \mathbb{N} \times \mathbb{R}_+$, with intensity measure:

$$l(ds) \otimes \left(\sum_{j \geq 1} p(j) \delta_j(dk) \right) \otimes \bar{D}(dy) \otimes \left(\sum_{j \geq 1} \delta_j(di) \right) \otimes d\theta,$$

where p and \bar{D} are given as in Assumption 1.

Moreover, we enlarge the original probability space $(\Omega, \mathcal{F}, \mathbb{P})$ to the filtered probability space $(\Omega, \mathcal{F}, \mathcal{F}_t, \mathbb{P})$, where $(\mathcal{F}_t)_{t \geq 0}$ is the canonical filtration generated by these processes.

We then introduce the following measures which are useful representing the main events as follows:

- Seeds maturing and becoming plants

$$\nu_{dep}(t) = \int_{[0, t] \times \mathbb{N}^* \times \mathbb{R}_+} (\delta_{y_i(s-)}, -\delta_{x_i(s-), y_i(s-)} \mathbb{1}_{\{i \leq N_s(s-)\}} \mathbb{1}_{\{\theta \leq \lambda(y_i(s-), z_i(s-))\}}) Q_{dep}(ds, di, d\theta).$$

- Plants dispersing new groups of seeds

$$\nu_{dis}(t) = \sum_k p(k) \int_{[0, t] \times \mathcal{X} \times \mathbb{N} \times \mathbb{R}_+} (0, k \delta_{x_i(s-), y}) \mathbb{1}_{\{i \leq N_p(s-)\}} \mathbb{1}_{\{\theta \leq D(x_i(s-), y)\}} Q_{dis}(ds, dy, di, d\theta).$$

We are now able to define our process as a solution of the following integral equation:

Definition A.2. An \mathcal{F}_t -adapted stochastic process $(\nu_t)_{t \geq 0} = (\nu_p(t), \nu_s(t) : t \geq 0)$ is called measure-valued GDM process, if we have that almost surely:

$$\langle \phi, \nu(t) \rangle = \langle \phi, \nu(0) \rangle + \langle \phi, \nu_{dep}(t) \rangle + \langle \phi, \nu_{dis}(t) \rangle + \int_0^t (\langle L\phi_{sp}, \nu(s) \rangle + \langle L\phi_s, \nu_s(s) \rangle) ds$$

(A.1)

$$+ \int_0^t \sum_{i=1}^{N_p(s-)} \sum_{j=1}^{N_s(s-)} \sigma(z_j(s-)) \nabla_z \phi_{s,p}(x_i(s), y_j(s-), z_j(s-)) \cdot dW_s^i + \int_0^t \sum_{j=1}^{N_s(s-)} \sigma(z_j(s-)) \nabla_z \phi_s(y_i(s-), z_i(s-)) dx \cdot dW_s^i,$$

for all $t \geq 0$, and all $\phi = \{\phi_p, \phi_s\}$.

The following proposition shows that the process given by (A.1) coincides with the process with infinitesimal generator \mathcal{L} .

Proposition A.3. Let $(\boldsymbol{\nu}_t)_{t \geq 0} = (\nu_p(t), \nu_s(t) : t \geq 0)$ be a solution of (A.1) such that for all $T > 0$

$$(A.2) \quad \mathbb{E} \left(\sup_{t \in [0, T]} (\langle 1, \nu_p(t) \rangle + \langle 1, \nu_s(t) \rangle)^2 \right) < \infty.$$

Then, under Assumption 1, the process $(\boldsymbol{\nu}_t)_{t \geq 0}$ is Markov with infinitesimal generator \mathcal{L} given by (1.4).

Proof. The proof of this proposition is standard. We refer the reader to [5] or [1] for a proof in similar contexts. \square

It remains to show that starting from controlled initial conditions at time zero, we can extend this control for later times.

Proposition A.4. Let $\boldsymbol{\nu}_0 = (\nu_p(0), \nu_s(0))$ be such that for some $p \geq 1$ we have

$$(A.3) \quad \mathbb{E} ((\langle 1, \nu_p(0) \rangle + \langle 1, \nu_s(0) \rangle)^p) < \infty.$$

Assume that the counting distribution has finite p -th moment:

$$(A.4) \quad \mu_p := \sum_{\kappa} p(\kappa) \kappa^p < \infty.$$

Then, under Assumption 1, the process $(\boldsymbol{\nu}_t)_{t \geq 0} = (\nu_p(t), \nu_s(t) : t \geq 0)$ satisfies

$$(A.5) \quad \mathbb{E} \left(\sup_{t \in [0, T]} (\langle 1, \nu_p(t) \rangle + \langle 1, \nu_s(t) \rangle)^p \right) < \infty.$$

Proof. Let us use a stopping time argument to show this proposition. Define the following sequence:

$$(A.6) \quad \tau_n = \inf\{t \geq 0 : \langle 1, \nu_p(t) \rangle + \langle 1, \nu_s(t) \rangle \geq n\}.$$

Let us also consider F_p given by the choice:

$$(A.7) \quad F_p(\boldsymbol{\nu}) = \langle 1, \nu_p \rangle^p + \langle 1, \nu_s \rangle^p.$$

Hence, by (A.1) we have:

$$(A.8) \quad \sup_{s \in [0, t \wedge \tau_n]} \langle 1, \nu_p(s) \rangle^p + \langle 1, \nu_s(s) \rangle^p \leq \langle 1, \nu_p(0) \rangle^p + \langle 1, \nu_s(0) \rangle^p + \sum_k p(k) \int_0^{t \wedge \tau_n} \int_{\mathcal{X} \times \mathbb{N} \times \mathbb{R}_+} [(\langle 1, \nu_s(s-) \rangle + k)^p - \langle 1, \nu_s(s-) \rangle^p] \mathbf{1}_{\{i \leq N_p(s-)\}} \mathbf{1}_{\{\theta \leq D(x_i(s-), y)\}} Q_{dis}(ds, dy, di, d\theta),$$

where the diffusion terms vanished due to the presences of derivatives of the constant function 1, and we have dropped the integral term related to maturation since it has a negative contribution.

Taking expectation we obtain

$$(A.9) \quad \mathbb{E} \left[\sup_{s \in [0, t \wedge \tau_n]} \langle 1, \nu_p(s) \rangle^p + \langle 1, \nu_s(s) \rangle^p \right] \leq \mathbb{E} [\langle 1, \nu_p(0) \rangle^p + \langle 1, \nu_s(0) \rangle^p] + C \mathbb{E} \left[\int_0^{t \wedge \tau_n} N_p(s-) \sum_k p(k) [(\langle 1, \nu_s(s-) \rangle + k)^p - \langle 1, \nu_s(s-) \rangle^p] ds \right],$$

where the constant C incorporates the bounds given by Assumption 1.

Using (A.4) and the simple inequality $(x+1)^p - x^p \leq C_p(1+x^{p-1})$ we obtain

$$(A.10) \quad \mathbb{E} \left[\sup_{s \in [0, t \wedge \tau_n]} \langle 1, \nu_p(s) \rangle^p + \langle 1, \nu_s(s) \rangle^p \right] \leq \mathbb{E} [\langle 1, \nu_p(0) \rangle^p + \langle 1, \nu_s(0) \rangle^p] + C \mathbb{E} \left[\int_0^t N_p(s \wedge \tau_n) [1 + \langle 1, \nu_s(s \wedge \tau_n) \rangle^{p-1}] ds \right],$$

where the constant C has changed its value again. Moreover, using the equality

$$(A.11) \quad N_p(s) = \langle 1, \nu_p(s) \rangle,$$

we obtain

$$(A.12) \quad \mathbb{E} \left[\sup_{s \in [0, t \wedge \tau_n]} \langle 1, \nu_p(s) \rangle^p + \langle 1, \nu_s(s) \rangle^p \right] \leq C_p \left(1 + \mathbb{E} \left[\int_0^t \langle 1, \nu_p(s) \rangle^p + \langle 1, \nu_s(s) \rangle^p ds \right] \right),$$

where again the constant C_p changed its value incorporating new constants.

By Gronwalls inequality we then have

$$(A.13) \quad \mathbb{E} \left[\sup_{s \in [0, t \wedge \tau_n]} \langle 1, \nu_p(s) \rangle^p + \langle 1, \nu_s(s) \rangle^p \right] \leq C_p e^{Bt},$$

for some constant B independent of n . From (A.13) we can deduce that τ_n goes to a.s. to infinity as $n \rightarrow \infty$. We can then apply Fatou's lemma to conclude. \square

For existence the well-definedness of the process ν_t one has to construct the process step by step, where the time steps are given by a sequence of jump instants T_n exponentially distributed with law

$$(A.14) \quad R(\nu_{n-1})e^{-R(\nu_{n-1})t},$$

and where the total jump rate $R(\nu)$ is bounded by Assumption 1. It is then enough to check that the sequence T_n goes to infinity almost surely. This follows from

$$\mathbb{E} \left(\sup_{t \leq T} \sum_{\alpha \in \mathcal{X}} \langle \nu_\alpha(t), 1 \rangle_{\mathcal{D}_\alpha}^p \right) < \infty,$$

which is a consequence of (A.5) when $p = 1$. However, for the martingale characterization of $\{\nu(t)\}_{t \geq 0}$ higher values of p are needed.

APPENDIX B. ESTIMATE ON THE PDE SYSTEM

In this section, we prove Proposition 4.1 and establish some estimates on the solution to the regularised problem. Let us start with the proof of Proposition 4.1.

Proof. Let us introduce the following sequence of functions $(f_n, g_n)_{n \in \mathbb{N}}$, where $(f_0(t, x), g_0(t, x)) = (f_0(x), 0)$, and for all $n > 0$, f_n and g_n are respectively solution of the following equations:

$$(B.1) \quad \partial_t f_n(t, x) = \int_{\mathcal{X}} \lambda(z, x) g_n(t, z, x) dz \quad \text{for all } t > 0, x \in \mathcal{X}$$

$$(B.2) \quad \partial_t g_n(t, x, y) = \mathcal{L}_\varepsilon[g_n](t, x, y) - \lambda(x, y) g_n(t, x, y) + \mu_1 D(x, y) f_{n-1}(t, x) \quad \text{for all } t > 0, (x, y) \in \mathcal{X}^2$$

$$(B.3) \quad \nabla_{x,u} g_n(t, x, y) \cdot \vec{n} = 0, \quad \text{for all } t > 0, (x, y) \in \partial \mathcal{X}^2$$

$$(B.4) \quad f_n(0, x) = f_0(x) \quad \text{for all } x \in \mathcal{X}$$

$$(B.5) \quad g_n(0, x, y) = 0 \quad \text{for all } (x, y) \in \mathcal{X}^2.$$

Since, for all $n > 0$, g_n satisfies a parabolic equation, the sequence (f_n, g_n) is then well defined for all t as soon as for all n , we show that $f_n(t, x) \in L^2(\mathcal{X})$ for all $t > 0$. We claim that

Claim B.1. For all $n \geq 0$ and $t > 0$, $f_n(t, x) \in L^2(\mathcal{X})$.

Proof. To prove such claim we will use a induction argument. Since by definition $f_0(t, x) = f_0(x)$ for all t the results is trivial for $n = 0$. Let us assume that for some $n \geq 0$, for all $t > 0$, $f_n(t, \cdot) \in L^2(\mathcal{X})$ and $f_n \geq 0$ let us prove that

for all $t > 0$, $f_{n+1}(t, \cdot) \in L^2(\mathcal{X})$ and $f_{n+1}(t, x) \geq 0$. Observe that since $f_n \in L^2(\mathcal{X})$ for all $t > 0$, and $\bar{D}(x) \in L^\infty(\mathcal{X})$ we have

$$\int_{\mathcal{X}^2} F_n^2(t, x, y) dx dy = \int_{\mathcal{X}^2} \mu_1^2 D^2(x, y) f_n^2(x) dy dx = \mu_1^2 \int_{\mathcal{X}} \left(f_n^2(t, x) \int_{\mathcal{X}} D^2(x, y) dy \right) dx \leq C \int_{\mathcal{X}} f_n^2(x) dx,$$

and therefore g_{n+1} satisfies a parabolic equation with a right hand side $F_n(t, x, y) \in L^2(\mathcal{X}^2)$. From standard parabolic theory [6, 9, 16], since \mathcal{X} is smooth, g_{n+1} is well defined for all $t > 0$ and $g_{n+1} \in C^1(\mathbb{R}^+, H^1(\mathcal{X}^2))$. Now observe that since $g_n \in C^1(\mathbb{R}^+, H^1(\mathcal{X}^2))$ and $\lambda \in L^\infty(\mathcal{X}^2)$, the function $h_{n+1}(t, x) := \int_{\mathcal{X}} \lambda(x, z) g_{n+1}(t, z, x) dz$ belongs to $C^1(\mathbb{R}^+, L^1(\mathcal{X}))$ and from the equation satisfied by $f_{n+1}(t, x)$, we straightforwardly get that f_{n+1} is well defined for all times $t > 0$ and $f_{n+1} \in C^1(\mathbb{R}^+, L^1(\mathcal{X}))$. Moreover, we have

$$(B.6) \quad f_{n+1}(t, x) = f_0(x) + \int_0^t \int_{\mathcal{X}} \lambda(z, x) g_{n+1}(s, z, x) dz ds.$$

By Young's and Jensen inequalities, we get that

$$f_{n+1}^2(t, x) \leq C \left(f_0^2(x) + \bar{\lambda} \int_0^t \int_{\mathcal{X}} g_{n+1}^2(s, z, x) ds dz \right),$$

and therefore $f_{n+1}(t, \cdot) \in L^2(\mathcal{X})$ for all $t > 0$. Note that as a side result of the claim we get that for all $n \geq 0$ ($f_n(t, x), g_n(t, x, y)$) is well defined for all $t > 0$, $x \in \mathcal{X}, y \in \mathcal{X}$ and moreover, for all n , $g_n \in C^1(\mathbb{R}^+, H^1(\mathcal{X}^2))$ and $f_n - f_0 \in C^2(\mathbb{R}^+, H^1(\mathcal{X}))$. \square

Now, to conclude to the existence of a solution, we need to check that $(f_n(t, x), g_n(t, x))_{n \in \mathbb{N}}$ converges in a certain sense. Before that, let us show that the sequences f_n and g_n are monotone increasing

Claim B.2. For all $n > 0$ and $t > 0$, $f_{n+1}(t, x) \geq f_n$ and $g_{n+1}(t, x, y) \geq g_n(t, x, y)$ for all $x, y \in \mathcal{X}$.

Proof. Observe by the relation (B.6) we get the monotone behaviour of the sequence $(f_n)_{n \in \mathbb{N}}$ directly from the one of $(g_n)_{n \in \mathbb{N}^*}$.

To obtain the behaviour of the sequences f_n, g_n , we will argue by induction. But before let us observe that thanks to (B.6) we have $f_1 \geq f_0$ if we can prove that $g_1 \geq 0$. The positivity of g_1 , is a straightforward consequence of the parabolic maximum principle. Indeed, since, f_0, D, λ and μ_1 are positive quantities, by definition of g_1 , it satisfies for all $t > 0$, $(x, y) \in \mathcal{X}^2$:

$$\begin{aligned} \partial_t g_1(t, x, y) &= \mathcal{L}_\varepsilon[g_1](t, x, y) - \lambda(x, y)g_1(t, x, y) + \mu_1 D(x, y)f_0(x) \geq \mathcal{L}_\varepsilon[g_1](t, x, y) - \lambda(x, y)g_1(t, x, y), \\ \nabla_{x, y} g_1(t, x, y) \cdot \vec{n} &= 0, \\ g_1(0, x, y) &= 0, \end{aligned}$$

which from the parabolic maximum principle entails that $g_1(t, x, y) \geq 0$ for all $t \geq 0$ and $(x, y) \in \mathcal{X}^2$.

As a consequence $f_1 \geq f_0$, we can check that $w = g_2 - g_1$ satisfies for all $t > 0$, $(x, y) \in \mathcal{X}^2$:

$$\begin{aligned} \partial_t w(t, x, y) &= \mathcal{L}_\varepsilon[w](t, x, y) - \lambda(x, y)w + \mu_1 D(x, y)(f_1(t, x) - f_0(x)) \geq \mathcal{L}_\varepsilon[w](t, x, y) - \lambda(x, y)w(t, x, y), \\ \nabla_{x, y} w(t, x, y) \cdot \vec{n} &= 0, \\ w(0, x, y) &= 0, \end{aligned}$$

which again by the parabolic maximum principle entails that $w(t, x, y) \geq 0$ for all $t \geq 0$ and $(x, y) \in \mathcal{X}^2$, meaning that $g_2 \geq g_1$ and by (B.6), $f_2 \geq f_1$.

Let us now argue by induction, and let us assume there exists $k \geq 2$ such that for all $0 < n \leq k$, $f_n \leq f_{n+1}$ and $g_n \leq g_{n+1}$. Let us prove that these inequality are still true for $k + 1$. Note that thanks to our induction assumption we only need to show that $f_{k+1} \leq f_{k+2}$ and $g_{k+1} \leq g_{k+2}$. Define now $w = g_{k+2} - g_{k+1}$. Thanks to the definition of

g_n and using that $f_{k+1} \geq f_k$ we deduce that for all $t > 0, (x, y) \in \mathcal{X}^2$

$$\begin{aligned} \partial_t w(t, x, y) &= \mathcal{L}_\varepsilon[w](t, x, y) - \lambda(x, y)w + \mu_1 D(x, y)(f_{k+1}(t, x) - f_k(x)) \geq \mathcal{L}_\varepsilon[w](t, x, y) - \lambda(x, y)w(t, x, y), \\ \nabla_{x, y} w(t, x, y) \cdot \vec{n} &= 0, \\ w(0, x, y) &= 0. \end{aligned}$$

By using once again the parabolic maximum principle, we get $w \geq 0$, for all $t \geq 0$ and $(x, y) \in \mathcal{X}^2$, meaning that $g_{k+2} \geq g_{k+1}$. We then ensure that $f_{k+2} \geq f_{k+1}$ by using (B.6).

Since from the above computation, the induction assumption is true for $k = 2$, we are done. \square

We are now in position to show that $(f_n(t, x), g_n(t, x))_{n \in \mathbb{N}}$ converges. From the monotone behaviour of the sequence, to ensure the convergence, we only need to obtain a uniform upper bound independent of n .

To do so, since $f_n \in L^2(\mathcal{X})$, let us multiply (B.1) by f_n and integrate over \mathcal{X} . We then get using Jensen inequality

$$\frac{1}{2} \frac{d}{dt} \int_{\mathcal{X}} f_n^2(t, x) dx = \int_{\mathcal{X}} \left(f_n(t, x) \int_{\mathcal{X}} \lambda(z, x) g_n(t, z, x) dz \right) dx \leq \bar{\lambda} \|f_n\|_2 \|g_n\|_2.$$

Therefore we get

$$\frac{d}{dt} \|f_n\|_2 \leq \bar{\lambda} \|g_n\|_2,$$

leading to

$$\|f_n\|_2 \leq \|f_0\|_0 + \bar{\lambda} \int_0^t \|g_n\|_2(s) ds.$$

Now let us multiply (B.2) by g_n and integrate over \mathcal{X}^2 , a short computation shows that

$$\frac{d}{dt} \|g_n\|_2 \leq C \left(\int_{\mathcal{X}} f_{n-1}^2(t, x) \left(\int_{\mathcal{X}} D^2(x, y) dy \right) dx \right)^{\frac{1}{2}} \leq C \|f_{n-1}\|_2.$$

Thus, we get

$$\|g_n\|_2(t) \leq C \int_0^t \|f_{n-1}\|_2(s) ds$$

leading to

$$\|f_n\|_2 \leq \|f_0\|_2 + \bar{\lambda} C \int_0^t \int_0^s \|f_{n-1}\|_2(\tau) d\tau.$$

By a recursive argument, for all $n \geq 0$ we achieve

$$\|f_n\|_2 \leq \|f_0\|_2 \sum_{i=0}^n \frac{(Ct^2)^i}{2i!},$$

and so we obtain the bound

$$\|f_n\|_2 \leq \|f_0\|_2 e^{Ct^2}.$$

The latter enforces also the following bound on g_n

$$\|g_n\|_2 \leq C \int_0^t \|f_{n-1}\|_2(s) ds \leq C \|f_0\|_2 \int_0^t e^{Cs^2} ds.$$

To obtain H^1 estimate, let us observe that since $\lambda, D \in W^{1, \infty}$, then we can check that $\nabla_x f_n$ and $\nabla_i g_n$ with $i = x$ or $i = y$ respectively satisfies

$$\begin{aligned}
\partial_t \nabla_y f_n(t, y) &= \int_{\mathcal{X}} \nabla_y \lambda(z, y) g_n(t, z, y) dz + \int_{\mathcal{X}} \lambda(z, y) (\nabla_y g_n)(t, z, y) dz && \text{for all } t > 0, y \in \mathcal{X} \\
\partial_t \nabla_i g_n(t, x, y) &= \mathcal{L}_\varepsilon[\nabla_i g_n](t, x, y) - \nabla_i(\lambda(x, y) g_n(t, x, y)) + \nabla_i(\mu_1 D(x, y) f_{n-1}(t, x)) && \text{for all } t > 0, (x, y) \in \mathcal{X}^2 \\
\nabla_y f_n(0, y) &= \nabla_y f_0(y) && \text{for all } y \in \mathcal{X} \\
\nabla_i g_n(0, x, y) &= 0 && \text{for all } (x, y) \in \mathcal{X}^2.
\end{aligned}$$

By multiplying the first equality by $\nabla_y f_n$ and the second by $\nabla_i g_n$ and integrate respectively on \mathcal{X} and \mathcal{X}^2 , we get after straightforward computations using the estimates on f_n, g_n above

$$\begin{aligned}
\partial_t \|\nabla f_n\|_2 &\leq C \|f_0\|_2 \int_0^t e^{Cs^2} ds + C \|\nabla_y g_n\|_2 && \text{for all } t > 0, \\
\partial_t \|\nabla_y g_n\|_2 &\leq C \|f_0\|_2 \left(e^{Ct^2} + \int_0^t e^{Cs^2} ds \right) && \text{for all } t > 0, \\
\partial_t \|\nabla_x g_n\|_2 &\leq C \|f_0\|_2 \left(e^{Ct^2} + \int_0^t e^{Cs^2} ds \right) + C \|\nabla f_{n-1}\|_2 && \text{for all } t > 0.
\end{aligned}$$

We then obtain the desired H^1 estimates by exploiting the above differential relations and the estimates on f_n and g_n .

Now, for all $t \leq T$ the sequence $(g_n, f_n - f_0)$ is uniformly bounded in $C^{1,1}((0, T), H^1(\mathcal{X}^2)) \times C^{1,1}((0, T), H^1(\mathcal{X}))$, so, up to extraction of a subsequence, there exists a convergent subsequence in $C^1((0, T), L^2(\mathcal{X}))$ to some functions (\bar{g}, \bar{f}) which is a weak solution to

$$\begin{aligned}
\partial_t f(t, x) &= \int_{\mathcal{X}} \lambda(z, x) g(t, z, x) dz && \text{for all } 0 < t < T, x \in \mathcal{X} \\
\partial_t g(t, x, y) &= \mathcal{L}_\varepsilon[g](t, x, y) - \lambda(x, y) g(t, x, y) + \mu_1 D(x, y) f(t, x) && \text{for all } 0 < t < T, (x, y) \in \mathcal{X}^2 \\
\nabla_{x,y} g(t, x, y) \cdot \bar{\mathbf{n}} &= 0, && \text{for all } 0 < t < T, (x, y) \in \partial \mathcal{X}^2 \\
f(0, x) &= f_0(x) && \text{for all } x \in \mathcal{X} \\
g(0, x, y) &= 0 && \text{for all } (x, y) \in \mathcal{X}^2.
\end{aligned}$$

Thanks to elliptic and parabolic regularity estimates, we can check that \bar{g}, \bar{f} is indeed a strong solution to the above equation. To complete the construction, we need to verify that \bar{g}, \bar{f} is non trivial which is straightforward, since $(f_n)_{n \in \mathbb{N}}$ and $(g_n)_{n \in \mathbb{N}}$ are monotone increasing sequences. \square

Next, we obtain some useful estimate and prove Proposition 4.3 that we recall below.

Proposition B.3. *Let \mathcal{X} be a smooth bounded domain, with at least a C^1 boundary. Assume further that $f_0 \in H^1(\mathcal{X})$, $f_0 \geq 0$, $\lambda, D \in W^{1,\infty}(\mathcal{X}^2)$ are non negative functions. Let $f_\varepsilon, g_\varepsilon$ be positive functions satisfying $f_\varepsilon \in C^1((0, T), H^1(\mathcal{X}))$, $g_\varepsilon \in C^1((0, T), H^1(\mathcal{X}) \times H^1(\mathcal{X}))$ and such that $(f_\varepsilon, g_\varepsilon)$ is a solution to (4.6)–(4.10).*

Then there exists positive constants $C_0, C_1, C_2, C_3, C_4, C_5$ independent of ε such that

$$\|f_\varepsilon\|_{H^1} \leq \|\nabla f_0\|_2 + \|f_0\|_2 \left(e^{C_0 t^2} + C_0 \int_0^t \int_0^\tau e^{C_0 s^2} ds d\tau + C_1 \int_0^t \int_0^\tau \int_0^s e^{C_0 \sigma^2} d\sigma ds d\tau \right),$$

and

$$\begin{aligned}
\|g_\varepsilon\|_{H^1} \leq C_1 \|\nabla f_0\|_2 t + \|f_0\|_2 \left(C_2 \int_0^t e^{C_0 \tau^2} d\tau + C_3 \int_0^t \int_0^\tau e^{C_0 s^2} ds d\tau \right. \\
\left. + C_4 \int_0^t \int_0^\tau \int_0^s e^{C_0 \sigma^2} d\sigma ds d\tau + C_5 \int_0^t \int_0^\tau \int_0^s \int_0^\sigma e^{C_0 \omega^2} d\omega d\sigma ds d\tau \right).
\end{aligned}$$

Moreover, for all $T > 0$, along any sequence $\varepsilon_n \rightarrow 0$ there exists a subsequence $(\varepsilon_{n_k})_{k \in \mathbb{N}}$ such that

$$f_{\varepsilon_{n_k}} \geq f_0 \quad \text{and} \quad \|g_{\varepsilon_{n_k}}\|_2(t) \geq \frac{1}{2}\|g_1\|_2(t) \quad \text{for} \quad 0 < t < T,$$

where g_1 is the unique solution of the ultra-parabolic equation:

$$\begin{aligned} \partial_t g_1(t, x, y) &= \Delta_y g_1(t, x, y) - \lambda(x, y)g_1(t, x, y) + \mu_1 D(x, y)f_0(x) && \text{for all } t > 0, (x, y) \in \mathcal{X}^2, \\ \nabla_y g_1(t, x, y) \cdot \vec{n} &= 0, && \text{for all } t > 0, x \in \mathcal{X}, y \in \partial\mathcal{X}, \\ g_1(0, x, y) &= 0 && \text{for all } (x, y) \in \mathcal{X}^2. \end{aligned}$$

Proof. To obtain now the estimate on the H^1 norm, we need to get uniform L^2 estimates on the $\nabla f, \nabla_x g$ and $\nabla_y g$. To do so, let us observe that since $\lambda, D \in W^{1,\infty}$, $f \in H^1(\mathcal{X})$ and $g \in H^1(\mathcal{X}^2)$, from (4.7) we can check that $\nabla_y g$ satisfies, in the sense of distribution,

$$\partial_t \nabla_y g(t, x, y) = \mathcal{L}_\varepsilon[\nabla_y g](t, x, y) - \nabla_y \lambda(x, y) \cdot g(t, x, y) - \lambda(x, y) \nabla_y g(t, x, y) + \mu_1 f(t, x) \nabla_y D(x, y).$$

By multiplying the above equation by $\nabla_y g$ and integrating over \mathcal{X}^2 , we get, using the coercivity of \mathcal{L}_ε and Cauchy-Schwarz inequality,

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int_{\mathcal{X}^2} |\nabla_y g(t, x, y)|^2 dx dy &= \langle \mathcal{L}_\varepsilon[\nabla_y g](t, x, y), \nabla_y g(t, x, y) \rangle - \int_{\mathcal{X}^2} g \nabla_y \lambda \nabla_y g - \int_{\mathcal{X}^2} \lambda |\nabla_y g|^2 + \mu_1 \int_{\mathcal{X}^2} f \nabla_y D \nabla_y g \\ &\leq \|\nabla_y g\|_2 \left(\|\nabla_y \lambda\|_\infty \|g\|_2 + \mu_1 \sqrt{\int_{\mathcal{X}} f^2(t, x) \left(\int_{\mathcal{X}} |\nabla_y D(x, y)|^2 dy \right) dx} \right), \end{aligned}$$

which after simplification by $\|\nabla_y g\|_2$, gives

$$\frac{d}{dt} \|\nabla_y g\|_2 \leq C_2 \|g\|_2 + C_3 \|f\|_2,$$

where $C_2 := \|\lambda\|_{1,\infty}$ and $C_3 := \mu_1 \sqrt{\sup_{x \in \mathcal{X}} \int_{\mathcal{X}} |\nabla_y D(x, y)|^2 dy}$.

Using the estimates (4.12) and (4.13), we achieve

$$\frac{d}{dt} \|\nabla_y g\|_2 \leq C_2 C_1 \|f_0\|_2 \int_0^t e^{C_0 \tau^2} d\tau + C_3 \|f_0\|_2 e^{C_0 t^2},$$

which after integration in time yields the L^2 bound

$$(B.7) \quad \|\nabla_y g\|_2 \leq C_2 C_1 \|f_0\|_2 \int_0^t \int_0^\tau e^{C_0 s^2} ds d\tau + C_3 \|f_0\|_2 \int_0^t e^{C_0 \tau^2} d\tau.$$

Let us now obtain a L^2 bound for ∇f . Since $\lambda \in W^{1,\infty}$, $f \in H^1(\mathcal{X})$ and $g \in H^1(\mathcal{X}^2)$, from (4.6) we can check that ∇f satisfies, in the sense of distribution,

$$\frac{d}{dt} \nabla f(t, y) = \int_{\mathcal{X}} \nabla_y \lambda(z, y) g(t, z, y) dy + \int_{\mathcal{X}} \lambda(z, y) \nabla_y g(t, z, y) dz.$$

By multiplying the later equality by ∇f and integrating over \mathcal{X} , it yields, using Cauchy-Schwarz inequality and $\lambda \in W^{1,\infty}$,

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int_{\mathcal{X}} |\nabla f(t, y)|^2 dy &= \int_{\mathcal{X}} \nabla f(t, y) \left(\int_{\mathcal{X}} \nabla_y \lambda(z, y) g(t, z, y) dy + \int_{\mathcal{X}} \lambda(z, y) \nabla_y g(t, z, y) dz \right) dy, \\ &\leq \|\nabla f\|_2 (\|\nabla_y \lambda\|_\infty \|g\|_2 + \|\lambda\|_\infty \|\nabla_y g\|_2) \\ &\leq \|\nabla f\|_2 C_2 (\|g\|_2 + \|\nabla_y g\|_2). \end{aligned}$$

After simplification by $\|\nabla f\|_2$, we get

$$\frac{d}{dt}\|\nabla f\|_2 \leq C_2(\|g\|_2 + \|\nabla_y g\|_2).$$

Integrating in time and using (4.13) and (B.7), then yields

$$(B.8) \quad \|\nabla f\|_2 \leq \|\nabla f_0\|_2 + C_2(C_3 + C_1)\|f_0\|_2 \int_0^t \int_0^\tau e^{C_0 s^2} ds d\tau + C_2 C_1 \|f_0\|_2 \int_0^t \int_0^\tau \int_0^s e^{C_0 \sigma^2} d\sigma ds d\tau.$$

Last we obtain a L^2 bound for $\nabla_x g$. As for $\nabla_y g$ from (4.7) we can check that $\nabla_x g$ satisfies, in the sense of distribution,

$$\partial_t \nabla_x g(t, x, y) = \mathcal{L}_\varepsilon[\nabla_x g](t, x, y) - \nabla_x \lambda(x, y) \cdot g(t, x, y) - \lambda(x, y) \nabla_x g(t, x, y) + \mu_1 f(t, x) \nabla_x D(x, y) + \mu_1 D(x, y) \nabla f(t, x).$$

By multiplying the above equation by $\nabla_x g$ and integrating over \mathcal{X}^2 , we get, using the coercivity of \mathcal{L}_ε and Cauchy-Schwarz inequality,

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int_{\mathcal{X}^2} |\nabla_x g|^2 dx dy &= \langle \mathcal{L}_\varepsilon[\nabla_x g], \nabla_x g \rangle - \int_{\mathcal{X}^2} g \nabla_x \lambda \nabla_x g - \int_{\mathcal{X}^2} \lambda |\nabla_x g|^2 + \mu_1 \int_{\mathcal{X}^2} f \nabla_x D \nabla_x g + \mu_1 \int_{\mathcal{X}^2} D \nabla f \nabla_x g \\ &\leq \|\nabla_x g\|_2 \left[\|\nabla_x \lambda\|_\infty \|g\|_2 + \mu_1 \sqrt{\int_{\mathcal{X}} f^2(t, x) \left(\int_{\mathcal{X}} |\nabla_x D(x, y)|^2 dy \right) dx} + \mu_1 \sqrt{\|\bar{D}\|_\infty \|\nabla f\|_2} \right] \\ &\leq \|\nabla_x g\|_2 \left[C_2 \|g\|_2 + \mu_1 \sqrt{\int_{\mathcal{X}} f^2(t, x) \left(\int_{\mathcal{X}} |\nabla_x D(x, y)|^2 dy \right) dx} + C_1 \|\nabla f\|_2 \right]. \end{aligned}$$

After simplification by $\|\nabla_x g\|_2$, and setting $C_4 := \mu_1 \sqrt{\sup_{x \in \mathcal{X}} \left(\int_{\mathcal{X}} |\nabla_x D(x, y)|^2 dy \right)}$ we get

$$\begin{aligned} \frac{d}{dt} \|\nabla_x g\|_2 &\leq \left[C_2 \|g\|_2 + \mu_1 \sqrt{\int_{\mathcal{X}} f^2(t, x) \left(\int_{\mathcal{X}} |\nabla_x D(x, y)|^2 dy \right) dx} + C_1 \|\nabla f\|_2 \right], \\ &\leq [C_2 \|g\|_2 + C_4 \|f\|_2 + C_1 \|\nabla f\|_2], \end{aligned}$$

which after integrating in time and using (4.13), (B.8), then yields

$$(B.9) \quad \|\nabla_x g\|_2 \leq C_1 \|\nabla f_0\|_2 t + C_4 \|f_0\|_2 \int_0^t e^{C_0 \tau^2} d\tau + C_2 C_1 \|f_0\|_2 \int_0^t \int_0^\tau e^{C_0 s^2} ds d\tau \\ + C_1 C_2 (C_3 + C_1) \|f_0\|_2 \int_0^t \int_0^\tau \int_0^s e^{C_0 \sigma^2} d\sigma ds d\tau + C_2 C_1^2 \|f_0\|_2 \int_0^t \int_0^\tau \int_0^s \int_0^\sigma e^{C_0 \omega^2} d\omega d\sigma ds d\tau.$$

Collecting (4.12), (B.8), (4.13), (B.7) and (B.9), then get the desired estimate.

To obtain the estimate from below, we just have to observe that since for all ε , g_ε is positive, the first assertion is then trivial. Now since $f_\varepsilon(t, x) \geq f_0$ for all ε , we have for all $t > 0$, $(x, y) \in \mathcal{X}^2$

$$\partial_t g_\varepsilon(t, x, y) = \mathcal{L}_\varepsilon[g_\varepsilon](t, x, y) - \lambda(x, y) g_\varepsilon(t, x, y) + \mu_1 D(x, y) f_\varepsilon(t, x) \geq \mathcal{L}_\varepsilon[g_\varepsilon](t, x, y) - \lambda(x, y) g_\varepsilon(t, x, y) + \mu_1 D(x, y) f_0(x),$$

and for all ε , we have by the parabolic comparison principle, $g_\varepsilon(t, x, y) \geq g_{1, \varepsilon}(t, x, y)$. To obtain the last part of the estimate from below, we need to characterised the convergence of $g_{1, \varepsilon}$ with respect to ε . Such characterisation follows as consequence of uniform H^1 bounds on $g_{1, \varepsilon}$ that can be obtained in a similar fashion as the one obtained for g_ε . The uniqueness of the limit solution follows also from a standard argument on the L^2 norm. \square

REFERENCES

- [1] Mario Ayala, Jerome Coville, and Raphael Forien. “A measure-valued stochastic model for vector-borne viruses”. In: *arXiv preprint arXiv:2211.04563* (2022).
- [2] Lyn C Branch, Diego Villarreal, and Gene S Fowler. “Recruitment, dispersal, and group fusion in a declining population of the plains vizcacha (*Lagostomus maximus*; Chinchillidae)”. In: *Journal of mammalogy* 74.1 (1993), pp. 9–20.

- [3] Nicolas Champagnat, Régis Ferrière, and Sylvie Méléard. “From Individual Stochastic Processes to Macroscopic Models in Adaptive Evolution”. In: *Stochastic Models* 24.sup1 (2008), pp. 2–44.
- [4] Nicolas Champagnat and Sylvie Méléard. “Invasion and Adaptive Evolution for Individual-Based Spatially Structured Populations”. In: *Journal of Mathematical Biology* 55.2 (2007), p. 147.
- [5] Nicolas Fournier and Sylvie Méléard. “A Microscopic Probabilistic Description of a Locally Regulated Population and Macroscopic Approximations”. In: *The Annals of Applied Probability* 14.4 (2004), pp. 1880–1919.
- [6] Avner Friedman. *Partial differential equations of parabolic type*. Courier Dover Publications, 2008.
- [7] Henry F Howe. “Scatter-and clump-dispersal and seedling demography: hypothesis and implications”. In: *Oecologia* 79.3 (1989), pp. 417–426.
- [8] Sylvie Méléard and Vincent Bansaye. *Stochastic Models for Structured Populations: Scaling Limits and Long Time Behavior*. Vol. 1. Springer, 2015.
- [9] Amnon Pazy. *Semigroups of linear operators and applications to partial differential equations*. Vol. 44. Springer Science & Business Media, 2012.
- [10] Marco A Pizo and Isaac Simão. “Seed deposition patterns and the survival of seeds and seedlings of the palm *Euterpe edulis*”. In: *Acta Oecologica* 22.4 (2001), pp. 229–233.
- [11] Steven B Reichling. “Group dispersal in juvenile *Brachypelma vagans* (Araneae, Theraphosidae)”. In: *The Journal of Arachnology* 28.2 (2000), pp. 248–250.
- [12] Sylvie Roelly-Coppoletta. “A criterion of convergence of measure-valued processes: application to measure branching processes”. In: *Stochastics: An International Journal of Probability and Stochastic Processes* 17.1-2 (1986), pp. 43–65.
- [13] Samuel Soubeyrand, Tomáš Mrkvicka, and Antti Penttinen. “A Nonstationary Cylinder-Based Model Describing Group Dispersal in a Fragmented Habitat”. In: *Stochastic Models* 30.1 (2014), pp. 48–67.
- [14] Samuel Soubeyrand et al. “Patchy patterns due to group dispersal”. In: *Journal of Theoretical Biology* 271.1 (2011), pp. 87–99.
- [15] Kazuaki Takahashi et al. “Seasonal variation in fleshy fruit use and seed dispersal by the Japanese black bear (*Ursus thibetanus japonicus*)”. In: *Ecological Research* 23.2 (2008), pp. 471–478.
- [16] Zhuoqun Wu, Jingxue Yin, and Chunpeng Wang. *Elliptic and parabolic equations*. World Scientific Publishing Company, 2006.