

Thermogravimetry and FT-IR spectroscopy: efficient approaches for organic amendments stability analysis?

Nadia Bennegadi-Laurent, Sabine Houot, Nathalie Damay, Jean-Baptiste Besnier, Karine Laval, Isabelle Trinsoutrot-Gattin

▶ To cite this version:

Nadia Bennegadi-Laurent, Sabine Houot, Nathalie Damay, Jean-Baptiste Besnier, Karine Laval, et al.. Thermogravimetry and FT-IR spectroscopy: efficient approaches for organic amendments stability analysis?. 17th INTERNATIONAL CONFERENCE RAMIRAN, Sep 2017, Wexford, Ireland. hal-04561826

HAL Id: hal-04561826 https://hal.science/hal-04561826

Submitted on 28 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Paper number : 180

Thermogravimetry and FT-IR spectroscopy: efficient approaches for organic amendments stability analysis?

Bennegadi-Laurent, N.1, Houot, S.2, Damay, N.3, Besnier, J.B.4, Laval, K.1, Trinsoutrot-Gattin, I.1

¹Institut Polytechnique LaSalle-Esitpa, campus Rouen, Research unit Agroecology Hydrogeochemistry Environment and Ressources (AGHYLE), 3 Rue du Tronquet - F-76130 Rouen Cedex, France ; 2 INRA, UMR 1091 Énvironnement et Grandes Cultures (EGC), 78850 Grignon, France ; ³ Laboratoire Départemental d'Analyses et de Recherche, Pôle du Griffon - 180 rue Pierre Gilles de Gennes, Barenton-Bugny F02007 Laon Cedex ; 4 Institut Polytechnique LaSalle-Esitpa, Transformation and Agroressources, 3 Rue du Tronquet - F-76130 Rouen Cedex- France.

nadia.laurent@unilasalle.fr

Introduction

- Spreading of exogenous organic material (EOMs) is un opportunity to provide nutrients for crops, improuve soil fertility and decrease chemical fertilizers
- EOMs have different physico-chemical characteristics, which depend on their origin and the processes of their treatments. These characteristics can strongly influence their behavior and nutrient release after spreading
- Objective : Develop technical solutions easy to implement and able to predic EOM behaviour in soils

Materials and Methods

Six organic materials representative of the large classe of EOMs ehold compost C/N = 19.5

Three analytical techniques and 4 replicates :


C/N= 9.9

FT-IR spectroscopy : 4000-400 cm⁻¹, resolution set to

EOMs were dried at 38°C and ground (1 mm)

cattle man C/N = 12.9

4 cm⁻¹ and 32 scans recorded




Poultry manure P C/N = 8.9

Van Soest biochemical fractionation ¹ & 3 days carbon mineralization in soil ² biological stability of EOMs (IROC indicator3) Thermogravimetric analysis (TG) : Oxidizing atmosphere (80% O₂/ 20% N₂) between

23-900°C. for calculated thermal stability index R1 (Fig 1)

Fig 1. Example of TG profile (TG: loss mass

curve,% ; DTG : first derivative, LM% : Loss



Results

FT-IR spectra :

EOMs show different compositions in biochemichal functions (Fig. 2), specifically in Alipahtic and Amide functions

TG analysis :

Two exothermal peaks of combustion: Aliphatic peak (268-300°C) as cellulose and lignocellulosic and aromatic peak (441-497°C) for recalcitrant compound (results not shown)





Fig 3.: Stability of OEMs with R1 and I_{ROC} indexes : sclae of stability

Mass



RAMIRAN

Similar classification of EOMS stability between R1 and I_{ROC} indexes (Fig 3) : Very strong (digestat, sludge compost); middle (manures), low (sludge, household compost)

Conclusions & Perspectives

- TG and IR spectra show they potential to provide a classification of EOMs according to their stability
 - FT-IR analysis could be further refined to develop an indicator such as R1 or I_{ROC}
 - As a perspective, these approaches should be compared with sequential chemical extractions coupling with 3D fluorescence spectroscopy 4



J. et al. (2014). Water Res. 50, 359-72 This work was supported by the National Research Programme of French Environment & Energy Managment Agency (ADEME)

Lashermes et al. 2009. Europ. J.of Soil Sci., 60, 297-310 ; ² AFNOR, 2009. French standard XPU 44-162 ; ³ AFNOR, 2009. French standard XPU 44-163 ; ⁴. Jimenez,

Fig 2. Infrared spectroscopic characteristics for EOMs