



HAL
open science

Inexact subgradient methods for semialgebraic functions

Jérôme Bolte, Tam Le, Éric Moulines, Edouard Pauwels

► **To cite this version:**

Jérôme Bolte, Tam Le, Éric Moulines, Edouard Pauwels. Inexact subgradient methods for semialgebraic functions. 2024. hal-04562371

HAL Id: hal-04562371

<https://hal.science/hal-04562371v1>

Preprint submitted on 29 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Inexact subgradient methods for semialgebraic functions

Jérôme Bolte* Tam Le † Eric Moulines ‡ Edouard Pauwels§

April 29, 2024

Abstract

Motivated by the widespread use of approximate derivatives in machine learning and optimization, we study inexact subgradient methods with non-vanishing additive errors and step sizes. In the nonconvex semialgebraic setting, under boundedness assumptions, we prove that the method provides points that eventually fluctuate close to the critical set at a distance proportional to ϵ^ρ where ϵ is the error in subgradient evaluation and ρ relates to the geometry of the problem. In the convex setting, we provide complexity results for the averaged values. We also obtain byproducts of independent interest, such as descent-like lemmas for nonsmooth nonconvex problems and some results on the limit of affine interpolants of differential inclusions.

1 Introduction

Let $f: \mathbb{R}^p \rightarrow \mathbb{R}$ be locally Lipschitz and consider the unconstrained global minimization problem

$$\min_{x \in \mathbb{R}^p} f(x).$$

An important tool for addressing such problems in modern high-dimensional nonsmooth settings, are subgradient algorithms, see e.g. [55, 56, 24, 49]. We focus here on the *inexact* or *biased subgradient method*, see [56]:

$$x_{k+1} \in x_k - \alpha_k [\partial^c f(x_k) + \bar{B}(0, \epsilon)], \quad x_0 \in \mathbb{R}^n,$$

where for all $k \in \mathbb{N}$, $\alpha_k > 0$ is a sequence of positive step sizes, $\epsilon > 0$ is an error or a bias level and ∂^c denotes the Clarke subdifferential [20].

*Toulouse School of Economics, Université Toulouse Capitole, Toulouse, France.

†Laboratoire Jean Kuntzmann, Université Grenoble Alpes, Grenoble, France.

‡CMAP, Ecole polytechnique, Paris and Mohamed Bin Zayed University of AI

§Toulouse School of Economics, Université Toulouse Capitole et IUF, Toulouse, France.

Let us first provide some motivations and some insights into this type of method, commenting in particular on two of its most distinctive features: inexact oracle evaluation and nonsmoothness.

The sources of error in the subgradient evaluation are very diverse. They may be generated by numerical errors or approximation techniques for derivatives: for instance the use of complex oracles for derivatives [34, 44, 13, 19], or the use of low precision computations, as the default 32 bits precision for training neural networks or more advanced techniques [25]. In recent decades, due to the increasing size of problems in both dimension and structural complexity—such as large sums, composite structures, intrinsic complexity of layers, differentiable programming, and federated learning—there has been a rapid development of what we could call “sketchy calculations”. In our case, the concept of a sketch calculus is employed to achieve cost-effective evaluation of subgradients, with the goal of saving memory, reducing computational overhead, and managing heterogeneous information. One can, for instance, think of subgradients sketched through mini-batching [39], sparsification [61], using sophisticated compression techniques [45] or incremental techniques [56, 46]. Federated and distributed learning, see e.g. [45], also provides many cases where subgradients estimates are inexact. The vast area of stochastic optimization provides many examples where one needs to deal with noisy measurement of sub-gradients, due to random subsampling of mini-batches, or Monte Carlo approximation errors; see [29, 47, 57, 32, 33, 38]. However, in this paper, we do not address stochastic approximation as it deserves a separate study.

Nonsmooth (and inexact) subgradient methods have deep roots in optimization with the many works of Shor [55], Ermoliev [29], Norkin [50], Nemirovskii [48] and also the pioneering work of Solodov-Zaidev [56]. Nonsmoothness is indeed prevalent for both convex and nonconvex problems. In the convex world, robustness questions naturally provide nonsmooth max problems [8], regularization techniques resort to nonsmooth regularizers [1, 59, 22, 7], while bundle methods rely on complex polyhedral oracles, see e.g. [44]. Modern machine learning provides a wealth of applications involving nonsmoothness. While this aspect has always been witnessed in deep learning through standard building blocks such as ReLU activations and MaxPooling, it appears in more recent contexts: for instance, sorting procedures may be used to promote sparsity [30], while optimization layers [3] may be used to refine learning abilities.

An original aspect of our work related to nonsmoothness is to deal with *nonvanishing stepsizes*. This makes the study considerably more difficult as steps are not used to mitigate the oscillation effects inherent to nonsmoothness and errors. One of our interests in this aspect is to be able to deal with the widespread use of schedulers in deep learning that do not systematically provide steps tending toward zero [51, 42, 21].

Let us now outline the main contributions of this article, highlighting the most original ones:

- The regularity assumptions on the cost f are weak and easily verifiable. Our results indeed apply to all local Lipschitz semialgebraic functions (or definable in an o-minimal structure), covering therefore a vast area of applications,

- We deal with constant step sizes and the endogenous bias they produce. As a byproduct, Lemma 13 provides a general result of independent interest for the ODE method, characterizing the limit points of small constant steps recursions.
- We show that *all the terms of the sequence eventually fluctuate around the critical sets*.
- We provide a bound for the fluctuation zone in $O(\epsilon^\rho)$, where ρ is related to the local geometry of the problem.
- In the convex case we do not make use of compactness or strong error bounds assumptions, we merely assume the function to be convex semialgebraic and coercive.

Our work is inscribed in a long series of reflections on nonsmooth and inexact (sub)gradient methods, for which we give a brief non-comprehensive overview.

The inexact subgradient algorithm with Clarke regular losses was studied in the pioneering [56]. There are many results for the convex case since the first studies within the Russian-Soviet school, they generally use demanding error bounds or compactness of the domain, see e.g. [46, 36], contrary to ours which resort uniquely on semialgebraicity and coercivity. Our convergence analysis is based on the comparison of sequences and trajectories of continuous time dynamical systems. It has become classical since the seminal work of [41, 10]. The ODE method for nonsmooth optimization with zero mean noise was considered in the pioneering [24], from which we borrow several ideas. Yet, we deal with non vanishing errors, and we go beyond vanishing steps. Previous studies on constant step sizes also use the ODE method [9, 31, 53, 11, 12], but our main theorems provide stronger results for an ubiquitous class of functions (semialgebraic, and even definable functions). In the nonconvex setting, the study of stochastic gradient algorithm with bias is not new [58, 27, 52, 49], but the results are either for smooth functions or based on strong assumptions, as sharpness or metric regularity. Let us mention [27] which contains results that are qualitatively similar to ours but for continuously differentiable functions. See [26] for a recent survey of this literature.

2 Preliminaries and statement of the main results

2.1 Notations.

For $f: \mathbb{R}^p \rightarrow \mathbb{R}$ locally Lipschitz, we denote its Clarke subdifferential by $\partial^c f$, and, for $\epsilon > 0$, we define

$$\begin{aligned} \text{crit}_\epsilon f &= \{x : \text{dist}(0, \partial^c f(x)) \leq \epsilon\}, \\ \text{vcrit}_\epsilon f &= f(\text{crit}_\epsilon f), \end{aligned} \tag{1}$$

the ϵ -critical set, which is closed and the set of ϵ -critical values. When $l \notin \text{vcrit}_\epsilon f$, it is called an ϵ -regular value. We also denote by $\text{crit} f$ the critical set and $\text{vcrit} f$ the set of critical values. The set of minimizers is $\text{argmin} f$.

For notational convenience, we write the algorithm as

$$x_{k+1} = x_k - \alpha_k v_\epsilon(x_k), \quad (2)$$

where $v_\epsilon : \mathbb{R}^p \rightarrow \mathbb{R}$ represent the biased oracle, i.e. satisfies $\text{dist}(v_\epsilon(x), \partial^c f(x)) \leq \epsilon$ for all $x \in \mathbb{R}^p$.

Throughout the next sections, we use the shorthand $[a \leq g \leq b]$, for real numbers a, b and function $g : \mathbb{R}^p \rightarrow \mathbb{R}$, to denote $\{x \in \mathbb{R}^p : a \leq g(x) \leq b\}$. We also use the same notations for $=$ and $<, >$. We denote the Euclidean norm by $\|\cdot\|$. For $A \subset \mathbb{R}^p$, we let $\|A\| = \sup_{a \in A} \|a\|$. For a subset, $A \subset \mathbb{R}^p$, \bar{A} is its closure, and A^c its complement. For $\epsilon > 0$, $\bar{B}(0, \epsilon)$ is the closed ball of center 0 and radius ϵ . We denote the distance of a point x to a compact set $A \subset \mathbb{R}^p$ by $\text{dist}(x, A) := \inf_{z \in A} \|x - z\|$. Given a set-valued map $Z : \mathbb{R}^p \rightrightarrows \mathbb{R}^q$, we denote the graph of Z by

$$\text{graph}[Z] := \{(x, y) \in \mathbb{R}^p \times \mathbb{R}^q : y \in Z(x)\}.$$

2.2 Main results

The nonconvex setting We will work under the following assumption.

Assumption 1. *The function $f : \mathbb{R}^p \rightarrow \mathbb{R}$ is L -Lipschitz, lower-bounded, semialgebraic with $\text{crit}_\epsilon f$ bounded for some $\epsilon > 0$.*

As stated in Lemma 7, Assumption 1 ensures that f is also coercive. Our first main result relates to vanishing step sizes.

Theorem 1 (Convergence for biased subgradient method with vanishing step size). *Under Assumption 1, there is $\bar{\epsilon} > 0$, $C > 0$ $\rho > 0$ such that for any $\epsilon < \bar{\epsilon}$, $x_0 \in \mathbb{R}^p$, there is $\bar{\alpha} > 0$, such that for any $(x_k)_{k \in \mathbb{N}}$ given by (2) with $0 < \alpha_k \leq \bar{\alpha}$ for all $k \in \mathbb{N}$, $\alpha_k \rightarrow 0$ and $\sum_{k=0}^{\infty} \alpha_k = \infty$, we have*

$$\begin{aligned} \lim_{k \rightarrow \infty} \text{dist}(f(x_k), \text{vcrit}_\epsilon f) &= 0, \\ \limsup_{k \rightarrow \infty} \text{dist}(x_k, \text{crit } f) &\leq C\epsilon^\rho. \end{aligned}$$

Our second main result relates to small constant step sizes.

Theorem 2 (Convergence for biased subgradient method with constant step size). *Under Assumption 1, there is $\bar{\epsilon} > 0$, $C > 0$ $\rho > 0$ such that for any $\epsilon < \bar{\epsilon}$, $x_0 \in \mathbb{R}^p$, and $(x_k(\alpha))_{k \in \mathbb{N}}$ given by (2) with $\alpha_k = \alpha > 0$ for all $k \in \mathbb{N}$, we have*

$$\begin{aligned} \lim_{\alpha \rightarrow 0^+} \limsup_{k \rightarrow \infty} \text{dist}(f(x_k(\alpha)), \text{vcrit}_\epsilon f) &= 0, \\ \limsup_{\alpha \rightarrow 0^+} \limsup_{k \rightarrow \infty} \text{dist}(x_k(\alpha), \text{crit } f) &\leq C\epsilon^\rho. \end{aligned}$$

The convex setting We complete these two results with an explicit estimate for the convex case. Our result is in the line of usual results, see, e.g., [46, 36], but does not use any compactness or strong error bounds.

For a coercive, convex, semialgebraic function $f: \mathbb{R}^p \rightarrow \mathbb{R}$, there exists $c > 0$ and $a \in (0, 1]$, such that

$$\frac{c}{2} ((f(x) - \min f)^a + (f(x) - \min f)) \geq \text{dist}(x, \text{argmin } f) \quad \forall x \in \mathbb{R}^p. \quad (3)$$

Theorem 3 (Biased subgradient complexity: convex case). *Let $f: \mathbb{R}^p \rightarrow \mathbb{R}$ be L -Lipschitz, semialgebraic, convex and coercive. Then with $a \in (0, 1]$ and $c > 0$, as in (3), for any sequence generated by (2), we have*

$$(2 - a - \epsilon c) \frac{\sum_{i=0}^k \alpha_i (f(x_i) - f^*)}{\sum_{i=0}^k \alpha_i} \leq (1 - a)(\epsilon c)^{\frac{1}{1-a}} + \frac{\|x_0 - x_*\|^2 + (L + \epsilon)^2 \sum_{i=0}^k \alpha_i^2}{\sum_{i=0}^k \alpha_i}.$$

We understand for $a = 1$, the value $(1 - a)(\epsilon c)^{\frac{1}{1-a}}$ as 0 if $\epsilon c < 1$ and undefined or infinite otherwise.

This allows to get convergence rates in value with various choices of step size. For example, if $\epsilon c < 1$, choosing $\alpha_i = \frac{1}{\sqrt{k+1}}$ for $i = 0, \dots, k$, leads to

$$\min_{i=1, \dots, k} f(x_k) - f^* \leq \frac{(1 - a)(\epsilon c)^{\frac{1}{1-a}}}{2 - a - \epsilon c} + \frac{\|x_0 - x_*\|^2 + (L + \epsilon)^2}{(2 - a - \epsilon c)\sqrt{k+1}}.$$

Note that the bound is vacuous if for example $a = 1$ and $\epsilon \geq 1/c$ and provides effective guarantees only for small values of ϵ . This is somewhat unavoidable: if f is the absolute value and $\epsilon > 1$, one completely loses control of the resulting biased subgradient sequence. This is the so called “low error” setting.

2.3 Reading keys and natural extensions

For the two first theorems, we examine the continuous-time limit of the recursion (2), which leads to a differential inclusion given in (4). A special Lyapunov mechanism is introduced in Section 3: Along the flow, the objective decreases only in certain regions of the space and may increase in other regions. This mechanism is used to describe the asymptotics of the system and provides explicit estimates in the Lipschitz coercive setting, under a Kurdyka-Lojasiewicz assumption for f and a metric regularity assumption for $\partial^c f$. This leads to an explicit estimate of the distance to $\text{crit } f$ for any positively invariant set A such that $f(A) \subset \text{vcrit}_\epsilon f$. The proof of Theorem 1 and Theorem 2 then consists of justifying the fact that relevant asymptotic sets fulfill this invariance property, which is done in Section 4. The main device is to use the small step limiting relation between the recursion (2) and its continuous-time counterpart for which the analysis was made in Section 3.

Theorem 3 is obtained by using standard Lyapunov functions in convex optimization and by applying an error bound condition that is automatically satisfied in the semialgebraic

case. These arguments are given in Section 5 and are completely independent of the arguments for Theorem 1 and Theorem 2, which make up the main technical part of this work.

These results could be readily extended in various ways.

- It can be seen directly that the same results apply beyond semialgebraicity to any polynomially bounded o-minimal structure, the prototypical example being the structure of globally subanalytic sets, see [60] for an overview.
- It is also directly possible to obtain qualitatively similar results, without the convergence rate estimates, in any o-minimal structure, replacing the exponent estimates by definable functions [60].
- While we consider the Clarke subdifferential, the main device used in the proof is the chain rule along Lipschitz curves, which is satisfied for the broader class of conservative gradient fields [17]. Examples are automatic differentiation oracles and generalized derivatives as in [28]. It is known that conservative gradients may induce spurious stationary points. The result in [16, Theorem 4.12] ensures that up to removing a zero measure meagre set of possible initialization $x_0 \in \mathbb{R}^p$ and a finite set of step sizes $\alpha \in \mathbb{R}$, one does actually manipulate the subdifferential oracle.
- We only consider the deterministic setting, it is possible to extend these results to the stochastic approximation setting, by adding zero mean stochastic perturbation terms which satisfy summability hypotheses for vanishing step sizes [10] or by resorting to the study of convergence of stationary measures in the constant step size setting [53].

3 The continuous-time system and auxiliary results

Let $f: \mathbb{R}^p \rightarrow \mathbb{R}$ be locally Lipschitz. Consider for $\epsilon > 0$, the differential inclusion

$$\dot{x}(t) \in -\partial^c f(x(t)) + \bar{B}(0, \epsilon), \quad (4)$$

for almost every $t \in [0, +\infty)$, where the solution is to be found among locally Lipschitz curves. Existence of solutions on maximal intervals follows from classical results, see [6, 4]. In particular, if f is Lipschitz, we may consider solutions defined on \mathbb{R}_+ . Equivalently, x is solution to (4) if for almost every t , $\text{dist}(\dot{x}(t), -\partial^c f(x(t))) \leq \epsilon$.

3.1 Asymptotics of the biased dynamics

Following Valadier and [17], a locally Lipschitz function f is called path-differentiable if for any Lipschitz curve $\gamma: \mathbb{R} \rightarrow \mathbb{R}^p$, we have for almost every $t \in \mathbb{R}$,

$$\frac{d}{dt}(f \circ \gamma)(t) = \langle v, \dot{\gamma}(t) \rangle, \quad \forall v \in \partial^c f(\gamma(t)).$$

Semialgebraic functions are path-differentiable [24, 17]. Under path-differentiability, we obtain the following results.

Lemma 4 (Descent properties). *Let f be locally Lipschitz and path-differentiable. Let $\epsilon > 0$ and $x : \mathbb{R}_+ \rightarrow \mathbb{R}^p$ be a solution to the differential inclusion (4) assumed to be defined on \mathbb{R}_+ (see remark below). Then,*

1. (Weak Lyapunov) *There is a measurable selection v_x of $\partial^c f(x(\cdot))$ such that,*

$$f(x(t_2)) - f(x(t_1)) \leq - \int_{t_1}^{t_2} \|v_x(t)\|(\|v_x(t)\| - \epsilon)dt. \quad \forall 0 \leq t_1 \leq t_2.$$

In particular, if $f(x(0)) \notin \text{vcrit}_\epsilon f$, there is $t > 0$ such that $f(x(s)) < f(x(0))$ for all $0 \leq s \leq t$.

2. (Decrease) *Let $l \notin \text{cl vcrit}_\epsilon f$ be such that $f(x(0)) \leq l$ then for all $t > 0$, $f(x(t)) < l$.*
3. (Asymptotics) *Either $\|x(t)\| \rightarrow +\infty$ as $t \rightarrow +\infty$, or we have both*

$$\begin{aligned} \liminf_{t \rightarrow \infty} \text{dist}(0, \partial^c f(x(t))) &\leq \epsilon \\ \lim_{t \rightarrow \infty} \text{dist}(f(x(t)), \text{vcrit}_\epsilon f) &= 0. \end{aligned}$$

4. (Quantitative estimates) *For any $0 \leq a < b$ and $\delta > \epsilon$, set $T = \frac{b-a}{\delta(\delta-\epsilon)}$ and assume that $f(x([0, T])) \subset [a, b]$, then there is $t \in [0, T]$ such that $\text{dist}(0, \partial^c f(x(t))) \leq \delta$.*
5. (ϵ -stationarity) *For any $a \in \mathbb{R}$ and $T > 0$, if $f(x([0, T])) = a$ then $x([0, T]) \subset \text{crit}_\epsilon f$.*

Remark 1. Actually Lemma 4 points 1, 2, 4, 5 hold on the interval of definition of the solution which does not need to be \mathbb{R}_+ . In our proofs, we will apply Lemma 4 to solutions defined on an interval $[0, T] \subset \mathbb{R}_+$.

Proof :

1. Consider a measurable selection v_x such that for almost every $t \geq 0$,

$$v_x(t) \in \text{argmin}_{v \in \partial^c f(x(t))} \|\dot{x}(t) + v\|.$$

Such a selection exists, see e.g. [2, 18.13]. Since $\dot{x}(t) \in -\partial^c f(x(t)) + \bar{B}(0, \epsilon)$, then $\|\dot{x}(t) + v_x(t)\| \leq \epsilon$ for almost every $t \geq 0$. Then, by path-differentiability of f , we have for almost every $t \geq 0$,

$$\begin{aligned} \frac{d}{dt}(f \circ x)(t) &= \langle v_x(t), \dot{x}(t) \rangle \\ &= \langle v_x(t), -v_x(t) + v_x(t) + \dot{x}(t) \rangle \\ &= -\|v_x(t)\|^2 + \langle v_x(t), v_x(t) + \dot{x}(t) \rangle \\ &\leq -\|v_x(t)\|(\|v_x(t)\| - \|\dot{x}(t) + v_x(t)\|) \\ &\leq -\|v_x(t)\|(\|v_x(t)\| - \epsilon). \end{aligned} \tag{5}$$

Integrating from t_1 to t_2 gives the desired inequality. If $f(x(0)) \notin \text{vcrit}_\epsilon f$, then $x(0) \notin \text{crit}_\epsilon f$ since $\text{crit}_\epsilon f$ is closed, there is a compact set U with $x(0) \in \text{int } U$ such that $U \cap \text{crit}_\epsilon f = \emptyset$ and f is Lipschitz on U . We may therefore choose $t > 0$ small enough such that $x(s) \in U$ for all $s \in [0, t]$ and the result follows.

2. We distinguish two cases.

Case 1. Assume that $f(x(0)) \notin \text{cl vcrit}_\epsilon f$. Let us show that $f(x(t)) < f(x(0))$ for all $t > 0$.

Since $(\text{cl vcrit}_\epsilon f)^c$ is open and f is locally Lipschitz, there exists $t^* > 0$ small enough such that $f(x(s)) \notin \text{cl vcrit}_\epsilon f$ for all $s \in [0, t^*]$. In this case, 1 gives us $f(x(s)) \leq f(x(0)) - \int_0^{t^*} \|v_x(s)\|(\|v_x(s)\| - \epsilon) ds < f(x(0))$ for all $s \in (0, t^*]$.

Now we show that for all $t \geq t^*$, $f(x(t)) \leq f(x(t^*))$. Assume toward a contradiction that there exists $t \geq t^*$ such that $f(x(t^*)) < f(x(t))$. Since $f(x(t^*)) \notin \text{cl vcrit}_\epsilon f$, we may choose t so that $[f(x(t^*)), f(x(t))] \subset (\text{cl vcrit}_\epsilon f)^c$ by openness of the latter. Now, consider $t^- = \max\{s : s \in [t^*, t], f(x(s)) \leq f(x(t^*))\}$, and $t^+ = \min\{s : s \in [t^-, t], f(x(s)) \geq f(x(t))\}$. By continuity of $f \circ x$, t^- and t^+ are well defined with $t^+ \geq t^-$, and they satisfy for all $s \in [t^-, t^+]$, $f(x(s)) \in [f(x(t^-)), f(x(t^+))] \subset (\text{cl vcrit}_\epsilon f)^c \subset (\text{vcrit}_\epsilon f)^c$, as well as $f(x(t^-)) = f(x(t^*))$ and $f(x(t^+)) = f(x(t))$. In particular, $f(x(t^+)) > f(x(t^-))$ hence $t^+ > t^-$. Let v_x be given by 1. Then we have

$$f(x(t^+)) - f(x(t^-)) \leq - \int_{t^-}^{t^+} \|v_x(s)\|(\|v_x(s)\| - \epsilon) ds < 0,$$

where the last inequality comes from $t^+ > t^-$ and $\|v_x(s)\| > \epsilon$ for almost every $s \in [t^-, t^+]$ since $f(x(s)) \notin \text{vcrit}_\epsilon f$. This yields a contradiction. We have shown that for all $t \geq t^*$, $f(x(t)) \leq f(x(t^*))$.

Finally, for $t \in (0, t^*]$, $f(x(t)) < f(x(0))$, and for $t > t^*$, $f(x(t)) \leq f(x(t^*)) < f(x(0))$ hence the desired result under the assumption that $f(x(0)) \notin \text{cl vcrit}_\epsilon f$.

Case 2. Now, assume that $f(x(0)) \in \text{cl vcrit}_\epsilon f$ and $f(x_0) \leq l$ for $l \notin \text{cl vcrit}_\epsilon f$. In this case we actually have $f(x(0)) < l$. Since $(\text{cl vcrit}_\epsilon f)^c$ is open, there is $l' \notin \text{cl vcrit}_\epsilon f$ such that $f(x(0)) < l' < l$. Then by continuity of $f \circ x$, either $f(x(t)) < l$ for all t or there exists $t > 0$ such that $f(x(t)) = l'$ and t can be chosen to be minimal since $[f \circ x = l']$ is closed and lower bounded. We have $f(x(s)) \leq l'$ for all $s \leq t$, and by 2.1, we have for all $s \geq t$, $f(x(s)) \leq l'$. Since $l' < l$, we have the desired result.

4. We now prove the fourth item, which does not depend on item 3 but is used to prove item 3. We have $a < b$. Assume toward a contradiction that $\text{dist}(0, \partial^c f(x(t))) > \delta$ for all $t \in [0, T]$. Since $s \rightarrow \text{dist}(0, \partial^c f(x(s)))$ is lower semi-continuous, there exists $\delta' > \delta$ such that $\text{dist}(0, \partial^c f(x(t))) \geq \delta'$ for all $t \in [0, T]$. In this case, 1 gives us $f(x(T)) \leq f(x(0)) - T\delta'(\delta' - \epsilon)$, hence

$$f(x(T)) \leq b - (b - a) \frac{\delta'(\delta' - \epsilon)}{\delta(\delta - \epsilon)} < b - (b - a) = a$$

This is a contradiction as we assumed that $f(x(T)) \geq a$.

3. Assume that $\|x(t)\|$ does not go to $+\infty$, this means that the trajectory has accumulation

points. So $f(x(t))$ also has finite accumulation values and in particular $f(x(t))$ does not diverge to $-\infty$ or $+\infty$.

— Using 1, we have that $\liminf_{t \rightarrow \infty} \text{dist}(0, \partial^c f(x(t))) > \epsilon$ implies that $f(x(t)) \rightarrow -\infty$ as $t \rightarrow \infty$ and we obtain the first limit.

— Denote by I all the accumulation points of $f(x(t))$ as $t \rightarrow \infty$. I is a nonempty interval by continuity of $f \circ x$ and by the fact that $f \circ x$ does not go to $+\infty$ or to $-\infty$. We distinguish two cases.

First if I has empty interior, then $I = \{l\}$ and $f(x(t)) \rightarrow l$ for some $l \in \mathbb{R}$. Since the trajectory x has accumulation points, we may find a diverging sequence $(t_k)_{k \in \mathbb{N}}$ in \mathbb{R}_+ such that $x(t_k) \rightarrow \bar{x} \in \mathbb{R}^p$ as $k \rightarrow \infty$, and we have $f(\bar{x}) = l$. By 4.2, for any $\delta > \epsilon$, set $a = l - \delta(\delta - \epsilon)$, $b = l + \delta(\delta - \epsilon)$ we have $f(x(t_k + \mathbb{R}_+)) \in [a, b]$ for k sufficiently large and therefore, there exists $s_k \in [t_k, t_k + 2]$ such that $\text{dist}(0, \partial^c f(x(s_k))) \leq \delta$. This can be repeated for smaller δ and we may find a sequence, $(s_k)_{k \in \mathbb{N}}$, such that for each k , $s_k \in [t_i, t_i + 2]$ for some $i \in \mathbb{N}$ and $\text{dist}(0, \partial^c f(x(s_k))) \rightarrow \epsilon$. The restricted trajectory $\{x([t_i, t_i + 2])\}_{i \in \mathbb{N}}$ remains in a bounded set, so up to a subsequence, we may assume that $x(s_k)$ converges to a point $\tilde{x} \in \text{crit}_\epsilon f$ with $f(\tilde{x}) = l$. This shows that $l \in \text{vcrit}_\epsilon f$.

Secondly, let us assume that I has nonempty interior. Assume toward a contradiction that there exists $l \in \text{int } I \cap (\text{cl } \text{vcrit}_\epsilon f)^c$. In this case, there is $u > 0$ such that $[l - u, l + u] \subset \text{int } I \cap (\text{cl } \text{vcrit}_\epsilon f)^c$ and t such that $f(x(t)) \in [l - u, l]$. By 2, we have $f(x(s)) < l$ for all $s > t$, but this is contradictory with the fact that $l + u \in \text{int } I$, because this implies that $t \mapsto f(x(t))$ has accumulation values strictly greater than $l + u$. So I is an interval such that $\emptyset \neq \text{int } I \subset \text{cl } \text{vcrit}_\epsilon f$ hence $I \subset \text{cl } \text{vcrit}_\epsilon f$. This means that $\sup_{v \in I} \text{dist}(v, \text{vcrit}_\epsilon f) = 0$, which is the desired result.

5. If $T > 0$ and $f \circ x$ is constant on $[0, T]$, by 1, we necessarily have $\text{dist}(0, \partial^c f(x(t))) \leq \epsilon$ for almost every $t \in (0, T)$, and $x([0, T]) \in \text{crit}_\epsilon f$ because $\text{crit}_\epsilon f$ is closed and x is continuous.

□

3.2 Estimates under the nonsmooth KL inequality and a metric subregularity condition

We will obtain more precise estimates under the following assumption.

Assumption 2. f is L -Lipschitz, path-differentiable, the set $\text{vcrit } f$ is non-empty finite, and there exists $\bar{\epsilon} \in (0, 1)$ such that for any $0 \leq \epsilon \leq \bar{\epsilon}$, $\text{vcrit}_\epsilon f$ is a finite union of segments.

Furthermore, there exists $c > 0$ and $\theta \in [0, 1)$, and $\beta > 0$ such that for all $x \in f^{-1}(\text{vcrit}_{\bar{\epsilon}} f)$

$$\text{dist}(f(x), \text{vcrit } f)^\theta \leq c \text{dist}(0, \partial^c f(x)) \quad (\text{KL})$$

$$\text{dist}(x, \text{crit } f) \leq c \text{dist}(0, \partial^c f(x))^\beta. \quad (\text{MR})$$

Property (KL) is some form of the nonsmooth Kurdyka-Lojasiewicz inequality [15], while (MR) is a form of metric sub-regularity of the subdifferential around the critical set, see

[40, Proposition 3.1] for a general result, but also [35, 5, 54, 43] for concrete applications in optimization.

In our context, Assumption 2 is essential to control trajectories of (4). Some previous works on biased algorithms relied as well on similar conditions. For instance, KL inequality for real analytic functions was used in [27] and metric regularity ($\beta = 1$) appears in [49]. Assumption 2 is actually satisfied for all semialgebraic functions:

Lemma 5 (Semialgebraicity implies regularity). *If Assumption 1 is satisfied, then Assumption 2 is also satisfied for some $\bar{\epsilon} > 0$.*

Proof : According to Sard’s theorem for semialgebraic functions [15], $\text{vcrit } f$ is finite. Assumption 1 ensures that there exists $\bar{\epsilon}$ such that $\text{crit}_\epsilon f$ is compact for every $0 \leq \epsilon \leq \bar{\epsilon}$. For $0 \leq \epsilon \leq \bar{\epsilon}$ the sets $\text{vcrit}_\epsilon f \subset \mathbb{R}$ are then compact and semialgebraic, i.e. they consist of a finite number of segments. Furthermore, by Lemma 7 (see next subsection), f is coercive so $f^{-1}(\text{vcrit}_\epsilon f)$ is compact. (KL) follows from Kurdyka-Lojasiewicz inequality for nonsmooth semialgebraic functions [14] and compactness. As for (MR), this is Hölder metric subregularity as given in [40, Proposition 3.1] on the compact set $f^{-1}(\text{vcrit}_\epsilon f)$. □

We will therefore prove Theorem 1 and Theorem 2 under Assumption 2.

Remark 2 (Beyond semialgebraicity). Semialgebraicity in Assumption 1 can be replaced by global subanalyticity and all results would hold true in the exact same form. This allows to include the logarithm and exponential function restricted to compact segments for example. More generally, our results hold provided that f is definable in a polynomially bounded o-minimal structure [23], for which semialgebraic sets and globally subanalytic sets are the main examples. Our results could also be extended if in Assumption 1 we assume that f is definable in an o-minimal structure (not necessarily polynomially bounded) instead of being semialgebraic. Under this assumption, we would obtain the same results in Theorem 1 and Theorem 2, but the power like estimates would be replaced by abstract non negative increasing definable functions continuous at 0 with value 0.

The next lemma generalizes the following fact, “for a locally Lipschitz continuous semialgebraic f and a subgradient curve x ($\dot{x}(t) \in -\partial^c f(x(t))$ for all $t \in \mathbb{R}_+$), the inclusion $f(x(\mathbb{R}_+)) \subset \text{vcrit } f$, implies $0 \in \partial^c f(x(t))$ for all $t \geq 0$ ”. Indeed, $\text{vcrit } f$ is made of a finite number of singletons, and if there is $t > 0$ such that $\text{dist}(0, \partial^c f(x(t))) > 0$, this would hold locally and result in a strict decrease by path-differentiability. We would thus have a time $t' > 0$ such that $f(x(t')) \notin \text{vcrit } f$. This fact, corresponding to the case when $\epsilon = 0$, can be extended to a general $\epsilon > 0$.

Lemma 6 (Approximate stationarity of near-critical curves). *Under Assumption 2, set $\rho := \min \left\{ \frac{(1-\theta)\beta}{\theta(1+\beta)}, \beta \right\}$. There exists $C > 0$, such that for any $\epsilon \in [0, \bar{\epsilon}]$ and any solution curve $x : \mathbb{R}_+ \rightarrow \mathbb{R}^p$ such that $f(x(\mathbb{R}_+)) \subset \text{vcrit}_\epsilon f$, we have for all $t \geq 0$,*

$$\text{dist}(x(t), \text{crit } f) \leq C e^{\rho t}. \tag{6}$$

Proof : Fix an arbitrary $\epsilon \leq \bar{\epsilon}$, and $x: \mathbb{R}_+ \rightarrow \mathbb{R}^p$ an arbitrary solution. Since $f(x(\mathbb{R}_+))$ is connected, $f(x(\mathbb{R}_+))$ is contained in a single connected component of $\text{vcrit}_\epsilon f$ of the form $[a, b] \subset \mathbb{R}$.

For all $z \in \text{crit}_\epsilon f$, $\text{dist}(0, \partial^c f(z)) \leq \epsilon$ implies $\text{dist}(f(z), \text{vcrit } f) \leq (c\epsilon)^{\frac{1}{\theta}}$ using (KL). Let $N \in \mathbb{N}$ such that $v_0 \leq v_1 \leq \dots \leq v_{N+1}$ are the ordered critical values in $[a, b]$ to which we added $v_0 = a$ and $v_{N+1} = b$. This defines $N + 1$ segments which cover $[a, b]$, one of them has length at least $\frac{b-a}{N+1}$. Consequently, there is an open segment (u, v) such that $v - u = \frac{b-a}{N+1}$ which does not contain any critical value. Therefore, we may choose $z \in \text{crit}_\epsilon f$ such that $f(z) = (u + v)/2$ and we have

$$\text{dist}(f(z), \text{vcrit } f) \geq \frac{v - u}{2} = \frac{b - a}{2(N + 1)}. \quad (7)$$

Combining (7) with (KL), we obtain

$$b - a \leq 2(N + 1)(c\epsilon)^{\frac{1}{\theta}} := K_1\epsilon^{\frac{1}{\theta}}. \quad (8)$$

We fix an arbitrary $\alpha > 0$ and $t \geq 0$. By Lemma 4.4 and (8), there is $t' \geq t$, such that

$$\text{dist}(0, \partial^c f(x(t'))) \leq \epsilon + \epsilon^\alpha \quad (9)$$

$$t' - t \leq \frac{b - a}{(\epsilon + \epsilon^\alpha)\epsilon^\alpha} \leq \frac{K_1\epsilon^{\frac{1}{\theta}}}{(\epsilon + \epsilon^\alpha)\epsilon^\alpha} \leq \frac{K_1\epsilon^{\frac{1}{\theta}}}{\epsilon^{1+\alpha}}. \quad (10)$$

It follows using the inclusion (4), the fact that f is L -Lipschitz so that its subgradient is bounded by L , and (10) that

$$\|x(t') - x(t)\| \leq \int_t^{t'} \|\dot{x}(s)\| ds \leq (t' - t)(L + \epsilon) \leq (L + \epsilon) \frac{K_1\epsilon^{\frac{1}{\theta}}}{\epsilon^{1+\alpha}} \quad (11)$$

Finally, using the estimates (11), (9) and (MR),

$$\begin{aligned} \text{dist}(x(t), \text{crit } f) &\leq \|x(t) - x(t')\| + \text{dist}(x(t'), \text{crit } f) \\ &\leq (L + \epsilon)K_1\epsilon^{\frac{1}{\theta}-1-\alpha} + c(\epsilon + \epsilon^\alpha)^\beta. \end{aligned} \quad (12)$$

Since $t \geq 0$ was arbitrary, the estimate (12) holds for all $t \geq 0$. Furthermore since $\alpha > 0$ was arbitrary, we may choose α freely. We distinguish two cases.

- If $\theta > \frac{1}{2+\beta}$, then we choose $\alpha = \frac{1-\theta}{\theta(1+\beta)} < 1$, and the exponent in (12) is of the form $C\epsilon^{\frac{(1-\theta)\beta}{\theta(1+\beta)}}$.
- If $\theta \leq \frac{1}{2+\beta}$ then we may choose $\alpha = \frac{1}{\theta} - \beta - 1 \geq 1$, and the exponent in (12) is of the form $C\epsilon^\beta$.

□

Remark 3 (On the initialization). It may be puzzling not to see a condition on the initialization in Lemma 6. This is actually hidden in the condition $f(x(\mathbb{R}_+)) \subset \text{vcrit}_\epsilon f$ which enforces x to start close enough to $\text{crit } f$ so that $f(x(t))$ cannot leave $\text{vcrit}_\epsilon f$ near $t = 0$.

Lemma 6 has the following direct consequence.

Corollary 1 (Invariant sets and biased dynamics). *Under Assumption 2, let $\epsilon \leq \bar{\epsilon}$, $S \subset \mathbb{R}^p$ be a positively invariant set, that is for any $z \in S$, there is $x: \mathbb{R}_+ \rightarrow \mathbb{R}^p$, solution to (4) such that $x(\mathbb{R}_+) \subset S$ and $x(0) = z$. If $f(S) \subset \text{vcrit}_\epsilon f$, then for any $z \in S$, $\text{dist}(z, \text{crit } f) \leq C\epsilon^\rho$, where C, ρ are given by Lemma 6.*

For the continuous time dynamics in (4), it is known that the set of accumulation points of a bounded solution trajectory form invariant set (see for example [10, Theorem 3.6, Lemma 3.5]). We obtain the following.

Corollary 2 (Asymptotics of biased dynamics). *Under Assumption 2, let $\epsilon \leq \bar{\epsilon}$, and $x: \mathbb{R}^+ \rightarrow \mathbb{R}^p$ be a bounded solution to the differential inclusion (4). For any $z \in \mathbb{R}^p$, accumulation point of the trajectory, we have $\text{dist}(z, \text{crit } f) \leq C\epsilon^\rho$, where C, ρ are given by Lemma 6.*

3.3 Coercivity and boundedness of ϵ critical points

Due to the importance of boundedness of curves in our results, we need to make a brief detour through coercivity properties and their link with the boundedness of $\text{crit}_\epsilon f$. Let us recall first:

Theorem 1 (Ekeland's variational principle). *Let X be a complete metric space with distance d . Let $f: X \mapsto \mathbb{R} \cup \{+\infty\}$ be lower semi-continuous, bounded below and finite at least at one point. Fix $\epsilon > 0$ and $x_0 \in X$ such that $f(x_0) \leq \epsilon + \inf_x f(x)$. Then for any $\lambda > 0$, there is $y_0 \in X$ such that*

$$f(y_0) \leq f(x_0) \quad d(x_0, y_0) \leq \lambda \quad f(x) + \frac{\epsilon}{\lambda}d(x, y_0) > f(y_0), \forall x \neq y_0.$$

The conclusion tells us that y_0 is a strict global minimizer of $g: x \rightarrow f(x) + \frac{\epsilon}{\lambda}d(x, y_0)$.

In our framework, \mathbb{R}^p is endowed with the Euclidean distance and f is locally Lipschitz, so using the sum rule with the Clarke subdifferential yields the following fact:

$$0 \in \partial^c g(y_0) \subset \partial^c f(y_0) + \bar{B}\left(0, \frac{\epsilon}{\lambda}\right) \quad (13)$$

This has the following consequences:

Lemma 7 (Boundedness of the ϵ -critical set implies coercivity). *Assume that $f: \mathbb{R}^p \rightarrow \mathbb{R}$ is locally Lipschitz, then for any $x \in \mathbb{R}^p$, $a \in (0, 1)$, there is $y \in \mathbb{R}^p$ such that*

$$\|y\| \geq \|x\| - \|x\|^a, \quad \text{dist}(0, \partial^c f(y))\|x\|^a \leq f(x) - \inf f.$$

In particular, if f is lower bounded and $\text{crit}_{\bar{\epsilon}} f$ is bounded for some $\bar{\epsilon} > 0$, then f is coercive.

Proof : Fix $x \in \mathbb{R}^p$, if $f(x) = \inf f$ there is nothing to prove. Choose $\epsilon = f(x) - \inf f$ (assumed finite, otherwise there is nothing to prove) and $\lambda = \|x\|^a$. By Ekeland's variational principle in Theorem 1 there is $y \in \mathbb{R}^p$ such that

$$\|x - y\| \leq \|x\|^a,$$

hence $\|y\| \geq \|x\| - \|x\|^a$. Moreover using (13), i.e., $0 \in \partial^c f(y) + \bar{B}\left(0, \frac{\epsilon}{\|x\|^a}\right)$, we get $\text{dist}(0, \partial^c f(y)) \leq \frac{\epsilon}{\|x\|^a}$.

If $\text{crit}_{\bar{\epsilon}} f$ is bounded for some $\bar{\epsilon} > 0$, and f was not coercive, we could choose a sequence $\|x_k\| = k + 1$ so that $f(x_k) - \inf f$ is bounded by some M , and obtain an unbounded sequence y_k in $\text{crit}_{\bar{\epsilon}} f$ for k large enough. \square

Remark 4 (Coercivity and critical points). (a) Of course $\text{crit} f$ bounded and nonempty does not imply coercivity, e.g., take $s \rightarrow (1 + s^2)^{-1}$.

(b) The fact that “ $\text{crit}_{\epsilon} f$ bounded (for some $\epsilon > 0$) implies f coercive” is a generalization of the classical result in convex analysis “ $\text{argmin} f$ bounded implies f coercive when f is convex”.

4 Main consequences for the biased subgradient method

4.1 Link between discrete and continuous-time

In this section, we repeatedly use the connection between the discrete and continuous-time systems in the limit of small step sizes. This is a classical approach which we state for a general set-valued map, see [18, Lemma 21]. We use the following assumptions which guarantee the existence of solutions to the differential inclusion [6, Chapter 2, Section 1, Theorem 3].

Assumption 3. Z is a set-valued map defined from \mathbb{R}^p into the subsets of \mathbb{R}^p ; it is assumed to be nonempty convex valued and locally bounded with closed graph.

Lemma 8 (Approximate solutions to differential inclusions). *Let Z be as in Assumption 3. For each $i \in \mathbb{N}$, let $T_i > 0$ and assume that $T_i \rightarrow T$ as $i \rightarrow \infty$ for some $T > 0$. For each $i \in \mathbb{N}$, let $\gamma_i: [0, T_i] \rightarrow \mathbb{R}^p$ be a Lipschitz curve. Assume that the sequence $(\gamma_i)_{i \in \mathbb{N}}$ converges to some bounded and Lipschitz curve $\gamma: [0, T] \rightarrow \mathbb{R}^p$, in the sense that $\sup_{t \in [0, \min(T_i, T)]} \|\gamma(t) - \gamma_i(t)\| \rightarrow 0$, and*

$$\lim_{i \rightarrow \infty} \int_0^{T_i} \text{dist}((\gamma_i(t), \dot{\gamma}_i(t)), \text{graph}[Z]) dt = 0. \quad (14)$$

Then $\dot{\gamma}(t) \in Z(\gamma(t))$ for almost every $t \in [0, T]$.

The following is classical and can be found for example in [18]. We will use it extensively to obtain discrete descent lemmas in the coming section.

Lemma 9 (Affine interpolants of discrete dynamics and their limit curves). *Consider $T > 0$ and sequences satisfying*

$$x_{k+1} - x_k \in \alpha_k Z(x_k), \quad k = 0, 1, \dots, K,$$

where K is such that $\sum_{k=0}^{K-1} \alpha_k \geq T$ and $0 < \alpha_k \leq \alpha$, $k = 0, \dots, K-1$. Let $\gamma_\alpha: [0, T] \rightarrow \mathbb{R}^p$ be the interpolation of $(x_k)_{k=0, \dots, K}$ defined as follows: for all $k = 0, 1, \dots, K$, $\gamma_\alpha \left(\sum_{i=0}^{k-1} \alpha_i \right) = x_k$, and γ_α is affine between these points. The curve γ_α satisfies

$$\int_0^T \text{dist}((\gamma_\alpha(t), \dot{\gamma}_\alpha(t)), \text{graph}[Z]) dt \leq \alpha T \sup_{0 \leq k \leq K-1} \|Z(x_k)\|.$$

In particular, if Z is bounded and we have a uniformly bounded family of such curves γ_α , up to a subsequence, as $\alpha \rightarrow 0$, the curves converge uniformly to a solution of the underlying differential inclusion.

Proof : For all $\alpha > 0$, the curve γ_α satisfies for any $t \in [0, T]$,

$$\text{dist}((\gamma_\alpha(t), \dot{\gamma}_\alpha(t)), \text{graph}[Z]) \leq \alpha \sup_{0 \leq k \leq K-1} \|Z(x_k)\|,$$

as $\dot{\gamma}_\alpha(t) = Z(x_k)$ where k is the closest point x_k on the curve corresponding to time smaller than t .

When Z is bounded by a constant L , one has a uniform bound since $\max_{0 \leq k \leq K} \|Z(x_k)\| \leq L$. In that case, the curves γ_α are L -Lipschitz continuous and thus equicontinuous.

Whence, by letting $\alpha \rightarrow 0$, if the curves are bounded, Arzelà-Ascoli theorem applies and provides a subsequence of γ_α that uniformly converges to solutions of the continuous-time differential inclusion, thanks to Lemma 8. \square

4.2 Descent lemmas

We now state some consequences of Lemma 8 which apply to our setting, see also [37] for similar arguments in the unbiased setting.

Lemma 10 (Quasi-descent Lemma). *Let f be locally Lipschitz and path-differentiable, $\epsilon > 0$ and $l \notin \text{vcrit}_\epsilon f$. Then for any $\eta > 0$ and $M > 0$, there is $\bar{\alpha} > 0$ such that for any $z \in \mathbb{R}^p$ satisfying $f(z) \leq l$, $\|z\| \leq M$, and for any sequence generated by (2), with $x_0 = z$ and $\alpha_k \leq \bar{\alpha}$, we have $f(x_k) \leq l + \eta$ for all $k < \inf\{i \in \mathbb{N} : \|x_i\| > M\}$.*

In particular, under Assumption 1, there is $M > 0$, such that for α small enough, all sequences generated by (2) initialized with $f(x_0) \leq l$ are bounded by M .

Proof : Toward a contradiction, we assume that there is $M > 0$ such that for any $\alpha > 0$, there exists $z = x_0$ such that $f(z) \leq l$ and a sequence $(\alpha_k)_{k \in \mathbb{N}}$ smaller than α such that there exists $K > 0$ satisfying $f(x_K) > l + \eta$ and $\|x_k\| \leq M$ for all $k = 0, \dots, K$. Observe that $(\alpha_k)_{k \in \mathbb{N}}$ depends on α , and denote by L a Lipschitz constant of f on the ball of radius M .

We fix a time horizon $T < \eta/(L^2 + L\epsilon)$ and we may assume that $\alpha < T$. We may then also find $k \in \mathbb{N}$ such that $f(x_k) \leq l$ and $f(x_i) > l$ for all $i = k + 1, \dots, K$. We have

$$l + \eta < f(x_K) \leq f(x_k) + L \sum_{j=k}^{K-1} \|x_{j+1} - x_j\| \leq l + L \sum_{j=k}^{K-1} \alpha_j(L + \epsilon) = l + (L^2 + L\epsilon) \sum_{j=k}^{K-1} \alpha_j.$$

We deduce that $\sum_{j=k}^{K-1} \alpha_j > T$. Let $K' \geq k$ be the smallest value such that $\sum_{j=k}^{K'} \alpha_j \geq T$, since $\alpha < T$ and by the argument above, we have $k < K' \leq K - 1$. We may then consider the truncated sequence $(x_i)_{i=k, \dots, K'+1}$, which is nonempty whenever $\alpha < T$. Consider the affine interpolation of this truncated sequence from $i = k$ to $i = K' + 1 \leq K$ and restricted to $[0, T]$ (which is possible because $\sum_{i=k}^{K'} \alpha_k \geq T$) as in Lemma 9, call it x_α , it is Lipschitz and satisfies:

- $\|x_\alpha(t)\| \leq M$ for all $t \in [0, T]$.
- $x_\alpha(0) = x_k$, $f(x_\alpha(0)) \leq l$ and $f(x_\alpha(t)) > l$ for some $t \in [0, \alpha]$.
- $l \leq f(x_\alpha(t)) \leq l + \eta + \alpha L(L + \epsilon)$ for all $t \in [\alpha, T]$.
- $\text{dist}((x_\alpha(t), \dot{x}_\alpha(t)), \text{graph}[-\partial^c f + \bar{B}(0, \epsilon)]) \leq \alpha(L + \epsilon)$ and $\|\dot{x}_\alpha(t)\| \leq L + \epsilon$ for almost every t .

Thus, these trajectories are uniformly bounded and equicontinuous. Applying Arzelà-Ascoli theorem allows to obtain a converging subsequence as $\alpha \rightarrow 0$. Lemma 8 ensures that the limit $x : [0, T] \rightarrow \mathbb{R}^p$ is a solution to (4) such that $f(x(0)) = l$ and $f(x([0, T])) \geq l$ which contradicts Lemma 4.1 (see Remark 1 for its validity for x defined on $[0, T]$).

The last remark follows because under Assumption 1, f is coercive (Lemma 7). Then it is Lipschitz on the (compact) sublevel set $l + \eta$, say with constant L . For a step size threshold α_0 , choose $M_0 = \max\{\|x\| + \alpha_0(L + \epsilon), f(x) \leq l + \eta\}$ and denote by \tilde{L} a Lipschitz constant of f on the ball or radius M_0 centered at zero. By coercivity, we may choose $M > 0$ large enough such that $\inf_{\|x\| \geq M} f(x) > l + \eta + \alpha_0 \tilde{L}(L + \epsilon)$. Applying the lemma for this choice of M , reducing the resulting $\bar{\alpha}$ if it is bigger than α_0 , for any admissible sequence, we have for all k such that $f(x_k) \leq l + \eta$, it holds that $\|x_{k+1} - x_k\| \leq \alpha_0(L + \epsilon)$ and $\|x_k\| + \alpha_0(L + \epsilon) \leq M_0$ so that both x_k and x_{k+1} are contained in the ball of radius M_0 . We deduce that $f(x_{k+1}) \leq l + \eta + \alpha_0 \tilde{L}(L + \epsilon)$ so $\|x_{k+1}\| \leq M$. We deduce that $k + 1 < \inf\{i \in \mathbb{N} : \|x_i\| > M\}$ and the main statement of the lemma ensures that, $f(x_{k+1}) \leq l + \eta$. By induction this holds true for all $k \in \mathbb{N}$. \square

Lemma 11 (ϵ -regular values are repulsive). *Let $f : \mathbb{R}^p \rightarrow \mathbb{R}$ be locally Lipschitz and path differentiable, $\epsilon > 0$, $l \notin \text{vcrit}_\epsilon f$, and $\eta := \text{dist}(l, \text{vcrit}_\epsilon f)/16 > 0$. For any $M > 0$, there is $\bar{\alpha} > 0$ such that for any sequence generated by (2), with $\alpha_k \leq \bar{\alpha}$ for all $k \in \mathbb{N}$, and $\sum_{k \in \mathbb{N}} \alpha_k = +\infty$, either $\{i \in \mathbb{N}, \|x_i\| > M\}$ is nonempty or $\liminf_{k \rightarrow \infty} f(x_k) > l + 2\eta$ or $\limsup_{k \rightarrow \infty} f(x_k) < l - 2\eta$.*

In particular under Assumption 1, there is $\bar{\alpha} > 0$ such that regardless of the initial condition for any sequence generated by (2), with $\alpha_k \leq \bar{\alpha}$, and $\sum_{k \in \mathbb{N}} \alpha_k = +\infty$ for all $k \in \mathbb{N}$, $\liminf_{k \rightarrow \infty} f(x_k) > l + 2\eta$ or $\limsup_{k \rightarrow \infty} f(x_k) < l - 2\eta$.

Proof : Fix $M > 0$ and denote by L a Lipschitz constant of f on the ball of radius M . By definition of η , we have that $[l - 8\eta, l + 8\eta] \subset (\text{vcrit}_\epsilon f)^c$. Set $\delta := \min\{\|v\| : v \in \partial^c f(x), x \in \mathbb{R}^p, \|x\| \leq M, |f(x) - l| \leq 8\eta\} > \epsilon$ and $T \geq 12\frac{\eta}{\delta(\delta - \epsilon)}$. Any solution $x : [0, T] \rightarrow \mathbb{R}^p$ to (4) bounded by M such that $f(x(0)) \leq l + 4\eta$ satisfies $f(x(T)) \leq l - 8\eta$ by Lemma 4.1.

1. Therefore there is $\bar{\alpha} > 0$, such that any sequence generated by (2), bounded by M , with $f(x_0) \leq l + 4\eta$ satisfies $f(x_k) \leq l - 4\eta$ for some $k \in \mathbb{N}$ and therefore has accumulation values below $l - 4\eta$. Indeed, if this was not the case, using Lemma 8 and the fact that $\sum_{k \in \mathbb{N}} \alpha_k = +\infty$, we could construct a solution to the differential inclusion satisfying $f(x(0)) \leq l + 4\eta$ satisfying $f(x(T)) \geq l - 4\eta$, which contradicts the previous statement.

2. Let us first observe that for all k , if $\|x_k\| \leq M$ and $\|x_{k+1}\| \leq M$, then

$$|f(x_{k+1}) - f(x_k)| \leq L\|x_{k+1} - x_k\| \leq \bar{\alpha}L(L + \epsilon),$$

so that we can assume, shrinking $\bar{\alpha}$ if necessary that this gap is less than η . Hence we may assume that for any sequence bounded by M such that $\limsup_{k \rightarrow \infty} f(x_k) \geq l - 2\eta$ and $\liminf_{k \rightarrow \infty} f(x_k) \leq l + 2\eta$, $(f(x_k))_{k \in \mathbb{N}}$ takes infinitely many value in $[l - 4\eta, l + 4\eta]$.

3. We may reduce $\bar{\alpha}$ as given by Lemma 10 so that any sequence initialized with $f(x_0) \leq l - 4\eta$ and bounded by M satisfies $f(x_k) \leq l - 3\eta$ for all $k \in \mathbb{N}$. The chosen $\bar{\alpha}$ satisfies the required properties.

Let us first summarize the properties of $\bar{\alpha}$. Note that the properties claimed above can be shifted by initializing a sequence at an arbitrary $K \in \mathbb{N}$ and by considering $k \geq K$. Below, $(x_k)_{k \in \mathbb{N}}$ is any sequence generated by (2) with $\alpha_k \leq \bar{\alpha}$ for all $k \in \mathbb{N}$, bounded by M , and $K \in \mathbb{N}$ is arbitrary.

1. If $f(x_K) \leq l + 4\eta$, there is $k \geq K$ such that $f(x_k) \leq l - 4\eta$.
2. If $\limsup_{k \rightarrow \infty} f(x_k) \geq l - 2\eta$ and $\liminf_{k \rightarrow \infty} f(x_k) \leq l + 2\eta$, then there is $k \in \mathbb{N}$ such that $f(x_k) \in [l - 4\eta, l + 4\eta]$.
3. If $f(x_K) \leq l - 4\eta$ then $f(x_k) \leq l - 3\eta$ for all $k \geq K$.

For the choice of $\bar{\alpha}$ as above, assume toward a contradiction that there exists a sequence generated by (2), bounded by M , with $\alpha_k \leq \bar{\alpha}$ for all $k \in \mathbb{N}$, satisfying $\limsup_{k \rightarrow \infty} f(x_k) \geq l - 2\eta$ and $\liminf_{k \rightarrow \infty} f(x_k) \leq l + 2\eta$. There would be $K \in \mathbb{N}$ such that $f(x_K) \in [l - 4\eta, l + 4\eta]$ by 2 and we deduce by 1 that there is $k \geq K$ such that $f(x_k) \leq l - 4\eta$. This implies by 3 that $\limsup_{k \rightarrow \infty} f(x_k) \leq l - 3\eta$ which is contradictory.

The last comment follows because by Lemma 10 reducing $\bar{\alpha}$ if necessary, there is $M > 0$ such that all sequences generated by (2) with $f(x_0) \leq l + 3\eta$ and $\alpha_k \leq \bar{\alpha}$ for all $k \in \mathbb{N}$ are bounded by M . If $\liminf_{k \rightarrow \infty} f(x_k) \leq l + 2\eta$, then there is $K > 0$ such that $f(x_K) \leq l + 3\eta$ and by considering $\tilde{x}_k = x_{k+K}$, bounded by M , the result follows. \square

Under Assumption 1, f is coercive and $\text{vcrit}_\epsilon f$ is closed. Combining the two previous lemma we obtain

Corollary 3 (Large-horizon descent Lemma). *Under Assumption 1, fix x_0 such that $f(x_0) \notin \text{vcrit}_\epsilon f$ and $\eta > 0$. There is $\bar{\alpha} > 0$ such that any sequence generated by (2) with $\alpha_k \leq \bar{\alpha}$ and $\sum_{k \in \mathbb{N}} \alpha_k = +\infty$ we have*

(i) $f(x_k) \leq f(x_0) + \eta$ for all $k \in \mathbb{N}$,

(ii) there is κ such that

$$f(x_k) \leq c + \eta, \quad \forall k \geq \kappa$$

where c is the first ϵ -critical value below $f(x_0)$.

Note that if $(1 + \rho)\eta < f(x_0) - c$ for some $\rho > 0$, the above implies

$$f(x_k) \leq f(x_0) - \rho\eta \text{ for } k \geq \kappa,$$

where κ is unknown.

4.3 Vanishing step sizes

For vanishing step sizes, the work of Benaim-Hofbauer-Sorin [10] ensures that the set of accumulation points is invariant. We additionally show that it has accumulation values in $\text{vcrit}_\epsilon f$. The main result stated in Theorem 1 is a consequence of the following result and Lemma 5.

Theorem 12 (Convergence for biased subgradient with vanishing step). *Under Assumption 2, let C, ρ be given by Lemma 6 and $\epsilon < \bar{\epsilon}$. Assume that $(x_k)_{k \in \mathbb{N}}$, is given by (2) with $\alpha_k \rightarrow 0$ and $\sum_{k=0}^{\infty} \alpha_k = \infty$. If $\sup_{k \in \mathbb{N}} \|x_k\| < \infty$ (which always holds if f is coercive for small enough step sizes),*

$$\begin{aligned} \lim_{k \rightarrow \infty} \text{dist}(f(x_k), \text{vcrit}_\epsilon f) &= 0, \\ \limsup_{k \rightarrow \infty} \text{dist}(x_k, \text{crit } f) &\leq C\epsilon^\rho. \end{aligned}$$

The boundedness condition always holds for small enough step sizes if f is coercive.

Proof : By assumption, there is $M > 0$ such that for all $k \in \mathbb{N}$, $\|x_k\| \leq M$. Denote by L_f the set of limit points of $f(x_k)$, it is an interval since $\alpha_k \rightarrow 0$. We first prove the first statement which is equivalent to the fact that L_f does not contain any value $l \notin \text{vcrit}_\epsilon f$. Suppose toward a contradiction that this is the case, then there exists a value $l \in (\text{vcrit}_\epsilon f)^c \cap L_f$. For such a value l , let $\eta > 0$ be given by Lemma 11 ($\text{vcrit}_\epsilon f$ is closed by Assumption 2). Recall that $(\alpha_k)_{k \in \mathbb{N}}$ is vanishing so that either $\liminf_{k \rightarrow \infty} f(x_k) > l + 2\eta$ or $\limsup_{k \rightarrow \infty} f(x_k) < l - 2\eta$, which is in contradiction with $l \in L_f$.

Denote by L the set of accumulation points of the sequence $(x_k)_{k \in \mathbb{N}}$. By the previous statement we have $f(L) \subset \text{vcrit}_\epsilon f$. By [10, Theorem 4.2 and 4.3], L is an internally chain transitive set. In particular, L is invariant for the inclusion (4) by [10, Lemma 3.5]. This means that for any $\bar{x} \in L$, there is a solution x to (4), such that $x(0) = \bar{x}$ and $x(\mathbb{R}) \in L$. The second statement follows by Corollary 1. \square

4.4 Constant step sizes

We start this section with a general result on constant step size discretization which is of independent interest.

Lemma 13 (Limits of accumulation points are invariant). *Let $M > 0$ be a bound and $x_0 \in \mathbb{R}^p$ be fixed. Under Assumption 3, for any $s > 0$, denote by $L(s)$ the set of accumulation points of sequences satisfying $x_{k+1}(s) = x_k(s) + sh(x_k(s))$ for each $k \in \mathbb{N}$, where h is a selection in Z . Assume that $L(s)$ is nonempty and bounded by M for all s small enough. Set*

$$L = \bigcap_{\alpha > 0} \text{cl} \left\{ \bigcup_{0 < s \leq \alpha} L(s) \right\}.$$

Then the set L is nonempty and invariant with respect to the flow induced by Z in the sense that for any $\bar{x} \in L$ there exists a solution to the differential inclusion $\dot{x}(t) \in Z(x(t))$ for almost every $t \in \mathbb{R}$ such that $x(\mathbb{R}) \subset L$ and $x(0) = \bar{x}$.

Proof : The assumption that $L(s)$ is bounded by a fixed M for small enough s ensures that all considered sequences are bounded and we may restrict all the asymptotics to happen in a fixed bounded set (for example of size $2M$), in particular, Z will be bounded by ζ on this set. This ensures that L is nonempty and this that all considered objects remain in a bounded set.

Let $\bar{x} \in L$. For a fixed $\alpha > 0$, $\bar{x} \in \text{cl} \cup_{s \leq \alpha} L(s)$ means that for any $e > 0$, there is $s \leq \alpha$ such that $\text{dist}(\bar{x}, L(s)) \leq e$. Therefore if $\bar{x} \in L$, we deduce that there exists a sequence $(s_j)_{j \in \mathbb{N}}$ which tends to 0 and a sequence $\bar{x}_j \in L(s_j)$ which tends to \bar{x} as $j \rightarrow \infty$.

Fix an arbitrary $T > 0$ and $j \in \mathbb{N}$. We set $K_j = \lceil T/s_j \rceil$ and consider for $k \geq K_j$, $X_{j,k} = (x_i(s_j))_{i=k-K_j}^{i=k+K_j}$. Up to a subsequence, as $k \rightarrow \infty$, $X_{j,k}$ converges to \bar{X}_j which contains $2K_j + 1$ accumulation points in $L(s_j)$, the central one chosen to be \bar{x}_j . The affine interpolation of the (ordered) collection of points in \bar{X}_j is denoted $\bar{\gamma}_j: [-T, T] \rightarrow \mathbb{R}^p$ with $\bar{\gamma}_j(0) = \bar{x}_j$.

This construction can be repeated for all $j \in \mathbb{N}$ and using Arzelà-Ascoli, there is a subsequence of $(\bar{\gamma}_j)_{j \in \mathbb{N}}$ which converges uniformly to $\bar{\gamma}: [-T, T] \rightarrow \mathbb{R}^p$. We have $\bar{\gamma}([-T, T]) \subset L$. Indeed let $t \in [-T, T]$. For any $e > 0$, we have $\|\bar{\gamma}_j(t) - \bar{\gamma}(t)\| \leq e$ for j high enough. By construction, $\bar{\gamma}_j(s_j \lceil t/s_j \rceil) \in L(s_j)$. Since $s_j \lceil t/s_j \rceil$ converges to t and $\bar{\gamma}_j$ is Lipschitz with same constant for all j , then $\|\bar{\gamma}(t) - \bar{\gamma}_j(s_j \lceil t/s_j \rceil)\| \leq 2e$ for all j high enough. Since e was arbitrary, this means $\bar{\gamma}(t) \in L$.

We claim that $\bar{\gamma}$ is a solution to (4).

Indeed, for any $j \in \mathbb{N}$, there is $k_j \in \mathbb{N}$ such that $\|X_{j,k_j} - \bar{X}_j\| \leq 1/j$. Let $\gamma_j: [-T, T] \rightarrow \mathbb{R}^p$ be the affine interpolation of X_{j,k_j} . Up to the Arzelà-Ascoli subsequence, $\gamma_j \rightarrow \bar{\gamma}$ uniformly on $[-T, T]$. Furthermore, for each j

$$\int_{-T}^T \text{dist}((\gamma_j(t), \dot{\gamma}_j(t)), \text{graph}[Z]) dt \leq 2Ts_j\zeta$$

By Lemma 8, this shows that $\bar{\gamma}$ is a solution to (4).

The function $\bar{\gamma}$ may be extended to $[-2T, 2T]$, $[-3T, 3T]$, ... by taking further subsequences for each j and further Arzelà-Ascoli subsequences. The resulting function is defined on \mathbb{R} and has to be a solution to (4). This concludes the proof. \square

The main result stated in Theorem 2 is a consequence of the following result and Lemma 5.

Theorem 14 (Convergence for biased subgradient with constant steps). *Under Assumptions 2, let C, ρ be given by Lemma 6 and $\epsilon < \bar{\epsilon}$. Set for all $\alpha > 0$, $(x_k(\alpha))_{k \in \mathbb{N}}$, as given by (2) with $\alpha_k = \alpha$, for all $k \in \mathbb{N}$. Then if $\limsup_{\alpha \rightarrow 0} \sup_{k \rightarrow \infty} \|x_k(\alpha)\| < \infty$ (which always holds if f is coercive),*

$$\begin{aligned} \lim_{\alpha \rightarrow 0} \limsup_{k \rightarrow \infty} \text{dist}(f(x_k(\alpha)), \text{vcrit}_\epsilon f) &= 0, \\ \limsup_{\alpha \rightarrow 0} \limsup_{k \rightarrow \infty} \text{dist}(x_k(\alpha), \text{crit } f) &\leq C\epsilon^\rho. \end{aligned}$$

Proof : By assumption, there is $M > 0$ such that for small enough α , $\|x_k(\alpha)\| \leq M$ for all $k \in \mathbb{N}$ large enough.

For each s denote by $L(s)$ the set of limit points of $x_k(s)$, which is closed. We set $L = \bigcap_{\alpha > 0} \text{cl } \bigcup_{s \leq \alpha} L(s)$ which is closed as an intersection of closed set, and bounded by M . Intuitively L is the set of accumulation points of accumulation points of $(x_k(\alpha))_{k \in \mathbb{N}}$ as $\alpha \rightarrow 0$. By Lemma 13, L is invariant. We will show that $f(L) \subset \text{vcrit}_\epsilon f$ from which the first statement follows. The second is then a consequence of Corollary 1.

For any $l \notin \text{vcrit}_\epsilon f$, since $\text{vcrit}_\epsilon f$ is closed by Assumption 2, Lemma 11 ensures that there is $\bar{\alpha}$ and η , such that for any $s \leq \bar{\alpha}$, $[l - 2\eta, l + 2\eta] \cap f(L(s)) = \emptyset$. We deduce that

$$\begin{aligned} [l - 2\eta, l + 2\eta] \cap f(\bigcup_{s \leq \bar{\alpha}} L(s)) &= \emptyset \\ [l - \eta, l + \eta] \cap f(\text{cl } \bigcup_{s \leq \bar{\alpha}} L(s)) &= \emptyset \quad (\text{use continuity}). \end{aligned}$$

Whence we see that for any $l \notin \text{vcrit}_\epsilon f$, $l \notin f(\bigcap_{\alpha > 0} \text{cl } \bigcup_{s \leq \alpha} L(s))$, and therefore $f(L) \subset \text{vcrit}_\epsilon f$ through complementation. This proves the first claim. \square

5 The convex case

Proof of Theorem 3: Consider a sequence of step sizes $(\alpha_k)_{k \in \mathbb{N}}$ and a iterates $(x_k)_{k \in \mathbb{N}}$ such that for all $k \in \mathbb{N}$,

$$x_{k+1} = x_k - \alpha_k(v_k + b_k),$$

for some $v_k \in \partial^c f(x_k)$ and $\|b_k\| \leq \epsilon$. We have for any $k \in \mathbb{N}$ and any $x^* \in X^*$,

$$\begin{aligned} \frac{1}{2}\|x_{k+1} - x^*\|_2^2 &= \frac{1}{2}\|x_k - \alpha_k(v_k + b_k) - x^*\|_2^2 \\ &= \frac{1}{2}\|x_k - x^*\|_2^2 + \alpha_k(v_k + b_k)^T(x^* - x_k) + \frac{\alpha_k^2}{2}\|v_k + b_k\|_2^2 \\ &\leq \frac{1}{2}\|x_k - x^*\|_2^2 - \alpha_k(f(x_k) - f^*) + \alpha_k\epsilon\|x_k - x^*\| + \frac{\alpha_k^2}{2}(L + \epsilon)^2. \end{aligned}$$

We deduce, using the error bound (3), that

$$\begin{aligned} &\frac{1}{2}\text{dist}(x_{k+1}, X^*)^2 \tag{15} \\ &\leq \frac{1}{2}\text{dist}(x_k, X^*)^2 - \alpha_k(f(x_k) - f^*) + \alpha_k\epsilon\text{dist}(x_k, X^*) + \frac{\alpha_k^2}{2}(L + \epsilon)^2 \\ &\leq \frac{1}{2}\text{dist}(x_k, X^*)^2 + \alpha_k \left(\epsilon \frac{c}{2} ((f(x_k) - f^*)^a + (f(x_k) - f^*)) - (f(x_k) - f^*) \right) + \frac{\alpha_k^2}{2}(L + \epsilon)^2. \end{aligned}$$

If $a = 1$: This means that we have a sharp function, then, as long as $\epsilon c < 1$, we obtain the same global rate as the classical subgradient algorithm, modulo a constant $(1 - \epsilon c)$, and all accumulation points in the argmin. Indeed, summing the previous inequality from $i = 0$ to k leads to

$$(1 - \epsilon c) \frac{\sum_{i=0}^k \alpha_i (f(x_i) - f^*)}{\sum_{i=0}^k \alpha_i} \leq \frac{\|x_0 - x^*\|_2^2 + (L + \epsilon)^2 \sum_{i=0}^k \alpha_i^2}{\sum_{i=0}^k \alpha_i}.$$

We remark that this is exactly the claimed formula for $a = 1$.

If $a < 1$: Then we can use the following lemma

Lemma 15. *Set $g(\delta) = s\delta^t - \delta$ for some $t \in (0, 1)$ and $s > 0$, then for all $\delta \in \mathbb{R}_+$*

$$g(\delta) \leq -(1 - t)(\delta - s^{\frac{1}{1-t}}).$$

Proof : The function g is concave on \mathbb{R}_+ , set $\delta_0 = s^{\frac{1}{1-t}}$, we have $g(\delta_0) = 0$ and $g'(\delta_0) = t - 1 < 0$ and the result follows by concavity. \square

We obtain setting $\delta = f(x_k) - f^*$ applying Lemma 15 to (15)

$$\begin{aligned} &\frac{1}{2}\text{dist}(x_{k+1}, X^*)^2 - \frac{1}{2}\text{dist}(x_k, X^*)^2 \\ &\leq \frac{\alpha_k}{2} \left(-(1 - a) \left((f(x_k) - f^*) - (\epsilon c)^{\frac{1}{1-a}} \right) + (\epsilon c - 1)(f(x_k) - f^*) \right) + \frac{\alpha_k^2}{2}(L + \epsilon)^2 \\ &= \frac{\alpha_k}{2} \left((\epsilon c + a - 2)(f(x_k) - f^*) + (1 - a)(\epsilon c)^{\frac{1}{1-a}} \right) + \frac{\alpha_k^2}{2}(L + \epsilon)^2 \end{aligned}$$

We deduce by summation

$$(2 - a - \epsilon c) \frac{\sum_{i=0}^k \alpha_i (f(x_i) - f^*)}{\sum_{i=0}^k \alpha_i} \leq (1 - a)(\epsilon c)^{\frac{1}{1-a}} + \frac{\|x_0 - x_*\|^2 + (L + \epsilon)^2 \sum_{i=0}^k \alpha_i^2}{\sum_{i=0}^k \alpha_i}.$$

□

Acknowledgements The authors acknowledge the support of Centre Lagrange. JB, TL, EP thank AI Interdisciplinary Institute ANITI funding, through the French “Investments for the Future – PIA3” program under the grant agreement ANR-19-PI3A0004, Air Force Office of Scientific Research, Air Force Material Command, USAF, under grant numbers FA8655-22-1-7012, ANR Chess, grant ANR-17-EURE-0010, ANR Regulia.

References

- [1] Compressed sensing. *IEEE Transactions on information theory*, 52(4):1289–1306, 2006.
- [2] C.D. Aliprantis and K.C. Border. *Infinite Dimensional Analysis*. Springer Berlin, Heidelberg, 2006.
- [3] Brandon Amos and J Zico Kolter. Optnet: Differentiable optimization as a layer in neural networks. In *International Conference on Machine Learning*, pages 136–145. PMLR, 2017.
- [4] F.M. Arscott and A.F. Filippov. *Differential Equations with Discontinuous Righthand Sides: Control Systems*. Mathematics and its Applications. Springer Netherlands, 1988.
- [5] Timo Aspelmeier, C Charitha, and D Russell Luke. Local linear convergence of the admm/douglas–rachford algorithms without strong convexity and application to statistical imaging. *SIAM Journal on Imaging Sciences*, 9(2):842–868, 2016.
- [6] J-P Aubin and Arrigo Cellina. *Differential inclusions: set-valued maps and viability theory*, volume 264. Springer Science & Business Media, 1983.
- [7] Francis Bach, Rodolphe Jenatton, Julien Mairal, and Guillaume Obozinski. Convex optimization with sparsity-inducing norms. *Optimization for Machine Learning*, pages 19–49, 2011.
- [8] Aharon Ben-Tal, Laurent El Ghaoui, and Arkadi Nemirovski. *Robust optimization*, volume 28. Princeton university press, 2009.
- [9] Michel Benaïm. Recursive algorithms, urn processes and chaining number of chain recurrent sets. *Ergodic Theory and Dynamical Systems*, 18(1):53–87, 1998.
- [10] Michel Benaïm, Josef Hofbauer, and Sylvain Sorin. Stochastic approximations and differential inclusions. *SIAM Journal on Control and Optimization*, 44(1):328–348, 2005.

- [11] Pascal Bianchi, Walid Hachem, and Adil Salim. Constant step stochastic approximations involving differential inclusions: stability, long-run convergence and applications. *Stochastics*, 91(2):288–320, 2019.
- [12] Pascal Bianchi, Walid Hachem, and Sholom Schechtman. Convergence of constant step stochastic gradient descent for non-smooth non-convex functions. *Set-Valued and Variational Analysis*, 30(3):1117–1147, 2022.
- [13] Mathieu Blondel, Quentin Berthet, Marco Cuturi, Roy Frostig, Stephan Hoyer, Felipe Llinares-López, Fabian Pedregosa, and Jean-Philippe Vert. Efficient and modular implicit differentiation. *Advances in neural information processing systems*, 35:5230–5242, 2022.
- [14] Jérôme Bolte, Aris Daniilidis, and Adrian Lewis. The Łojasiewicz inequality for non-smooth subanalytic functions with applications to subgradient dynamical systems. *SIAM Journal on Optimization*, 17(4):1205–1223, 2007.
- [15] Jérôme Bolte, Aris Daniilidis, Adrian Lewis, and Masahiro Shiota. Clarke subgradients of stratifiable functions. *SIAM Journal on Optimization*, 18(2):556–572, 2007.
- [16] Jérôme Bolte, Tam Le, and Edouard Pauwels. Subgradient sampling for nonsmooth nonconvex minimization. *SIAM Journal on Optimization*, 33(4):2542–2569, 2023.
- [17] Jérôme Bolte and Edouard Pauwels. Conservative set valued fields, automatic differentiation, stochastic gradient methods and deep learning. *Mathematical Programming*, 188(1):19–51, 2021.
- [18] Jérôme Bolte, Edouard Pauwels, and Rodolfo Rios-Zertuche. Long term dynamics of the subgradient method for lipschitz path differentiable functions. *Journal of the European Mathematical Society*, 2022.
- [19] Jérôme Bolte, Edouard Pauwels, and Samuel Vaiter. One-step differentiation of iterative algorithms. *Advances in Neural Information Processing Systems*, 36, 2023.
- [20] Frank H Clarke. *Optimization and nonsmooth analysis*. SIAM, 1983.
- [21] Jeremy Cohen, Simran Kaur, Yuanzhi Li, J Zico Kolter, and Ameet Talwalkar. Gradient descent on neural networks typically occurs at the edge of stability. In *International Conference on Learning Representations*, 2020.
- [22] Patrick L Combettes and Jean-Christophe Pesquet. Proximal splitting methods in signal processing. In *Fixed-point algorithms for inverse problems in science and engineering*, pages 185–212. Springer, 2011.
- [23] Michel Coste. *An introduction to ϕ -minimal geometry*. Istituti editoriali e poligrafici internazionali Pisa, 2000.
- [24] Damek Davis, Dmitriy Drusvyatskiy, Sham Kakade, and Jason D Lee. Stochastic subgradient method converges on tame functions. *Foundations of computational mathematics*, 20(1):119–154, 2020.

- [25] Christopher De Sa, Matthew Feldman, Christopher Ré, and Kunle Olukotun. Understanding and optimizing asynchronous low-precision stochastic gradient descent. In *Proceedings of the 44th annual international symposium on computer architecture*, pages 561–574, 2017.
- [26] Aymeric Dieuleveut, Gersende Fort, Eric Moulines, and Hoi-To Wai. Stochastic approximation beyond gradient for signal processing and machine learning. *IEEE Transactions on Signal Processing*, 2023.
- [27] A Doucet and V Tadic. Asymptotic bias of stochastic gradient search. *Annals of Applied Probability*, 27(6), 2017.
- [28] Yu M Ermol’ev and VI Norkin. Stochastic generalized gradient method for nonconvex nonsmooth stochastic optimization. *Cybernetics and Systems Analysis*, 34(2):196–215, 1998.
- [29] Yuri M. Ermoliev. Stochastic quasigradient methods and their application to system optimization. *Stochastics*, 9:1–36, 1983.
- [30] William Fedus, Barret Zoph, and Noam Shazeer. Switch transformers: Scaling to trillion parameter models with simple and efficient sparsity. *Journal of Machine Learning Research*, 23(120):1–39, 2022.
- [31] Jean-Claude Fort and Gilles Pages. Asymptotic behavior of a markovian stochastic algorithm with constant step. *SIAM journal on control and optimization*, 37(5):1456–1482, 1999.
- [32] Saeed Ghadimi and Guanghui Lan. Stochastic first-and zeroth-order methods for nonconvex stochastic programming. *SIAM Journal on Optimization*, 23(4):2341–2368, 2013.
- [33] Saeed Ghadimi, Guanghui Lan, and Hongchao Zhang. Mini-batch stochastic approximation methods for nonconvex stochastic composite optimization. *Mathematical Programming*, 155(1):267–305, 2016.
- [34] Andreas Griewank and Christèle Faure. Piggyback differentiation and optimization. In *Large-scale PDE-constrained optimization*, pages 148–164. Springer, 2003.
- [35] Robert Hesse and D Russell Luke. Nonconvex notions of regularity and convergence of fundamental algorithms for feasibility problems. *SIAM Journal on Optimization*, 23(4):2397–2419, 2013.
- [36] Woonghee Tim Huh and Paat Rusmevichientong. Online sequential optimization with biased gradients: theory and applications to censored demand. *INFORMS Journal on Computing*, 26(1):150–159, 2014.
- [37] Cédric Josz and Lexiao Lai. Global stability of first-order methods for coercive tame functions. *Mathematical Programming*, pages 1–26, 2023.

- [38] Guanghui Lan and Zhiqiang Zhou. Algorithms for stochastic optimization with function or expectation constraints. *Computational Optimization and Applications*, 76(2):461–498, 2020.
- [39] Yann LeCun, Bernhard Boser, John S Denker, Donnie Henderson, Richard E Howard, Wayne Hubbard, and Lawrence D Jackel. Backpropagation applied to handwritten zip code recognition. *Neural computation*, 1(4):541–551, 1989.
- [40] Jae Hyoung Lee and Tien-Son Pham. Openness, hölder metric regularity, and hölder continuity properties of semialgebraic set-valued maps. *SIAM Journal on Optimization*, 32(1):56–74, 2022.
- [41] Lennart Ljung. Analysis of recursive stochastic algorithms. *IEEE transactions on automatic control*, 22(4):551–575, 1977.
- [42] Ilya Loshchilov and Frank Hutter. SGDR: Stochastic gradient descent with warm restarts. In *International Conference on Learning Representations*, 2017.
- [43] D Russell Luke, Marc Teboulle, and Nguyen H Thao. Necessary conditions for linear convergence of iterated expansive, set-valued mappings. *Mathematical Programming*, 180:1–31, 2020.
- [44] Marko Mäkelä. Survey of bundle methods for nonsmooth optimization. *Optimization methods and software*, 17(1):1–29, 2002.
- [45] Konstantin Mishchenko, Eduard Gorbunov, Martin Takáč, and Peter Richtárik. Distributed learning with compressed gradient differences. *arXiv preprint arXiv:1901.09269*, 2019.
- [46] Angelia Nedić and Dimitri P Bertsekas. The effect of deterministic noise in subgradient methods. *Mathematical programming*, 125(1):75–99, 2010.
- [47] Arkadi Nemirovski, Anatoli Juditsky, Guanghui Lan, and Alexander Shapiro. Robust stochastic approximation approach to stochastic programming. *SIAM Journal on optimization*, 19(4):1574–1609, 2009.
- [48] Arkadij Semenovič Nemirovskij and David Borisovich Yudin. Problem complexity and method efficiency in optimization. 1983.
- [49] Nhu Nguyen and George Yin. Stochastic approximation with discontinuous dynamics, differential inclusions, and applications. *The Annals of Applied Probability*, 33(1):780–823, 2023.
- [50] V. I. Norkin. Nonlocal minimization algorithms of nondifferentiable functions. *Cybernetics*, 14(5):704–707, Sep 1978.
- [51] Alec Radford, Karthik Narasimhan, Tim Salimans, Ilya Sutskever, et al. Improving language understanding by generative pre-training. 2018.

- [52] Arunselvan Ramaswamy and Shalabh Bhatnagar. Analysis of gradient descent methods with nondiminishing bounded errors. *IEEE Transactions on Automatic Control*, 63(5):1465–1471, 2017.
- [53] Grégory Roth and William H Sandholm. Stochastic approximations with constant step size and differential inclusions. *SIAM Journal on Control and Optimization*, 51(1):525–555, 2013.
- [54] D Russell Luke, Nguyen H Thao, and Matthew K Tam. Quantitative convergence analysis of iterated expansive, set-valued mappings. *Mathematics of Operations Research*, 43(4):1143–1176, 2018.
- [55] Naum Zuselevich Shor. *Minimization methods for non-differentiable functions*, volume 3. Springer Science & Business Media, 2012.
- [56] Mikhail V Solodov and SK Zavriev. Error stability properties of generalized gradient-type algorithms. *Journal of Optimization Theory and Applications*, 98:663–680, 1998.
- [57] James C Spall. Stochastic optimization. *Handbook of computational statistics: Concepts and methods*, pages 173–201, 2012.
- [58] Vladislav B Tadić and Arnaud Doucet. Asymptotic bias of stochastic gradient search. In *2011 50th IEEE Conference on Decision and Control and European Control Conference*, pages 722–727. IEEE, 2011.
- [59] Robert Tibshirani. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 58(1):267–288, 1996.
- [60] Lou van den Dries. *Tame topology and o-minimal structures*, volume 248. Cambridge university press, 1998.
- [61] Jianqiao Wangni, Jialei Wang, Ji Liu, and Tong Zhang. Gradient sparsification for communication-efficient distributed optimization. *Advances in Neural Information Processing Systems*, 31, 2018.