



HAL
open science

On the link between Dempster's rule of combination of evidence and fuzzy set intersection

Henri Prade

► **To cite this version:**

Henri Prade. On the link between Dempster's rule of combination of evidence and fuzzy set intersection. Bulletin pour les sous-ensembles flous et leurs applications, 1981, 8, pp.60–64. hal-04563922

HAL Id: hal-04563922

<https://hal.science/hal-04563922>

Submitted on 30 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On the link between Dempster's rule of combination of evidence and
fuzzy set intersection

Henri Prade

Laboratoire Langages & Systèmes Informatiques
Université Paul Sabatier
118 route de Narbonne - 31062 Toulouse Cedex

0. First, basic results concerning the representation of belief or plausibility functions are recalled together with the expression of Dempster's rule of combination of evidence in terms of basic probability assignments and in terms of commonality functions. Then, the normalization of belief and plausibility functions is **discussed**. Lastly, in case of consonant belief functions, it is stressed that Dempster's rule of combination can be interpreted as a fuzzy set intersection based on product, taking care of the normalization question.

1. Let S be a finite set : $S = \{s_1, \dots, s_n\}$. Let $\mathcal{P}(S)$ denote the set of subsets of S . Shafer [6] has shown that any belief (or credibility) function Bel is uniquely defined through the specification of a mapping m from $\mathcal{P}(S)$ to $[0, 1]$ called 'basic probability assignment', satisfying the conditions

$$m(\emptyset) = 0 \quad (1)$$

$$\text{and} \quad \sum_{A \subseteq X} m(A) = 1 \quad (2)$$

then, we have

$$\forall A \subseteq S, \quad Bel(A) = \sum_{B \subseteq A} m(B) \quad (3)$$

Conversely, m is obtained from Bel by

$$\forall A \subseteq S, \quad m(A) = \sum_{B \subseteq A} (-1)^{|A-B|} Bel(B) \quad (4)$$

where $|A|$ denotes the cardinality of the set A .

The set function Pl defined by duality

$$\forall A \subseteq S, \quad Pl(A) = 1 - Bel(\bar{A}) \quad (5)$$

is called a plausibility function (\bar{A} denotes the complement of A). Note that

$$\forall A \subseteq S, \quad Pl(A) \geq Bel(A)$$

We have

$$\forall A \subseteq S, \quad Pl(A) = \sum_{B \cap A \neq \emptyset} m(B) \quad (6)$$

[N.B.: Originally, Dempster [1] introduced plausibility and belief functions (under the names 'upper and lower probabilities') by considering a multiva_

lued mapping Γ from a probability space $(X, \mathcal{P}(X), \mu)$ to a set S , and the following subsets of X induced by Γ : $A^* = \{x \in X, \Gamma(x) \cap A \neq \emptyset\}$ and $A_* = \{x \in X, \Gamma(x) \neq \emptyset, \Gamma(x) \subseteq A\}$ where $A \subseteq S$; in this framework, $Pl(A) = \mu(A^*)/\mu(S^*)$ and $Bel(A) = \mu(A_*)/\mu(S_*)$ where $S^* = S_* = \{x \in X, \Gamma(x) \neq \emptyset\}$. Thus, Pl and Bel are completely determined by the 2^n quantities $\mu(\{x \in X, \Gamma(x) = B\})$ where B is any subset of S ; note that the subsets $\{x \in X, \Gamma(x) = B\}$ determine a partition of X , thus $\sum_{B \subseteq S} \mu(\{x \in X, \Gamma(x) = B\}) = 1$; as pointed out by Goodman [3] and Nguyen [5], these 2^n quantities determine a random set on S . We have

$$\forall B \subseteq S, B \neq \emptyset, \quad m(B) = \frac{\mu(\{x \in X, \Gamma(x) = B\})}{1 - \mu(\{x \in X, \Gamma(x) = \emptyset\})} \quad (7)$$

The set $\mathcal{F} = \{B \subseteq S, m(B) > 0\}$ may be thought as the set of possible localizations of the truth, probabilistically weighted by m . If Bel is a probability measure, \mathcal{F} contains only singletons: the possible localizations of the truth are then precise. The elements of \mathcal{F} are called 'focal elements'.

A so-called commonality function Q can be defined from m by

$$\forall A \subseteq S, \quad Q(A) = \sum_{\substack{B \subseteq S \\ A \subseteq B}} m(B) \quad (8)$$

Note that $Q(\{s\}) = Pl(\{s\})$; $Q(\emptyset) = 1$. Bel can be expressed directly in terms of Q :

$$\forall A \subseteq S, \quad Bel(A) = \sum_{B \subseteq \bar{A}} (-1)^{|B|} Q(B) \quad (9)$$

Thus, Q provides another way of specifying a belief or a plausibility function.

Given two basic probability assignments m_1 and m_2 , Dempster's rule ([1], [6]) enables to build a basic probability assignment m representing the combination of the two evidences. We have

$$m(\emptyset) = 0; \quad \forall A \subseteq S, A \neq \emptyset \quad m(A) = \frac{\sum_{\substack{i,j \\ A_i \cap B_j = A}} m_1(A_i) \cdot m_2(B_j)}{1 - \sum_{\substack{i,j \\ A_i \cap B_j = \emptyset}} m_1(A_i) \cdot m_2(B_j)} \quad (10)$$

It is easy to check that $\sum_{A \subseteq S} m(A) = 1$. Thus, Dempster's rule corresponds to the intersection of two statistically independent random sets of the same base space, after renormalization.

N.B.: In Dempster's original framework, the combination of $((X_1, \mathcal{P}(X_1), \mu_1), \Gamma_1)$ and $((X_2, \mathcal{P}(X_2), \mu_2), \Gamma_2)$ yields $((X, \mathcal{P}(X), \mu), \Gamma)$ defined by $X = X_1 \times X_2$, $\mu = \mu_1 \times \mu_2$ and $\Gamma(x) = \Gamma_1(x) \cap \Gamma_2(x)$.

As pointed out by Dempster [1] and Shafer [6], the combination is more easily expressed in terms of commonality functions since we have

$$\forall A \subseteq S, \quad Q(A) = \frac{Q_1(A) \cdot Q_2(A)}{K} \quad (11)$$

where K denotes the denominator of fraction (10).

2. Zadeh [8] has questioned the validity of the normalization in (10). It is worth noticing that the problem of normalization vanishes as soon as we

admit that a basic probability assigned to the empty set \emptyset may be non-zero and we define the "intersection" of m_1 and m_2 by

$$m^*(A) = \sum_{i,j} m_1(A_i) \cdot m_2(B_j) \quad (12)$$

It is easy to check we have $\sum_{A \subseteq S} m^*(A) = 1$, even if $m_1(\emptyset) \neq 0$ or $m_2(\emptyset) \neq 0$. From now, the normalization condition of a basic probability assignment will be only given by (2); (1) is dropped. Keeping definition (8) unchanged, (11) becomes

$$\forall A \subseteq S, \quad Q(A) = Q_1(A) \cdot Q_2(A) \quad (13)$$

Noticing that (3) can be written

$$\forall A \subseteq S, \quad \text{Bel}(A) = \sum_{B \subseteq S} m(B) \cdot [\inf_{s \in B} \chi_A(s)] \quad (14)$$

where χ_A is the characteristic function of the subset A . Thus, $\text{Bel}(A)$ appears to be the expectation that the truth lies certainly in A , computed from the set of probabilistic weights $\{m(B), B \subseteq S \text{ and } B \neq \emptyset\}$ attached to the possible localizations of the truth since $\inf \chi(s) = 1$ if $B \subseteq A$ and is zero otherwise.

According to this interpretation in terms of expectation, when $m(\emptyset)$ may be non-zero, (3) must be modified into

$$\forall A \subseteq S, \quad \text{Bel}(A) = \sum_{\substack{B \subseteq A \\ B \neq \emptyset}} m(B) \quad (15)$$

Similarly, (6) can be written

$$\forall A \subseteq S, \quad \text{Pl}(A) = \sum_{B \subseteq S} m(B) \cdot (\sup_{s \in B} \chi_A(s)) \quad (16)$$

Thus, $\text{Pl}(A)$ is the expectation that the truth lies possibly in A . (6) remains unchanged, but (5) becomes

$$\forall A \subseteq S, \quad \text{Pl}(A) + \text{Bel}(\bar{A}) = 1 - m(\emptyset) \quad (17)$$

However, we keep $\forall A \subseteq S, \text{Pl}(A) \geq \text{Bel}(A)$. As soon as $m(\emptyset) \neq 0$, Bel and Pl are no longer normalized:

$$\text{Bel}(S) < 1 \quad ; \quad \text{Pl}(S) < 1 \quad (18)$$

We still have $\text{Bel}(\emptyset)$ and $\text{Pl}(\emptyset)$ equal to zero.

If the empty subset of S has a non-zero probabilistic weight, it means that a possible localization of the truth is outside of S , then it may seem natural that the credibility and the plausibility that the truth lies in S are strictly less than 1; then, (17) means that 'A non plausible' does not entail that \bar{A} is completely credible.

[N.B.: (7) is modified into $\forall B \subseteq S, m(B) = \mu(\{x \in X, \Gamma(x) = B\})$; i.e. $m(\emptyset) = 0$ if and only if the set of points in X which are not mapped to S by Γ has a measure equal to zero.

However, we have from (13)

$$\forall s \in S, \quad Q(\{s\}) = Q_1(\{s\}) \cdot Q_2(\{s\}) \quad (25)$$

i.e. since $Q_i(\{s\}) = Pl_i(\{s\}) = \pi_i(s)$, $i=1, 2$, (25) expresses that $s \mapsto Q(\{s\})$ is the membership function of a fuzzy set which is the intersection (performed using product) of the fuzzy sets whose membership functions are respectively π_1 and π_2 .

Consonance is lost and moreover $\{Q(\{s\}), s \in S\}$ does not represent all the information. Indeed the number of unknowns is generally greater than the number of equations in the following

$$\begin{cases} Q(\{s\}) = \sum_{\substack{A \in S \\ A \ni s}} m(A) \\ 1 = \sum_{A \in S} m(A) \end{cases} \quad (26)$$

4. The two intended purposes of this short note were

1) to question the normalization condition $m(\emptyset) = 0$ of a basic probability assignment m and to study its implications for Dempster's rule of combination especially.

2) to show in what way Dempster's rule of combination is related to fuzzy set intersection based on product. In that perspective a natural question is raised: What combination rule would be related to fuzzy set intersection based on min operation (for instance)? Normalization of fuzzy sets and normalization of basic probability assignments appear to be clearly related.

References:

- [1] Dempster, A.P. Upper and lower probabilities induced by a multivalued mapping. Ann. Math. Statist. Vol. 38 pp 325-339. 1967.
- [2] Dubois, D., Prade, H. On several representations of an uncertain body of evidence. Submitted.
- [3] Goodman, I.R. Identification of fuzzy sets with a class of canonically induced random sets. Proc. 19th IEEE Conference on Decision and Control, Albuquerque, New Mexico, Dec. 10-12, 1980, pp 353-357.
- [4] Kampé de Fériet, J. Une interprétation des mesures de plausibilité et de crédibilité au sens de G. Shafer et de la fonction d'appartenance définissant un ensemble flou de L. Zadeh. Pub. IRMA - Lille (France) Vol. 2, Fasc. 6 n°2, pp II-01, II-22, 1980.
- [5] Nguyen, H.T. On random sets and belief functions. J. Math. Anal. & Appl. Vol. 65 pp 531-542. 1978.
- [6] Shafer, G. A mathematical theory of evidence. Princeton University Press 1976.
- [7] Zadeh, L.A. Fuzzy sets as a basis for a theory of possibility. Fuzzy Sets & Systems Vol. 1 n°1 pp 3-28. 1978.
- [8] Zadeh, L.A. On the validity of Dempster's rule of combination of evidence. Memo UCB/ERL n°79/24, Univ. of California, Berkeley 1979.