



HAL
open science

Spotting method as a high throughput alternative to the conventional spread plating method conventional spread plating method for bacterial time kill experiments

Vincent Aranzana-Climent, Laure Prouvensier, Tom Collet, Sandrine Marchand, Nicolas Grégoire, Julien M Buyck

► To cite this version:

Vincent Aranzana-Climent, Laure Prouvensier, Tom Collet, Sandrine Marchand, Nicolas Grégoire, et al.. Spotting method as a high throughput alternative to the conventional spread plating method conventional spread plating method for bacterial time kill experiments. 34th European Congress of Clinical Microbiology and Infectious Diseases (ECCMID), Apr 2024, Barcelona, Spain. hal-04571076

HAL Id: hal-04571076

<https://hal.science/hal-04571076>

Submitted on 7 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Spotting method as a high throughput alternative to the conventional spread plating method for bacterial time-kill experiments.

V. Aranzana-Climent¹, L. Prouvensier^{1,2}, T. Collet¹, S. Marchand^{1,2}, N. Grégoire^{1,2}, J.M. Buyck¹

Contact : vincent.aranzana.climent@univ-poitiers.fr

¹Université de Poitiers, INSERM U1070, France

²CHU de Poitiers, Laboratoire de Toxicologie et de Pharmacocinétique, France

Introduction

In vitro time-kill experiments are commonly performed to obtain information about the dynamics of a drug antimicrobial activity. It is however labor intensive due to the conventional spread plating method used to count bacteria. In this work we compare a spotting method with the conventional spread plating method.

Materials and Methods




Figure 1: Experimental protocol.

Correction of colony counts to account for overcrowding

- Colony forming unit counts were converted to bacterial densities after correcting for overcrowding (Figure 2) by the « most probable number » method from Martini *et al.* (1)
- Under this method, for a given measurement, the probability of overcrowding is higher when the number of counted colonies (Figure 3) is close to the maximal number of countable colonies.

Validation of the spotting method

- To validate the spotting method against the conventional plating method we applied correlated bivariate least square regression developed by Francq and Govaerts (2) on \log_{10} transformed bacterial densities after exclusion of measurements below 10^3 CFU/mL (limit of detection of the spotting method)

Mathematical methods and software

- Optimization of the likelihood function of the « most probable number » (1) method was performed using the R function `optim` (3). Correlated bivariate least square regression was performed using the `BivRegBLS` package (4). Data wrangling and graphing was performed with the `tidyverse` package (5)




Figure 2: Illustration of the overcrowding problem.
A. Number of bacteria at time of plating.
B. Number of colonies after incubation

Figure 3: Illustration of how the maximal number of countable colonies was determined.
A. Conventional plating B. Spotting

Results




Figure 4: Bland-Altman plot (n=551).

Table 1: Performance of the two plating methods

	Conventional plating	Spotting
Number of time-kill per week	8-12	48-96
Mean maximal number of colonies per plating	675	7.84
Limit of detection (CFU/mL)	$4 \cdot 10^1$	10^3
Average bias (\log_{10} CFU/mL)		$6.6 \cdot 10^{-4}$
Repeatability standard deviation (\log_{10} CFU/mL)	0.105	0.445

- As seen in Table 1 spotting method was accurate but exhibited lower precision than the conventional plating method.
- The accuracy was confirmed over the whole range of measurements as evidenced by the red regression line on Figure 4.
- The two measurement methods were considered in agreement when the difference was between -1 and +1 \log_{10} CFU/mL (Figure 4 green area). In our experiment they agreed since the 95% prediction interval of the differences (Figure 4 purple lines) lied inside the agreement interval (Figure 4 green area).
- This agreement is qualitatively shown by Figure 5 where typical time-kill profiles obtained by plating (Figure 5A) and by spotting (Figure 5B) are comparable.




Figure 5: Typical time-kill results. A. Conventional plating B. Spotting

Conclusion

The spotting method yields bacterial counting results comparable with the conventional plating method while being significantly higher throughput.

References

- Martini *et al.* [10.1101/2023.05.18.541301](https://doi.org/10.1101/2023.05.18.541301)
- Francq and Govaerts, [10.1002/sim.6872](https://doi.org/10.1002/sim.6872)
- R Core Team (2023) www.R-project.org
- Francq *et al.* [10.1002/sim.8709](https://doi.org/10.1002/sim.8709)
- Wickham *et al.* [10.21105/joss.01686](https://doi.org/10.21105/joss.01686)



This work was supported by the Agence Nationale de recherche as part of its "Investissement d'Avenir" research program grant n°ANR-20-PAMR-0010

34th ECCMID EUROPEAN CONGRESS OF CLINICAL MICROBIOLOGY AND INFECTIOUS DISEASES

Barcelona, Spain
27 – 30 April 2024