



HAL
open science

Bidimensional empirical mode decomposition and dynamic differential microscopy reveals multiscale dynamics of critical fluctuations

Ana Oprisan, Sorinel Oprisan, Carole Lecoutre, Yves Garrabos, Daniel Beysens

► To cite this version:

Ana Oprisan, Sorinel Oprisan, Carole Lecoutre, Yves Garrabos, Daniel Beysens. Bidimensional empirical mode decomposition and dynamic differential microscopy reveals multiscale dynamics of critical fluctuations. 21st Symposium on Thermophysical Properties, National Institute of Standards and Technology; Joint ASME-AIChE Committee on Thermophysical Properties, Jun 2021, Visioconférence, United States. hal-04571379

HAL Id: hal-04571379

<https://hal.science/hal-04571379>

Submitted on 7 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Bidimensional Empirical Mode Decomposition and Dynamic Differential Microscopy Reveals Multiscale Dynamics of Critical Fluctuations

Ana Oprisan^{C, S} and Sorinel Oprisan
Department of Physics, College of Charleston, Charleston, SC, U.S.A.
oprisana@cofc.edu

Carole Lecoutre-Chabot and Yves Garrabos
Bordeaux INP, ICMCB, CNRS, Université de Bordeaux, Pessac, France

Daniel Beysens
*Physique et Mécanique des Milieux Hétérogènes, Sorbonne Université, CNRS, ESPCI Paris - PSL University,
Paris, France*

Critical fluids have diverging compressibility, which hampers the study of critical phenomena due to convection, sedimentation, and buoyancy. Microgravity environments on the International Space Station (ISS) eliminate the experimental limitations due to compressibility near the critical point. We used a multiscale approach to investigate the dynamics of fluctuations near the critical point of sulfur hexafluoride (SF₆) in microgravity. The Dynamic Differential Microscopy (DDM) has been extensively used for extracting the structure factor and the intermediate scattering function (ISF) of critical fluctuations. Oftentimes, the ISF has a complicated structure with multiple overlapping relaxation processes. Instead of fitting the ISF with multiple weighted exponentials, we first separated different spatial scales from the original images using the Bidimensional Empirical Mode Decomposition (BEMD) technique. Each spatial scale is represented by an Intrinsic Mode Function (IMF) image. We computed the ISF and the structure factor of each IMF. We found over 90% correlation between the ISF of the first IMF and the original image. The ISF determines the correlation time of fluctuations, which providing thus thermal diffusivity coefficient above the critical temperature and effective diffusion coefficients below the critical temperature very close in magnitude. The relaxation time associated with the distinguishable structures observed in the second IMF could be attributed to the fractal nature of fluctuations and light scattering at low wavenumber during the stationary behavior and the transient evolution of the critical fluid cell, which are not easy to detect in the original image.