



**HAL**  
open science

# Copper-based grape pest management has impacted wine aroma

Irene de Guidi, Virginie Galeote, Bruno Blondin, Jean-Luc Legras

## ► To cite this version:

Irene de Guidi, Virginie Galeote, Bruno Blondin, Jean-Luc Legras. Copper-based grape pest management has impacted wine aroma. *Scientific Reports*, 2024, 14 (1), pp.10124. <10.1038/s41598-024-60335-9>. <hal-04572656>

**HAL Id: hal-04572656**

**<https://hal.science/hal-04572656v1>**

Submitted on 11 May 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



OPEN

## Copper-based grape pest management has impacted wine aroma

Irene De Guidi, Virginie Galeote, Bruno Blondin &amp; Jean-Luc Legras✉

Despite the high energetic cost of the reduction of sulfate to  $H_2S$ , required for the synthesis of sulfur-containing amino acids, some wine *Saccharomyces cerevisiae* strains have been reported to produce excessive amounts of  $H_2S$  during alcoholic fermentation, which is detrimental to wine quality. Surprisingly, in the presence of sulfite, used as a preservative, wine strains produce more  $H_2S$  than wild (oak) or wine velum (*flor*) isolates during fermentation. Since copper resistance caused by the amplification of the sulfur rich protein Cup1p is a specific adaptation trait of wine strains, we analyzed the link between copper resistance mechanism, sulfur metabolism and  $H_2S$  production. We show that a higher content of copper in the must increases the production of  $H_2S$ , and that  $SO_2$  increases the resistance to copper. Using a set of 51 strains we observed a positive and then negative relation between the number of copies of *CUP1* and  $H_2S$  production during fermentation. This complex pattern could be mimicked using a multicopy plasmid carrying *CUP1*, confirming the relation between copper resistance and  $H_2S$  production. The massive use of copper for vine sanitary management has led to the selection of resistant strains at the cost of a metabolic tradeoff: the overproduction of  $H_2S$ , resulting in a decrease in wine quality.

The most ancient traces of wine making have been discovered in Georgia<sup>1</sup> and have been dated as 6000 BC. Since that ancient time, cultivation of grapevine and winemaking knowledge spread progressively all over the world<sup>2</sup>. All along this period, winemaking practices have evolved, especially with the discovery of the use of sulfite to limit the growth of undesired microorganisms to protect wine from oxygen and to preserve aroma profile. Similarly, the cultivation of *Vitis vinifera* has faced changes, especially with the development of grafting and the spray of chemical compounds required to face the import in Europe of three major pests for vine: phylloxera, powdery mildew and downy mildew. Among chemicals sprayed on vines, copper has been intensively used in vineyards to control the development of *Plasmopara viticola*. This intensive use of copper in vineyards has translated into high copper in grape musts<sup>3</sup>.

Wine fermentation is mainly achieved by the yeast species *Saccharomyces cerevisiae* which is found also in many fermented products: sake, bread, cheese and more<sup>4–7</sup>, as well as in natural biotopes such as forests<sup>8,9</sup>. *S. cerevisiae* strains display specific physiological properties associated to the different ecological niches they live in, as result of several domestication events<sup>4,10–12</sup>.

One of the most remarkable and contrasting adaptation events can be seen in fermenting wine strains and in wine velum isolates (*flor* yeasts). *S. cerevisiae* velum strains have developed a specialized aerobic lifestyle, highly different from the one of fermenting wine strains<sup>13</sup>. Since they colonize the wine when fermentation is concluded, velum strains develop the ability to grow in media depleted for nitrogen, vitamins, glucose and fructose.

Wine fermentation poses a challenging environment for *S. cerevisiae*. Different genomic features have been identified as traces of adaptation to the wine environment, in line with its domestication<sup>14</sup>. The first and best described adaptation of *S. cerevisiae* to the wine environment is the resistance to sulfite, obtained from several translocation events resulting in a high expression of the sulfite export pump Ssu1<sup>15–19</sup>. Another example of adaptation to the grape and must environments can be seen in the selection of strains carrying multiple copies of the *CUP1* gene. This gene amplification leads to an enhanced protein abundance/synthesis, providing resistance to high concentrations of copper in the grape must, resulting from the massive use of copper as fungicide<sup>20</sup>. Cup1p is among the ten sulfur richest yeast proteins<sup>21</sup> and some *S. cerevisiae* strains can harbor up to 79 copies<sup>18,20,22</sup>. Therefore, the high synthesis of Cup1p caused by its amplification requires a high availability of sulfur containing amino acids methionine and cysteine that are scarce in grape musts. These amino acids can be synthesized by yeast through the sulfur assimilation pathway (SAP), which reduces inorganic sulfate into hydrogen sulfide ( $H_2S$ ) with the consumption of 7 mol of NADPH and 4 of ATP per mole of S-amino acid<sup>23</sup>. Consequently, the

SPO, INRAE, Institut Agro, Université de Montpellier, 34060 Montpellier, France. ✉email: jean-luc.legras@inrae.fr

biosynthesis of the sulfur amino acids has a significant impact on the yeast redox and energy balances. A high diversity in the production of  $H_2S$  during alcoholic fermentation has been described for wine strains<sup>24</sup>, and because its content is detrimental to wine aroma, different studies have deciphered its genetic bases and found allelic variations in *MET10*, *SKP2*, *MET2*, *TUM1*<sup>25–28</sup>, genes involved in the sulfur assimilation pathway or its regulation. Some of these findings have been patented and have led to the improvement of industrial winemaking starters. Surprisingly, no investigation has been carried out to understand the biological meaning of such overproduction, nor to evaluate a potential relation with different ecological niches. Interestingly, for wine *S. cerevisiae*,  $SO_2$  and copper tolerance have been found negatively associated<sup>29</sup>. Transcriptional and proteomic analysis in sulfur-limited medium, demonstrated that *SSU1* over-expression induced sulfur limitation during exposure to copper and provoked an increased sensitivity to copper<sup>30</sup>.


Because the production of  $H_2S$  is so costly to the cell<sup>23</sup>, we wondered why some wine strains were overproducing it. Comparing three groups of strains: isolated from velum and wine, two contrasted anthropogenic environments, and oak, as a natural environment, we show that the total content of  $H_2S$  produced during alcoholic fermentation depends on the ecological niche, and that exposure of yeast cells to copper enhances  $H_2S$  production. We evaluated how the amplification of *CUP1* may explain such variation, using a set of strains with variable number of copies of *CUP1* or strains carrying a plasmid overproducing *CUP1*. Last, we measured the impact of sulfites availability in the media on copper resistance.

## Results

### Strain variability in $H_2S$ production during alcoholic fermentation

To assess the variability of the production of  $H_2S$  during alcoholic fermentation of *Saccharomyces cerevisiae*, we evaluated 33 strains isolated from three ecological niches: wine (n = 10), wine velum (n = 14), and from oak trees (n = 9), as a wild reference. Because  $SO_2$  is an intermediate of the sulfur assimilation pathway, and used in most wine fermentations as an additive for its antimicrobial, antioxidant and anti-oxidizing activities, we compared the  $H_2S$  production of the mentioned ecological groups in a synthetic grape must in the absence or presence of sulfites. The variability in  $H_2S$  production among strains of these three groups is presented in Fig. 1 and Supplementary Fig. 1. A two-way ANOVA revealed that the origin of the strain has a significant effect on  $H_2S$  produced ( $F_{2,128} = 36.31$ ,  $p\_value = 3.24 \times 10^{-13}$ ) as well as the addition of  $SO_2$  to the must ( $F_{1,128} = 59.19$ ,  $p\_value = 3.36 \times 10^{-12}$ ). A significant interaction between the effects of the two factors (i.e.  $SO_2$  and origin) on  $H_2S$  produced during alcoholic fermentation ( $F_{2,128} = 14.5$ ,  $p\_value = 2.20 \times 10^{-6}$ ) was detected.

Tukey multiple comparisons of means at 95% family-wise confidence level showed that  $H_2S$  production between the independent origins was significant when the must contained sulfite (Fig. 1b). Wine strains produced more  $H_2S$  compared to velum and oak ( $p\_value = 1.50 \times 10^{-6}$  and 0.014 respectively). Oak isolates also produced more  $H_2S$  than velum strains in the presence of sulfite ( $p\_value = 1.50 \times 10^{-6}$ ). This difference was not noticeable when the must did not contain sulfites (Fig. 1a).



**Figure 1.** Difference in the average of cumulate  $H_2S$  production during alcoholic fermentation of velum, oak and wine strains in absence (A) or presence (B) of  $SO_2$ .

Comparing Fig. 1a,b, it was clear that strains isolated from different origins did not respond with the same amplitude to the sulfite treatment. Velum strains displayed a remarkable low H<sub>2</sub>S production even in the presence of sulfite, in comparison to wine and oak strains ( $p_{\text{value}} = 1.50 \times 10^{-6}$  and  $3.34 \times 10^{-5}$  respectively). This explains the significant interaction detected by the model.

### Influence of copper content on H<sub>2</sub>S production

The copper content of the grape must or wine can result from the traces left with sanitary treatment performed on vine, and also from treatments aimed at reducing H<sub>2</sub>S production. Indeed, thiols functions reduce Cu<sup>2+</sup> and produce a Cu<sup>+</sup> that binds to -SH functions<sup>31</sup>. We evaluated the effect of copper concentrations of the must on H<sub>2</sub>S production of two industrial winemaking starters: VL1, a wine strain, identified as low H<sub>2</sub>S producer in the first experiment, and LMD17, a high H<sub>2</sub>S producer wine strain<sup>32</sup>. The analysis of variance revealed a significant effect of both factors: strain ( $F_{1,14} = 57$ ,  $p_{\text{value}} = 2.7 \times 10^{-6}$ ), and copper ( $F_{2,14} = 10.6$ ,  $p_{\text{value}} = 0.002$ ). The results presented in Fig. 2 show a clear increase of H<sub>2</sub>S production with an increase in copper content in the synthetic grape must at concentrations compatible with those encountered in winemaking<sup>3</sup>. For both strains, higher concentrations provoke the formation of a black precipitate that hampers H<sub>2</sub>S measurement with our method, and suggests higher production (Supplementary Fig. 2).

### Evaluating the relation between H<sub>2</sub>S production and *CUP1* copy number


Besides inducing H<sub>2</sub>S production, the copper content in the growth medium controls the expression of *CUP1*<sup>33</sup> that is involved in its detoxification. In addition, we observed that *CUP1* is one of the proteins with the highest sulfur containing aminoacid content (21.31%), just after *MNC1* (25.76%), another membrane protein that is upregulated by toxic concentrations of heavy metal ions<sup>34</sup>.

In a first approach aimed at exploring the effect of *CUP1* copy number on H<sub>2</sub>S production in strains of the same three niches analysed above, we increased the number of strains to test (+ 18 wine isolates, total  $n = 51$ ), in order to include strains with 2 to 71 *CUP1* copy number. Surprisingly, we observed a non-linear relation between *CUP1* copy number and total H<sub>2</sub>S production.


As shown in Fig. 3A, strains with 1 to 10 *CUP1* copies exhibit an increasing total H<sub>2</sub>S production, whereas for more copies, it progressively decreases until it reaches almost null values. A 3rd degree polynomial model described well the H<sub>2</sub>S production in relation to the *CUP1* copy number of the strains (black line in Fig. 3), displaying a bell shape, that remained even after the removal of the two highest values for H<sub>2</sub>S production (red line in Fig. 3A). A complex polynomial relation between *CUP1* copy number and H<sub>2</sub>S production is noticeable when the model was built with wine strains only (Fig. 3C).

This bell shape is conserved and amplified in the presence of sulfite in the media, except for strains with a high copy number of *CUP1* (Fig. 3B,D).

Notably, the increase in H<sub>2</sub>S concentration with the copy number of *CUP1* within the range 1–10 copies is similar to the response caused by the increase in copper content of the grape must observed for VL1 and LMD17.



**Figure 2.** Effect of copper content in synthetic must without SO<sub>2</sub> on total H<sub>2</sub>S production during alcoholic fermentation by wine strains VL1 and LMD17.  $p$  values refer to two-way ANOVA.



**Figure 3.**  $H_2S$  production (in synthetic must without  $SO_2$ ) distribution as function of *CUP1* copy number of each strain. (A) Whole set of studied strains, in synthetic must without  $SO_2$ ; (B) whole set of studied strains, in synthetic must supplemented with  $SO_2$ ; (C) wine strains, in synthetic must without  $SO_2$ ; (D) wine strains in synthetic must supplemented with  $SO_2$ ; Isolation origins are described by colours as in Fig. 1. Black solid line: polynomial model describing the relation between  $H_2S$  production and *CUP1* copy number; red line: same model excluding the two highest  $H_2S$  producers.

### Impact of the modulation of *CUP1* copy number on $H_2S$ production

In order to validate the effect of *CUP1* copy number on  $H_2S$  production, we tried to manipulate the number of *CUP1* copies per cell. With this aim, we built a multicopy yeast episomal plasmid (YE<sub>p</sub>) expressing *CUP1* under the control of the strong promoter from the translational elongation factor EF-1 alpha (*TEF1*). Three strains with different number of genomic copies of *CUP1* were transformed with this plasmid and tested in a media containing a low copper concentration (0.25 mg/L).


First, the overexpression of *CUP1* in the oak strain OAK-Rom 3\_2, a low  $H_2S$  producer with one copy of *CUP1*, led to a significant increase of  $H_2S$  production ( $F_{2,6} = 9.61$ ,  $p\_value = 0.013$ , Fig. 4), without affecting the growth. In contrast to the oak strain, the overexpression of *CUP1* in the wine strain LMD17, a high  $H_2S$  producer, with 11 copies of *CUP1*, decreased  $H_2S$  production by half ( $F_{2,5} = 17.49$ ,  $p\_value = 0.006$ , Fig. 4), with no impact on fermentation kinetic. Last, the overexpression of *CUP1* in L1374, which carries 36 copies of *CUP1* and was ranked among the lowest  $H_2S$  producers, did not change its production ( $F_{2,6} = 0.2$ ,  $p\_value = 0.824$ , Fig. 4). The responses displayed by these three constructions are in agreement to the experimental data presented in Fig. 3, reproducing the “bell-shape” trend of  $H_2S$  production.

### Impact of sulfite addition in the culture media on copper resistance



Because sulfite is required for the synthesis of sulfur containing amino acids essential for *CUP1* synthesis, it was logical to test how the deregulation of the pathway by exogenous sulfite might affect copper resistance. Three strains were tested: oak strain Oakrom 3.2, and wine strains LMD17 and L1374, that have 1, 10 and 34 copies of *CUP1* respectively. The overexpression of *CUP1* in Oakrom 3.2, increased the resistance to copper in the control media, which was not the case for wine strains LMD17 and L1374. However, yeast growth was improved for all strains overexpressing *CUP1* on copper-supplemented media when  $SO_2$  was present (Fig. 5A,B).

### Discussion

Our findings, obtained under conventional winemaking conditions, involving the presence of sulfites, demonstrate for the first time that the production of hydrogen sulfide ( $H_2S$ ) by *Saccharomyces cerevisiae* during alcoholic fermentation varies among natural and two distinct groups of domesticated strains. Surprisingly, wine populations exhibited the highest  $H_2S$  production levels when sulfites are added to the grape must. This observation is unexpected, given the widely acknowledged undesirability of  $H_2S$  in winemaking processes, caused by its unpleasant smell of rotten egg. Indeed, high residual concentrations of  $H_2S$  require specific treatment to eliminate this off-flavour. The increase in  $H_2S$  production induced by sulfite addition is explicable due to its role as an intermediate metabolite in the sulfur assimilation pathway. However, the significant differences we observed between wine and oak strains are intriguing, especially considering that wine strains harbour several



**Figure 4.** Effect of *CUP1* overexpression on total H<sub>2</sub>S released during alcoholic fermentation (in synthetic must without SO<sub>2</sub>) in three *S. cerevisiae* strains. Different lowercase letters indicate statistically significant differences between the molecular modifications (control wild-type strain, empty vector or *CUP1* overexpressing vector) for each strain separately, after Tukey multiple comparison of means at 95% family-wise confidence level.



**Figure 5.** Effect of SO<sub>2</sub> addition in the media on copper resistance. (A) Resistance of oak strain Oak-Rom 3.2 to copper (0 to 1 mM) in the absence (control) and presence of 40 mg/L SO<sub>2</sub>. (B) Resistance of wine strain LMD17 and L1374 to copper (0 to 12 mM) in the absence (control) and presence of SO<sub>2</sub> (60 mg/L).

types of translocations leading to a higher expression of the sulfite efflux pump *SSU1*. This apparently contradicts the antagonistic role of *SSU1* in copper resistance<sup>30</sup>, suggesting a limitation in the sulfur assimilation pathway. However, it should be noted that Onetto et al.'s study was conducted in the absence of added sulfites and indeed, we could show that the introduction of sulfites into the grape must, as commonly practiced by winemakers, increases copper resistance, including for strains with a high *CUP1* copy number. We therefore propose that exogenous sulfite may exceed the expulsion capacity of the sulfite export transporter, thereby increasing H<sub>2</sub>S, and hence sulfur containing amino acids and *CUP1* synthesis in wine strains.

We also demonstrate that the presence of copper in grape must increases H<sub>2</sub>S production. Our results align with previous expression data<sup>35,36</sup>, which revealed that copper exposure triggers a higher expression of genes encoding the two subunits of the sulfite reductase *MET5*, *MET10* (i.e., the main enzyme of the sulfur assimilation pathway), and the protein responsible for copper resistance/detoxification *CUP1* following. Furthermore, the metallothionein protein Cup1p, which has one of the highest contents of sulfur-containing amino acids methionine and cysteine in the *S. cerevisiae* proteome, requires the availability of these amino acids for its synthesis. The differences in H<sub>2</sub>S production among oak, wine, and velum yeast strains may be attributed to variations in the number of *CUP1* copies in their genomes. However, we describe a complex relationship between the number of *CUP1* copies and H<sub>2</sub>S production. Moderate amplification of *CUP1* (up to approximately 10 copies) leads to an increase in H<sub>2</sub>S production, whereas higher copy numbers result in a lower fraction of H<sub>2</sub>S being stripped by the CO<sub>2</sub> generated during fermentation. One possibility of this curve could result from an increasing activation of SAP to support *CUP1* production when up to at least 10 gene copies are present, whereas for higher number of copies, the higher requirement of sulfur amino acids could exceed maximum activity of the SAP, leading to an increased use of H<sub>2</sub>S for the synthesis and a lower release. The increased resistance obtained in the presence of SO<sub>2</sub> supports this hypothesis. In this case, we propose that this reflects a higher utilization of H<sub>2</sub>S for amino acid synthesis.

Lastly, our results also shed light on the specific behaviour of flor strains. Unlike wine strains, velum strains exhibit very low H<sub>2</sub>S production, and have a lower number of *CUP1* copies. Velum strains grow at the surface of wine, after alcoholic fermentation, which significantly reduces the copper content of wine. It is likely that the selection pressure for copper-resistant strains has been less intense for flor strains compared to wine yeast. Another possible explanation for the lower H<sub>2</sub>S production in velum strains is the reduced activity of the pentose phosphate pathway, which provides NADPH, in comparison to oak, bread and wine strains<sup>37</sup>. This observation elucidates the divergent domestication trajectories of wine and flor strains, reflecting their distinct lifestyles<sup>13</sup>.

## Conclusion

The long-term exposure of yeast to copper, used for vine pest management over 150 years, has led to their adaptation by selecting strains with multiple copies of *CUP1*. Our results suggest that this adaptation involves a significant trade-off: increased resistance to copper, but also high H<sub>2</sub>S production by the yeast, which is detrimental to wine quality. This increased H<sub>2</sub>S production is further exacerbated in the presence of sulfite, another common additive in winemaking. Given the energetic cost of H<sub>2</sub>S production, its impact on the global yeast metabolism should be evaluated. Although many projects and techniques have been dedicated to understanding and limiting H<sub>2</sub>S production<sup>26,27,28</sup>, none have investigated the potential role of copper use in causing this phenotype. Therefore, the yeast *CUP1* background should be considered when selecting wine yeast for low H<sub>2</sub>S production. However, diversity data suggests that the amplification of *CUP1* likely is not the sole mechanism explaining variations in H<sub>2</sub>S production, which requires further investigations.

## Materials and methods

### Strains

Fifty-one *Saccharomyces cerevisiae* from different geographical areas were characterized for their H<sub>2</sub>S production during alcoholic fermentation. The genetic group, identified in previous works indicated in the references of Supplementary Table 1, reflected the colonized ecological niche: 28 belong to the “wine” clade, 14 to “velum” group and 9 to the “oak” one. Strains were selected from our laboratory collection and maintained on solid medium (agar YPD: 2% glucose, 1% yeast extract, 2% bactopectone, 2% agar) at 4 °C.

### Fermentation conditions and H<sub>2</sub>S quantification

Fermentation experiments were conducted using synthetic must (SM), designed to mimic the characteristics of a natural grape must<sup>38</sup>. It contained a 200 g/L equimolar glucose and fructose content, and 200 mg/L assimilable nitrogen, 3.8 mg/L phytosterol, and 0.25 mg/L of Cu<sup>2+</sup>. The pH was adjusted to 3.3 with sodium hydroxide solution.

One colony of each strain was grown in 5 ml of liquid YPD at 28 °C for 24 h and then diluted 100 times in SM. After 24 h at 28 °C, cells were counted with an electronic particle counter (Multisizer 3 counter; Beckman Coulter) and 250 mL of SM, supplemented with 60 mg/L of SO<sub>2</sub> when the impact of sulfite was evaluated, were inoculated to 1 × 10<sup>6</sup> cells/mL. Fermentations were carried out at 28 °C, under permanent stirring (280 rpm) and they were followed daily by weight loss, until the theoretical percentage of sugar consumed reached 95% (87.4 g CO<sub>2</sub>/L produced). Total H<sub>2</sub>S produced during alcoholic fermentation was collected with a zinc-based trap system and quantified with sulfide specific fluorescent probe, as described before<sup>32</sup>.

When the impact of the overexpression of *CUP1* was in study, SM was supplemented with Geneticin (G418—Sigma A1720-5G) to maintain the plasmid allowing the overexpression itself. Suitable antibiotic concentrations were defined for each strain (100 µg/mL for wine strains, 40 µg/mL for the oak one), to simultaneously allow the maintenance of the plasmid and a good fermentation rate, but prevent the growth of the sensitive strain (i.e. the wild-type strain without the plasmid).

When assessing the impact of copper concentration on H<sub>2</sub>S production, SM was supplemented with copper sulfate to reach 1 or 2 mg/L of copper; control copper concentration was 0.25 mg/L in all the experiments. More details about the experiments are given in the “[Experimental design and statistical analyses](#)” section.

### Drop test on copper and sulfite supplemented media

Copper resistance in presence or absence of SO<sub>2</sub>, was assessed by a drop-test for three wild type strains with different *CUP1* copy number in their genome (Oak-Rom 3-2, LMD17 and L1374), and their counterpart engineered to over-express *CUP1* (see below). Triplicates of these strains were grown overnight at 28 °C in 5 mL of YPD. Cells were then counted with an electronic particle counter (Multisizer 3 counter; Beckman Coulter), washed with PBS and resuspended in sterile PBS to obtain 10<sup>7</sup> cells/mL. Three successive 1/10 dilutions were prepared and 1.5 µL of each dilution was spotted on synthetic must having the same composition of the one used for the fermentations, gelled with 20 g/L agar. According to the tested modalities, copper (0, 0.5, 1, 6, 12 mM) and sulfite (0, 40, 60 mg/L SO<sub>2</sub>) were added to the media to evaluate their effect. Agar plates were incubated at 28 °C for 72 h and growth was assessed by visual examination.

### CUP1 copy number evaluation

For most of the strains, *CUP1* copy number was estimated from their genome sequence, obtained from previous works or from sequencing performed in this study. To obtain the values, the median sequencing depth measured at SNPs encountered between coordinates 212,500 and 213,000, and between 214,500 and 215,000 on Chromosome VIII was divided by the median sequencing depth over the entire genome (excluding mitochondria and 2 microns). For Italian strains, *CUP1* copy number data had been already quantified by Real Time PCR<sup>18</sup>.

### Genomic DNA extraction for sequencing

Genomic DNA was isolated from liquid yeast cultures in stationary phase, with a classical phenol–chloroform method, as described before<sup>39</sup>, with an additional purification step based on the use of silica-coated magnetic beads (GMG-252-A-100 mL—PerkinElmer), as follows. Cells were broken mechanically by shaking them in the presence of 600 µm diameter glass beads, lysis buffer (Tris 50 mM pH 8, EDTA 50 mM, NaCl 100 mM, Triton 2%, SDS 1.25%) and phenol chloroform isoamyl alcohol 25:24:1. DNA was precipitated with isopropanol and ethanol, dried, resuspended in TE (Tris 10 mM, EDTA 1 mM) and treated with RNase A. Samples were mixed with the DNA absorption solution (for one sample: 50 µL 5 M NaCl, 15 µL magnetic beads (GMG-252-A-100 mL—PerkinElmer), 250 µL 7.8 M guanidium chloride, 800 µL isopropanol), after which metal beads with DNA absorbed on their silica surface were recovered using the DynaMag™-2 Magnet tube holder (12321D-DynaMag-2—Invitrogen) and washed twice with AMMLAV/E buffer (10 mM Tris pH 8, 0.1 mM EDTA, 60 mM potassium acetate, 65% ethanol) and twice with ethanol 75%. DNA was then desorbed and in aqueous solution.

DNA purity was checked from the 260 nm/280 nm and 260 nm/230 nm OD ratio measured with NanoDrop 1000 (ThermoScientific). The DNA was quantified by fluorescence using the QuantiFluor kit, dsDNA system (Promega) and then stored at –20 °C.

### Genome sequence and analysis

DNA samples were processed to generate libraries of 500 bp inserts. After passing quality control, the libraries were sequenced with DNBseq technology using BGISEQ-500 platform, generating paired-end reads of 2 × 150 bp.

For each library, low-quality reads were processed and filtered using the FASTX Toolkit v0.0.13.2 and TRIM-MOMATIC v0.36<sup>40</sup> with the following parameters (LEADING:10 HEADCROP:5 SLIDINGWINDOW:4:15 MINLEN:50).

Reads were then mapped to the S288C reference genome with BWA v0.6.2 with default parameters<sup>41</sup> and genotyping made with samtools v1.11 to obtain a variant file including the sequencing depth of each variant position. Sequence positions were afterwards filtered for quality criteria: sufficient coverage position as well as genotyping and mapping quality (MQ > –20) were kept.

### Plasmid construction and yeast transformation

*CUP1* was inserted via Gibson assembly method<sup>42</sup> between TEF promoter and terminator in a high copy Yeast Episomal plasmid (YEp352), modified to confer geneticin resistance (YEp352-G418) to the host cell. In detail, the backbone was amplified with primers P1 and P2, designed to replace the original *URA3* copy of YEp with *CUP1*, since the strains used were not auxotrophic and the selection had been made by antibiotic. Therefore, the backbone contained a 2 um replication origin (multicopy), AmpR, ColE1, pPGK and G418 resistance cassette. *CUP1* was amplified from OakGri7\_1, a strain previously sequenced by our laboratory<sup>10</sup>, with a single metallothionein copy and the same sequence as laboratory reference strain S288C, used to design primers (P5–P6). TEF promoter and terminator were amplified from pCfB2312<sup>43</sup> with primers P3–P4 and P7–P8, respectively. Primer sequences are listed in Table 1.

Proper fragment insertion was verified by enzymatic digestion (NarI, ClaI, PacI—New England Biolabs). To assure that the phenotype was related to the overexpression of *CUP1*, a YEp352-G418 plasmid without *CUP1* was used as control. PCRs were performed with Phusion™ High-Fidelity DNA Polymerase and validated by gel electrophoresis. *Escherichia coli* strain DH5α was used to maintain and amplify the plasmid; cells were selected on LB medium with ampicillin (100 µg/mL) and grown at 37 °C. Yeasts (Oak-Rom 3-2, LMD17 and L1374) were transformed with the lithium acetate method<sup>44</sup> and strains containing the recombinant plasmids were selected on YPD agar with 200 µg/mL geneticin (G418—Sigma A1720-5G).

Primer name	Primer sequence (uppercase letters are homologous to the template to amplify, lowercase letters are homologous to the adjacent fragment to assemble)
P1	acgccgcatccagtgctgaACATATGCGTATATATACCAATCTAAG
P2	cggcgggacaaggcaagctAGGAGAAAATACCGCATCAG
P3	ctgatgctgtattttctctAGCTTGCCTGTCCCCGCCG
P4	ttaattaatcgctgaacatGGTTGTTTATGTTTCGGATGTGATGTGAGAA
P5	acatccgaacataacaaccATGTTTCAGCGAATTAATTAACCTC
P6	ctttttatgtcagtactgaTCATTTCCAGAGCAGCATG
P7	catgctgctctgggaatgaTCAGTACTGACAATAAAAAGATTCTGTGTTTCAAGA
P8	tggtatataatcgcatgtTCGACACTGGATGGCGCGTTA

**Table 1.** Primers used in this work.

## Experimental design and statistical analyses

### *Experiment 1: impact of the origin of the isolate and sulfites on H<sub>2</sub>S production*

33 strains were selected randomly from our laboratory collection (Supplementary Table 1, Dataset 1). Alcoholic fermentations were performed in absence or presence of SO<sub>2</sub>, in duplicate for each strain and each condition.

The factors accounting for the variation of H<sub>2</sub>S were analyzed with the following analysis of variance model:

$$Y_{ijk} = \mu + \alpha_i + \beta_j + \gamma_{ij} + \varepsilon_{ijk},$$

where  $Y_{ijk}$  is the H<sub>2</sub>S production,  $\mu$  the overall grand mean,  $\alpha_i$  is the fixed strain group effect,  $\beta_j$  is the fixed SO<sub>2</sub> effect,  $\gamma_{ij}$  is their interaction effect, and  $\varepsilon_{ijk}$  the residual error.

The analysis of the residuals showed that three values were distant from the global distribution. Since results of the statistical analysis did not change after removing all the observations of the three outlier strains, the complete dataset was kept as the method is sufficiently robust to mild deviations.

### *Experiment 2: impact of copper content of the media on H<sub>2</sub>S production*

Fermentations were performed without SO<sub>2</sub> in triplicate, for each strain (VL1 and LMD17) and each condition (0.25–1 and 2 mg/L of copper).

To evaluate the effect of copper and strains on H<sub>2</sub>S production, ANOVA was performed, after checking for the equality of variance with a Levene test. The most parsimonious model was kept after checking of the absence of interaction between strain and the copper content:

$$Y_{ijk} = \mu + \alpha_i + \beta_j + \varepsilon_{ijk},$$

where  $Y_{ijk}$  is the H<sub>2</sub>S production,  $\mu$  the overall grand mean,  $\alpha_i$  is the fixed strain effect,  $\beta_j$  is the fixed copper effect and  $\varepsilon_{ijk}$  the residual error.

### *Experiment 3: impact of CUP1 copy number on H<sub>2</sub>S production*

To the strains evaluated in experiment 1, we added 18 wine strains (total strains analyzed = 55), some known to harbor a high number of copies of *CUP1*, and some commercial strains known to be high H<sub>2</sub>S producers (Supplementary Table 1, Dataset 2). Alcoholic fermentations were performed in absence of SO<sub>2</sub>, in duplicate for each strain.

Different polynomial models were used to describe the interaction between H<sub>2</sub>S production and *CUP1* copy number (first-, second- and third-degree polynomial models); ANOVA was used to assess the significance of these models.

### *Experiment 4: impact of the overexpression of CUP1 on H<sub>2</sub>S production*

Fermentations were performed without SO<sub>2</sub> and with standard copper content (0.25 mg/L) in triplicate, for each strain (OAK\_ROM 1–3, LMD17 and L1374) and each condition (wild-type strain, strain with the empty vector, strain with the *CUP1* overexpressing vector).

ANOVA was performed to test the effect of the genetic modification in each strain. The model used was:

$$Y_{ij} = \mu + \alpha_i \cdot \varepsilon_{ijk},$$

where  $Y_{ij}$  is the H<sub>2</sub>S production,  $\mu$  the overall grand mean,  $\alpha_i$  is the fixed genetic modification effect, and  $\varepsilon_{ijk}$  the residual error.


Figure 6 summarizes the experimental design.

For all the experiments, when the impact of one (or more) factor was significant, differences between modalities were evaluated by post-hoc testing (Tukey's HSD multiple-comparison test,  $p < 0.05$ ).

Statistical analyses were performed in the R environment (R version 4.0.2 (2020-06-22)<sup>45</sup>).

## Compliance with international and national regulation

Yeast strains were available from culture collection, or gifted by other authors, or provided by the company Lal-lemmand. The yeast collection and use was in accordance with all the relevant guidelines.



**Figure 6.** Experimental design.

### Data availability

The datasets generated and analysed during the current study are available at Data Gouv (<https://entrepot.reche.reche.data.gouv.fr/>) with the following <https://doi.org/10.57745/5ECVDJ>. Genome sequences were deposited at EBI (<https://www.ebi.ac.uk/>) and corresponding accession numbers are given in Supplementary Table 1 Strain-dataset.xlsx.

Received: 15 July 2023; Accepted: 22 April 2024

Published online: 02 May 2024

### References

1. McGovern, P. *et al.* Early neolithic wine of Georgia in the South Caucasus. *Proc. Natl. Acad. Sci. U.S.A.* **114**(48), E10309–E10318. [https://doi.org/10.1073/PNAS.1714728114/SUPPL\\_FILE/PNAS.1714728114.SD03.XLSB](https://doi.org/10.1073/PNAS.1714728114/SUPPL_FILE/PNAS.1714728114.SD03.XLSB) (2017).
2. Pretorius, I. S. Tailoring wine yeast for the new millennium: Novel approaches to the ancient art of winemaking. *Yeast* **16**(8), 675–729. [https://doi.org/10.1002/1097-0061\(20000615\)16:8%3c675::AID-YEA585%3e3.0.CO;2-B](https://doi.org/10.1002/1097-0061(20000615)16:8%3c675::AID-YEA585%3e3.0.CO;2-B) (2000).

3. Darriet, P. *et al.* Effects of copper fungicide spraying on volatile thiols of the varietal aroma of Sauvignon blanc, Cabernet Sauvignon and Merlot wines. *Vitis* **40**(2), 93–99 (2001).
4. Barbosa, R. *et al.* Multiple rounds of artificial selection promote microbe secondary domestication—The case of Cachaça yeasts. *Genome Biol. Evol.* **10**(8), 1939–1955. <https://doi.org/10.1093/gbe/evy132> (2018).
5. Cromie, G. A. *et al.* Genomic sequence diversity and population structure of *Saccharomyces cerevisiae* assessed by RAD-seq. *G3 Genes Genomes Genet.* **3**(12), 2163–2171. <https://doi.org/10.1534/g3.113.007492> (2013).
6. Fay, J. C. & Benavides, J. A. Evidence for domesticated and wild populations of *Saccharomyces cerevisiae*. *PLoS Genet.* **1**(1), 0066–0071. <https://doi.org/10.1371/journal.pgen.0010005> (2005).
7. Ludlow, C. L. *et al.* Independent origins of yeast associated with coffee and cacao fermentation. *Curr. Biol.* **26**(7), 965–971. <https://doi.org/10.1016/j.cub.2016.02.012> (2016).
8. Wang, Q. M., Liu, W. Q., Liti, G., Wang, S. A. & Bai, F. Y. Surprisingly diverged populations of *Saccharomyces cerevisiae* in natural environments remote from human activity. *Mol. Ecol.* **21**(22), 5404–5417. <https://doi.org/10.1111/j.1365-294X.2012.05732.X> (2012).
9. Lee, T. J. *et al.* Extensive sampling of *Saccharomyces cerevisiae* in Taiwan reveals ecology and evolution of predomesticated lineages. *Genome Res.* **32**(5), 864–877. <https://doi.org/10.1101/GR.276286.121/-/DC1> (2022).
10. Legras, J.-L. *et al.* Adaptation of *S. cerevisiae* to fermented food environments reveals remarkable genome plasticity and the footprints of domestication. *Mol. Biol. Evol.* **35**(7), 1712–1727. <https://doi.org/10.1093/molbev/msy066> (2018).
11. Gonçalves, M. *et al.* Distinct domestication trajectories in top-fermenting beer yeasts and wine yeasts. *Curr. Biol.* **26**(20), 2750–2761. <https://doi.org/10.1016/j.cub.2016.08.040> (2016).
12. De Guidi, L., Legras, J. L., Galeote, V. & Sicard, D. Yeast domestication in fermented food and beverages: Past research and new avenues. *Curr. Opin. Food Sci.* **51**, 101032. <https://doi.org/10.1016/j.cofs.2023.101032> (2023).
13. Coi, A. L. *et al.* Genomic signatures of adaptation to wine biological ageing conditions in biofilm-forming flor yeasts. *Mol. Ecol.* **26**(7), 2150–2166. <https://doi.org/10.1111/mec.14053> (2017).
14. García-Ríos, E. & Guillamón, J. M. Genomic adaptations of *Saccharomyces* genus to wine niche. *Microorganisms* **10**(9), 1811. <https://doi.org/10.3390/microorganisms10091811> (2022).
15. Pérez-Ortín, J. E., Querol, A., Puig, S. & Barrio, E. Molecular characterization of a chromosomal rearrangement involved in the adaptive evolution of yeast strains. *Genome Res.* **12**(10), 1533–1539. <https://doi.org/10.1101/gr.436602> (2002).
16. Zimmer, A. *et al.* QTL dissection of lag phase in wine fermentation reveals a new translocation responsible for *Saccharomyces cerevisiae* adaptation to sulfite. *PLoS ONE* **9**(1), e86298. <https://doi.org/10.1371/journal.pone.0086298> (2014).
17. García-Ríos, E., Nuévalos, M., Barrio, E., Puig, S. & Guillamón, J. M. A new chromosomal rearrangement improves the adaptation of wine yeasts to sulfite. *Environ. Microbiol.* **21**(5), 1771–1781. <https://doi.org/10.1111/1462-2920.14586> (2019).
18. Crosato, G. *et al.* The impact of CUP1 gene copy-number and XVI-VIII/XV-XVI translocations on copper and sulfite tolerance in vineyard *Saccharomyces cerevisiae* strain populations. *FEMS Yeast Res.* **20**(4), 28. <https://doi.org/10.1093/femsyr/foaa028> (2020).
19. Brion, C., Ambroset, C., Sanchez, I., Legras, J.-L. & Blondin, B. Differential adaptation to multi-stressed conditions of wine fermentation revealed by variations in yeast regulatory networks. *BMC Genom.* **14**(1), 1–21. <https://doi.org/10.1186/1471-2164-14-681> (2013).
20. Fogel, S., Welch, J. W., Cathala, G. & Karin, M. Gene amplification in yeast: CUP1 copy number regulates copper resistance. *Curr. Genet.* **7**(5), 347–355. <https://doi.org/10.1007/BF00445874> (1983).
21. Bateman, A. *et al.* UniProt: The universal protein knowledgebase in 2021. *Nucleic Acids Res.* **49**(D1), D480–D489. <https://doi.org/10.1093/NAR/GKAA1100> (2021).
22. Strobe, P. K. *et al.* The 100-genomes strains, an *S. cerevisiae* resource that illuminates its natural phenotypic and genotypic variation and emergence as an opportunistic pathogen. *Genome Res.* **125**(5), 762–774. <https://doi.org/10.1101/GR.185538.114> (2015).
23. Celton, M. *et al.* A comparative transcriptomic, fluxomic and metabolomic analysis of the response of *Saccharomyces cerevisiae* to increases in NADPH oxidation. *BMC Genom.* **13**(1), 317. <https://doi.org/10.1186/1471-2164-13-317> (2012).
24. Spiropoulos, A., Tanaka, J., Flerianos, I. & Bisson, L. F. Characterization of hydrogen sulfide formation in commercial and natural wine isolates of *Saccharomyces*. *Am. J. Enol. Viticult.* **51**(3), 233–248 (2000).
25. Huang, C., Roncoroni, M. & Gardner, R. C. MET2 affects production of hydrogen sulfide during wine fermentation. *Appl. Microbiol. Biotechnol.* **98**(16), 7125–7135. <https://doi.org/10.1007/s00253-014-5789-1> (2014).
26. Noble, J., Sanchez, I. & Blondin, B. Identification of new *Saccharomyces cerevisiae* variants of the MET2 and SKP2 genes controlling the sulfur assimilation pathway and the production of undesirable sulfur compounds during alcoholic fermentation. *Microb. Cell Factor.* **14**(1), 1–16. <https://doi.org/10.1186/s12934-015-0245-1> (2015).
27. Huang, C. W. *et al.* The yeast TUM1 affects production of hydrogen sulfide from cysteine treatment during fermentation Dawes I, editor. *FEMS Yeast Res.* **16**(8), 100. <https://doi.org/10.1093/femsyr/fow100> (2016).
28. Linderholm, A., Dietzel, K., Hirst, M. & Bisson, L. F. Identification of MET10-932 and characterization as an allele reducing hydrogen sulfide formation in wine strains of *Saccharomyces cerevisiae*. *Appl. Environ. Microbiol.* **76**(23), 7699–7707. <https://doi.org/10.1128/AEM.01666-10> (2010).
29. Schmidt, S. A., Kolouchova, R., Forgan, A. H. & Borneman, A. R. Evaluation of *Saccharomyces cerevisiae* wine yeast competitive fitness in enologically relevant environments by barcode sequencing. *G3 Genes Genomes Genet.* **10**(2), 591–603. <https://doi.org/10.1534/G3.119.400743> (2020).
30. Onetto, C. A. *et al.* SO<sub>2</sub> and copper tolerance exhibit an evolutionary trade-off in *Saccharomyces cerevisiae*. *PLoS Genet.* **19**(3), e1010692. <https://doi.org/10.1371/JOURNAL.PGEN.1010692> (2023).
31. Kreitman, G. Y., Danilewicz, J. C., Jeffery, D. W. & Elias, R. J. Reaction mechanisms of metals with hydrogen sulfide and thiols in model wine. Part 2: Iron- and copper-catalyzed oxidation. *J. Agric. Food Chem.* **64**(20), 4105–4113. <https://doi.org/10.1021/ACS.JAFC.6B00642> (2016).
32. De Guidi, I., Farines, V., Legras, J.-L. & Blondin, B. Development of a new assay for measuring H<sub>2</sub>S production during alcoholic fermentation: Application to the evaluation of the main factors impacting H<sub>2</sub>S production by three *Saccharomyces cerevisiae* wine strains. *Fermentation* **7**, 1–14 (2021).
33. Peña, M. M. O., Koch, K. A. & Thiele, D. J. Dynamic regulation of copper uptake and detoxification genes in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **18**(5), 2514–2523. <https://doi.org/10.1128/mcb.18.5.2514> (1998).
34. Andreeva, N. *et al.* Transcriptome profile of yeast reveals the essential role of PMA2 and uncharacterized gene YBR056W-A (MNC1) in adaptation to toxic manganese concentration. *Metallomics* **9**(2), 175–182. <https://doi.org/10.1039/C6MT00210B> (2017).
35. Oc, S., Eraslan, S. & Kirdar, B. Dynamic transcriptional response of *Saccharomyces cerevisiae* cells to copper. *Sci. Rep.* **10**(1), 1–13. <https://doi.org/10.1038/s41598-020-75511-w> (2020).
36. Yasokawa, D. *et al.* Mechanisms of copper toxicity in *Saccharomyces cerevisiae* determined by microarray analysis. *Environ. Toxicol.* **23**(5), 599–606. <https://doi.org/10.1002/TOX.20406> (2008).
37. Nidelet, T., Brial, P., Camarasa, C. & Dequin, S. Diversity of flux distribution in central carbon metabolism of *S. cerevisiae* strains from diverse environments. *Microb. Cell Factor.* **15**(1), 1–13. <https://doi.org/10.1186/S12934-016-0456-0/FIGURES/7> (2016).
38. Bely, M., Sablayrolles, J. M. & Barre, P. Description of alcoholic fermentation kinetics: Its variability and significance. *Am. J. Enol. Viticult.* **41**(4), 319–324 (1990).
39. Saubin, M. *et al.* Investigation of genetic relationships between *Hanseniaspora* species found in grape musts revealed interspecific hybrids with dynamic genome structures. *Front. Microbiol.* **10**, 1. <https://doi.org/10.3389/FMICB.2019.02960/FULL> (2020).

40. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for illumina sequence data. *Bioinformatics* **30**(15), 2114–2120. <https://doi.org/10.1093/bioinformatics/btu170> (2014).
41. Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows–Wheeler transform. *Bioinformatics* **26**(5), 589–595. <https://doi.org/10.1093/bioinformatics/btp698> (2010).
42. Gibson, D. G. *et al.* Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat. Methods* **6**(5), 343–345. <https://doi.org/10.1038/nmeth.1318> (2009).
43. Jessop-Fabre, M. M. *et al.* EasyClone-MarkerFree: A vector toolkit for marker-less integration of genes into *Saccharomyces cerevisiae* via CRISPR-Cas9. *Biotechnol. J.* **11**(8), 1110–1117. <https://doi.org/10.1002/biot.201600147> (2016).
44. Gietz, R. D. & Schiestl, R. H. High-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG method. *Nat. Protoc.* **2**(1), 31–34. <https://doi.org/10.1038/nprot.2007.13> (2007).
45. Team R Core. *R: A Language and Environment for Statistical Computing*. <https://www.r-project.org/> (2020).

## Acknowledgements

The authors thank Jessica Noble (Lallemand SAS), Prof Joseph Schacherer (University of Strasbourg) and Prof Viviana Corich (University of Padova) for providing the strains analyzed in this work. All the authors are grateful to Thérèse Marlin for providing her expertise with the Gibson assembly strategy and for the molecular cloning of *CUP1*, to Dr Thibault Nidelet for the polynomial models and Dr Delphine Sicard for the statistical counseling and the proofreading of the manuscript.

## Author contributions

BB and JLL conceived the study and supervised the work; IDG carried out the experiments and wrote the manuscript with support from JLL and BB; VG and IDG conceived the molecular biology experiments. All authors discussed the results and contributed to the final manuscript. All authors have read and agreed to the published version of the manuscript.

## Funding

This work has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie Actions (Grant Agreement No 764364) and from the French National Research Agency (Funding Agreement No ANR-21-PRRD-0008-01).

## Competing interests

The authors declare no competing interests.

## Additional information

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1038/s41598-024-60335-9>.

**Correspondence** and requests for materials should be addressed to J.-L.L.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Publisher’s note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2024