



HAL
open science

Dealing with the irregular in inverse and imaging problems

Gabriel Claret, Michael Hinz, Anna Rozanova-Pierrat, Alexander Teplyaev

► **To cite this version:**

Gabriel Claret, Michael Hinz, Anna Rozanova-Pierrat, Alexander Teplyaev. Dealing with the irregular in inverse and imaging problems. Congrès des Jeunes Chercheurs en Mathématiques et Applications 2023, Sep 2023, Gif-sur-Yvette, France. . hal-04573038

HAL Id: hal-04573038

<https://hal.science/hal-04573038v1>


Submitted on 13 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Description of the problem

In medical imaging, cancer tumours are known to have irregular boundaries, which are best modelled by **non-Lipschitz shapes such as fractals**. We consider an imaging problem set on a domain Ω of \mathbb{R}^n , $n \geq 2$, inside which is an inclusion D characterized by its conductivity k :



$$\begin{cases} \nabla \cdot \left((1 + (k-1)\mathbb{1}_D) \nabla u \right) = 0 & \text{on } \Omega, \\ \frac{\partial_i u}{\partial \nu} \Big|_{\partial \Omega} = g, \\ \int_{\Omega} u \, dx = 0. \end{cases} \quad (1)$$

Here,

- Ω is a bounded domain,
- $\partial \Omega$ is connected,
- $D \subset \subset \Omega$ is a domain,
- $k > 0$ and $k \neq 1$,
- $\mathbb{1}_D$ is the indicator function of D .

We wish to **identify the inclusion D from boundary measurement on $\partial \Omega$** when Ω (and D) can be an irregular or fractal domain.

• **Domain regularity:** Ω is a **(two-sided) extension domain**:

- Ω is an H^1 -extension domain: \exists a linear continuous extension operator $H^1(\Omega) \rightarrow H^1(\mathbb{R}^n)$,
 - $\Omega^c := \mathbb{R}^n \setminus \bar{\Omega}$ is an H^1 -extension domain,
 - the Hausdorff dimension of $\partial \Omega$ is (locally) in $[n-1, n]$.
- ▶ $\partial \Omega$ can have a **changing Hausdorff dimension**.
 ▶ Ω cannot have a cusp (conditions (i) and (ii)).

2 Trace and weak normal derivatives

• **Trace operators** – Dirichlet boundary conditions [2]:

$$\text{Tr}_i u(x) = \lim_{r \rightarrow 0} \frac{1}{\lambda^{(n)}(\Omega \cap B_r(x))} \int_{\Omega \cap B_r(x)} u \, dx, \quad u \in H^1(\Omega), \quad x \in \partial \Omega \text{ q.e.},$$

is the **interior trace operator**, where $\lambda^{(n)}$ is Lebesgue's measure on \mathbb{R}^n .

- ▶ $\text{Ker}(\text{Tr}_i) = H_0^1(\Omega) = \overline{C_0^\infty(\Omega)}^{\|\cdot\|_{H^1(\Omega)}}$
- ▶ $\text{Ker}(\text{Tr}_i)^\perp$ is the space of **1-harmonic functions on Ω** .
- ▶ $\mathcal{B}(\partial \Omega) := \text{Tr}_i(H^1(\Omega))$ is a Hilbert space for the norm:

$$\|f\|_{\mathcal{B}(\partial \Omega)} := \min \{ \|v\|_{H^1(\Omega)} \mid \text{Tr}_i v = f \} = \|\tilde{v}\|_{H^1(\Omega)},$$

where \tilde{v} is 1-harmonic on Ω and $\text{Tr}_i \tilde{v} = f$.

Similarly, we define the **exterior trace operator** $\text{Tr}_e : H^1(\Omega^c) \rightarrow \mathcal{B}(\partial \Omega)$.

• **Weak normal derivatives** – Neumann boundary conditions [3]: for $u \in H^1(\Omega)$ with $\Delta u \in L^2(\Omega)$, denoting $\mathcal{B}'(\partial \Omega) := \mathcal{L}(\mathcal{B}(\partial \Omega), \mathbb{R})$,

$$\forall v \in H^1(\Omega), \quad \left\langle \frac{\partial_i u}{\partial \nu}, \text{Tr}_i v \right\rangle_{\mathcal{B}'(\partial \Omega), \mathcal{B}(\partial \Omega)} = \int_{\Omega} (\Delta u) v \, dx + \int_{\Omega} \nabla u \cdot \nabla v \, dx,$$

is the **weak interior normal derivative** of u .

Similarly, we define the **weak exterior normal derivative** $\frac{\partial_e u}{\partial \nu} \in \mathcal{B}'(\partial \Omega)$ for $u \in H^1(\Omega^c)$ with $\Delta u \in L^2(\Omega^c)$.

3 Layer potential operators

$$\text{Transmission problem: } \begin{cases} -\Delta u = 0 & \text{on } \mathbb{R}^n \setminus \partial \Omega, \\ \text{Tr}_i u - \text{Tr}_e u = -f & \text{on } \partial \Omega, \\ \frac{\partial_i u}{\partial \nu} - \frac{\partial_e u}{\partial \nu} = g & \text{on } \partial \Omega. \end{cases} \quad (2)$$

• If Ω is **Lipschitz**, $f, g \in L^2(\partial \Omega, \sigma)$ with σ Lebesgue's measure on $\partial \Omega$ and $\int_{\partial \Omega} g \, d\sigma = 0$, then $u = \mathcal{S}_{\partial \Omega} g - \mathcal{D}_{\partial \Omega} f$ [4], where:

$$\begin{aligned} \mathcal{S}_{\partial \Omega} g(x) &= \int_{\partial \Omega} G(x-y) g(y) \, \sigma(dy), \quad x \in \mathbb{R}^n, \\ \mathcal{D}_{\partial \Omega} f(x) &= \int_{\partial \Omega} \frac{\partial G}{\partial \nu_y}(x-y) f(y) \, \sigma(dy), \quad x \in \mathbb{R}^n \setminus \partial \Omega, \end{aligned}$$

with G Green's function for $-\Delta$ on \mathbb{R}^n , are the **single** and **double layer potential operators** respectively.

The Neumann-Poincaré operator is the singular integral:

$$\mathcal{K}_{\partial \Omega} f(x) = \frac{1}{\omega_n} \text{p.v.} \int_{\partial \Omega} \frac{\langle y-x, \nu_y \rangle}{|x-y|^n} f(y) \, \sigma(dy),$$

with ω_n the area of the unit sphere and ν_y the outward normal to $\partial \Omega$.

It holds:

$$\mathcal{K}_{\partial \Omega} = \frac{1}{2} (\text{Tr}_i + \text{Tr}_e) \circ \mathcal{D}_{\partial \Omega}. \quad (3)$$

• If Ω is an **extension domain**, $f \in \mathcal{B}(\partial \Omega)$, $g \in \mathcal{B}'(\partial \Omega)$ with $\langle g, 1 \rangle_{\mathcal{B}', \mathcal{B}} = 0$ (i.e., $g \in \mathcal{B}'_0(\partial \Omega)$), then $u = \mathcal{S}_{\partial \Omega} g - \mathcal{D}_{\partial \Omega} f$, where

$$\begin{aligned} \mathcal{S}_{\partial \Omega} : g \in \mathcal{B}'_0(\partial \Omega) &\mapsto u_{\mathcal{S}} \text{ weak sol. to (2) for } f = 0, \\ \mathcal{D}_{\partial \Omega} : f \in \mathcal{B}(\partial \Omega) &\mapsto u_{\mathcal{D}} \text{ weak sol. to (2) for } g = 0, \end{aligned}$$

are the **generalized layer potential operators**.

The Neumann-Poincaré operator is defined as $\mathcal{K}_{\partial \Omega} : \mathcal{B}(\partial \Omega) \rightarrow \mathcal{B}(\partial \Omega)$ by (3).

4 Imaging results

We generalize imaging results given in [1] in the Lipschitz case.

• **Representation formula:** let Ω be an **extension domain** and $g \in \mathcal{B}'_0(\partial \Omega)$. Then, the unique weak solution to (1) is:

$$u = H - \mathcal{S}_{\partial D} \varphi \quad \text{with} \quad H = \mathcal{S}_{\partial \Omega} g - \mathcal{D}_{\partial \Omega} f,$$

where $f = \text{Tr}_i u|_{\partial \Omega} \in \mathcal{B}(\partial \Omega)$ and $\varphi \in \mathcal{B}'_0(\partial D)$ such that:

$$\left(\frac{k+1}{2k-1} I + \mathcal{K}_{\partial D}^* \right) \varphi = \frac{\partial H}{\partial \nu} \Big|_{\partial D}.$$

• **Monotonous identification:** let $D_1 \subset D_2 \subset \subset \Omega$ be **extension domains** of \mathbb{R}^n . If for some $g \in \mathcal{B}'_0(\partial \Omega) \setminus \{0\}$, the solutions u_1 and u_2 to problem (1) for $D = D_1$ and $D = D_2$ resp. satisfy $\text{Tr}_i u_1|_{\partial \Omega} = \text{Tr}_i u_2|_{\partial \Omega}$, then $D_1 = D_2$.

• **Identification of disks:** let Ω be an **extension domain of \mathbb{R}^2** and $D_1, D_2 \subset \subset \Omega$ be **disks**. If for some $g \in \mathcal{B}'_0(\partial \Omega) \setminus \{0\}$, the solutions u_1 and u_2 to problem (1) for $D = D_1$ and $D = D_2$ resp. satisfy $\text{Tr}_i u_1|_{\partial \Omega} = \text{Tr}_i u_2|_{\partial \Omega}$, then $D_1 = D_2$.

References

- [1] H. AMMARI AND H. KANG, *Reconstruction of Small Inhomogeneities from Boundary Measurements*, Springer Science & Business Media, springer-verlag ed., Aug. 2004.
- [2] M. BIEGERT, *On Traces of Sobolev Functions on the Boundary of Extension Domains*, Proceedings of the American Mathematical Society, 137 (2009), pp. 4169–4176. Publisher: American Mathematical Society.
- [3] M. R. LANCIA, *A Transmission Problem with a Fractal Interface*, Zeitschrift für Analysis und ihre Anwendungen, 21 (2002), pp. 113–133.
- [4] G. VERCHOTA, *Layer potentials and regularity for the Dirichlet problem for Laplace's equation in Lipschitz domains*, Journal of Functional Analysis, 59 (1984), pp. 572–611.