



HAL
open science

Tooth wear analyses track niche partitioning at Gerakarou, a 1.8 Ma old site from Greece

Evangelia Alifieri, Emilie Berlioz, Anastasia G Gkeme, Dimitris S Kostopoulos, Gildas Merceron

► **To cite this version:**

Evangelia Alifieri, Emilie Berlioz, Anastasia G Gkeme, Dimitris S Kostopoulos, Gildas Merceron. Tooth wear analyses track niche partitioning at Gerakarou, a 1.8 Ma old site from Greece. *Quaternary Science Reviews*, 2024, 334, pp.108712. 10.1016/j.quascirev.2024.108712 . hal-04573203

HAL Id: hal-04573203

<https://hal.science/hal-04573203>

Submitted on 13 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License



Tooth wear analyses track niche partitioning at Gerakarou, a 1.8 Ma old site from Greece

Evangelia Alifieri^a, Emilie Berlioz^{b,*}, Anastasia G. Gkeme^a, Dimitris S. Kostopoulos^a, Gildas Merceron^c

^a Laboratory of Geology and Palaeontology, School of Geology, Aristotle University of Thessaloniki, 54124, Thessaloniki, Greece

^b EVOADAPTA, Universidad de Cantabria, Santander, Spain

^c Laboratoire PALEVOPRIM, UMR 7262, CNRS & Université de Poitiers, France

ARTICLE INFO

Handling editor: Mira Matthews

Keywords:

Geometric morphometrics
Mesowear
Dental microwear texture analysis (DMTA)
Early Pleistocene
Villafranchian
Ungulates

ABSTRACT

The climatic instability of the Early Pleistocene induced regional paleoenvironmental changes, which impacted faunal composition and dynamics. It coincides with the *Homo* dispersal out of Africa. The role played by the Eastern Balkans in this dispersal is at the heart of current scientific debates. The rich fossil assemblage of Gerakarou-1 (2–1.8 Ma; northern Greece) can provide important insight into the environmental conditions of that time and area. In this study, we provide insights into the dietary habits of the herbivore guild of Gerakarou-1 at both short and long-term scales based on two complementary dental proxies, mesowear and microwear textures applied on 70 specimens and we infer the local paleoecology and paleoenvironment. As diet varies between and among populations within the same species, we conduct a population-level study, including a reference dataset of 144 extant individuals belonging to six populations of ruminants with well-known ecology. As we aim to contribute to the collective effort of the community to improve the mesowear methodology, we propose and discuss here the benefits of an alternative, quantitative Geometric Morphometric approach. We show that the occlusal relief, as calculated in the classical mesowear approach, is a variable of little relevance and that the quantitative approach allows better discrimination between dietary categories. By proposing for the first time a multi-scale and multi-proxies study of the entire herbivore guild at Gerakarou-1, we nuance and clarify the paleoecology and niche-partitioning of the ungulates from this site and improve past paleoenvironmental interpretations.

1. Introduction

In the Early Pleistocene of the Northern Hemisphere, alternating glacial and interglacial cycles led to a gradual global cooling and habitat opening (Bonifay and Brugal, 1996; Gibbard and Cohen, 2008; Kahlke et al., 2011; Lisiecki and Raymo, 2005). Large-scale global changes have impacts on local environments as they significantly affect floral and faunal assemblages, and thus ecological and biogeographical aspects (Kahlke et al., 2011 and references therein). This topic is of prime interest as the Early Pleistocene coincides with the dispersal of the genus *Homo* from Africa, as documented in the locality of Dmanisi, Georgia (1.85–1.77 Ma; Ferring et al., 2011). The diversity of habitats available during the dispersal of early humans throughout Europe is at the heart of the discussions (Bartolini-Lucenti et al., 2022; Spassov, 2016; van der

Made and Mateos, 2010). Due to its location, Eastern Balkans may have played a crucial role in the early *Homo* dispersal into Europe (Harvati and Roksandic, 2017), although there are currently no known human fossil records in Europe before the Middle Pleistocene (Tourloukis and Karkanas, 2012). The Early Pleistocene site of Gerakarou-1 (Mygdonia Basin, Greece) is biochronologically estimated to be around 2–1.8 Ma (Konidaris et al., 2015), making it roughly contemporaneous with the first occurrence of the genus *Homo* out of Africa.

Being the most durable organs and thus best-preserved remains found in paleontological sites, teeth of herbivorous mammals are often used to infer paleoecology and paleoenvironmental conditions (Merceron et al., 2023). Here, we extract two dental proxies that constitute complementary temporal windows over the dietary habits of past ungulates (Davis and Pineda-Munoz, 2016). Numerous studies on wild

* Corresponding author.

E-mail addresses: evi0812@yahoo.gr (E. Alifieri), emilie.berlioz@unican.es (E. Berlioz), sia2006@yahoo.com (A.G. Gkeme), dkostop@geo.auth.gr (D.S. Kostopoulos), gildas.merceron@univ-poitiers.fr (G. Merceron).

<https://doi.org/10.1016/j.quascirev.2024.108712>

Received 6 November 2023; Received in revised form 3 May 2024; Accepted 4 May 2024

Available online 9 May 2024

0277-3791/© 2024 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC license (<http://creativecommons.org/licenses/by-nc/4.0/>).

extant populations with long-term records of ecological preferences (e.g. Berlioz et al., 2017; Kubo et al., 2017; Kubo and Fujita, 2021; Kubo and Yamada, 2014; Merceron et al., 2021) and controlled experimental settings (e.g. Louail et al., 2021; Winkler et al., 2020) have shown the relations between food properties and resulting dental microwear textures. Besides, molar mesowear is known to inform feeding habits on a much longer scale (Ackermans et al., 2020).

Here, by combining dental microwear texture analysis with a new protocol of molar mesowear analysis using geometric morphometrics, set up on a comparative dataset of extant species (Fig. 1), we analyze 70 specimens of five meso-herbivores from Gerakarou-1 to decipher dietary preferences, niche partitioning, and habitats available in this Early Pleistocene site.

2. Material and methods

2.1. Material

Gerakarou-1 is one of the 12 known fossil sites from the Mygdonia basin (Koufos and Kostopoulos, 2016), which covers the Lower Pleistocene (2.6–0.8 Ma; Gibbard and Head, 2009). Based on typical late Villafranchian faunal elements and more recent taxa, Gerakarou-1 is attributed to MNQ 18 (Koufos, 2001) before the Olduvai event (1.92–1.78 Ma; Cohen and Gibbard, 2022). In Gerakarou-1, almost 60% of the faunal assemblage comprises ungulates while carnivore species account for around 30% and rodents around 10% (Koufos, 2001).

Only adults with erupted UM3/lm3 were considered for the analyses. We excluded teeth at the severely worn stage or unworn. As the first molars are often too worn, and the third molars are frequently not worn enough, we work preferentially on the second molars. This is the tooth most studied in Dental Microwear Texture (DMT) studies, which facilitates inter-team comparisons. We analysed 70 molars belonging to *Equus altidens* (N = 8), *Antilope koufosi* (N = 3), *Gazella bouvrainae* (N = 30), *Croizetoceros ramosus gerakarensis* (hereafter *C. ramosus*; N = 26), and

Eucladoceros tegulensis (N = 3) (Table S1). The few specimens of *Leptobos* cf. *etruscus* and an undetermined Bovidae were excluded. All fossil specimens are housed in the Museum of Geology, Paleontology, Palaeoanthropology (LGPOT) of the Aristotle University of Thessaloniki, Greece.

As diet varies between populations within a species (Gebert and Verheyden-Tixier, 2001) depending on age, sex, physiological condition, and season, we conducted a population-level study. When considered among herbivore ungulates, phylogenetic relationships and masticatory movements are known to have little impact on the two dental proxies used in this study (Calandra and Merceron, 2016; DeSantis et al., 2018; Schulz et al., 2010). Therefore, in the absence of strict present-day representatives, our comparative material comprises 144 individuals from six European populations of extant species of ruminants (Table 1; Table S1) with well-known ecology, from biomes comparable with Early Pleistocene European landscapes. The sympatric *Ovis ammon* (N = 21), *Rupicapra rupicapra* (N = 19), *Cervus elaphus* (N = 19), and *Capreolus capreolus* (N = 18) samples come from the Bauges Natural Regional Park (French Alps; Redjadj et al., 2014). *Alces alces* (N = 28) are from the Biebrza marshes (Eastern Poland; Kuijper et al., 2016), and *Bos taurus* (N = 39) are from a Camargue sample (Rhône delta, France; Blaise, 2005; Duncan and D’Herbes, 1982). Previous dental microwear texture (DMT) studies have focused on these extant populations (Berlioz et al., 2022; Hermier et al., 2020; Merceron et al., 2021). All dental molds are stored at the PALEVOPRIM lab (University of Poitiers, France).

In our comparative material, the population that is the most engaged in grazing is *Bos taurus* (Duncan and D’Herbes, 1982), and the most browsing is the moose *Alces alces* (Kuijper et al., 2016). The red deer *Cervus elaphus*, the chamois *Rupicapra rupicapra*, and the mouflon *Ovis ammon* are known to be mixed feeders (Gebert and Verheyden-Tixier, 2001; Marchand et al., 2013; Pérez-Barbería and Nores, 1994) while the roe deer *Capreolus capreolus* is a selective browser prone to select the most nutritive items (Tixier et al., 1997).

2.2. Methods

Regarding dental facets for DMT analysis, we followed recommendations by Ramdarshan et al. (2017): we focused on the mesio-lingual facet of the protocone on upper molars and the disto-labial facet of the protoconid on lower molars. For the upper teeth of equids, we adapted the location by molding the lingual facet of the protocone. Cleaning and molding procedures are detailed here: <https://zenodo.org/doi/10.5281/zenodo.7305566>. For scanning with “Trident”, the Leica DCM8 (Leica Microsystems) microscope of PALEVOPRIM Lab has been used; for pre-treating and analysing the data we used Leica MAP v. 8.2Sfrax software (Surfract, www.surfract.com), following procedures by Merceron et al. (2016) and in supplementary information therein; appendix A1). We applied Scale Sensitive Fractal Analysis and extracted complexity (Asfc), heterogeneity of complexity (HASfc; nine cells), and anisotropy (epLsar, measured following the traditional method; Calandra et al., 2022; Scott et al., 2006). Whenever the sample size was large enough (N ≥ 5; Table 1, Table S1), we also calculated inter-individual disparities for Asfc and epLsar (Disp-Asfc and Disp-epLsar, respectively) following Levene’s equations (Plavcan and Cope, 2001; Disp-X = $\Sigma(\ln(X_i) - \text{Med}_i) >_n(\ln(X_i)) / n$). This approach allows the assessment of inter-sample variation, which has interesting ecological implications in terms of diet breadth, social behaviours, and habitat use (Burgman et al., 2016; Merceron et al., 2021).

Regarding molar mesowear analysis, we followed the procedures by Rivals and Semprebon (2006). The coding of the occlusal relief (OR) and the apex shape (CS) generates a mesowear score (Ms) for each individual. A lower Ms corresponds to sharper cusps with higher relief, while a higher Ms indicates rounder and lower cusps. Our study is part of an ongoing collective effort to refine the link between mesowear and dietary ecology and improve the discriminatory power of the approach




Fig. 1. Combination of long-term (Mesowear) and short-term (Microwear) approaches for palaeoecological inferences. Top of the figure: Geometric Morphometric MesoWear (GMMW) is based on the occlusal relief as a continuous variable and the apex cusp shape determined by an outline digitalized with 2-D landmarks and semi-landmarks. DMT (lower part): Dental microwear texture analysis allows the quantification of different texture parameters using 3D cloud points here represented as a photosimulation and a false-colour elevation map for the lower second molars of *Eucladoceros tegulensis* (GER 136; middle of the figure); and *Equus altidens* (GER 33; bottom of the figure).

Table 1

Summary statistics for the molar mesowear and dental microwear textures for both extant and extinct species (for populations with N>5). N: sample size; SE: standard error; SD: standard deviation; ORR: Occlusal Relief Ratio; ACS: Apex Cusp Shape using Geometric Morphometric Mesowear Method; Ms: Mesowear score (Rivals and Semperebon, 2006); Asfc: complexity; ePLsar: anisotropy; Disp-Asfc: disparity of complexity; HAsfc: heterogeneity of complexity (9 cells). Regarding DMT, only variables displaying significant variations are shown.

| Variables | | Extant Species | | | | | | Fossil Species | | |
|-----------|------|------------------|---------------------|-----------------|-------------------|---------------------|-----------------|----------------------|-------------------|--------------------|
| | | <i>B. taurus</i> | <i>R. rupicapra</i> | <i>O. ammon</i> | <i>C. elaphus</i> | <i>C. capreolus</i> | <i>A. alces</i> | <i>G. bouvrainae</i> | <i>C. ramosus</i> | <i>E. altidens</i> |
| | N | 39 | 19 | 21 | 19 | 18 | 28 | 30 | 26 | 8 |
| ORR | Mean | 0.59 | 0.56 | 0.46 | 0.44 | 0.39 | 0.33 | 0.32 | 0.33 | 0.12 |
| | SE | 0.02 | 0.03 | 0.02 | 0.01 | 0.02 | 0.01 | 0.02 | 0.02 | 0.02 |
| | SD | 0.12 | 0.11 | 0.08 | 0.04 | 0.08 | 0.07 | 0.09 | 0.09 | 0.05 |
| ACS (PC1) | Mean | 0.06 | -0.01 | 0.02 | -0.04 | -0.01 | -0.05 | -0.03 | -0.01 | 0.12 |
| | SE | 0.01 | 0.02 | 0.02 | 0.02 | 0.03 | 0.01 | 0.02 | 0.02 | 0.03 |
| | SD | 0.09 | 0.09 | 0.11 | 0.07 | 0.11 | 0.07 | 0.10 | 0.10 | 0.07 |
| Ms | Mean | 0.92 | 0.89 | 0.81 | 0.84 | 0.72 | 0.57 | 0.87 | 0.92 | 2.25 |
| | SE | 0.04 | 0.07 | 0.09 | 0.09 | 0.11 | 0.10 | 0.17 | 0.12 | 0.25 |
| | SD | 0.27 | 0.32 | 0.40 | 0.37 | 0.46 | 0.50 | 0.94 | 0.63 | 0.71 |
| Asfc | Mean | 1.59 | 1.85 | 2.04 | 1.38 | 2.61 | 4.63 | 2.01 | 1.33 | 0.96 |
| | SE | 0.16 | 0.23 | 0.19 | 0.14 | 0.59 | 0.72 | 0.20 | 0.13 | 0.11 |
| | SD | 0.99 | 1.00 | 0.89 | 0.60 | 2.49 | 3.82 | 1.11 | 0.65 | 0.30 |
| epLsar | Mean | 5.29 | 4.29 | 3.80 | 5.89 | 6.05 | 1.87 | 4.51 | 6.39 | 3.88 |
| | SE | 0.34 | 0.43 | 0.57 | 0.62 | 0.55 | 0.24 | 0.34 | 0.44 | 0.63 |
| | SD | 2.11 | 1.88 | 2.61 | 2.72 | 2.34 | 1.25 | 1.84 | 2.25 | 1.78 |
| Disp-Asfc | Mean | 0.21 | 0.23 | 0.25 | 0.19 | 0.43 | 0.39 | 0.27 | 0.21 | 0.14 |
| | SE | 0.04 | 0.05 | 0.04 | 0.03 | 0.09 | 0.07 | 0.04 | 0.03 | 0.02 |
| | SD | 0.24 | 0.21 | 0.16 | 0.15 | 0.38 | 0.35 | 0.20 | 0.16 | 0.06 |
| HAsfc | Mean | 0.41 | 0.39 | 0.33 | 0.58 | 0.36 | 0.69 | 0.41 | 0.34 | 0.24 |
| | SE | 0.05 | 0.05 | 0.02 | 0.12 | 0.03 | 0.07 | 0.02 | 0.03 | 0.02 |
| | SD | 0.28 | 0.24 | 0.11 | 0.50 | 0.14 | 0.39 | 0.13 | 0.17 | 0.07 |

(Ackermans, 2020; Alifieri et al., 2021, 2022; Mühlbachler et al., 2023). This is why we also proposed a Geometric Morphometric MesoWear method (GMMW). We photographed the dental rows of all specimens, focusing preferentially on second molars (UM2/lm2; Table S1), using an Olympus TG-3 and a Nikon D5200 digital cameras and a professional photographic table equipped with a graduated arm and lighting. For the upper molars, the picture was taken on the labial side, and for the lower molars, on the lingual side. We favored the right molars, but, when these were unusable, we photographed the left ones, and mirrored the picture along the X-axis. We orientated the samples by placing them on foam pads, so that the walls of the molars were perfectly perpendicular to the camera lens. Each photo was taken with the same scale and the same distance between the lens and the specimen. We then digitally computed the Occlusal Relief Ratio (ORR; corresponding to H/IC, as illustrated in Fig. 1) with tpsDig (v. 2.32; Rohlf, 2021). The Apex Cusp Shape (ACS; Fig. 1) is digitalized by 40 2-D semi-landmarks from the mesial to the distal side of the anterior cusp (or posterior whenever the anterior was damaged; Table S1) with TpsUtil (v 1.78; Rohlf, 2021). Then we selected the upper-25% semi-landmarks reflecting the shape of the apex of the cusp and performed generalized least squares procrustes superimpositions with MorphoJ (v 107a; Klingenberg, 2011) to remove scale, rotation, and translation differences and work in the same morphological space (Kendall, 1977; Slice, 2007; Webster and Sheets, 2010).

Then, we generated an eco-dietary space by conducting a Principal Component Analysis on extant specimens. In this PCA, we secondarily included the fossil specimens, using PC scores from the PCA on extant. Analyses focus on PC1, which determines the Apex Cusp Shape (56.93% of the variance; Figs. 1 and 2, Table S3, Fig. S1).

After rank-transformations (Conover and Iman, 1981), we performed one-way ANOVAs (Table S2) to determine whether the DMT and GMMW significantly varied, followed by Tukey (HSD) and Fisher (LSD) post-hoc tests to identify inter-group significant differences (Table 2).

3. Results

3.1. Dental microwear texture analysis

The ANOVA detected significant differences in complexity, disparity

of complexity, heterogeneity of complexity, and anisotropy (Table S2). Among extant populations, *Alces alces* differs from all other extant species in having a higher complexity and lower anisotropy (Tables 1 and 2, Fig. 2). It also has a higher heterogeneity of complexity than all species, but *C. elaphus*. *A. alces* also has a higher disparity of complexity than *B. taurus* and *C. elaphus*. *Alces alces* differs significantly from *C. capreolus* in having higher complexity, heterogeneity of complexity, and lower anisotropy. However, these cervids share a high inter-individual disparity of complexity (Table 1) compared to the other extant ruminants. *O. ammon*, *R. rupicapra*, and *C. elaphus*, known as mixed feeders, and *B. taurus*, known as grazer, all show significant differences with *A. alces*. It is worth noting that there are no significant differences between *R. rupicapra* and *B. taurus* and that *C. elaphus* has the lowest complexity among all the extant species (Tables 1 and 2, Fig. 2).

Looking at the fossil samples, *Equus altidens* differs from all extant species in having lower complexity (except *C. elaphus*) and lower heterogeneity of complexity (except *O. ammon*; Tables 1 and 2, Fig. 2). It also differs in having a lower anisotropy than *C. elaphus* and *C. capreolus* and a lower disparity of complexity than *C. capreolus* and *A. alces*. *Gazella bouvrainae* has a higher complexity than *B. taurus* and *C. elaphus*. It has lower anisotropy than *C. capreolus* and *C. elaphus* but higher than *A. alces*. *Gazella bouvrainae* differs from *R. rupicapra* and *O. ammon* in having a higher heterogeneity of complexity (but lower than that of *A. alces*). *Croizetoceros ramosus* has significantly lower complexity than *A. alces*, *C. capreolus*, and *O. ammon* and lower heterogeneity of complexity than *C. capreolus* and *A. alces*. All but one of the *Antelope koufosi* and *Eucladoceros tegulensis* (Fig. 2) plot far from *A. alces*. However, the low sample size (N = 3 in both cases) for these two species forbids us to apply statistical tests and discuss further the differences with extant species.

3.2. Molar mesowear analysis

The occlusal relief (OR), one of the two variables used to calculate the mesowear score (Ms), is high for 100% of the individuals used as a reference, and for 78% of the fossil individuals (Table S1). The extant *A. alces* has the lowest mesowear score (Ms; Table 1, Fig. S2), and *B. taurus* the highest (Table 1; Fig. S2). Other species plot between these




Fig. 2. Microwear and Mesowear results for extant and extinct populations. Mean and confidence interval of the mean at 95% of anisotropy (epLsar) and complexity (Asfc, A and B) and of occlusal relief ratio (ORR) and apex cusp shape calculated with the GMMW approach (PC1, C and D) for extant and extinct species.

Table 2

Results of post hoc tests for between samples differences. Significant differences supported by Fisher's Least Significant Difference tests (LSD), which are underlined when these differences are also supported by Tukey's Honestly Significant Difference tests (HSD). Analyses were done only whenever $N \geq 5$ specimens.

| | Extant Species | | | | | | Fossil Species | | |
|---------------------|--------------------------|---------------------|-----------------|-----------------------------|---------------------|---|-----------------------------|------------------------------------|---|
| | <i>B. taurus</i> | <i>R. rupicapra</i> | <i>O. ammon</i> | <i>C. elaphus</i> | <i>C. capreolus</i> | <i>A. alces</i> | <i>G. bouvrinae</i> | <i>C. ramosus</i> | <i>E. altidens</i> |
| <i>B. taurus</i> | / | | Asfc epLsar | HAsfc | Disp-Asfc | <u>Asfc epLsar</u> Disp- Asfc <u>HAsfc</u> | Asfc | | Asfc HAsfc |
| <i>R. rupicapra</i> | ACS | / | | epLsar HAsfc | epLsar | <u>Asfc epLsar</u> <u>HAsfc</u> | | <u>epLsar</u> | Asfc HAsfc |
| <i>O. ammon</i> | <u>ORR</u> | ORR | / | Asfc <u>epLsar</u> HAsfc | <u>epLsar</u> | <u>Asfc epLsar</u> <u>HAsfc</u> | HAsfc | Asfc <u>epLsar</u> | <u>Asfc</u> |
| <i>C. elaphus</i> | <u>ORR</u> <u>ACS</u> | ORR | ACS | / | Disp-Asfc HAsfc | <u>Asfc epLsar</u> Disp- <u>Asfc</u> | Asfc epLsar | <u>HAsfc</u> | epLsar <u>HAsfc</u> |
| <i>C. capreolus</i> | <u>ORR</u> <u>ACS</u> | <u>ORR</u> | <u>ORR</u> | ORR | / | <u>Asfc epLsar</u> <u>HAsfc</u> | epLsar | Asfc | Asfc epLsar Disp- Asfc HAsfc |
| <i>A. alces</i> | <u>ORR</u> <u>ACS</u> | <u>ORR</u> | <u>ORR</u> ACS | <u>ORR</u> | ORR | / | <u>Asfc epLsar</u> HAsfc | <u>Asfc epLsar</u> <u>HAsfc</u> | <u>Asfc epLsar</u> Disp- Asfc <u>HAsfc</u> |
| <i>G. bouvrinae</i> | <u>ORR</u> <u>ACS</u> | <u>ORR</u> | <u>ORR</u> | <u>ORR</u> | ORR | / | / | Asfc HAsfc <u>epLsar</u> | <u>Asfc</u> <u>HAsfc</u> epLsar |
| <i>C. ramosus</i> | <u>ORR</u> <u>ACS</u> | <u>ORR</u> | <u>ORR</u> | <u>ORR</u> | ORR | ACS | / | / | epLsar |
| <i>E. altidens</i> | <u>ORR</u> | <u>ORR</u> ACS | <u>ORR</u> ACS | <u>ORR</u> ACS | <u>ORR</u> ACS | <u>ORR</u> ACS | <u>ORR</u> ACS | <u>ORR</u> ACS | / |

two end-members with significant overlap. When it comes to fossils, the mesowear score of *Equus altidens* is the only one that is different from all extant and extinct ruminants in having the highest values (Tables 1 and S4, Fig. S2).

When considering the occlusal relief ratio as a continuous variable (ORR; Tables 1 and 2, Fig. 2), it occurs that *A. alces* has the lowest relief and *B. taurus* the highest. Other species plot between them (Fig. 2). The post-hoc test shows that *A. alces* differs from other extant species (but marginally with *C. capreolus*) in having a significantly lower occlusal relief ratio (Tables 1 and 2; Fig. 2). *Capreolus capreolus* also differs from *B. taurus*, *R. rupicapra*, *O. ammon* and marginally from *C. elaphus* in having a significantly lower occlusal relief ratio (Tables 1 and 2; Fig. 2). *Cervus elaphus* has a significantly lower occlusal relief ratio than *B. taurus* and marginally lower than *R. rupicapra*. *Ovis ammon* also has a lower occlusal relief ratio than *B. taurus*.

The Apex Cusp Shape (Table S3; Fig. S1) significantly varies along the PC1, explaining 56.93 % of the variance. PC2 does not reflect mesowear but is rather related to cusp inclination (mesial or distal) and explains 32.32% of the variance (Table S3; Fig. S1). *Bos taurus* significantly differs from all species but *O. ammon* in having higher values evidencing more rounded cusp apices (PC1, Tables 1 and 2; Fig. 2). *Ovis ammon* also differs from *C. elaphus* and *A. alces* in having higher PC1, meaning rounder cusps (Tables 1 and 2; Fig. 2). PC1 of the Model PCA based on extant populations and setting our interpretive eco-dietary space explains 58.78% of the variance for the fossil dataset (Table S3). When considering the occlusal relief ratio, equids show the lowest value, significantly different from all extant and extinct ruminants. *Equus altidens* displays the highest coordinates along PC1 (Fig. 2, Table 1), being not different from *B. taurus*. The occlusal relief ratio of *G. bouvrinae* and *C. ramosus* are similar to the ones of *A. alces* and *C. capreolus*. They show significant differences in occlusal relief ratio with the three extant bovids and *C. elaphus*. Their coordinates along PC1 have lower values than *B. taurus*, confirming sharper cusps. *Eucladoceros tegulensis* and *A. koufosi* have occlusal relief ratio similar to *A. alces*. *Eucladoceros tegulensis* displays low values along the PC1 supporting sharp cusps, while two of the three specimens of *A. koufosi* have high values, the third one plotting near the mean values for the three extant cervids (Fig. 2).

4. Discussion

Molar mesowear bears a long-term dietary signal, at least on a yearly scale (Ackermans, 2020). In our study, the occlusal relief on mandibular cheek teeth of mid-latitude C₃ plant-eating ruminants does not discriminate between extant populations well-known to forage on different resources (OR; Tables S1 and S4). This finding can be generalized to a larger geographical scale since data from a recent study show similar results (Mihlbachler et al., 2023). The cusp shape is by then the only parameter that significantly contributes to the mesowear scores (Ms). It results that, in our study, all six extant ruminants share low mesowear scores, overlapping with the ranges of extant browsers, mixed feeders, and C₃ grazers in the comparative datasets from Fortelius and Solounias (2000; Fig. S2). In addition to the need for improving the measurement variables, such results highlight the need to set up *ad hoc* comparative models considering the fine-scale ecology. Indeed, large datasets built with species represented by few individuals occupying different biomes dispersed at different latitudes and sampled at different years, or even decades, might be, at best, non-adapted and, at worst, a source of misinterpretation regarding dietary habits (Fortelius and Solounias, 2000; Mihlbachler et al., 2023).

We calculated an occlusal relief ratio and modeled Apex Cusp Shape to overcome these limitations, thus treating these variables as continuous. They significantly discriminate extant populations. It is worth noting that the highest occlusal relief ratio is not found for browsing species but for *Bos taurus*, the most grazing species of our extant dataset, challenging the traditional view of a positive correlation between low

occlusal relief and the consumption of monocots. One could interpret the low occlusal relief ratio of the two most browsing species (*Alces alces* and *Capreolus capreolus*) as the result of a diet rich in lignin components, requiring a higher chewing stroke to comminute these food items. That said, the Apex Cusp Shape results are consistent with former studies (Solounias and Semprebon, 2002): sharper apices were found for browsing species and blunter apices for grazing ones. The molar mesowear (GMMW) proposed here clearly shows better dietary discrimination.

Combined molar mesowear and dental microwear textures provide complementary long-term and short-term signals offering a more comprehensive view into the paleoecology of species and populations. The analysis of the herbivore guild of the Greek faunal assemblage of Gerakarou-1 emphasizes food preferences and ecological niche partitioning. *Gazella bouvrinae* had a mixed-feeding diet similar to modern *R. rupicapra* and *O. ammon*, as the similar short-term DMT (complexity and anisotropy) and long-term Apex Cusp Shape of the three species demonstrate. *Croizerceros ramosus* likely fed on resources more similar to that of the population of the extant *C. elaphus*. This small-size extinct deer likely consumed less lignified tissues and more herbaceous vegetation than *G. bouvrinae* as it displays higher texture anisotropy than *G. bouvrinae*. *Equus altidens* composed the third most important sample in the Gerakarou-1 community. Its diet was mainly based on herbaceous vegetation, as indicated by its Apex Cusp Shape and microwear values. The two other species investigated in this study, *E. tegulensis* and *A. koufosi*, are inadequately represented, so making interpretations is limited. However, we can pinpoint that the molar mesowear depicts *E. tegulensis* as a browser while the microwear texture supports mixed feeders. Observing contrasting results between the two approaches is not inconsistent: mesowear reflects an average diet, while microwear texture provides information on the last few weeks of the animal's life. On the other hand, *A. koufosi* was more engaged in grazing.

Considering that a taphocenosis does not represent a biocenosis, but an undetermined portion of it (Soligo and Andrews, 2005) and that the extinct species composing the taphocenosis might have originated from different habitats in the surrounding of the site (Merceron et al., 2021), the dietary reconstruction of the herbivorous species draws a more accurate picture of the habitat availabilities at a local scale. Our results exclude the occurrence of wooded landscapes, as none of the species were browsers. They all include a significant proportion of herbaceous resources, whose the development would have been severely restricted by a dense tree canopy. Indeed, *E. tegulensis* is the only species that could be classified as a mixed-feeding ruminant engaged in browsing (Berlitz et al., 2018). While the feeding habits of the ruminants support the presence of open habitats with woody vegetation present either as patches or shrub/bushland, the feeding habits of the equid support that it occupied the habitats providing the most abundant herbaceous resources. By proposing for the first time a multi-scale study of the entire herbivore guild at Gerakarou-1, we enhance and clarify previous paleoenvironmental interpretations, whether based on the faunal assemblage (Kahlke et al., 2011) or exclusively on the study of the ecology of Gerakarou-1 deer (Berlitz, 2017; Berlitz et al., 2018).

5. Conclusions

The present study investigates the feeding habits of the guild of the meso-herbivores found at Gerakarou-1, an Early Pleistocene faunal assemblage in northern Greece. The combination of dental microwear texture analysis and an improved version of molar mesowear analysis using geometric morphometry first applied on reference extant populations with known differences in diet, enables us to take a further look into the paleoecology, niche partitioning, and paleoenvironment of the ungulates from this site. *Equus altidens* is the most grazing species from this assemblage. The other species, all ruminants, were mixed feeders, with *Eucladoceros tegulensis* most likely the most engaged in browsing. Our results support the presence of open habitats with a low abundance of

woody patches or shrub/bushlands. Gerakarou-1 is one of the oldest sites in the Mygdonian basin. With a succession of sedimentary formations, including several fossil sites (Konidaris et al., 2015; Kostopoulos et al., 2023; Koufos, 2001), the Mygdonian basin provides a unique opportunity to decipher fine-scale changes in the ecological niches of the meso-herbivores based on complementary dental proxies, as a bridge to assess environmental variations throughout Early Pleistocene times.

The study also shows the need to continue improving the mesowear tool, particularly by increasing repeatability and moving towards a more effective quantitative (but still affordable and easy to use) approach.

Funding

This study was funded by the CNRS, France [IEA- PICS08245; 2019–2021: « *Exploring Paradolichopithecus from Greece and the early Pleistocene primate evolution* », PIs: G. Merceron and D. S. Kostopoulos], and partly funded by the French Research Agency [ANR-13-JSV7-0008-01 : TRIDENT, PI: G. Merceron]. The first author was granted an ERASMUS + fellowship (2020-0043).

Author contributions

GM and DSK conceptualized the study in the framework of the master thesis of EA. EB-, GM, and DSK trained EA and supervised her work. EA and AG sampled the specimens. EA conducted all analyses and interpreted the results under the supervision of EB and GM. EA, GM, and EB wrote the manuscript. All co-authors collaborated on reviewing and editing the manuscript. All co-authors agreed to the published version of the manuscript.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgments

The authors are thankful to the colleagues who contributed to the field excavations, to Christos Plastiras and Jérôme Surault for their help in the data acquisition, and to Corentin Gibert for discussions on the statistics.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.quascirev.2024.108712>.

References

- Ackermans, N.L., 2020. The history of mesowear: a review. *PeerJ* 8, e8519.
- Ackermans, N.L., Martin, L.F., Codron, D., Hummel, J., Kircher, P.R., Richter, H., Kaiser, T.M., Clauss, M., Hatt, J.-M., 2020. Mesowear represents a lifetime signal in sheep (*Ovis aries*) within a long-term feeding experiment. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 553, 109793.
- Alfieri, E., Berlioz, E., Kostopoulos, D.S., Merceron, G., 2022. Toward quantifying mesowear proxies: geometric morphometric mesowear (GMMW) method. In: 2nd Virtual Conference for Women Archaeologists and Paleontologists.
- Alfieri, E., Berlioz, E., Kostopoulos, D.S., Merceron, G., 2021. Feeding ecology and paleoenvironment of the early Pleistocene Gerakarou herbivore mammal community (Mygdonia basin, Greece) as inferred by DMTA analysis. In: 3rd Palaeontological Virtual Conference.
- Bartolini-Lucenti, S., Cirilli, O., Pandolfi, L., Bernor, R.L., Bukhsianidze, M., Carotenuto, F., Lordkipanidze, D., Tsikaridze, N., Rook, L., 2022. Zoogeographic significance of Dmanisi large mammal assemblage. *J. Hum. Evol.* 163, 103125.
- Berlioz, E., 2017. *Écologie alimentaire et paléoenvironnements des cervidés européens du Pléistocène inférieur: le message des textures de micro-usure dentaire* (PhD Thesis). Université de Poitiers, France.
- Berlioz, E., Azorit, C., Blondel, C., Tellado Ruiz, M.S., Merceron, G., 2017. Deer in an arid habitat: dental microwear textures track feeding adaptability. *Hystrix* 28, 222–230.
- Berlioz, É., Kostopoulos, D.S., Blondel, C., Merceron, G., 2018. Feeding ecology of *Eucladoceros ctenoides* as a proxy to track regional environmental variations in Europe during the early Pleistocene. *Comptes Rendus Palevol* 17, 320–332.
- Berlioz, E., Leduc, C., Hofman-Kamińska, E., Bignon-Lau, O., Kowalczyk, R., Merceron, G., 2022. Dental microwear foraging ecology of a large browsing ruminant in Northern Hemisphere: the European moose (*Alces alces*). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 586, 110754 <https://doi.org/10.1016/j.palaeo.2021.110754>.
- Blaise, E., 2005. L'élevage au Néolithique final dans le sud-est de la France: éléments de réflexion sur la gestion des troupeaux. *Anthropozoologica* 40, 191–216.
- Bonifay, M.-F., Brugal, J.-P., 1996. Biogéographie et biostratigraphie des grandes faunes du Pléistocène inférieur et moyen en Europe du Sud: apport des gisements français. *Paléo* 8, 19–29.
- Burgman, J.H., Leichliter, J., Avenant, N.L., Ungar, P.S., 2016. Dental microwear of sympatric rodent species sampled across habitats in southern Africa: implications for environmental influence. *Integr. Zool.* 11, 111–127.
- Calandra, I., Bob, K., Merceron, G., Blateyron, F., Hildebrandt, A., Schulz-Kornas, E., Souron, A., Winkler, D.E., 2022. Surface texture analysis in Toothfrax and MountainsMap® SSFA module: different software packages, different results? *Peer Community Journal* 2.
- Calandra, I., Merceron, G., 2016. Dental microwear texture analysis in mammalian ecology. *Mamm Rev.* 46, 215–228.
- Cohen, K., Gibbard, P., 2022. Global Chronostratigraphical Correlation Table for the Last 2.7 Million Years v.2019 (Poster Version) vol. 5.
- Conover, W.J., Iman, R.L., 1981. Rank transformations as a bridge between parametric and nonparametric statistics. *Am. Statistician* 35, 124–129.
- Davis, M., Pineda-Munoz, S., 2016. The temporal scale of diet and dietary proxies. *Ecol. Evol.* 6, 1883–1897.
- DeSantis, L., Fortelius, M., Grine, F.E., Janis, C., Kaiser, T.M., Merceron, G., Purnell, M. A., Schulz-Kornas, E., Saarinen, J., Teaford, M., 2018. The phylogenetic signal in tooth wear: what does it mean? *Ecol. Evol.* 8, 11359–11362.
- Duncan, P., D'Herbes, J.-M., 1982. The use of domestic herbivores in the management of wetlands for waterbirds in the Camargue, France. In: *Proc. Of the 3rd Techn. Meeting on West. Pal. Migr. Bird Manag. Munster*, pp. 51–67.
- Ferring, R., Oms, O., Agustí, J., Berna, F., Nioradze, M., Shelia, T., Tappen, M., Vekua, A., Zhvania, D., Lordkipanidze, D., 2011. Earliest human occupations at Dmanisi (Georgian caucasus) dated to 1.85–1.78 Ma. *Proc. Natl. Acad. Sci. U.S.A.* 108, 10432–10436. <https://doi.org/10.1073/pnas.1106638108>.
- Fortelius, M., Solounias, N., 2000. Functional characterization of ungulate molars using the abrasion-attrition wear gradient: a new method for reconstructing paleodiets. *Am. Mus. Novit.* 3301, 1–36.
- Gebert, C., Verheyden-Tixier, H., 2001. Variations of diet composition of red deer (*Cervus elaphus* L.) in Europe. *Mamm Rev.* 31, 189–201.
- Gibbard, P., Cohen, K.M., 2008. Global chronostratigraphical correlation table for the last 2.7 million years. *Episodes* 31, 243–247.
- Gibbard, P.L., Head, M.J., 2009. IUGS ratification of the quaternary system/period and the Pleistocene series/epoch with a base at 2.58 Ma. *Quaternaire. Revue de l'Association française pour l'étude du Quaternaire* 20, 411–412.
- Harvati, K., Roksandic, M. (Eds.), 2017. *Paleoanthropology of the Balkans and Anatolia: Human Evolution and its Context*. Springer.
- Hermier, R., Merceron, G., Kostopoulos, D.S., 2020. The emblematic eurasian villafranchian antelope *Gazellospira* (Mammalia: Bovidae): new insights from the lower Pleistocene Dafnero fossil sites (northern Greece). *Geobios* 61, 11–29.
- Kahlke, R.-D., García, N., Kostopoulos, D.S., Lacomart, F., Lister, A.M., Mazza, P.P., Spassov, N., Titov, V.V., 2011. Western Palaeoartctic palaeoenvironmental conditions during the Early and early Middle Pleistocene inferred from large mammal communities, and implications for hominin dispersal in Europe. *Quat. Sci. Rev.* 30, 1368–1395.
- Kendall, D.G., 1977. The diffusion of shape. *Adv. Appl. Probab.* 9, 428–430.
- Klingenberg, C.P., 2011. MorphoJ: an integrated software package for geometric morphometrics. *Molecular ecology resources* 11, 353–357.
- Konidaris, G.E., Tourloukis, V., Kostopoulos, D.S., Thompson, N., Giusti, D., Michailidis, D., Koufos, G.D., Harvati, K., 2015. Two new vertebrate localities from the early Pleistocene of Mygdonia basin (Macedonia, Greece): preliminary results. *Comptes Rendus Palevol* 14, 353–362.
- Kostopoulos, D.S., Konidaris, G.E., Amanatidou, M., Chitoglou, K., Fragkioudakis, E., Gerakakis, N., Giannakou, V., Gkeme, A., Kalaitzi, C., Tsakalidis, C., Tsatsalis, V., 2023. The new fossil site Krimni-3 in Mygdonia Basin and the first evidence of a giant ostrich in the Early Pleistocene of Greece. *PalZ* 97, 147–161. <https://doi.org/10.1007/s12542-022-00632-8>.
- Koufos, G.D., 2001. The Villafranchian mammalian faunas and biochronology of Greece. *Bollettino-Società Paleontologica Italiana* 40, 217–224.
- Koufos, G.D., Kostopoulos, D.S., 2016. The Plio-Pleistocene large mammal record of Greece: implications for early human dispersals into Europe. In: *Paleoanthropology of the Balkans and Anatolia*. Springer, pp. 269–280.
- Kubo, M.O., Fujita, M., 2021. Diets of Pleistocene insular dwarf deer revealed by dental microwear texture analysis. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 562, 110098.
- Kubo, M.O., Yamada, E., 2014. The inter-relationship between dietary and environmental properties and tooth wear: comparisons of mesowear, molar wear rate, and hypsodonty index of extant sika deer populations. *PLoS One* 9.

- Kubo, M.O., Yamada, E., Kubo, T., Kohno, N., 2017. Dental microwear texture analysis of extant sika deer with considerations on inter-microscope variability and surface preparation protocols. *Biosurface and Biotribology* 3, 155–165.
- Kuijper, D.P.J., Devriendt, K., Bormans, M., Van Diggelen, R., 2016. Do moose redistribute nutrients in low-productive fen systems? *Agric. Ecosyst. Environ.* 234, 40–47.
- Lisiecki, L.E., Raymo, M.E., 2005. A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records. *Paleoceanography* 20, 1–17.
- Louail, M., Neaux, D., Ferchaud, S., Merceron, G., 2021. Investigating the Relationship between Diet and Mandibular Morphology: New Insights from a Controlled-Feeding Experiment on Domestic Pigs.
- Marchand, P., Redjadj, C., Garel, M., Cugnasse, J., Maillard, D., Loison, A., 2013. Are mouflon *Ovis gmelini musimon* really grazers? A review of variation in diet composition. *Mamm Rev.* 43, 275–291. <https://doi.org/10.1111/mam.12000>.
- Merceron, G., Berlioz, E., Vonhof, H., Green, D., Garel, M., Tütken, T., 2021. Tooth tales told by dental diet proxies: an alpine community of sympatric ruminants as a model to decipher the ecology of fossil fauna. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 562, 110077 <https://doi.org/10.1016/j.palaeo.2020.110077>.
- Merceron, G., Ramdarshan, A., Blondel, C., Boisserie, J.-R., Brunetiere, N., Francisco, A., Gautier, D., Milhet, X., Novello, A., Pret, D., 2016. Untangling the environmental from the dietary: dust does not matter. *Proceeding of the Royal Society B* 283, 1–8.
- Merceron, G., Tütken, T., Scott, R., 2023. Editorial preface to special issue: understanding dental proxies of ancient diets. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 111589.
- Mihlbachler, M.C., Barrón-Ortiz, C.I., Rankin, B.D., Theodor, J.M., 2023. Sharpening the mesowear tool: geometric morphometric analysis of cusp shape and diet in ruminants. *Frontiers in Ecology and Evolution* 11, 950463.
- Pérez-Barbería, F.J., Nores, C., 1994. Seasonal variation in group size of Cantabrian chamois in relation to escape terrain and food. *Acta Theriol.* 39, 295–305.
- Plavcan, J.M., Cope, D.A., 2001. Metric variation and species recognition in the fossil record. *Evol. Anthropol.: Issues, News, and Reviews: Issues, News, and Reviews* 10, 204–222.
- Ramdarshan, A., Blondel, C., Gautier, D., Surault, J., Merceron, G., 2017. Overcoming sampling issues in dental tribology: insights from an experimentation on sheep. *Palaeontol. Electron.* 20, 1–19.
- Redjadj, C., Darmon, G., Maillard, D., Chevrier, T., Bastianelli, D., Verheyden, H., Loison, A., Saïd, S., 2014. Intra- and interspecific differences in diet quality and composition in a large herbivore community. *PLoS One* 9, e84756.
- Rivals, F., Semperebon, G.M., 2006. A comparison of the dietary habits of a large sample of the Pleistocene pronghorn *Stockoceros onusrosagris* from the Papago Springs Cave in Arizona to the modern *Antilocapra americana*. *J. Vertebr. Paleontol.* 26, 495–500.
- Rohlf, F.J., 2021. *TpsDig2*. <https://www.sbmorphometrics.org/index.html>.
- Schulz, E., Calandra, I., Kaiser, T.M., 2010. Applying tribology to teeth of hoofed mammals. *Scanning* 32, 162–182.
- Scott, R.S., Ungar, P.S., Bergstrom, T.S., Brown, C.A., Childs, B.E., Teaford, M.F., Walker, A., 2006. Dental microwear texture analysis: technical considerations. *J. Hum. Evol.* 51, 339–349.
- Slice, D.E., 2007. Geometric morphometrics. *Annu. Rev. Anthropol.* 36, 261–281.
- Soligo, C., Andrews, P., 2005. Taphonomic bias, taxonomic bias and historical non-equivalence of faunal structure in early hominin localities. *J. Hum. Evol.* 49, 206–229.
- Solounias, N., Semperebon, G., 2002. Advances in the reconstruction of ungulate ecomorphology with application to early fossil equids. *Am. Mus. Novit.* 3366, 1–49.
- Spassov, N., 2016. Southeastern Europe as a route for the earliest dispersal of *Homo* toward Europe: ecological conditions and the timing of the first human occupation of Europe. In: *Paleoanthropology of the Balkans and Anatolia*. Springer, pp. 281–290.
- Tixier, H., Duncan, P., Sechovic, J., Yant, A., Gleizes, M., Lila, M., 1997. Food selection by European roe deer (*Capreolus capreolus*): effects of plant chemistry, and consequences for the nutritional value of their diets. *J. Zool.* 242, 229–245.
- Tourloukis, V., Karkanas, P., 2012. The Middle Pleistocene archaeological record of Greece and the role of the Aegean in hominin dispersals: new data and interpretations. *Quat. Sci. Rev.* 43, 1–15.
- van der Made, J., Mateos, A., 2010. Longstanding biogeographic patterns and the dispersal of early *Homo* out of Africa and into Europe. *Quaternary International, Oldest Human Expansions in Eurasia: Favouring and Limiting Factors* 223–224, 195–200. <https://doi.org/10.1016/j.quaint.2009.11.015>.
- Webster, M., Sheets, H.D., 2010. A practical introduction to landmark-based geometric morphometrics. *Quantitative Methods in Paleobiology* 16, 168–188.
- Winkler, D.E., Schulz-Kornas, E., Kaiser, T.M., Codron, D., Leichliter, J., Hummel, J., Martin, L.F., Clauss, M., Tütken, T., 2020. The turnover of dental microwear texture: testing the “last supper” effect in small mammals in a controlled feeding experiment. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 557, 109930.