



HAL
open science

Response to commentary "The importance of assessing left ventricular longitudinal function in presence of increased afterload".

Hugues De Courson, Alexandre Loiseau, Grégoire Chadeaux, Matthieu Biais

► To cite this version:

Hugues De Courson, Alexandre Loiseau, Grégoire Chadeaux, Matthieu Biais. Response to commentary "The importance of assessing left ventricular longitudinal function in presence of increased afterload".. Critical Care, 2024, 28 (1), pp.103. 10.1186/s13054-024-04886-3 . hal-04573530

HAL Id: hal-04573530

<https://hal.science/hal-04573530>

Submitted on 13 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

MATTERS ARISING

Open Access



Response to commentary “The importance of assessing left ventricular longitudinal function in presence of increased afterload”

Hugues de Courson^{1,2}, Alexandre Loiseau¹, Grégoire Chadefaux¹ and Matthieu Biais^{1,3,4*}

Dear Editor,

We very much appreciate Dr. Santonocito’s thoughtful comments on our research, titled Myocardial dysfunction assessed by speckle-tracking in good-grade subarachnoid hemorrhage patients (WFNS 1–2): a prospective observational study [1].

The authors raise two valuable points:

Threshold for defining left ventricular damage We agree that the chosen threshold may have been too high for our specific population. We addressed this issue in our paper by presenting and discussing results for a lower threshold of – 17%. However, it’s important to note that the research questioning this common threshold in critical care patients wasn’t published at the time our protocol design and clinical trial registration (NCT03761654) were finalized.

Accounting for high afterload We agree that high afterload could influence the results. Note that the S’ wave

results presented were for the right ventricle. However, we were able to re-analyse the ultrasound images to obtain the value of the lateral S’ wave at the mitral level. So, we performed, as asked, further analysis to assess the concordance and correlation between global longitudinal strain and mitral S’ wave, providing a more comprehensive picture of left ventricular function in this context. The correlation between S’ wave and SLG was very low and not statistically significant ($r = -0.023$; $p = 0.875$). Using a threshold of 6.8 according to the work of Park et al. [2], the concordance rate between GLS and S’ wave was 45% for a pathological GLS threshold of – 20% and 79% for a pathological GLS threshold of – 17% (Fig. 1).

These results therefore suggest that mitral S’ wave analysis is not a better surrogate for GLS in this population.

Abbreviations

GLS Global longitudinal strain
WFNS World Federation of Neurologic Surgeons

Acknowledgements

Authors thank Catherine Bigotto and Laure Estève for technical support

Author contributions

HDC: recruited patients, collected the data, performed statistical analysis and wrote the manuscript. AL: recruited patients, collected the data and helped in writing the manuscript. GC: recruited patients, analyzed the data and helped in writing the manuscript. MB: designed the study, helped analyze the data and helped in writing the manuscript. All authors approved the final version.

Funding

This study was supported by institutional and departmental funds.

Availability of data and materials

All data generated or analyzed during this study are included in this published article.

This comment refers to the article available online at <https://doi.org/10.1186/s13054-023-04738-6>.

This reply refers to the comment available online at <https://doi.org/10.1186/s13054-024-04801-w>.

*Correspondence:

Matthieu Biais
matthieu.biais@chu-bordeaux.fr

¹ Department of Anesthesiology and Critical Care, Bordeaux University Hospital, Place Amélie Raba-Léon, Hôpital Tripode, Troisième Étage Aile 1, 33000 Bordeaux, France

² INSERM, BPH, U1219, Univ. Bordeaux, 33000 Bordeaux, France

³ U1034, Biology of Cardiovascular Diseases, INSERM, 33600 Pessac, France

⁴ University of Bordeaux, Bordeaux, France



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 1 Relationship between left ventricular ejection fraction (LVEF) and Mitral S' velocity. **A** Strain threshold of -20 . **B** Strain threshold of -17

Declarations

Ethics approval and consent to participate

This study was approved by the Ethics Committee (Ile de France Research Subjects Protection Committee VI—ID RCB: 2018-A02434-51, November 21st, 2018). This study was registered on Clinicaltrials.gov: NCT03761654.

Consent for publication

The manuscript has been read and its submission approved by all co-authors. According to the French law, the patients and/or next of kin were informed about the inclusion of their anonymized data in the database, and none declined participation.

Competing interests

M.B received honoraria from Edwards Lifesciences and Pulsion Medical System as a lecturer. The other authors have no competing interests.

Received: 14 March 2024 Accepted: 20 March 2024

Published online: 29 March 2024

References

1. De Courson H, Chadeaux G, Loiseau A, Georges D, Biais M. Myocardial dysfunction assessed by speckle-tracking in good-grade subarachnoid hemorrhage patients (WFNS 1–2): a prospective observational study. *Crit Care*. 2023;27(1):455.

2. Park YS, Park J-H, Ahn KT, Jang WI, Park HS, Kim JH, Lee JH, Choi SW, Jeong JO, Seong IW. Usefulness of Mitral annular systolic velocity in the detection of left ventricular systolic dysfunction: comparison with three dimensional echocardiographic data. *J Cardiovasc Ultrasound*. 2010;18(1):1.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.