

Using ultrasound shear wave elastography to characterize peripheral nerve mechanics: a systematic review on the normative reference values in healthy individuals

Tiago Neto, Johanna Johannsson, Ricardo Andrade

▶ To cite this version:

Tiago Neto, Johanna Johannsson, Ricardo Andrade. Using ultrasound shear wave elastography to characterize peripheral nerve mechanics: a systematic review on the normative reference values in healthy individuals. Ultrasonography, 2024, 43 (3), pp.169-178. 10.14366/usg.23211. hal-04577759

HAL Id: hal-04577759 https://hal.science/hal-04577759

Submitted on 28 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



Using ultrasound shear wave elastography to characterize peripheral nerve mechanics: a systematic review on the normative reference values in healthy individuals

Tiago Neto¹, Johanna Johannsson², Ricardo J. Andrade^{3,4,5}

¹Luxembourg Institute of Research in Orthopedics, Sports Medicine and Science, Luxembourg, Luxembourg; ²Department of Physiotherapy, LUNEX International University of Health, Exercise and Sports, Differdange, Luxembourg; ³Nantes University, Movement -Interactions - Performance, MIP, UR 4334, 44000, Nantes, France; ⁴Menzies Health Institute Queensland, Griffith University, Brisbane and Gold Coast, Australia; ⁵School of Health Sciences and Social Work, Griffith University, Brisbane and Gold Coast, Australia

Ultrasound shear wave elastography (SWE) is an emerging non-invasive imaging technique for peripheral nerve evaluation. Shear wave velocity (SWV), a surrogate measure of stiffness, holds promise as a biomarker for various peripheral nerve disorders. However, to maximize its clinical and biomechanical value, it is important to fully understand the factors that influence nerve SWV measurements. This systematic review aimed to identify the normal range of SWV for healthy sciatic and tibial nerves and to reveal the factors potentially affecting nerve SWV. An electronic search yielded 17 studies eligible for inclusion, involving 548 healthy individuals (age range, 17 to 72 years). Despite very good reliability metrics, the reported SWV values differed considerably across studies for the sciatic (1.9-9.9 m/s) and tibial (2.3-9.1 m/s) nerves. Factors such as measurement proximity to joint regions, limb postures inducing nerve axial stretching, and transducer alignment with nerve fiber orientation were associated with increased SWV. These findings suggest regional-specific nerve mechanical properties, non-linear elastic behaviour, and marked mechanical anisotropy. The impact of age and sex remains unclear and warrants further investigation. These results emphasize the importance of considering these factors when assessing and interpreting nerve SWE. While increased SWV has been linked to pathological changes affecting nerve tissue mechanics, the significant variability observed in healthy nerves highlights the need for standardized SWE assessment protocols. Developing guidelines for enhanced clinical utility and achieving a comprehensive understanding of the factors that influence nerve SWE assessments are critical in advancing the field.

Keywords: Ultrasound shear wave elastography; Ultrasonography; Sciatic neuropathy;

Peripheral nervous system; Nerve biomechanics; Non-invasive mechanics;

Diagnostic imaging

Key points: Nerve shear wave velocity, as measured by ultrasound shear wave elastography, is a surrogate measure of nerve stiffness and holds promise as a non-invasive biomarker of peripheral nerve disorders. The findings highlight that, despite very good reliability, nerve shear wave velocity is highly influenced by factors such as nerve region, limb postures, and transducer orientation. Standardized protocols for shear wave elastography assessment are crucial to enhance clinical utility and facilitate comparisons across studies, ultimately advancing the diagnosis and monitoring of nerve disorders.



REVIEW ARTICLE

https://doi.org/10.14366/usg.23211 eISSN: 2288-5943 Ultrasonography 2024;43:169-178

Received: November 14, 2023 Revised: February 9, 2024 Accepted: February 14, 2024

Correspondence to:

Ricardo J. Andrade, MSc, PhD, University of Nantes, Movement -Interactions - Performance (MIP), 23 Rue Recteur Schmitt Bat F0, BP 92235 44322, Nantes, France

Tel. +33251837241 Fax. +33251837045 E-mail: ricardo.andrade@univ-nantes.fr, r.andrade@griffith.edu.au

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (http://creativecommons.org/ licenses/by-nc/4.0/) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2024 Korean Society of Ultrasound in Medicine (KSUM)



How to cite this article:

Neto T, Johannsson J, Andrade RJ. Using ultrasound shear wave elastography to characterize peripheral nerve mechanics: a systematic review on the normative reference values in healthy individuals. Ultrasonography. 2024 May;43(3):169-178.

Introduction

Peripheral neuropathies are among the most common neurological disorders [1]. Nerve conduction studies, the current gold standard for detecting and monitoring peripheral nerve disorders, provide objective insights into the functioning of large $A\beta$ nerve fibers [2]. However, they have several limitations. For instance, they are less sensitive at detecting pathological changes in small sensory $A\delta$ or C nerve fibers [3], and they are typically limited to distal nerves, which are often severely damaged and unexcitable in length-dependent neuropathies [4]. A multimodal testing protocol, together with subjective patient-reported outcome measures, has therefore been suggested as the optimal peripheral nerve assessment strategy for several peripheral nerve disorders [5]. In this context, non-invasive imaging techniques, such as nerve high-resolution ultrasonography and ultrasound shear wave elastography (SWE), are emerging adjunct assessment modalities to both electrophysiological and clinical evaluations [6,7].

High-resolution ultrasonography is a rapidly evolving field, offering valuable clinical insights not only into nerve size and morphology [8], longitudinal excursion [9,10] and echogenicity [11] but also into the localization of nerve damage and associated pathological processes [12,13]. However, B-mode ultrasonography does not have the sensitivity required to detect all forms of peripheral nerve disorders and fails to provide information about nerve mechanical properties such as stiffness [7]. Indeed, abnormalities in nerve mechanical properties have been proposed as potential biomarkers for various peripheral nerve disorders [14,15]. For example, animal models of both focal neuropathies and polyneuropathies, including small and large nerve fiber neuropathies, have shown greater peripheral nerve stiffness regardless of the etiology [16-18]. Notably, SWE provides guantitative, real-time, and high spatial resolution elasticity mapping of biological soft tissues by measuring the propagation velocity of shear waves generated remotely using focused ultrasonic beams [19,20]. In peripheral nerves, the shear wave velocity has been shown to be highly sensitive to actual elasticity changes [21], making SWE a promising non-invasive method for evaluating changes in peripheral nerve mechanical properties associated with altered tissue structure and composition [6].

The diagnostic utility of ultrasound SWE has recently been demonstrated. For example, preliminary studies in individuals with carpal [22,23] and ulnar [24] tunnel syndromes, and distal symmetric diabetic polyneuropathy [25,26] have shown an increase in nerve shear wave velocity with the progression and severity of the neuropathy. Studies using ultrasound SWE in lower limb nerves have considerably increased in recent years, either to evaluate clinical conditions such as lumbar radicular pain [27,28] and

diabetic polyneuropathy [29,30], or to elucidate nerve biomechanics in healthy individuals [31,32]. Despite its recent popularity and reported validity, variables such as limb positioning, transducer orientation, or the region where the nerve elasticity is assessed, have been suggested to significantly influence the shear wave velocity measures of peripheral nerves [6]. However, it is not fully understood how these, and potentially other, variables may indeed affect the shear wave velocity of the most examined peripheral nerves: the sciatic and tibial nerves. A deeper understanding and mapping of the factors that may influence shear wave propagation within peripheral nerves would be essential for establishing and standardizing SWE assessment protocols, which can be useful for an early diagnosis and the reliable monitoring of nerve-related disorders. The objectives of this systematic review were to summarize the normative shear wave velocity values for the sciatic and tibial nerves in healthy individuals, and to describe the factors that may influence such measurements.

Materials and Methods

An electronic search for experimental or observational studies, written in any language (with at least an English title and abstract), was conducted on PubMed, Embase, and Web of Science. Due to resource constraints for translating articles in non-English languages and a commitment to ensuring accurate analysis, non-English studies were not included in this research. The following keywords, and respective combinations, were used during the search: peripheral nerve, sciatic or tibial nerve, SWE, nerve stiffness, and shear wave velocity. The electronic search was further complemented by reviewing the references of previous reviews and of the included studies. To be included, studies had to use ultrasound SWE to assess the shear wave velocity or shear modulus or Young's modulus of the sciatic and/or tibial nerves in healthy individuals of any age or sex. Studies in clinical populations reporting shear wave velocity values for healthy controls were also included. The exclusion criteria were as follows: reviews, studies in animals, studies in human cadavers, and studies in clinical populations without healthy control groups.

After selecting the studies, the following data were extracted and summarized in Table 1 [25,26,29,30,32–44]: demographic information of the participants, ultrasound scanner manufacturer, participant's posture during ultrasound scans, scanned nerve region, transducer orientation used for nerve scans, reliability outcomes, and the shear wave velocity values. To facilitate comparisons, peripheral nerve stiffness values reported in shear modulus (kPa) were transformed into shear wave velocity (m/s), using the following equation: $\mu = \rho V_s^2$, where ρ represents the density of soft tissues (approximately 1,000 kg/m³), and V_s represent shear wave speed [20]. Soft tissue density is generally modelled as a constant in

Table 1. Summary of the included studies

Study	Participants	Nerve	SWE equipment	SWV measurement	ICC (95% confidence interval)	Results for nerve SWV
Andrade et al. (2016) [32]	10 Male participants (25 years)	SN	Aixplorer, version 6.1 Probe: 10–2 MHz	Prone SN measured at the subgluteal region, with the knee flexed at 90° or fully extended, and the ankle in 40° of plantarflexion	Knee 90°: 0.98 (0.93–1.00) Knee 0°: 0.97 (0.90–0.99)	Knee at 90°: 5.07±1.2 m/s (25.7 kPa) Knee at 0°: 6.8±0.9 m/s (46.2 kPa) ^{a)}
Andrade et al. (2018) [33]	15 Participants (13 M, 2F), 22 years	SN	Aixplorer, version 6.1 Probe: 10–2 MHz	Prone SN measured at the subgluteal region, with full knee extension, during ankle passive motion: 0% of dorsiflexion (i.e., 40° plantarflexion) to 100% (i.e., 20° dorsiflexion)		0% Dorsiflexion: 5.43±1.3 m/s (29.5 kPa) 100% Dorsiflexion: 6.73±1.6 m/s (45.3 kPa) ^{b)}
Andrade et al. (2022) [34]	60 Participants (31 M, 29 F), 21 years	SN and TN	Aixplorer, version 6.1 Probe: 10–2 MHz	SWV measured in 2 hip positions, neutral (prone) and flexed at 90° (long sitting position). SN measured at the subgluteal region (i.e., proximal) and at the nerve bifurcation (i.e., distal). TN measured at the popliteal fossa (i.e., proximal), at 50% of the distance between the lateral femoral condyle and tibial malleolus (i.e., intermediate), and at the tibial malleolus (i.e., distal).		Hip neutral SN_p: 4.64±1.2 m/s (21.6 kPa) SN_d: 5.67±1.2 m/s (32.2 kPa) TN_p: 6.50±1.5 m/s (42.3 kPa) TN_int: 5.8±1.1 m/s (33.6 kPa) TN_d: 6.20±1.0 m/s (38.4 kPa) Hip flexed SN_p: 7.06±1.2 m/s (49.8 kPa) ^{c1} SN_d: 8.47±1.6 m/s (71.8 kPa) ^{c2} TN_p: 9.42±2.0 m/s (88.8 kPa) ^{c2} TN_int: 8.29±1.7 m/s (68.7 kPa) ^{c2} TN_d: 9.13±1.5 m/s (83.3 kPa) ^{c2}
Anegawa et al. (2023) [35]	21 Participants (10 M, 11 F), 21 years	TN	Canon Aplio300 Probe: 10 MHz	Participants' trunk and neck in the intermediate position, hip at 90° flexion and the knee at 30° flexion. TN measured above the medial malleolus at 25% (i.e., -16°) and 75% (i.e., 8°) of ankle passive dorsiflexion.	At 25%: 0.96 (0.77–0.99) At 75%: 0.95 (0.67–0.99)	25% Dorsiflexion: 5.5±1.3 m/s (30.3 kPa) 75% Dorsiflexion: 7.4±0.7 m/s (54.8 kPa) ^{b)}
Aslan et al. (2019) [25]	32 Participants (20 F, 12 M), 17 years	TN	Mindray Resona 7 Probe: L14–5WU ComboWave linear	Supine, ankle in slight plantarflexion TN measured above the medial malleolus	0.99 (0.99–1.00)	Transverse axis: 2.37 m/s Longitudinal axis: 4.17±0.9 m/s (17.4 kPa) ^{d)}
Bedewei et al. (2021) [36]	36 Participants (21 F, 25 M), 33 years	TN	Aixplorer, Mach 30 Probe: 18—5 MHz	Prone TN measured at the popliteal fossa	0.78	Transverse axis: 23.3±7.2 kPa (4.83 m/s) Longitudinal axis: 26.1±6.4 kPa (5.11 m/s)
Chen et al. (2020) [26]	19 Female participants (52 years)	TN	Aixplorer Probe: 4–15 MHz	Supine, ankle in slight plantarflexion TN measured above the medial malleolus		L: 21.3±5.0 kPa (4.65 m/s) R: 20.3±5.2 kPa (4.51 m/s)
Dikici et al. (2017) [29]	20 Participants (11 F, 9 M), 58 years	TN	Aixplorer Probe: 4–15 MHz	Supine, ankle in slight plantarflexion TN measured above the medial malleolus	0.96 (0.93–0.98)	L: 31.3±11.6 kPa (5.59 m/s) R: 30.3±7.6 kPa (5.50 m/s)
Greening and Dilley (2017) [37]	26 Participants: 11 M (38 years) and 15 F (39 years)	TN	Siemens, ACUSON S2000 Probe: 4–9 MHz	Supine TN measured above the medial malleolus, with knee flexed at 90° and fully extended (0°)	Knee 90°: 0.37 (0.10–0.64) Knee 0°: 0.53 (0.29–0.74)	Knee 90°: 3.25±0.1 m/s (10.6 kPa) Knee 0°: 5.16±0.2 m/s (26.6 kPa) ^{a)}
He et al. (2019) [38]	40 Participants (16 F, 24 M), 55 years	TN	Aixplorer Probe: 4–15 MHz	Supine, ankle in slight plantarflexion TN measured above the medial malleolus		L: 3.67±0.5 m/s (13.5 kPa) R: 3.64±0.5 m/s (13.2 kPa)

Continued

Table 1. Continued

Study	Participants	Nerve	SWE equipment	SWV measurement	ICC (95% confidence	Results for nerve SWV
					interval)	
Hirata et al. (2020) [39]	40 Male participants: 20 young adults (22 years) 20 older adults (72 years)	SN	Siemens, ACUSON S2000 Probe: 4–9 MHz	Prone SN measured at the mid-thigh region, with full knee extension, during ankle passive motion (i.e., 30° plantarflexion to 15° dorsiflexion)	Young adults: 0.98 (0.94–0.99) Older adults: 0.99 (0.98–1.00)	30° Plantarflexion Young: 2.66±0.6 m/s (9.6 kPa) ^{e)} Older: 1.92±0.3 m/s (4.4 kPa) 15° Dorsiflexion Young: 3.77±1.7 m/s (14.2 kPa) ^{b),e)} Older: 2.47±0.4 m/s (6.1 kPa) ^{b)}
Jiang et al. (2019) [30]	20 Participants (10 F, 10 M), 58 years	TN	Aixplorer Probe: 4-15 MHz	Supine, ankle in slight plantarflexion TN measured above the medial malleolus	0.95 (0.93–0.97)	Average of both limbs: 32.8±7.1 kPa (5.72 m/s)
Kawanishi et al. (2022) [40]	20 Participants (14 M, 6 F), 24 years	TN	Canon Aplio300 Probe: 10 MHz	Participants' position unknown. TN measured above the medial malleolus during ankle passive motion: 0% of dorsiflexion (i.e., 18° plantarflexion) to 100% (i.e., 17° dorsiflexion)	>0.85 (0.85-0.99)	0% Dorsiflexion: 4.5±1.7 m/s (20.3 kPa) 100% Dorsiflexion: 7.5±0.7 m/s (56.3 kPa) ^{b)}
Neto et al. (2017) [41]	14 Participants (11 M, 3 F), 30 years	SN	Aixplorer, version 10.0 Probe: 10–2 MHz	Prone SN measured at the subgluteal region, with full knee extension, during ankle passive motion: 0% of dorsiflexion (i.e., 40° plantarflexion) to 100% (i.e., 20° dorsiflexion)	0.93 (0.79–0.98)	0% Dorsiflexion: L: 7.77±1.5 m/s (60.4 kPa) R: 7.91±1.4 m/s (62.6 kPa) 100% Dorsiflexion: L: 9.65±1.8 m/s (93.0 kPa) R: 9.88±2.0 m/s (97.6 kPa)
Shang et al. (2023) [42]	50 Participants (23 M, 27 F), 40 years	TN	Mindray Resona 8 Probe: 4–15 MHz linear	Supine, ankle in slight plantarflexion TN measured above the medial malleolus	Interobserver 0.70 (0.58–0.79) Intraobserver 0.75 (0.65–0.82)	L: 6.06±2.4 m/s (36.7 kPa) R: 6.08±2.3 m/s (37.0 kPa) Males: 6.04±2.4 m/s (36.4 kPa) Females: 6.10±2.3 m/s (37.2 kPa) <40 years: 6.02±2.3 m/s (36.2 kPa) >40 years: 6.15±2.4 m/s (37.8 kPa)
Tang et al. (2021) [43]	105 Participants (49 M, 56 F), 49 years	TN	Aixplorer, version 12.0 Probe: 4–15 MHz	Supine, ankle in slight plantarflexion TN measured above the medial malleolus		All participants: 3.87–0.8 m/s (15.0 kPa) Males: 4.00–0.65 m/s (16.0 kPa) Females: 3.72–0.78 m/s (13.8 kPa) (values presented as median with IQR)
Wei and Ye (2020) [44]	20 Participants (12 F, 8 M), 57 years	TN	Siemens, ACUSON S2000 Probe: 4–9 MHz	Prone, ankle in neutral position TN measured at the popliteal fossa	0.86 (0.80–0.91)	L: 2.27±0.2 m/s (5.2 kPa) R: 2.38±0.2 m/s (5.7 kPa)

SWE, shear wave elastography; SWV, shear wave velocity; ICC, intraclass correlation coefficient; SN, sciatic nerve; SN_p, sciatic nerve proximal; SN_d, sciatic nerve distal; TN, tibial nerve; TN_p, tibial nerve proximal; TN_int, tibial nerve intermediate; TN_d, tibial nerve distal; F, female; M, male; L, left; R, right; IQR, interquartile range. ^{a)}Significantly higher than knee at 90°. ^{b)}Significantly higher than ankle plantarflexion. ^{c)}Significantly higher than hip neutral. ^{d)}Significantly higher than transverse axis. ^{e)}Significantly higher than older adult.

ultrasound machines, which simplifies the conversion of shear modulus to shear wave velocity. The risk of bias was assessed independently by two authors using the Joanna Briggs Institute Critical Appraisal Checklist for Analytical Cross-Sectional Studies [45].

The search strategy and its results are summarized in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram [46] (Fig. 1). The protocol of this review was registered in the International Prospective Register of Systematic Reviews (PROSPERO) with the number CRD42022302947.

Results

Characteristics of the Studies

Seventeen studies were included in the review and are summarized in Table 1. The tibial nerve was assessed in 12 studies, while the sciatic nerve was measured in four studies; one study assessed both nerves.

Tibial nerve shear wave velocity measurements were predominantly conducted with participants in a supine position, with the knee fully



Fig. 1. PRISMA flow diagram showing the results of the search strategy. SWE, shear wave elastography.

extended and ankle in plantarflexion (resting angle); however, an alternative prone position was utilized in three other studies.

For the sciatic nerve, all studies employed a prone position for participants, characterized by full knee extension and a resting ankle angle during shear wave velocity scans.

In most studies, the tibial nerve was assessed above the medial malleolus. In two studies [36,44], the tibial nerve was assessed in the popliteal fossa, and in another study [34], it was measured in three different locations from the popliteal fossa down to the medial malleolus. For the sciatic nerve, three studies [32,33,41] measured nerve shear wave velocity in the subgluteal region, one study measured it in the mid-thigh region [39], and another study [34] measured the sciatic shear wave velocity in two different regions: proximal (i.e., subgluteal) and distal (i.e., proximal to the bifurcation of common sciatic into tibial and fibular nerves).

Risk of Bias Assessment

The included studies showed an overall low risk of bias as the majority of the eight items of the Joanna Briggs Institute appraisal checklist for analytical cross-sectional studies were met. For the studies using a single group of healthy individuals, only items #3 and #4 were not met (i.e., without "exposure" or "condition" to be assessed).

Reliability of the Shear Wave Velocity Measurements

Thirteen studies reported reliability scores for nerve SWE assessments (Table 1). Overall, the use of SWE to measure sciatic and tibial shear wave velocity showed very good reliability, with intraclass correlation coefficient (ICC [2,1]) values ranging from 0.70 to 0.99. The exception is the study of Greening and Dilley [37],

which reported only moderate values of reliability (ICC=0.37-0.53). However, the authors suggested that these values may be attributed to methodological disparities, such as variations in the scanner used and the sequence of limb positioning, when compared to other studies of a similar nature.

Shear Wave Velocity of the Sciatic and Tibial Nerves

Table 1 shows the results for nerve shear wave velocities. The reported values show high variability for both the tibial (range, 2.3 to 9.1 m/s) and the sciatic nerves (range, 1.9 to 9.9 m/s). Most of the studies included young adults, while three studies [39,42] compared nerve shear wave velocity between younger and older participants. Hirata et al. [39] showed significantly lower sciatic nerve shear wave velocities in older adults than in younger adults. As for the tibial nerve, conflicting results were observed, with one study reporting higher tibial nerve shear wave velocities in older participants than younger participants [42], while another study [37] found a significant, moderate, and negative correlation (r=-0.58) between the tibial nerve shear wave velocity and age.

Posture-induced changes in nerve shear wave velocity were also investigated by using different lower limb postures through various combinations of hip, knee, and/or ankle angles. For example, one study [34] observed significantly higher sciatic and tibial nerve shear wave velocities (i.e., +54%) with hip flexion, compared to a hipneutral position. Postures involving full knee extension also resulted in significantly higher nerve shear wave velocities, compared to having the knee flexed at 90°, for both the sciatic [32] and tibial [37] nerves. Similar findings were reported for the influence of the ankle position in the shear wave velocities of the sciatic nerve, with higher shear wave velocities observed with the ankle positioned at higher

dorsiflexion angles compared to ankle plantarflexion [33,35,39-41].

Two studies [25,36] assessed the effect of transducer orientation in the shear wave velocity measures. Aslan et al. [25] reported significantly higher shear wave velocities (i.e., with a difference up to 1.8 m/s) obtained with longitudinal transducer orientation (parallel to the nerve fibers) compared to transverse orientation. One study investigated the spatial variation of shear wave velocity along sciatic and tibial nerves [34]. It was observed that distal nerve regions tend to show greater shear wave velocities. For example, an average difference of 1.41 m/s between proximal and distal sciatic has been reported. No studies reported significant left versus right limb differences in sciatic or tibial nerve shear wave velocities.

Discussion

In recent years, there has been a growing interest in the application of SWE to assess peripheral nerve mechanics in both clinical [6] and biomechanical [47] contexts. Shear wave velocity is not only a surrogate measure of nerve stiffness [21], but also a good indicator of tensile loading (i.e., nerve axial stretching) [32] and a promising biomarker for peripheral nerve disorders [6]. The popularity of SWE, especially in contexts of peripheral nerve evaluation, can be further attributed to its non-invasive and painless nature, and its capacity to estimate the mechanical properties of nerves in real time [20,48]. The observed high reliability, as evidenced by the ICC values reported in most of the studies, particularly in the context of intraexaminer measurements, highlights the clinical applicability of SWE. Nonetheless, the findings also point to substantial variability in the normal shear wave velocity values for both sciatic and tibial nerves across studies, which can be attributed to several confounding variables that will be discussed hereunder.

Spatial Variations in Nerve Shear Wave Velocity

The findings of this study indicate that both the sciatic and tibial nerves were assessed at multiple and non-standardized nerve regions across studies. This makes comparisons between studies difficult and may partly explain the large variability in shear wave velocities that has been observed for both peripheral nerves across studies. We identified only five studies [26,29,30,38,43] where a similar assessment protocol was employed, including the same scanner and limb positioning to measure the tibial nerve shear wave velocity in the longitudinal view. The shear wave velocity ranged from 3.9 m/s to 5.7 m/s (mean difference of 1.8 m/s), providing a more realistic notion of the expected variability when using SWE to assess peripheral nerves' stiffness. Importantly, these normal values for healthy tibial nerves are still considerably lower than the cutoff value of 51.05 kPa (approximately 7.1 m/s) for diagnosing diabetic

neuropathy with high sensitivity and specificity [29].

More recently, Andrade et al. [34] found spatial-dependent differences in shear wave velocities in the sciatic and tibial nerves. For instance, shear wave velocity tends to increase in a proximal to distal pattern at both upper and lower leg segments, being greatest in locations near the knee and ankle joints. These observations could be due to regional variations in sciatic and tibial nerves' ultrastructure, such as thinner and more numerous nerve fascicles wrapped by a thicker perineurium near joints [49], and an increased stiff epineural tissue content in joint regions [50]. Proximity to joint regions has also been associated with distinct nerve material properties in animal models [51]. Moreover, no significant correlations have been reported between nerve shear wave velocity and nerve thickness for both sciatic and tibial nerves [34], suggesting that it is still unclear to what extent nerve geometry influences shear wave velocity measures. However, it should be noted that for thin and stiff tissues such as tendons, the propagation of the shear waves can be guided [52]. Specifically, the shear waves can experience successive reflections on the nerve boundaries, and this phenomenon has the potential to affect the relationship between shear modulus and shear wave velocity [52]. This phenomenon cannot be totally excluded when assessing relatively thin peripheral nerves, and it can bias the quantification of nerve shear wave velocity, especially in neurological disorders known to significantly alter nerve cross-sectional area.

Influence of Limb Posture on Nerve Shear Wave Velocity

Limb posture is an important factor to consider when evaluating nerve shear wave velocity, as it can directly influence the nerve length and, thus, the tensile loading (i.e., the amount of axial stretching). Shear wave velocity is a good indicator of passive forces acting on biological soft tissues, such as muscles [53] and nerves [21]. The findings show that combining knee full extension with hip flexion [34,37] and knee full extension with ankle dorsiflexion [32,39,41] results in a considerable increase in sciatic and tibial nerve shear wave velocity. These observations are consistent with human cadaveric studies reporting an increase in nerve strain in similar lower limb postures [54]. They can be explained by the elastic nature of peripheral nerves, including their inherent layers of connective tissue, when subjected to tensile loads [14]. The elongation of peripheral nerves also results in increased endoneural pressure, which can further contribute to increased nerve stiffness [55]. To minimize variability in shear wave velocity measurements associated with nerve tensile loading during clinical evaluations, it is advisable to opt for scanning postures that mechanically unload the sciatic and tibial nerves, such as maintaining a neutral hip position, flexing the knee, and keeping a resting plantar flexion angle. This would make it possible to isolate true changes in nerve stiffness from potential nerve tensioning caused by postures that subject nerves to stretching, thereby increasing nerve wave propagation velocity.

Mechanical Anisotropy

Assessing peripheral nerves using SWE can be challenging due to their complex structural, mechanical, and geometric characteristics. Peripheral nerves, like other soft tissues in the body, are characterized by their heterogeneous and anisotropic nature, displaying non-linear and viscoelastic properties that lead to stiffening under substantial strains and exhibiting stress relaxation behaviour when axially stretched [14]. In addition to the elastic, heterogeneous, and thin characteristics discussed in earlier paragraphs, it is important to highlight that peripheral nerves exhibit a complex structural organization, which leads to strong anisotropic mechanics. Indeed, the effect of transducer orientation (i.e., along or across the nerve fibers) on shear wave velocity has been well described in anisotropic tissues [56], including peripheral nerves [21]. Measurements of the tibial nerve [25,36] performed with the transducer aligned parallel to the nerve fibers (i.e., longitudinal view) yielded considerably higher shear wave velocities compared to those performed perpendicular to the main direction of the nerve fibers (i.e., transverse scans) [25]. This expected mechanical anisotropy effect should be taken into consideration as a significant factor influencing nerve shear wave velocity outcomes. Ultrasound transducer positioning needs to be well controlled when using SWE to assess peripheral nerve's integrity. It is important to note, however, that mechanical anisotropy may also be an interesting biomarker for peripheral nerve disorders characterized by structural disorganization.

Influence of Sex and Age on Nerve Shear Wave Velocity

There is limited evidence regarding the influence of age and sex on peripheral nerve shear wave velocity, including that of the sciatic and tibial nerves. One study measuring the shear wave velocity of the median and tibial nerves observed that male participants had significantly higher shear wave velocities for the median nerve than female participants, but no differences were observed regarding the tibial nerve [43]. Similar findings were reported in another study [42] that observed no differences in the stiffness of the tibial nerve between male and female participants. Conflicting results have been reported on the effect of ageing on nerve shear wave velocity. One study found higher values of stiffness in the tibial nerve of older participants compared to younger participants. However, the sample in this study was relatively young, with only three participants above 60 years old, versus 47 participants below 60 years, which limits the validity of conclusions about the influence of ageing on nerve stiffness. Other studies observed the opposite tendency, with shear wave velocity decreasing with age, both for the tibial [37] and the sciatic [39] nerves. This may be explained by age-related reductions in the thickness and in the content of collagen fibril of the sciatic nerve's perineurium, as shown in animal studies [57]. Further studies are required to investigate the influence of demographic variables more comprehensively on peripheral nerve shear wave velocity, including their possible interactions with nerve regions and different tensile loads.

Additional Factors Influencing Nerve Shear Wave Velocity

The type of ultrasound scanner and the measurement depth are additional variables that can further impact normal shear wave velocity values and introduce variability when comparing across studies. For example, one study [58] reported significant differences in shear wave velocity obtained from ultrasound equipment from different manufacturers, using the same type of transducer and measuring at similar depths in an elasticity phantom. Another study reported different shear wave velocities from different ultrasound equipment when a soft elasticity phantom was used, but not with a harder phantom [59]. These small variations might be attributed to differences in the methodological procedures or computations used to estimate shear wave velocity among different manufacturers, potentially impacting the establishment of generalized cutoff values for various nerve-related disorders. Moreover, measuring shear wave velocity at depths ranging from 2 to 5 cm has also been shown to produce different results when using the same SWE scanner and transducer [58,60], with a trend for shear wave velocity to decrease in deeper regions [58]. As these results were mostly obtained from elasticity phantoms, it is important to determine whether these observations extend to the measurement of peripheral nerves, to improve the validity of nerve SWE assessments and its diagnostic value. However, the sciatic and tibial nerves are generally assessed at depths inferior to 3.5 cm, which might be covered by the focal area of linear probes typically used for these assessments.

Limitations of the Review

The current review presents some limitations. Other lower limb nerves, such as the fibular, femoral, obturator, and saphenous nerves, were not considered. However, elastography-based studies evaluating these nerves are scarce or even non-existent. Therefore, the primary focus of this review was on the most frequently assessed nerves—the sciatic and the tibial. Another potential limitation is the non-inclusion of clinical populations, where nerve mechanical properties have been shown to be considerably altered. This review exclusively considered healthy individuals, as we believe

that it is crucial to first establish a comprehensive understanding of shear wave velocity in structurally and functionally intact tibial and sciatic nerves. By taking this approach, it is possible to carefully examine the potential factors that may influence these shear wave velocity measurements before thoroughly investigating their clinical applicability in serial and multi-center settings. Finally, the high variability in the methods used to measure nerve stiffness made it impossible to pool the results, leading us to the decision not to perform a meta-analysis.

Conclusion

Ultrasound SWE has been recognized as a valid and accurate non-invasive imaging technique for evaluating the mechanical properties of peripheral nerves, with shear wave velocity serving as a surrogate marker for nerve stiffness. In this systematic review, we compiled a comprehensive range of normal shear wave velocity values for both the sciatic and tibial nerves in healthy adults. This contributes to summarizing normative reference values to which shear wave velocity from diseased nerves can be compared. Importantly, we also identified the main factors directly influencing SWE assessments in peripheral nerves. Better understanding and accounting for spatial variations in nerve shear wave velocity, the intrinsic mechanical anisotropy of nerves, and the implication of limb postures on axial stretching is paramount to enhance the clinical applicability of SWE and compare outcomes across research studies. Standardized assessment protocols are required and need to take into consideration variables such as scanning postures, transducer orientation, and the nerve regions under assessment, as these factors are likely to directly and significantly affect nerve shear wave velocity. The pursuit of such standardization holds substantial promise for the advancement of clinical diagnosis and the monitoring of peripheral neuropathies using SWE. Consensus among expert practitioners (e.g., a Delphi study) is required to guide decision-making in the acquisition, analysis, and interpretation of SWE in peripheral nerves in experimental and clinical settings.

ORCID: Tiago Neto: https://orcid.org/0000-0001-6696-5372; Johanna Johannsson: https://orcid.org/0000-0002-8927-1741; Ricardo J. Andrade: https://orcid.org/0000-0001-5316-7531

Author Contributions

Conceptualization: Neto T, Johannsson J, Andrade RJ. Data acquisition: Neto T, Johannsson J, Andrade RJ. Data analysis or interpretation: Neto T, Johannsson J, Andrade RJ. Drafting of the manuscript: Neto T, Johannsson J, Andrade RJ. Critical revision of the manuscript: Neto T, Johannsson J, Andrade RJ. Approval of the final version of the manuscript: all authors.

Conflict of Interest

No potential conflict of interest relevant to this article was reported.

Acknowledgments

RJA was supported by a Griffith University Postdoctoral Fellowship Scheme award.

References

- Watson JC, Dyck PJ. Peripheral neuropathy: a practical approach to diagnosis and symptom management. Mayo Clin Proc 2015;90:940-951.
- 2. Jones LK Jr. Nerve conduction studies: basic concepts and patterns of abnormalities. Neurol Clin 2012;30:405-427.
- Malik RA, Veves A, Tesfaye S, Smith G, Cameron N, Zochodne D, et al. Small fibre neuropathy: role in the diagnosis of diabetic sensorimotor polyneuropathy. Diabetes Metab Res Rev 2011;27:678-684.
- 4. Latov N. Diagnosis and treatment of chronic acquired demyelinating polyneuropathies. Nat Rev Neurol 2014;10:435-446.
- Argyriou AA, Park SB, Islam B, Tamburin S, Velasco R, Alberti P, et al. Neurophysiological, nerve imaging and other techniques to assess chemotherapy-induced peripheral neurotoxicity in the clinical and research settings. J Neurol Neurosurg Psychiatry 2019;90:1361-1369.
- Wee TC, Simon NG. Ultrasound elastography for the evaluation of peripheral nerves: a systematic review. Muscle Nerve 2019;60:501-512.
- 7. Gasparotti R, Padua L, Briani C, Lauria G. New technologies for the assessment of neuropathies. Nat Rev Neurol 2017;13:203-216.
- 8. Frost LR, Brown SH. Neuromuscular ultrasound imaging in low back pain patients with radiculopathy. Man Ther 2016;21:83-88.
- 9. Hough AD, Moore AP, Jones MP. Reduced longitudinal excursion of the median nerve in carpal tunnel syndrome. Arch Phys Med Rehabil 2007;88:569-576.
- 10. Ridehalgh C, Moore A, Hough A. Sciatic nerve excursion during a modified passive straight leg raise test in asymptomatic participants and participants with spinally referred leg pain. Man Ther 2015;20:564-569.
- 11. Holzapfel K, Ghosh T, Krischak S, Naumann M. Nerve echogenicity: changes in high-resolution nerve ultrasound in carpal tunnel syndrome after surgery. Ultrasound Med Biol 2022;48:223-227.
- 12. Erel E, Dilley A, Greening J, Morris V, Cohen B, Lynn B. Longitudinal sliding of the median nerve in patients with carpal tunnel syndrome. J Hand Surg Br 2003;28:439-443.
- 13. van Doesburg MH, Henderson J, Mink van der Molen AB, An KN, Amadio PC. Transverse plane tendon and median nerve motion in

the carpal tunnel: ultrasound comparison of carpal tunnel syndrome patients and healthy volunteers. PLoS One 2012;7:e37081.

- 14. Topp KS, Boyd BS. Structure and biomechanics of peripheral nerves: nerve responses to physical stresses and implications for physical therapist practice. Phys Ther 2006;86:92-109.
- Yayama T, Kobayashi S, Nakanishi Y, Uchida K, Kokubo Y, Miyazaki T, et al. Effects of graded mechanical compression of rabbit sciatic nerve on nerve blood flow and electrophysiological properties. J Clin Neurosci 2010;17:501-505.
- 16. Zhu Y, Jin Z, Luo Y, Wang Y, Peng N, Peng J, et al. Evaluation of the crushed sciatic nerve and denervated muscle with multimodality ultrasound techniques: an animal study. Ultrasound Med Biol 2020;46:377-392.
- 17. Bober BG, Shah SB. Paclitaxel alters sensory nerve biomechanical properties. J Biomech 2015;48:3559-3567.
- Layton BE, Sastry AM, Wang H, Sullivan KA, Feldman EL, Komorowski TE, et al. Differences between collagen morphologies, properties and distribution in diabetic and normal biobreeding and Sprague-Dawley rat sciatic nerves. J Biomech 2004;37:879-888.
- 19. Bercoff J, Tanter M, Fink M. Supersonic shear imaging: a new technique for soft tissue elasticity mapping. IEEE Trans Ultrason Ferroelectr Freq Control 2004;51:396-409.
- 20. Gennisson JL, Deffieux T, Fink M, Tanter M. Ultrasound elastography: principles and techniques. Diagn Interv Imaging 2013;94:487-495.
- 21. Schrier V, Lin J, Gregory A, Thoreson AR, Alizad A, Amadio PC, et al. Shear wave elastography of the median nerve: a mechanical study. Muscle Nerve 2020;61:826-833.
- 22. Kantarci F, Ustabasioglu FE, Delil S, Olgun DC, Korkmazer B, Dikici AS, et al. Median nerve stiffness measurement by shear wave elastography: a potential sonographic method in the diagnosis of carpal tunnel syndrome. Eur Radiol 2014;24:434-440.
- 23. Cingoz M, Kandemirli SG, Alis DC, Samanci C, Kandemirli GC, Adatepe NU. Evaluation of median nerve by shear wave elastography and diffusion tensor imaging in carpal tunnel syndrome. Eur J Radiol 2018;101:59-64.
- 24. Paluch L, Noszczyk B, Nitek Z, Walecki J, Osiak K, Pietruski P. Shearwave elastography: a new potential method to diagnose ulnar neuropathy at the elbow. Eur Radiol 2018;28:4932-4939.
- Aslan M, Aslan A, Emeksiz HC, Candan F, Erdemli S, Tombul T, et al. Assessment of peripheral nerves with shear wave elastography in type 1 diabetic adolescents without diabetic peripheral neuropathy. J Ultrasound Med 2019;38:1583-1596.
- 26. Chen R, Wang XL, Xue WL, Sun JW, Dong XY, Jiang ZP, et al. Application value of conventional ultrasound and real-time shear wave elastography in patients with type 2 diabetic polyneuropathy. Eur J Radiol 2020;126:108965.
- 27. Neto T, Freitas SR, Andrade RJ, Vaz JR, Mendes B, Firmino T, et al. Noninvasive measurement of sciatic nerve stiffness in patients with chronic low back related leg pain using shear wave elastography. J

Ultrasound Med 2019;38:157-164.

- Celebi UO, Burulday V, Ozveren MF, Dogan A, Akgul MH. Sonoelastographic evaluation of the sciatic nerve in patients with unilateral lumbar disc herniation. Skeletal Radiol 2019;48:129-136.
- 29. Dikici AS, Ustabasioglu FE, Delil S, Nalbantoglu M, Korkmaz B, Bakan S, et al. Evaluation of the tibial nerve with shear-wave elastography: a potential sonographic method for the diagnosis of diabetic peripheral neuropathy. Radiology 2017;282:494-501.
- 30. Jiang W, Huang S, Teng H, Wang P, Wu M, Zhou X, et al. Diagnostic performance of two-dimensional shear wave elastography for evaluating tibial nerve stiffness in patients with diabetic peripheral neuropathy. Eur Radiol 2019;29:2167-2174.
- 31. Zhu B, Yan F, He Y, Wang L, Xiang X, Tang Y, et al. Evaluation of the healthy median nerve elasticity: Feasibility and reliability of shear wave elastography. Medicine (Baltimore) 2018;97:e12956.
- 32. Andrade RJ, Nordez A, Hug F, Ates F, Coppieters MW, Pezarat-Correia P, et al. Non-invasive assessment of sciatic nerve stiffness during human ankle motion using ultrasound shear wave elastography. J Biomech 2016;49:326-331.
- 33. Andrade RJ, Freitas SR, Hug F, Le Sant G, Lacourpaille L, Gross R, et al. The potential role of sciatic nerve stiffness in the limitation of maximal ankle range of motion. Sci Rep 2018;8:14532.
- 34. Andrade RJ, Freitas SR, Hug F, Coppieters MW, Sierra-Silvestre E, Nordez A. Spatial variation in mechanical properties along the sciatic and tibial nerves: an ultrasound shear wave elastography study. J Biomech 2022;136:111075.
- Anegawa K, Kawanishi K, Nakamura M, Izumi M, Tsutsumi M, Kudo S. Tibial nerve dynamics during ankle dorsiflexion: the relationship between stiffness and excursion of the tibial nerve. J Biomech 2023;155:111646.
- 36. Bedewi MA, Elsifey AA, Alfaifi T, Kotb MA, Abdelgawad MS, Bediwy AM, et al. Shear wave elastography of the tibial nerve in healthy subjects. Medicine (Baltimore) 2021;100:e23999.
- 37. Greening J, Dilley A. Posture-induced changes in peripheral nerve stiffness measured by ultrasound shear-wave elastography. Muscle Nerve 2017;55:213-222.
- He Y, Xiang X, Zhu BH, Qiu L. Shear wave elastography evaluation of the median and tibial nerve in diabetic peripheral neuropathy. Quant Imaging Med Surg 2019;9:273-282.
- 39. Hirata K, Yamadera R, Akagi R. Associations between range of motion and tissue stiffness in young and older people. Med Sci Sports Exerc 2020;52:2179-2188.
- 40. Kawanishi K, Nariyama Y, Anegawa K, Tsutsumi M, Kudo S. Changes in tibial nerve stiffness during ankle dorsiflexion according to invivo analysis with shear wave elastography. Medicine (Baltimore) 2022;101:e29840.
- 41. Neto T, Freitas SR, Andrade RJ, Gomes J, Vaz J, Mendes B, et al. Sciatic nerve stiffness is not changed immediately after a slump neurodynamics technique. Muscles Ligaments Tendons J

2017;7:583-589.

- 42. Shang S, Yan W, Guo Y, Guo H, Chen R, Cong S, et al. The normal value and influencing factors of shear wave elastography in healthy tibial nerves: a cross-sectional study. Health Sci Rep 2023;6:e1412.
- 43. Tang X, Zhu B, Tian M, Guo R, Huang S, Tang Y, et al. Preliminary study on the influencing factors of shear wave elastography for peripheral nerves in healthy population. Sci Rep 2021;11:5582.
- 44. Wei M, Ye X. Feasibility of point shear wave elastography for evaluating diabetic peripheral neuropathy. J Ultrasound Med 2020;39:1135-1141.
- 45. Moola S, Munn Z, Tufanaru C, Aromataris E, Sears K, Currie M, et al. Checklist for analytical cross sectional studies: critical appraisal checklist for analytical cross sectional studies 2. Adelaide: Joanna Briggs Institute, 2017.
- 46. Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. PLoS Med 2009;6:e1000097.
- 47. Ellis R, Carta G, Andrade RJ, Coppieters MW. Neurodynamics: is tension contentious? J Man Manip Ther 2022;30:3-12.
- 48. Rossetto G, Lopomo NF, Shaikh SZ. Longitudinal movements and stiffness of lower extremity nerves measured by ultrasonography and ultrasound elastography in symptomatic and asymptomatic populations: a systematic review with meta-analysis. Ultrasound Med Biol 2023;49:1913-1929.
- 49. Hadzic A. Hadzic's textbook of regional anesthesia and acute pain management. 2nd ed. New York: McGraw Hill, 2017.
- 50. Sunderland S. The connective tissues of peripheral nerves. Brain 1965;88:841-854.
- 51. Phillips JB, Smit X, De Zoysa N, Afoke A, Brown RA. Peripheral nerves in the rat exhibit localized heterogeneity of tensile properties during limb movement. J Physiol 2004;557:879-887.

- 52. Brum J, Bernal M, Gennisson JL, Tanter M. In vivo evaluation of the elastic anisotropy of the human Achilles tendon using shear wave dispersion analysis. Phys Med Biol 2014;59:505-523.
- Hug F, Tucker K, Gennisson JL, Tanter M, Nordez A. Elastography for muscle biomechanics: toward the estimation of individual muscle force. Exerc Sport Sci Rev 2015;43:125-133.
- 54. Coppieters MW, Alshami AM, Babri AS, Souvlis T, Kippers V, Hodges PW. Strain and excursion of the sciatic, tibial, and plantar nerves during a modified straight leg raising test. J Orthop Res 2006;24:1883-1889.
- 55. Millesi H, Zoch G, Reihsner R. Mechanical properties of peripheral nerves. Clin Orthop Relat Res 1995;(314):76-83.
- Gennisson JL, Deffieux T, Mace E, Montaldo G, Fink M, Tanter M. Viscoelastic and anisotropic mechanical properties of in vivo muscle tissue assessed by supersonic shear imaging. Ultrasound Med Biol 2010;36:789-801.
- 57. Esquisatto MA, de Aro AA, Feo HB, Gomes L. Changes in the connective tissue sheath of Wistar rat nerve with aging. Ann Anat 2014;196:441-448.
- 58. Shin HJ, Kim MJ, Kim HY, Roh YH, Lee MJ. Comparison of shear wave velocities on ultrasound elastography between different machines, transducers, and acquisition depths: a phantom study. Eur Radiol 2016;26:3361-3367.
- 59. Dillman JR, Chen S, Davenport MS, Zhao H, Urban MW, Song P, et al. Superficial ultrasound shear wave speed measurements in soft and hard elasticity phantoms: repeatability and reproducibility using two ultrasound systems. Pediatr Radiol 2015;45:376-385.
- 60. Chang S, Kim MJ, Kim J, Lee MJ. Variability of shear wave velocity using different frequencies in acoustic radiation force impulse (ARFI) elastography: a phantom and normal liver study. Ultraschall Med 2013;34:260-265.