



HAL
open science

Ulrich ranks of Veronese varieties and equivariant instantons

Daniele Faenzi, Victor Do Valle Pretti

► **To cite this version:**

Daniele Faenzi, Victor Do Valle Pretti. Ulrich ranks of Veronese varieties and equivariant instantons. 2024. hal-04580438

HAL Id: hal-04580438

<https://hal.science/hal-04580438>

Preprint submitted on 20 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ULRICH RANKS OF VERONESE VARIETIES AND EQUIVARIANT INSTANTONS

DANIELE FAENZI AND VICTOR PRETTI

ABSTRACT. We construct Ulrich bundles on Veronese threefolds of arbitrary degree as generic deformations of symmetric squares of equivariant instanton bundles on the projective space, thus classifying the rank of Ulrich bundles on such varieties and proving a conjecture of Costa and Miró-Roig.

1. INTRODUCTION

Given an n -dimensional closed subscheme $X \subset \mathbb{P}^N$ polarized by a very ample divisor class D , an Ulrich sheaf of X is a non-zero coherent sheaf \mathcal{U} such that $H^*(\mathcal{U}(-tD)) = 0$ for $1 \leq t \leq n$. Ulrich sheaves attracted a lot of attention in view of their multiple links to other topics such as Boij-Söderberg theory, Chow forms, matrix factorisations and so on. As general reference on Ulrich bundles we use [CMRPL21]. We define the set of Ulrich ranks of X as

$$\mathrm{Ur}(X) = \{r \in \mathbb{N} \mid \text{there exists an Ulrich sheaf } \mathcal{U} \text{ on } X \text{ with } \mathrm{rk}(\mathcal{U}) = r\}.$$

In general, it is very difficult to determine $\mathrm{Ur}(X)$, except when $n = 1$ where one easily finds $\mathrm{Ur}(X) = \mathbb{N}^*$. A fundamental conjecture of [ES03] states that $\mathrm{Ur}(X)$ should be always non empty, while the main point of [BES17] is to compute the lower bound of $\mathrm{Ur}(X)$, called the Ulrich complexity of X . For a few surfaces, we understand $\mathrm{Ur}(X)$ completely, for instance we know $\mathrm{Ur}(X) = 2\mathbb{N}^*$ for K3 surfaces of Picard number one, see [Fae19, AFO17]; for the d -th Veronese surface, we know $\mathrm{Ur}(X) = 2\mathbb{N}^*$ if d is even and $\mathrm{Ur}(X) = \mathbb{N}^* \setminus \{1\}$ if d is odd, see [CMR18]. For $n \geq 3$ and for an arbitrary polarization, typically we have no clue of what $\mathrm{Ur}(X)$ should be, even though some results are available for minimal polarizations, e.g. [CFK23].

In this note, we focus on Veronese varieties of dimension $n \geq 3$ and mostly on $n = 3$, where the determination of $\mathrm{Ur}(X)$ is open. Our main result gives a complete answer for Veronese threefolds of arbitrary degree, answering a conjecture of Costa and Miró-Roig and giving the first example of varieties of dimension $n > 2$ where $\mathrm{Ur}(X)$ is completely understood, relatively to an arbitrary polarization. Actually $\mathrm{Ur}(X)$ is easy to determine also for d -th Veronese n -folds when $n!$ divides d , as Lemma 4 below shows, but when d modulo $n!$ is arbitrary this gets much harder. Here we look at $n = 3$, the most interesting cases being when d is congruent to 1, 3 or 5 modulo 6.

Let us formulate the precise result. Given integers $d, n \geq 1$ we set $N_d^n = \binom{n+d}{n} - 1$ and consider the d -th Veronese n -fold X_d^n in $\mathbb{P}^{N_d^n}$. We omit n from the notation when $n = 3$.

Theorem 1. *Let $d \geq 2$ and let X_d be the d -th Veronese threefold in \mathbb{P}^{N_d} . Set $\bar{d} \in \{0, \dots, 5\}$ for the remainder of the division of d by 6. We have:*

- i) if $\bar{d} = 0$, then $\mathrm{Ur}(X_d) = 6\mathbb{N}^*$;*
- ii) if $\bar{d} \in \{1, 5\}$, then $\mathrm{Ur}(X_d) = \mathbb{N}^* \setminus \{1\}$;*
- iii) if $\bar{d} \in \{2, 4\}$, then $\mathrm{Ur}(X_d) = 2\mathbb{N}^*$;*
- iv) if $\bar{d} = 3$, then $\mathrm{Ur}(X_d) = 3\mathbb{N}^*$.*

This proves [CMR21, Conjecture 1.1], where of course $r = 1$ has to be excluded from the statement. Let us say a word on the proof. First, non-existence outside the range mentioned in the theorem is easily dealt with by looking at the Hilbert polynomial of an Ulrich bundle on X_d . So the point is about constructing Ulrich bundles in the prescribed ranks.

2020 *Mathematics Subject Classification.* 14J60, 14F06.

Key words and phrases. Ulrich bundles, Veronese variety, instanton bundles, equivariant bundles.

D. F. partially supported by FanoHK ANR-20-CE40-0023, SupToPhAG/EIPHI ANR-17-EURE-0002, Bridges ANR-21-CE40-0017, Capes/COFECUB Ma 926/19. V.P. partially supported by CAPES 88887.647097/2021-00, FAPESP 2022/12883-3. Both authors thank Departamento de Matemática de Belo Horizonte for the warm hospitality in October 2023.

About existence, one first notes that Ulrich bundles of rank r are available when $n!$ divides r , due to a simple and smart construction appearing in [ES09, Theorem 6.1]. On the other hand, finding Ulrich bundles of rank $r < n!$ may be challenging and actually impossible for low r . However Costa and Miró-Roig observed that, if 3 divides $d^2 - 1$ and \mathcal{E} is a sufficiently general instanton bundle of rank 2 and charge $\frac{1}{3}(d^2 - 1)$, then $\mathcal{E}(2(d - 1))$ is an Ulrich bundle on X_d . This relies on a difficult result of Hartshorne and Hirschowitz, actually the main result of [HH82], to the effect that \mathcal{E} has natural cohomology, meaning that for all $t \in \mathbb{Z}$ there is at most one value of $i \in \mathbb{N}$ giving $H^i(\mathcal{E}(t)) \neq 0$. This, combined with Riemann-Roch, implies the vanishing of cohomology required for $\mathcal{E}(2(d - 1))$ to be Ulrich.

So in order to complete the classification, the main obstacle is constructing Ulrich bundles of rank 3 in the prescribed range, which is precisely the main goal of this note. The rather surprising fact is that, instead of searching for general bundles relying on the usual techniques based on constructing bundles from curves (typically unions of lines and conics) combined with deformation theory, we achieve this using very special bundles, in some sense the most special we could think of, namely symmetric squares of equivariant instanton bundles for the action of SL_2 on \mathbb{P}^3 by binary cubics.

These instantons were examined in the framework of spaces of equivariant matrices of constant rank in [BFL22]. They had been previously constructed and classified in [Fae07] and actually much earlier, to the first author's astonishment, see [BS97]. The outcome of the classification is that, for any integer $m \geq 1$, there is one and only one instanton bundle \mathcal{E}_m of rank 2 on \mathbb{P}^3 which is SL_2 -equivariant for this action and having $c_2(\mathcal{E}) = \binom{m+1}{2}$. These are precisely the numerical invariants needed to get Ulrich bundles after taking symmetric squares.

To explain how these sheaves are useful in the present setting, let us state a second result. We write $\mathrm{MI}(r, k)$ for the coarse moduli space of rank- r stable instanton bundles having second Chern class kH^2 , where H is the hyperplane class of \mathbb{P}^3 . According to [JV14], the space $\mathrm{MI}(2, k)$ is a smooth variety which is moreover irreducible by [Tik12, Tik13]. We recall that Le Potier defined in [LP83] an effective divisor that we denote by $\mathrm{LP}(k) \subset \mathrm{MI}(2, k)$ consisting of instanton bundles \mathcal{E} such that $H^*(S^2\mathcal{E}(-2)) \neq 0$. We also consider $\mathrm{MI}_{\mathrm{Sp}}(r, k)$ and $\mathrm{MI}_{\mathrm{SO}}(r, k)$, the moduli spaces of symplectic and orthogonal instantons, namely pairs $([\mathcal{F}], [\eta])$, where $[\mathcal{F}]$ lies in $\mathrm{MI}(r, k)$ and $[\eta]$ is the proportionality class of $\eta : \mathcal{F} \rightarrow \mathcal{F}^\vee$, a skew-symmetric or symmetric duality. Hence actually $\mathrm{MI}(2, k) \simeq \mathrm{MI}_{\mathrm{Sp}}(2, k)$ for any $k \geq 1$. The following theorem can be thought of as an incarnation of the classical isomorphism of Lie algebras $\mathfrak{sl}_2(\mathbb{C}) \simeq \mathfrak{so}_3(\mathbb{C})$.

Theorem 2. *For any $k \geq 1$, the assignment $[\mathcal{E}] \mapsto [S^2\mathcal{E}]$ gives an étale map onto an open subset:*

$$\varphi : \mathrm{MI}(2, k) \setminus \mathrm{LP}(k) \rightarrow \mathrm{MI}_{\mathrm{SO}}(3, 4k).$$

For $e \in \{0, 1, 2\}$ and $h \geq 0$, with $(h, e) \neq (0, 0)$, write $d = 6h + 2e + 1$, $m = 3h + e$, $k = \binom{m+1}{2}$. Then, for $[\mathcal{E}]$ sufficiently general in $\mathrm{MI}(2, k)$, $S^2\mathcal{E}(2(d - 1))$ is an Ulrich bundle on X_d .

The crux of the argument is that, when \mathcal{E} is the SL_2 -equivariant instanton bundle \mathcal{E}_m mentioned above, we may show that $\mathcal{F}_m = S^2\mathcal{E}_m$ satisfies $H^*(\mathcal{F}_m(-d - 2)) = 0$. This uses a remarkable sheafified minimal graded free SL_2 -equivariant resolution of \mathcal{E}_m , obtained in [Fae07] and recovered in [BFL22], that looks like:

$$(1) \quad 0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2m - 1) \rightarrow V_{3m} \otimes \mathcal{O}_{\mathbb{P}^3}(-m - 1) \rightarrow V_{3m+1} \otimes \mathcal{O}_{\mathbb{P}^3}(-m) \rightarrow \mathcal{E}_m \rightarrow 0.$$

Here, for $p \geq 0$, $V_p = S^p V_1$ is the irreducible SL_2 -representation of weight p .

With this being done, one observes that $H^*(\mathcal{F}_m(-d - 2)) = 0$ follows from Serre duality simply because \mathcal{F}_m is orthogonal. Then, choosing $[\mathcal{E}] \notin \mathrm{LP}(k)$ in a neighborhood of $[\mathcal{E}_m]$ will turn out in $S^2\mathcal{E}(2(d - 1))$ being an Ulrich bundle. In the very last passage, we use the irreducibility of $\mathrm{MI}(2, k)$. We could actually bypass it by showing $H^*(\mathcal{F}_m(-2)) = 0$, which we checked for several values of m but failed to prove in general. Note that \mathcal{E}_m itself is far from having natural cohomology, as the resolution (1) shows, but $S^2\mathcal{E}_m$ should have natural cohomology, at least we know it does for low values of m .

Remark 3. For $n \geq 4$ and arbitrary d , it seems difficult to determine $\mathrm{Ur}(X_n^d)$. Here are some comments.

- i) According to arithmetic conditions on d and n , we may determine $\mathrm{Ur}(X_n^d)$, for instance if $n!$ divides d then $\mathrm{Ur}(X_n^d) = n!\mathbb{N}^*$, see Lemma 4.
- ii) For most values of d modulo $n!$, we may not have a full picture of $\mathrm{Ur}(X_n^d)$. For instance, for $n = 5$ and $d = 2$, for any Ulrich bundle \mathcal{U} of rank r the discussion of Lemma 4 gives $8 \mid r$. Letting Sp_6 act linearly on \mathbb{P}^5 , we may consider the Sp_6 -bundle \mathcal{F} associated with the irreducible representation of

maximal weight $\varpi_2 + \varpi_3$, where $\varpi_1, \varpi_2, \varpi_3$ are the fundamental weights for Sp_6 . Then Borel-Weil ensures that $\mathcal{U} = \mathcal{F}(1)$ is an Ulrich bundle on X_2^5 . We have $\mathrm{rk}(\mathcal{U}) = 16$. But for any odd number $k < 15$, we don't know if X_2^5 supports Ulrich bundles of rank $8k$.

- iii) For $n \geq 4$ and $d \geq 2$, according to recent work of Lopez and Raychaudhury, see [LR24], X_d^n does not carry Ulrich bundles of rank ≤ 3 . This indicates that the situation for X_d^3 is very special.

2. PROOF OF THE MAIN RESULTS

Let us go through the proof of the results announced in the introduction. In the next subsection we recall some down-to-earth arithmetics about Ulrich bundles and classify Ulrich ranks on Veronese varieties in the simplest situation. Then we show a rather unexpected lemma stating that the SL_2 equivariant instantons considered here, in spite of being far from having natural cohomology, have a symmetric square that (almost) does. In the following subsection we interpret the symmetric square in terms of moduli spaces and show the fact bundles obtained this way fill an open dense subset of a component of their moduli space. This is surprising at first sight, but turns out to be very natural by looking at adjoint bundles. The last subsection summarizes the proof on Theorem 1.

2.1. Reminder on arithmetics of Ulrich bundles. Let \mathcal{U} be an Ulrich bundle of rank r on X_d^n . Then, the Hilbert polynomial of \mathcal{U} looks as follows

$$\chi(\mathcal{U}(t)) = \frac{r}{n!} \prod_{1 \leq j \leq n} (dj + t).$$

In particular, evaluating at the above expression for $t = 1$, we get:

$$(2) \quad n! \chi(\mathcal{U}(1)) = r \prod_{1 \leq j \leq n} (dj + 1).$$

Lemma 4. *We have $n! \mathbb{N}^* \subset \mathrm{Ur}(X_d^n)$. Moreover, if $n!$ divides d , then $\mathrm{Ur}(X_d^n) = n! \mathbb{N}^*$.*

Proof. This follows from the construction of [ES09, Theorem 6.1]. Indeed, we consider the product Y of n copies of \mathbb{P}^1 as a subvariety of degree $n!$ in $\mathbb{P}^{2^n - 1}$. Let π be a generic projection of Y to \mathbb{P}^n , so that π is a finite map of degree $n!$. Choose $s \in \mathbb{N}^*$. Then, with obvious notation, $\mathcal{U} = \pi_*(\mathcal{O}_Y(d-1, \dots, nd-1)^{\oplus s})$ is an Ulrich bundle on X_d^n of rank $n!s$. Therefore $n! \mathbb{N}^* \subset \mathrm{Ur}(X_d^n)$.

On the other hand, assuming that $n!$ divides d , we get that $\prod_{1 \leq j \leq n} (dj + 1)$ is prime to $n!$. So, given an Ulrich bundle \mathcal{U} on X_d^n of rank r , formula (2) shows that $n!$ divides r . Hence $\mathrm{Ur}(X_d^n) \subset n! \mathbb{N}^*$. \square

2.2. Cohomology vanishing in the positive quadrant. Let us consider the action of SL_2 on \mathbb{P}^3 by binary cubics, namely we identify \mathbb{P}^3 with $\mathbb{P}(V_3)$, in other words SL_2 acts linearly on \mathbb{P}^3 with the irreducible representation of weight 3. We use the notion of *instanton bundle* on \mathbb{P}^3 , which is an H -stable locally free sheaf \mathcal{E} of rank $r > 0$, such that $c_1(\mathcal{E}) = c_3(\mathcal{E}) = 0$ and $H^*(\mathcal{E}(-2)) = 0$. Writing $c_2(\mathcal{E}) = kH^2$, we say that k is the *charge* of \mathcal{E} . Note that stability is not always part of the definition of instanton bundle, however we do require it here. Also, note that $H^0(S^2\mathcal{E}) = 0$ for any instanton bundle \mathcal{E} of rank 2. Indeed, \mathcal{E} is of rank 2 with $c_1(\mathcal{E}) = 0$ so $\wedge^2 \mathcal{E} \simeq \mathcal{O}_{\mathbb{P}^3}$. Also, \mathcal{E} is stable, hence simple, so $\mathrm{Hom}(\mathcal{E}, \mathcal{E}^\vee) \simeq \mathrm{Hom}(\mathcal{E}, \mathcal{E})$ is one-dimensional, hence generated by the skew-symmetric duality responsible for the isomorphism $\wedge^2 \mathcal{E} \simeq \mathcal{O}_{\mathbb{P}^3}$. So there is no non-zero symmetric map $\mathcal{E} \rightarrow \mathcal{E}^\vee$.

Now, recall that according to [Fae07], for all $m \geq 1$ there is one and only one instanton bundle \mathcal{E}_m of rank 2 on \mathbb{P}^3 which is SL_2 -equivariant for the action of binary cubics and has charge $\binom{m+1}{2}$. Also, a rank-2 instanton bundle which is SL_2 -equivariant for the action of binary cubics is necessarily of the form \mathcal{E}_m .

Lemma 5. *For $e \in \{0, 1, 2\}$ and $h \geq 0$, with $(h, e) \neq (0, 0)$, write $d = 6h + 2e + 1$, $m = 3h + e$, $k = \binom{m+1}{2}$. Set $\mathcal{F}_m = S^2 \mathcal{E}_m$. Then we have $H^*(\mathcal{F}_m(d-2)) = 0$.*

Proof. We have $d = 2m + 1$ and a decomposition $\mathcal{E}_m \otimes \mathcal{E}_m$ as $\mathcal{F}_m \oplus \mathcal{O}_{\mathbb{P}^3}$, so

$$H^k(\mathcal{F}_m(t)) = H^k(\mathcal{E}_m \otimes \mathcal{E}_m(t)), \quad \text{for all } t \in \mathbb{Z} \text{ if } k \in \{1, 2\}.$$

We saw in the previous paragraph that $H^0(\mathcal{F}_m) = 0$, hence $H^0(\mathcal{F}_m(-t)) = 0$ for any $t \geq 0$. By Serre duality, since \mathcal{F}_m is self-dual, we get $H^3(\mathcal{F}_m(t-4)) = 0$ for any $t \geq 0$, in particular $H^3(\mathcal{F}_m(d-2)) = 0$.

We write ψ for the map $V_{3m} \otimes \mathcal{O}_{\mathbb{P}^3}(-m-1) \rightarrow V_{3m+1} \otimes \mathcal{O}_{\mathbb{P}^3}(-m)$ appearing in the resolution (1) of \mathcal{E}_m and consider such resolution as a two-step extension of \mathcal{E}_m and $\mathcal{O}_{\mathbb{P}^3}(-2m-1)$ given by an element

$$\zeta \in \text{Ext}_{\mathbb{P}^3}^2(\mathcal{E}_m, \mathcal{O}_{\mathbb{P}^3}(-2m-1)).$$

For all integers k and any locally free \mathcal{G} , we look at the Yoneda map:

$$\Upsilon_{\mathcal{G},k} : H^k(\mathcal{G} \otimes \mathcal{E}_m) \otimes \text{Ext}_{\mathbb{P}^3}^2(\mathcal{E}_m, \mathcal{O}_{\mathbb{P}^3}(-2m-1)) \rightarrow H^{k+2}(\mathcal{G}(-2m-1)).$$

This is functorial with respect to morphisms $\lambda : \mathcal{G} \rightarrow \mathcal{G}'$, namely, these give rise to commutative diagrams:

$$(3) \quad \begin{array}{ccc} H^k(\mathcal{G} \otimes \mathcal{E}_m) \otimes \text{Ext}_{\mathbb{P}^3}^2(\mathcal{E}_m, \mathcal{O}_{\mathbb{P}^3}(-2m-1)) & \xrightarrow{\Upsilon_{\mathcal{G},k}} & H^{k+2}(\mathcal{G}(-2m-1)) \\ \downarrow H^k(\lambda \otimes \text{id}) \otimes \text{id} & & \downarrow H^{k+2}(\lambda \otimes \text{id}) \\ H^k(\mathcal{G}' \otimes \mathcal{E}_m) \otimes \text{Ext}_{\mathbb{P}^3}^2(\mathcal{E}_m, \mathcal{O}_{\mathbb{P}^3}(-2m-1)) & \xrightarrow{\Upsilon_{\mathcal{G}',k}} & H^{k+2}(\mathcal{G}'(-2m-1)) \end{array}$$

We again use the resolution (1) to observe that cupping with ζ induces isomorphisms:

$$\begin{aligned} \Upsilon_{\mathcal{O}_{\mathbb{P}^3}(m-2),1}(- \otimes \zeta) &: H^1(\mathcal{E}_m(m-2)) \rightarrow H^3(\mathcal{O}_{\mathbb{P}^3}(-m-3)), \\ \Upsilon_{\mathcal{O}_{\mathbb{P}^3}(m-1),1}(- \otimes \zeta) &: H^1(\mathcal{E}_m(m-1)) \rightarrow H^3(\mathcal{O}_{\mathbb{P}^3}(-m-2)). \end{aligned}$$

We denote both such isomorphisms by $\wedge \zeta$.

Consider the resolution of $\mathcal{E}_m \otimes \mathcal{E}_m(d-2)$ obtained by tensoring (1) by $\mathcal{E}_m(d-2)$, that is,

$$0 \rightarrow \mathcal{E}_m(-2) \rightarrow V_{3m} \otimes \mathcal{E}_m(m-2) \rightarrow V_{3m+1} \otimes \mathcal{E}_m(m-1) \rightarrow \mathcal{E}_m \otimes \mathcal{E}_m(d-2) \rightarrow 0.$$

We already know that $H^*(\mathcal{E}_m(-2)) = 0$ since \mathcal{E}_m is an instanton bundle. Again from (1) we see

$$H^k(\mathcal{E}_m(m-1)) = H^k(\mathcal{E}_m(m-2)) = 0, \quad \text{if } k \neq 1.$$

Therefore $H^2(\mathcal{F}_m(d-2)) = H^2(\mathcal{E}_m \otimes \mathcal{E}_m(d-2)) = 0$. Moreover we get a map

$$f : V_{3m} \otimes H^1(\mathcal{E}_m(m-2)) \rightarrow V_{3m+1} \otimes H^1(\mathcal{E}_m(m-1)),$$

obtained as $f = H^1(\psi \otimes \mathcal{E}_m(2m-1))$, with the property that

$$(4) \quad H^0(\mathcal{E}_m \otimes \mathcal{E}_m(d-2)) \simeq \ker(f), \quad H^1(\mathcal{F}_m(d-2)) = H^1(\mathcal{E}_m \otimes \mathcal{E}_m(d-2)) \simeq \text{coker}(f).$$

Then applying (3) to $\psi \otimes \mathcal{O}_{\mathbb{P}^3}(2m-2)$ yields a commutative diagram induced by cupping with ζ :

$$\begin{array}{ccc} V_{3m} \otimes H^1(\mathcal{E}_m(m-2)) & \xrightarrow{V_{3m} \otimes \wedge \zeta} & V_{3m} \otimes H^3(\mathcal{O}_{\mathbb{P}^3}(-m-3)) \\ \downarrow f = H^1(\psi \otimes \mathcal{E}_m(2m-1)) & & \downarrow H^3(\psi \otimes \mathcal{O}_{\mathbb{P}^3}(-2)) \\ V_{3m+1} \otimes H^1(\mathcal{E}_m(m-1)) & \xrightarrow{V_{3m+1} \otimes \wedge \zeta} & V_{3m+1} \otimes H^3(\mathcal{O}_{\mathbb{P}^3}(-m-2)) \end{array}$$

The horizontal arrows are isomorphisms and the vertical arrow on the right is surjective since $H^*(\mathcal{E}_m(-2)) = 0$. Hence the vertical arrow on the left is also surjective. Also, the kernel of $H^3(\psi \otimes \mathcal{O}_{\mathbb{P}^3}(-2))$ is identified with $H^3(\mathcal{O}_{\mathbb{P}^3}(-2m-3)) = H^3(\mathcal{O}_{\mathbb{P}^3}(-d-2))$. Therefore, (4) gives:

$$\begin{aligned} H^1(\mathcal{F}_m) &= H^1(\mathcal{E}_m \otimes \mathcal{E}_m(d-2)) = 0 \\ H^0(\mathcal{F}_m(d-2)) \oplus H^0(\mathcal{O}_{\mathbb{P}^3}(d-2)) &= H^0(\mathcal{E}_m \otimes \mathcal{E}_m(d-2)) \simeq H^3(\mathcal{O}_{\mathbb{P}^3}(-d-2)). \end{aligned}$$

Hence $H^0(\mathcal{F}_m(d-2)) = 0$. This finishes the proof that $H^*(\mathcal{F}_m(d-2)) = 0$. \square

2.3. The symmetric square as map to the space of orthogonal instantons. Consider an element $([\mathcal{F}], [\eta])$ of $\text{MI}_{\text{SO}}(r, k)$ or $\text{MI}_{\text{Sp}}(r, k)$. Then \mathcal{F} is a stable bundle, isomorphic to \mathcal{F}^\vee via η . If $\eta' : \mathcal{F} \rightarrow \mathcal{F}^\vee$ is not proportional to η , then in the pencil $\lambda\eta + \lambda'\eta'$, for $[\lambda : \lambda'] \in \mathbb{P}^1$, there must be a morphism $\mathcal{F} \rightarrow \mathcal{F}^\vee$ having trivial determinant. However, \mathcal{F} and \mathcal{F}^\vee are stable of the same Hilbert polynomial, so this cannot happen unless η and η' are proportional. So $[\eta]$ is actually determined uniquely by $[\mathcal{F}]$ and we will suppress $[\eta]$ from the notation of the points in $\text{MI}_{\text{SO}}(r, k)$ and $\text{MI}_{\text{Sp}}(r, k)$.

Also, since any $[\mathcal{E}] \in \text{MI}(2, k)$ is equipped with skew-symmetric duality $\eta_{\mathcal{E}}$ which is canonical up to a non-zero scalar, $\mathcal{F} = S^2\mathcal{E}$ is equipped with the canonical class $[\eta_{\mathcal{F}}]$ of the symmetric duality $\eta_{\mathcal{F}} = S^2\eta_{\mathcal{E}}$.

Lemma 6. *Let $k \geq 1$ be an integer. Then the assignment $[\mathcal{E}] \rightarrow [S^2\mathcal{E}]$ gives an étale map*

$$\varphi : \text{MI}(2, k) \setminus \text{LP}(k) \rightarrow \text{MI}_{\text{SO}}(3, 4k)$$

and $\varphi(\text{MI}(2, k))$ is a smooth irreducible Zariski-dense open subset of a component of $\text{MI}_{\text{SO}}(3, 4k)$.

Proof. Let \mathcal{E} be an instanton bundle in $\text{MI}(2, k)$. We know from [JV14] that $\text{MI}(2, k)$ is smooth at \mathcal{E} . More precisely, the tangent space and the obstruction space to $\text{MI}(2, k)$ at the point $[\mathcal{E}]$ corresponding to \mathcal{E} are described as:

$$\mathcal{T}_{[\mathcal{E}]} \text{MI}(2, k) = H^1(S^2\mathcal{E}), \quad \text{with } h^1(S^2\mathcal{E}) = 8k - 3, \quad \mathcal{O}b_{[\mathcal{E}]} \text{MI}(2, k) = H^2(S^2\mathcal{E}) = 0.$$

Now, assuming $[\mathcal{E}]$ to lie away from the Le Potier divisor $\text{LP}(k)$, by definition $\mathcal{F} = S^2\mathcal{E}$ verifies $H^*(\mathcal{F}(-2)) = 0$. Moreover \mathcal{F} is slope-semistable since $S^2\mathcal{E}$ is and actually \mathcal{F} is slope-stable since $h^0(\mathcal{F}) = h^0(\mathcal{F}^\vee) = 0$. Therefore, computing Chern classes we see that \mathcal{F} is an instanton bundle corresponding to a point $[\mathcal{F}]$ of $\text{MI}_{\text{SO}}(3, 4k)$.

Now recall that, for an orthogonal bundle \mathcal{F} , the adjoint bundle $\text{ad}(\mathcal{F})$ is defined over every fibre of \mathcal{F} by the adjoint representation $\text{SO}(3) \rightarrow \text{ad}(\text{SO}(3))$, see for instance [Ram75]. For a given 3-dimensional vector space F , the associated adjoint representation is isomorphic to the exterior square $\wedge^2 F$, so $\text{ad}(\mathcal{F}) \simeq \wedge^2 \mathcal{F}$. Moreover, since \mathcal{F} has rank 3 and trivial determinant, we have $\wedge^2 \mathcal{F} \simeq \mathcal{F}^\vee$. However \mathcal{F} being orthogonal, it is self-dual, namely $\mathcal{F} \simeq \mathcal{F}^\vee$. Therefore, the tangent space and the obstruction space of $\text{MI}_{\text{SO}}(3, 4k)$ at $[\mathcal{F}]$ are described as

$$\mathcal{T}_{[\mathcal{F}]} \text{MI}_{\text{SO}}(3, 4k) = H^1(\text{ad}(\mathcal{F})) \simeq H^1(\mathcal{F}), \quad \mathcal{O}b_{[\mathcal{F}]} \text{MI}_{\text{SO}}(3, 4k) = H^2(\text{ad}(\mathcal{F})) \simeq H^2(\mathcal{F})$$

Therefore we get the following information:

$$\begin{aligned} \mathcal{T}_{[\mathcal{E}]} \text{MI}(2, k) &= H^1(S^2\mathcal{E}) \simeq H^1(\mathcal{F}) = \mathcal{T}_{[\mathcal{F}]} \text{MI}_{\text{SO}}(3, 4k), \\ 0 = \mathcal{O}b_{[\mathcal{E}]} \text{MI}(2, k) &= H^2(S^2\mathcal{E}) \simeq H^2(\wedge^2 \mathcal{F}) = \mathcal{O}b_{[\mathcal{F}]} \text{MI}_{\text{SO}}(3, 4k), \end{aligned}$$

where the isomorphism of tangent spaces is the differential of φ at $[\mathcal{E}]$. This proves that $\text{MI}_{\text{SO}}(3, 4k)$ is smooth at $[\mathcal{F}]$ and that φ is a smooth morphism of relative dimension 0, hence étale. Recall that $\text{MI}(2, k)$ is irreducible by [Tik12, Tik13]. Then, the image of φ is a smooth Zariski-dense subset of an irreducible component of $\text{MI}_{\text{SO}}(3, 4k)$. \square

2.4. Proof of the main theorems. All that remains to finish the proof of Theorem 2 is to combine the previous lemmas, the semicontinuity of the dimension of the cohomology groups and the result in [CMR21, Theorem 3.6] for the case of $r = 3$. So that we just have to construct a rank 3 instanton bundle of natural cohomology with charge $k = \binom{m+1}{2}$ for some integer $m \geq 1$ and $d = 2m + 1$.

Proof of Theorem 2. The first part of the statement is done in Lemma 6. To conclude we have to prove the existence of an Ulrich bundle of the form $S^2\mathcal{E}(2(d-1))$ on X_d for some $[\mathcal{E}]$ sufficiently general in $\text{MI}(2, k)$. For any $t \in \mathbb{Z}$, we consider the open subset W_t of $\text{MI}_{\text{SO}}(3, 4k)$ defined as

$$W_t = \{[\mathcal{F}] \in \text{MI}_{\text{SO}}(3, 4k) \mid H^*(\mathcal{F}(t-2)) = 0\}.$$

The intersection $U = W_d \cap W_{-d} \cap \varphi(\text{MI}(2, k))$ is non-empty because the class of $\mathcal{F}_m = S^2\mathcal{E}_m$ belongs to U . Indeed, Lemma 5 applies to guarantee that $[\mathcal{F}_m] \in W_d$. Moreover, since \mathcal{F}_m is orthogonal, hence self-dual, using Serre duality we see that $[\mathcal{F}_m] \in W_{-d}$, that is:

$$H^j(\mathcal{F}_m(-d-2))^\vee \simeq H^{3-j}(\mathcal{F}_m(d-2)) = 0, \quad \text{for all } j \in \mathbb{N}.$$

Then U is an open subset of an irreducible component of $\text{MI}_{\text{SO}}(3, 4k)$ by Lemma 6. Furthermore, the intersection of U with $\varphi(\text{MI}(2, k) \setminus \text{LP}(k))$ is also non-empty, being an intersection of non-empty open subsets inside the irreducible variety $\varphi(\text{MI}(2, k))$.

A representative of a class in the intersection $U \cap \varphi(\text{MI}(2, k) \setminus \text{LP}(k))$ is a rank 3 instanton bundle \mathcal{F} with charge $(d^2 - 1)/2$, satisfying $H^*(\mathcal{F}(-2 - td)) = 0$ for $t \in \{-1, 0, 1\}$. Therefore $\mathcal{F}(2d - 2)$ is an Ulrich bundle on X_d . \square

Proof of Theorem 1. Let us separate the various four cases of \bar{d} . Up to taking direct sums, it suffices to find Ulrich bundles of rank 2 and/or 3 in each case.

- i) This case is proven in Lemma 4.

- ii) When $\bar{d} \in \{1, 5\}$, then both 2 and 3 divide $d^2 - 1$. Note that $\bar{d} = 1$ and $\bar{d} = 5$ correspond respectively to the choices $e = 0$ and $e = 2$ of Theorem 2. From Theorem 2 and [CMR21, Proposition 3.7], we conclude the existence of Ulrich bundles of rank 2 and 3 and consequently any rank $r \geq 2$.
- iii) If $\bar{d} \in \{2, 4\}$ then 2 divides $d^2 - 1$ but 3 does not. Then we only need to apply [CMR21, Proposition 3.7] to conclude this case.
- iv) For $\bar{d} = 3$, we have that 3 divides $d^2 - 1$ but 2 does not. This corresponds to the case $e = 1$ in Theorem 2. Applying Theorem 2 affords the existence of Ulrich bundles of rank 3 and thereby the existence of Ulrich bundles of rank $3\mathbb{N}^*$.

□

REFERENCES

- [AFO17] Marian Aprodu, Gavril Farkas, and Angela Ortega, *Minimal resolutions, Chow forms and Ulrich bundles on K3 surfaces*, J. Reine Angew. Math. **730** (2017), 225–249.
- [BES17] Markus Bläser, David Eisenbud, and Frank-Olaf Schreyer, *Ulrich complexity*, Differential Geom. Appl. **55** (2017), 128–145.
- [BS97] Gil Bor and Jan Segert, *Symmetric instantons and the ADHM construction*, Comm. Math. Phys. **183** (1997), no. 1, 183–203.
- [BFL22] Ada Boralevi, Daniele Faenzi, and Paolo Lella, *A construction of equivariant bundles on the space of symmetric forms*, Rev. Mat. Iberoam. **38** (2022), no. 3, 761–782.
- [CFK23] Ciro Ciliberto, Flaminio Flamini, and Andreas Leopold Knutsen, *Ulrich bundles on Del Pezzo threefolds*, J. Algebra **634** (2023), 209–236.
- [CMR18] Laura Costa and Rosa Maria Miró-Roig, *Ulrich bundles on Veronese surfaces*, Singularities, algebraic geometry, commutative algebra, and related topics, Springer, Cham, 2018, pp. 375–381.
- [CMR21] ———, *Instanton bundles vs Ulrich bundles on projective spaces*, Beitr. Algebra Geom. **62** (2021), no. 2, 429–439.
- [CMRPL21] Laura Costa, Rosa María Miró-Roig, and Joan Pons-Llopis, *Ulrich bundles—from commutative algebra to algebraic geometry*, De Gruyter Studies in Mathematics, vol. 77, De Gruyter, Berlin, [2021] ©2021.
- [ES03] David Eisenbud and Frank-Olaf Schreyer, *Resultants and Chow forms via exterior syzygies*, J. Amer. Math. Soc. **16** (2003), no. 3, 537–579, With an appendix by Jerzy Weyman.
- [ES09] ———, *Betti numbers of graded modules and cohomology of vector bundles*, J. Amer. Math. Soc. **22** (2009), no. 3, 859–888.
- [Fae07] Daniele Faenzi, *Homogeneous instanton bundles on \mathbb{P}^3 for the action of $SL(2)$* , J. Geom. Phys. **57** (2007), no. 10, 2146–2157.
- [Fae19] ———, *Ulrich bundles on K3 surfaces*, Algebra Number Theory **13** (2019), no. 6, 1443–1454.
- [HH82] Robin Hartshorne and André Hirschowitz, *Cohomology of a general instanton bundle*, Ann. Sci. École Norm. Sup. (4) **15** (1982), no. 2, 365–390.
- [JV14] Marcos Jardim and Misha Verbitsky, *Trihyperfähler reduction and instanton bundles on $\mathbb{C}\mathbb{P}^3$* , Compos. Math. **150** (2014), no. 11, 1836–1868.
- [LP83] Joseph Le Potier, *Sur l'espace de modules des fibrés de Yang et Mills*, Mathematics and physics (Paris, 1979/1982), Progr. Math., vol. 37, Birkhäuser Boston, Boston, MA, 1983, pp. 65–137.
- [LR24] Angelo Felice Lopez and Debaditya Raychaudhury, *Non-existence of low rank Ulrich bundles on Veronese varieties*, In preparation, available at <http://ricerca.matfis.uniroma3.it/users/lopez/Veronese.pdf>, 2024.
- [Ram75] Annamalai Ramanathan, *Stable principal bundles on a compact Riemann surface*, Math. Ann. **213** (1975), 129–152.
- [Tik12] Alexander S. Tikhomirov, *Moduli of mathematical instanton vector bundles with odd c_2 on projective space*, Izv. Ross. Akad. Nauk Ser. Mat. **76** (2012), no. 5, 143–224.
- [Tik13] ———, *Moduli of mathematical instanton vector bundles with even c_2 on projective space*, Izv. Ross. Akad. Nauk Ser. Mat. **77** (2013), no. 6, 139–168.

DANIELE FAENZI, INSTITUT DE MATHÉMATIQUES DE BOURGOGNE, UMR 5584 CNRS, UNIVERSITÉ DE BOURGOGNE, 9 AVENUE ALAIN SAVARY, F-21000, DIJON, FRANCE
Email address: daniele.faenzi@u-bourgogne.fr

VICTOR PRETTI, INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, 1010 RUA DO MATÃO, SÃO PAULO, BRAZIL
Email address: edu-vp.zosxx@simplelogin.com