

## MAKAHO: An interactive cartographic Tool for Trend Analysis of hydrological extremes in France

Louis Héraut, Michel Lang, Benjamin Renard, Eric Sauquet, Jean-Philippe

Vidal

#### ▶ To cite this version:

Louis Héraut, Michel Lang, Benjamin Renard, Eric Sauquet, Jean-Philippe Vidal. MAKAHO: An interactive cartographic Tool for Trend Analysis of hydrological extremes in France. EGU24, Apr 2024, Wien, Austria, France. . hal-04582254

### HAL Id: hal-04582254 https://hal.science/hal-04582254

Submitted on 21 May 2024  $\,$ 

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Louis Héraut<sup>1</sup>, Michel Lang<sup>1</sup>, Benjamin Renard<sup>2</sup>, Éric Sauquet<sup>1</sup> and Jean-Philippe Vidal<sup>1</sup> <sup>1</sup>INRAE, UR RiverLy, Villeurbanne, France, <sup>2</sup>INRAE, UMR RECOVER, Aix-en-Provence, France



**232 stations** belonging to the French Reference Hydrometric Network (*Giuntoli et al.*, 2013). These stations show a **high measurement quality**, time series with a historical depth of over 40 years, and they crucially gauge near-natural catchments. The statistical test used for trend detection is a variant of the Mann-Kendall test accounting for first-order autocorrelation. The trend slope is provided by the **Theil-Sen estimator**.

Giuntoli, I., Renard, B., Vidal, J.-P., and Bard, A. (2013). Low flows in france and their relationship to largescale climate indices. Journal of Hydrology, 482:105–118. https:/doi.org/10.1016/j.jhydrol.2012.12.038

# MAKAHO : An interactive cartographic Tool for Trend Analysis of hydrological extremes in France

> The duration of low flows has significantly increased in all the country with an average of more than **10 days per decade**, reaching almost a month in extreme cases in the southwest.





between the northern upward trends and the southern downward trends in mean discharge.

> Nearly **10 % of stations** show an **upward significant trend** in the **annual maximum daily** streamflow at the 90 % confidence level with an average increase of 13 % per decade. Almost all of these stations are located in the northern part of the country.

> Mainly located in the south, around 40 % of stations show a decreasing trend in the **annual minimum monthly discharge**, with an average intensity of about 11 % per decade.





The definition of the **hydrological year** is fully customizable and an optimal hydrological year is proposed based on hydrological considerations about the variable. The **analysis period** can be adjust as well as the **significance level** of the Mann-Kendall test. 1.76 x 10<sup>-2</sup> [m<sup>3</sup>.s<sup>-1</sup>.vear<sup>-1</sup>] -0.56 [%.vear 20 - in the million will be will will will be the will be will be the will be the will be the will be the will be

A click on a station displays the **time series** for the variable of interest.

The daily discharge series are stored in the form of a table and are entirely processed by the **EXstat** R package. Initially, the aggregation of daily data into an annual variable



	Bestionn Bassin h Superfic Altitude X = 8973 Y = 6402	rafre : DVFAL Rinder-Alpes vidrographique : Rhône cie : 227 [km <sup>2</sup> ] Date de début : 01/01/1963 Date de fin : 14/02/2021 Jay [m ; Lambert93] Taux de lacunes : 0.4 [%] JFMAM J J ASOND
	125 -	1970 1980 1990 2000 2010 2
ċ.	100 - 75 -	
<u> </u>	50 - 25 -	
	0 -	ikas, and to population and an and an an an an an an an and an and an and a state of the second and an and an a In the second and an
		-1.42 x 10 <sup>-1</sup> [m <sup>1</sup> .s <sup>-1</sup> .an <sup>-1</sup> ] -0.40 [%.an <sup>-1</sup> ]
let _	100 -	
/31 jui	75 -	
01 août	50 -	
,	25 -	
	0 -	
2	iuin -	
01 août / 31 juillet	avril -	
	févr.	
	déc	
5000	oct	
		-1.09 x 10 <sup>-3</sup> [jour.an <sup>-2</sup> ]
		•••
ljour.ari 0t/31 tuillet	0.2 -	
	0.1-	
6 <sup>6</sup>	0.11	
	0.0-	
		-5.03 x 10 <sup>-4</sup> [jour.an <sup>-3</sup> ]
_	0.15	· · ·
juillet	0.10 -	
noût/31	0.05	
6 <sup>10</sup>	0.05 -	
	0.00 -	
	0.04 -	0.00 x 10 <sup>-1</sup> [jour.an <sup>-2</sup> ]
- *	0.03 -	•
'31 juillé	0.02 -	· · · · · · · · · · · · · · · · · · ·
1 août /	0.01 -	
. 0	0.00	
-	0.00 -	



Shiny web hosting by SK8 INRAE project

## Datasheets

Each hydrometric station has **technical** sheets summarizing all trend analysis results for a selection of variables. These datasheets are directly downloadable from the website.

louis.heraut@inrae.fr

