



HAL
open science

THE L_p -DUAL SPACE OF A SEMISIMPLE LIE GROUP

Bachir Bekka

► To cite this version:

| Bachir Bekka. THE L_p -DUAL SPACE OF A SEMISIMPLE LIE GROUP. 2024. hal-04582339

HAL Id: hal-04582339

<https://hal.science/hal-04582339>

Preprint submitted on 21 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE L_p -DUAL SPACE OF A SEMISIMPLE LIE GROUP

BACHIR BEKKA

ABSTRACT. Let G be a semisimple Lie group. We describe the irreducible representations of G by linear isometries on L_p -spaces for $p \in (1, +\infty)$ with $p \neq 2$. More precisely, we show that, for every such representation π , there exists a parabolic subgroup Q of G such that π is equivalent to the natural representation of G on $L_p(G/Q)$ twisted by a unitary character of Q . When G is of real rank one, we give a complete classification of the possible irreducible representations of G on an L_p -space for $p \neq 2$, up to equivalence.

1. INTRODUCTION

Let G be a locally compact topological group and E a Banach space. Denote by $\mathcal{B}(E)$ the algebra of bounded operators on E , equipped with the strong operator topology, that is, the weakest topology for which the map $\mathcal{B}(E) \rightarrow E, T \mapsto Tv$ is continuous for every $v \in E$. An *isometric representation* (for short, a *representation*) of G on E is a continuous group homomorphism $\pi : G \rightarrow \text{Iso}(E)$, where $\text{Iso}(E)$ is the subgroup of $GL(E)$ consisting of the linear surjective isometries on E .

In the context of semisimple Lie groups, there is strong evidence (see [HC53], [God52], [War72], [Fel65]) that the right notions of irreducibility and equivalence of Banach representations are defined as follows. Recall that to a representation $\pi : G \rightarrow \text{Iso}(E)$ of G there is associated an algebra homomorphism $C_c(G) \rightarrow \mathcal{B}(E)$, denoted by π again, defined by

$$\pi(f) = \int_G f(g)\pi(g)d\mu_G(g) \quad \text{for all } f \in C_c(G),$$

where $C_c(G)$ is the convolution algebra of continuous functions on G with compact support and μ_G is a left Haar measure on G .

Definition 1. (i) A representation $\pi : G \rightarrow \text{Iso}(E)$ is *completely irreducible* if the algebra $\pi(C_c(G))$ is dense in $\mathcal{B}(E)$ for the strong operator topology.

(ii) Two representations $\pi_1 : G \rightarrow \text{Iso}(E_1)$ and $\pi_2 : G \rightarrow \text{Iso}(E_2)$ on Banach spaces E_1 and E_2 are *Naimark equivalent* (for short, *equivalent*) if, for $i \in \{1, 2\}$, there exists a dense subspace V_i of E_i which are $\pi_i(C_c(G)$ -invariant and a closed injective linear map $T : V_1 \rightarrow V_2$ such that

$$\pi_2(f)Tv = T\pi_1(f)v \quad \text{for all } v \in V_1, f \in C_c(G).$$

(iii) Let \mathcal{E} be a class of Banach spaces. The \mathcal{E} -*dual* of G , denoted by $\widehat{G}_{\mathcal{E}}$, is the set of equivalence classes of completely irreducible representations of G on some Banach space $E \in \mathcal{E}$;

When E is a Hilbert space, a representation $G \rightarrow \text{Iso}(E)$ is traditionally called a unitary representation; a unitary representation π is completely irreducible if and only if π is irreducible (that is, $\{0\}$ and E are the only G -invariant closed subspaces

Date: May 21, 2024.

2000 Mathematics Subject Classification. 22E46; 47L10; 37A40.

The author acknowledges the support by the ANR (French Agence Nationale de la Recherche) through the project Labex Lebesgue (ANR-11-LABX-0020-01) .

of E). Moreover, unitary representations π_1 and π_2 are Naimark equivalent if and only if π_1 and π_2 are unitarily equivalent (see [War72, 4.3]); in particular, if \mathcal{E} denotes the class of Hilbert spaces, then $\widehat{G}_{\mathcal{E}}$ coincides with the usual unitary dual \widehat{G} of G .

Let G a noncompact connected semisimple linear Lie group. For such a group, we will be concerned with \widehat{G}_{L_p} where L_p is the class of L_p -spaces for $p \in [1, +\infty[$. By an L_p -space, we mean the usual space $L_p(X, \mathcal{B}, \mu)$ of equivalence classes (modulo null sets) of measurable p -integrable functions $f : X \rightarrow \mathbf{C}$, where μ is a positive σ -finite measure defined on a standard Borel space (X, \mathcal{B}) . The unitary dual space $\widehat{G} = \widehat{G}_{L_2}$ is of course a classical much studied object (see for instance [Kna01], [War72]). We will deal here with \widehat{G}_{L_p} for $p \neq 2$.

Let θ be a Cartan involution on the Lie algebra \mathfrak{g} of G and \mathfrak{k} (respectively \mathfrak{p}) the eigenspace for the eigenvalue 1 (respectively -1) of θ . Let \mathfrak{a} be a maximal commutative subspace of \mathfrak{p} and let $\Sigma \subset \mathfrak{a}^*$ be the corresponding root system, where $\mathfrak{a}^* = \text{Hom}_{\mathbf{R}}(\mathfrak{a}, \mathbf{R})$ is the real dual space of \mathfrak{a} . Let Σ^+ be the set of positive roots for an ordering of Σ and \mathfrak{n} the sum of the root spaces \mathfrak{g}_{α} for $\alpha \in \Sigma^+$. Let K , A , and N be the subgroups of G with Lie algebras \mathfrak{k} , \mathfrak{a} and \mathfrak{n} , respectively. Then K is a maximal compact, A an abelian and N a nilpotent subgroup; moreover, we have an *Iwasawa decomposition* $G = KAN$.

Let M be the centralizer of A in K . The group MAN is a minimal parabolic subgroup of G . Recall (see [Kna01, Section V.5] or [War72, Chap. I, 1.2]) that a *parabolic subgroup* of G is a closed subgroup containing a conjugate of MAN . Every such group is conjugate to a *standard parabolic subgroup* $Q = Q_F$, which is parametrized by a subset F of the set Δ of simple positive roots; more precisely, let \mathfrak{a}_F be the intersection of all $\ker \alpha$ for $\alpha \in F$ and \mathfrak{n}_F the sum of the root spaces \mathfrak{g}_{α} for $\alpha \in \Sigma^+ \setminus \text{span}(F)$. We have a *Langlands decomposition*

$$Q = M_Q A_Q N_Q,$$

where M_Q is reductive and contains M , $A_Q = \exp(\mathfrak{a}_F)$ and $N_Q = \exp(\mathfrak{n}_F)$. Moreover, M_Q and A_Q commute, and both normalize N_Q . In particular, the standard minimal parabolic subgroup is given by $Q_{\emptyset} = MAN$.

Fix a parabolic subgroup $Q = Q_F$ of G and a real number $p \in [1, +\infty[$. We are going to define the L_p analog of the (unitary) principal series representations associated to Q .

Let $K_Q := K \cap M_Q$ and choose a Borel fundamental domain $\Omega_Q \subset K$ for the coset space K/K_Q . Then Ω_Q is also a fundamental domain for G/Q , since $G = KQ$ and $K \cap Q = K_Q$. Every $g \in G$ has a unique decomposition

$$g = \kappa(g)\mu(g)\exp(H(g))n(g) \quad \text{for } \kappa(g) \in \Omega_Q, \mu(g) \in M_Q, H(g) \in \mathfrak{a}_F, n(g) \in N_Q.$$

Let μ_Q be the unique quasi-invariant Borel probability measure on G/Q which is K -invariant. We identify G/Q with Ω_Q and transfer μ_Q and the natural G -action on G/Q to Ω_Q . Let $\rho_{Q,p} \in \mathfrak{a}^*$ be defined by

$$\rho_{Q,p} := \frac{1}{p} \sum_{\alpha \in \Sigma^+ \setminus \text{span}(F)} \dim(\mathfrak{g}_{\alpha})\alpha.$$

Observe that $\rho_{MAN,2}$ is the familiar half-sum of positive roots.

Fix a real linear form $\lambda \in \mathfrak{a}^*$ and a unitary character

$$\chi : M_Q \rightarrow \mathbf{S}^1 = \mathbf{U}(1)$$

of the reductive group M_Q . We define a representation $\pi(Q, \chi, \lambda, p)$ of G on

$$L_p(G/Q, \mu_Q) = L_p(\Omega_Q, \mu_Q)$$

by

$$\pi(Q, \chi, \lambda, p)(g)f(x) = \overline{\chi(\mu(g^{-1}x))} e^{-(i\lambda + \rho_{Q,p})H(g^{-1}x)} f(\kappa(g^{-1}x))$$

for $f \in L_p(\Omega_Q, \mu_F)$, $x \in \Omega_Q$, $g \in G$. It is easily verified (see Section 2) that $\pi(Q, \chi, \lambda, p)$ is indeed an isometric representation of G on $L_p(G/Q, \mu_Q)$.

We denote by $\widehat{M_Q^{\text{Ab}}}$ the group of unitary characters of M_Q . Here is our main result.

Theorem A. *Let G be a noncompact connected semisimple linear Lie group, and let $p \in (1, +\infty)$ with $p \neq 2$. Let π be a completely irreducible representation of G on an L_p -space. Then there exists a parabolic subgroup Q of G , a real linear form $\lambda \in \mathfrak{a}^*$, and a unitary character $\chi \in \widehat{M_Q^{\text{Ab}}}$ such that π is equivalent to the representation $\pi(Q, \chi, \lambda, p)$ on $L_p(G/Q, \mu_Q)$ described above.*

In order to study the irreducibility of a representation $\pi(Q, \chi, \lambda, p)$ as in the previous theorem, we consider its associated Harish-Chandra (\mathfrak{g}, K) -module. For a unitary representation σ of M_Q and for $\nu \in \mathfrak{a}_{\mathbf{C}}^* = \text{Hom}_{\mathbf{R}}(\mathfrak{a}, \mathbf{C})$, following [Wal88, 5.2], we denote by $I_{Q, \sigma, \nu}$ the associated infinitesimal nonunitary principal series of G , that is, the (\mathfrak{g}, K) -module underlying the induced representation

$$\text{Ind}_Q^G(\sigma \otimes \nu \otimes 1_{N_Q}) = \text{Ind}_{M_Q A_Q N_Q}^G(\sigma \otimes \nu \otimes 1_{N_Q}).$$

Proposition 2. (i) *Let $\pi(Q, \chi, \lambda, p)$ be a representation of G on $L_p(G/Q, \mu_Q)$ as in Theorem A. Then $\pi(Q, \chi, \lambda, p)$ is completely irreducible if and only if the principal series representation $I_{Q, \chi, \nu}$ is (algebraically) irreducible, where*

$$\nu = i\lambda + \delta_p \rho_{Q,2}$$

for $\delta_p = \frac{2}{p} - 1 \in (-1, 1)$.

(ii) *For $i = 1, 2$, let $\pi(Q_i, \chi_i, \lambda_i, p_i)$ be representations of G on $L_{p_i}(G/Q_i, \mu_{Q_i})$ as in Theorem A. Then $\pi(Q_1, \chi_1, \lambda_1, p_1)$ and $\pi(Q_2, \chi_2, \lambda_2, p_2)$ are equivalent if and only if the corresponding principal series representations I_{Q_1, χ_1, ν_1} and I_{Q_2, χ_2, ν_2} as in (i) are (algebraically) equivalent.*

Principal series representations $I_{Q, \sigma, \nu}$ are of course central objects in the representation theory of semisimple Lie groups; their structure has been much studied, especially in the case where Q is minimal parabolic, and it has been shown that “most” of them are irreducible (see [GN57], [Bru56], [PRRV67], [Kos69], [Wal71], [LW73]).

Using known irreducibility and equivalence results from the literature, we can settle the case where G is a simple Lie group with real rank one. So, G belongs to one of the series $\text{SO}_0(n, 1)$, $\text{SU}(n, 1)$, $\text{Sp}(n, 1)$ or $\text{Sp}(n, 1)$ for $n \geq 2$ or G is the exceptional Lie group $F_{4(-20)}$. We set $P := MAN$.

Theorem B. *Let $p \in (1, +\infty)$ with $p \neq 2$.*

- (i) *Let $G = \text{SO}_0(n, 1)$ for $n \geq 2$. Then $\widehat{M^{\text{Ab}}} = 1_M$ for $n \neq 3$ and $\widehat{M^{\text{Ab}}} \cong \mathbf{Z}$ for $n = 3$. The representation $\pi(P, \chi, \lambda, p)$ is irreducible for every $\lambda \in \mathfrak{a}^*$ and every $\chi \in \widehat{M^{\text{Ab}}}$.*
- (ii) *Let $G = \text{SU}(n, 1)$ for $n \geq 2$. Then $\widehat{M^{\text{Ab}}} \cong \mathbf{Z}$.*
 - (ii1) *The representation $\pi(P, \chi, \lambda, p)$ is irreducible for every $\lambda \in \mathfrak{a}^* \setminus \{0\}$ and every $\chi \in \widehat{M^{\text{Ab}}}$.*
 - (ii2) *After an appropriate identification of $\widehat{M^{\text{Ab}}}$ with \mathbf{Z} , the representation $\pi(P, m, 0, p) = \pi(P, \chi, 0, p)$ for $m \in \mathbf{Z}$ is not irreducible if and only if*

p belongs to the finite set

$$\left\{ \frac{2n}{k} : k \in \{1, \dots, 2n-1\}, k \equiv m \pmod{2}, k \neq m, k \neq 2n+m \right\}.$$

- (iii) Let $G = \mathrm{Sp}(n, 1)$ for $n \geq 2$. Then $\widehat{M^{\mathrm{Ab}}} = 1_M$.
 - (iii1) The representation $\pi(P, 1_M, \lambda, p)$ is irreducible for every $\lambda \in \mathfrak{a}^* \setminus \{0\}$
 - (iii2) The representation $\pi(P, 1_M, 0, p)$ is not irreducible if and only if $p = \frac{2n+1}{2n}$ or $p = 2n+1$.
- (iv) Let $G = \mathrm{F}_{4(-20)}$. Then $\widehat{M^{\mathrm{Ab}}} = 1_M$.
 - (iv1) The representation $\pi(P, 1_M, \lambda, p)$ is irreducible for every $\lambda \in \mathfrak{a}^* \setminus \{0\}$
 - (iv2) The representation $\pi(P, 1_M, 0, p)$ is not irreducible if and only if p belongs to the set $\left\{ \frac{11}{10}, \frac{11}{9}, \frac{11}{8}, \frac{11}{3}, \frac{11}{2}, 11 \right\}$.

Corollary C. Let G be one of the groups as in Theorem B.

- (i) Let $p \in (1, +\infty)$ with $p \neq 2$. Every completely irreducible representation of G on an L_p -space is equivalent either to the trivial representation 1_G or to one of the irreducible representations of the form $\pi(P, \chi, \lambda, p)$ from Theorem B.
- (ii) Two irreducible representations $\pi(P, \chi, \lambda, p)$ and $\pi(P, \chi', \lambda', p')$ from Theorem B with $(\chi', \lambda', p') \neq (\chi, \lambda, p)$ are equivalent if and only if

$$(\chi', \lambda', p') = (\bar{\chi}, \lambda, p) \quad \text{or} \quad (\chi', \lambda', p') = (\bar{\chi}, -\lambda, q),$$
 where q is the conjugate exponent of p .

The paper is organized as follows. The proofs of Theorem A, Proposition 3, and Theorem B are given in Sections 2, 3, and 4 respectively. As an example, the case of the group $G = \mathrm{SL}_2(\mathbf{R})$, which is a twofold cover of $\mathrm{SO}_0(2, 1)$, is treated in Section 5.

2. PROOF OF THEOREM A

Let G be a noncompact connected semisimple linear Lie group, Let (X, \mathcal{B}, μ) be a standard Borel space equipped with a σ -finite measure μ on \mathcal{B} . Let

$$\pi : G \rightarrow \mathrm{Iso}(L_p(X, \mu))$$

be a completely irreducible representation of G by linear isometries on $L_p(X, \mu)$ for $p \in (1, +\infty)$ with $p \neq 2$. Assume that π is not the trivial representation 1_G of G .

Observe that $L_p(X, \mu)$ is isometrically isomorphic to $L_p(X, \mu')$ for every σ -finite measure μ' on X which is equivalent to μ ; indeed, if $\varphi = \frac{d\mu'}{d\mu}$ denotes the Radon-Nikodym derivative of μ' with respect to μ , then

$$f \mapsto f\varphi^{1/p}$$

is a bijective linear isometry from $L_p(X, \mu')$ to $L_p(X, \mu)$. So, upon choosing a probability measure in the measure class of μ , we can and will assume in the sequel that μ is a probability measure.

• *First step.* We claim that there exists a measure-class preserving measurable action

$$G \times X \rightarrow X, \quad (g, x) \mapsto gx$$

of G on X and a measurable map $c : G \times X \rightarrow \mathbf{S}^1$ such that, for $f \in L_p(X, \mu)$, $g \in G$, and $x \in X$ we have

$$(1) \quad \pi(g)f(x) = c(g^{-1}, x) \left(\frac{dg^{-1}(\mu)}{d\mu}(x) \right)^{1/p} f(g^{-1}x),$$

where \mathbf{S}^1 is the set of complex numbers of modulus 1 and $g_*(\mu)$ is the image of μ under the map $x \mapsto gx$; moreover c satisfies the cocycle relation

$$(2) \quad c(g_1g_2, x) = c(g_1, g_2x)c(g_2, x) \quad \text{for all } g_1, g_2 \in G, \text{ for almost all } x \in X.$$

Indeed, let $g \in G$. By the Banach-Lamperti theorem (see [Ban93], [Lam58]), there exists a measure-class preserving measurable map $\varphi(g) : X \rightarrow X$ and a measurable map $c(g) : X \rightarrow \mathbf{S}^1$ such that

$$\pi(g)f(x) = c(g)(x) \left(\frac{d\varphi(g)_*(\mu)}{d\mu}(x) \right)^{1/p} f(\varphi(g)(x))$$

for $f \in L_p(X, \mu)$, $g \in G$, and $x \in X$. Since $\pi(g_1g_2) = \pi(g_1)\pi(g_2)$ for all $g_1, g_2 \in G$ and since $g \mapsto \pi(g)f$ is measurable for every $f \in L_p(X, \mu)$, it is readily checked that $(g, x) \mapsto \varphi(g^{-1})x$ is a measure-class preserving G -action on X and that

$$c : (g, x) \rightarrow c(g^{-1})x$$

is a measurable map satisfying Equation (2).

Recall that another measurable cocycle $c' : G \times X \rightarrow \mathbf{S}^1$ is said to be cohomologous to c if there exists a measurable map $b : X \rightarrow \mathbf{S}^1$ such that

$$c'(g, x) = b(gx)c(g, x)b(x)^{-1} \quad \text{for all } g \in G, \text{ for almost all } x \in X.$$

Let π' be the representation of G on $L_p(X, \mu)$ defined by the same formula (1), with c' instead of c . Then π' is equivalent to π ; indeed, the map $U : L_p(X, \mu) \rightarrow L_p(X, \mu)$, defined by

$$Uf(x) = b(x)f(x) \quad \text{for all } f \in L_p(X, \mu), x \in X,$$

is a bijective isometry which intertwines π' and π .

• *Second step.* We claim that the measure μ is continuous (that is, μ has no atoms).

Indeed, let A be the (at most countable) set of atoms of μ . The decomposition of μ into its atomic and continuous parts is $\mu = \mu_a + \mu_c$, where $\mu_a = \mu|_A$ and $\mu_c = \mu - \mu_a$. We can write

$$L_p(X, \mu) = L_p(X, \mu_c) \oplus \ell_p(A)$$

and it follows from the Banach-Lamperti theorem that $L_p(X, \mu_c)$ and $\ell_p(A)$ are $\pi(G)$ -invariant and so define subrepresentations of π . Since π is irreducible, it follows that either μ is purely atomic or μ is continuous.

Assume by contradiction that μ is purely atomic, that is, $L_p(X, \mu) = \ell_p(A)$. Let $G \curvearrowright A$ be the action of G on A and $c : G \times A \rightarrow \mathbf{S}^1$ the cocycle as above. Since π is continuous, the map

$$G \rightarrow \ell_p(A), \quad g \mapsto c(g, x)\delta_{g \cdot x}$$

is continuous for every $x \in A$. As G is connected, this implies that G fixes pointwise A . By irreducibility of π , it follows that A is a singleton $\{x_0\}$. Equation (2) shows that then that $g \mapsto c(g, x_0)$ is homomorphism from G to \mathbf{S}^1 . Since G is semisimple, G has a trivial abelianization and hence $c(g, x_0) = 1$ for all $g \in G$. So, $\pi = 1_G$; this is a contradiction and the claim is proved.

As is well-known (see [Zim84, Theorem 2.1.19]), the measurable G -system (X, μ, G) admits a continuous compact model, that is, there exists a compact metric space Y on which G acts continuously, a probability measure ν on the Borel subsets of Y and a G -equivariant Borel isomorphism $\Phi : Y \rightarrow X$ with $\Phi_*(\nu) = \mu$.

Let $c' : G \times Y \rightarrow \mathbf{S}^1$ be defined by $c'(g, y) = c(g, \Phi(y))$. The representation π' , defined on $L_p(Y, \nu)$ by the formula

$$\pi'(g)f(y) = c'(g^{-1}, y) \left(\frac{dg_*^{-1}(\nu)}{d\nu}(y) \right)^{1/p} f(g^{-1}y),$$

is equivalent to π . So, without loss of generality, we may assume that X is a compact metric space on which G acts continuously and that μ is a quasi-invariant probability measure on the Borel subsets of X .

Next, let K be a maximal compact subgroup of G and denote by dk the normalized Haar measure on K . Then $\mu' = \int_K k_*(\mu)dk$, defined by

$$\mu'(A) = \int_K \mu(kA)dk$$

for all Borel subsets A of X , is a probability measure on X which is K -invariant; moreover, μ' is equivalent to μ . So, we can and will assume in the sequel that μ is K -invariant.

• *Third step.* We claim that there are countably many K -orbits $\mathcal{O}_1, \mathcal{O}_2, \dots$, in X such that

$$\mu \left(\bigcup_{i \geq 1} \mathcal{O}_i \right) = 1.$$

Indeed, let $Y := X/K$ be the space of K -orbits, equipped with the quotient topology structure given by the quotient map $r : X \rightarrow Y$ and with the probability measure $\nu := r_*(\mu)$.

Since K is compact, Y is a Hausdorff space. So, Y is a second countable compact space and is therefore a standard Borel space. Let $\nu = \nu_a + \nu_c$ be the decomposition of ν into its atomic and continuous parts. The claim will be proved if we show that $\nu_c = 0$.

By a well-known disintegration lemma (see e.g. Lemma 11.1 [Mac52] or Theorem 5.14 in [EW11]), there exists a map

$$\theta : Y \rightarrow \text{Prob}(X), y \mapsto \theta_y,$$

where $\text{Prob}(X)$ is the set of Borel probability measures on X , with the following properties:

- (a) for every $y \in Y$, we have $\theta_y(r^{-1}(y)) = 1$;
- (b) for every $f \in L_1(X, \mu)$, the map $y \mapsto \int_X f(x)d\theta_y(x)$ is measurable and we have

$$\int_X f d\mu = \int_Y \left(\int_X f(x)d\theta_y(x) \right) d\nu(y).$$

Moreover, if a second map $\theta' : Y \rightarrow \text{Prob}(X)$ satisfies (a) and (b), then $\theta_y = \theta'_y$ for ν -almost every $y \in Y$.

Since μ is K -invariant, it follows from the uniqueness of θ that θ_y is the unique K -invariant probability measure on the K -orbit $y \subset X$, for ν -almost every y .

For $f \in C(X)$ and $y \in Y$, let $f|_y \in C(y)$ denote the restriction of f to y . The map $f \mapsto (y \mapsto f|_y)$ extends to an isometric isomorphism

$$S : L_p(X, \mu) \rightarrow \int_Y^\oplus L_p(y, \theta_y) d\nu(y),$$

where $\int_Y^\oplus L_p(y, \theta_y) d\nu(y)$ is the L_p -direct integral of the family $(L_p(y, \theta_y))_{y \in Y}$ in the sense of [dJR17]. Let π_y denote the representation of K on $L_p(y, \theta_y)$ given by

$$\pi_y(k)f(x) = c(k^{-1}, x)f(k^{-1}x) \quad \text{for all } f \in L_p(y, \theta_y), k \in K, x \in y.$$

The direct integral $\int_Y^\oplus \pi_y d\nu(y)$ of the π_y 's is a representation of K defined on $\int_Y^\oplus L_p(y, \theta_y) d\nu(y)$ and the map S intertwines $\pi|_K$ with $\int_Y^\oplus \pi_y d\nu(y)$; for this, see [dJR17, Theorem 4.9] (the proof given there extends to representations twisted by a cocycle as in our situation).

For $\delta \in \widehat{K}$ (recall that \widehat{K} is the unitary dual of K), set

$$\chi_\delta = d(\delta) \text{ch}_\delta,$$

where $\text{ch}_\delta = \text{Tr} \circ \delta$ is the usual character of δ , and define

$$P_\pi(\delta) := \pi(\overline{\chi_\delta}) = \int_K \overline{\chi_\delta}(k) \pi(k) dk.$$

Then $P_\pi(\delta)$ is a continuous projection of $L_p(X, \mu)$ onto the isotypical K -submodule of $L_p(X, \mu)$ of type δ . Similarly, for every $y \in Y$,

$$P_{\pi_y}(\delta) := \pi_y(\overline{\chi_\delta}) = \int_K \overline{\chi_\delta}(k) \pi_y(k) dk$$

is a continuous projection of $L_p(y, \theta_y)$ onto the isotypical K -submodule of $L_p(y, \theta_y)$ of type δ .

The field $y \rightarrow P_{\pi_y}(\delta)$ defines a decomposable operator $\int_Y^\oplus P_{\pi_y}(\delta) d\nu(y)$ on $\int_Y^\oplus L_p(y, \theta_y) d\nu(y)$ and, since S intertwines $\pi|_K$ and $\int_Y^\oplus \pi_y d\nu(y)$, we have

$$(3) \quad SP_\pi(\delta)S^{-1} = \int_Y^\oplus P_{\pi_y}(\delta) d\nu(y)$$

Let Ω be a Borel subset of Y . Denote by V_Ω the range of the multiplication operator on $\int_Y^\oplus L_p(y, \theta_y) d\nu(y)$ by $\mathbf{1}_\Omega$, that is,

$$\begin{aligned} V_\Omega &= \left\{ f = (f_y)_{y \in Y} \in \int_Y^\oplus L_p(y, \theta_y) d\nu(y) : \mathbf{1}_\Omega f = f \right\} \\ &= \left\{ f = (f_y)_{y \in Y} \in \int_Y^\oplus L_p(y, \theta_y) d\nu(y) : f_y = 0 \text{ for } \nu\text{-almost every } y \notin \Omega \right\}. \end{aligned}$$

Observe that

$$(4) \quad \int_Y^\oplus P_{\pi_y}(\delta) d\nu(y) \text{ leaves invariant } V_\Omega.$$

Assume, by contradiction, that the continuous part ν_c of ν is non-zero; so, denoting by Y_c the complement in Y of set of atoms of ν , we have $\nu(Y_c) > 0$ and the measure $\nu|_{Y_c}$ is continuous.

For $\delta \in \widehat{K}$, the subset

$$D_\delta := \{y \in Y : P_{\pi_y}(\delta) = 0\}$$

of Y is measurable. Indeed, let $(f_n)_n$ be a dense sequence in $C(X)$ for the uniform convergence; then, by Urysohn lemma, $(f_n|_y)_n$ is dense in $C(y)$ and hence in $L_p(y, \theta_y)$ for every $y \in Y$, and so

$$D_\delta = \bigcap_{n \in \mathbf{N}} \{y \in Y : P_{\pi_y}(\delta)(f_n|_y) = 0\}.$$

Since $y \mapsto P_{\pi_y}(\delta)(f_n|_y)$ belongs to $\int_Y^\oplus L_p(y, \theta_y) d\nu(y)$ and is therefore a measurable field, D_δ is measurable.

Observe that \widehat{K} is countable, since K is a separable compact group. So, the set

$$D := \bigcap_{\delta \in \widehat{K}} D_\delta$$

is measurable. We claim that

$$(5) \quad \nu(D) = 0.$$

Indeed, assume, by contradiction that $\nu(D) > 0$. Then $V_D \neq \{0\}$. However, for every $(f_y)_{y \in Y} \in V_D$, we have

$$P_{\pi_y}(\delta)(f_y) = 0 \quad \text{for all } \delta \in \widehat{K}$$

and hence $f_y = 0$ for every $y \in D$. This contradiction shows that (5) holds.

It follows from (5) that there exists $\delta \in \widehat{K}$ such that $\nu(Y_c \cap N_\delta) < \nu(Y_c)$. So, for

$$\Omega := Y_c \setminus N_\delta$$

we have $\nu(\Omega) > 0$. Since $\nu|_{Y_c}$ is continuous, we can therefore find a partition

$$\Omega = \coprod_{n \in \mathbf{N}} \Omega_n$$

of Ω in countably many Borel subsets Ω_n with $\nu(\Omega_n) > 0$ for all $n \in \mathbf{N}$. We have

$$\left(\int_Y^{\oplus} P_{\pi_y}(\delta) d\nu(y) \right) V_{\Omega_n} \neq \{0\} \quad \text{for all } n \in \mathbf{N}.$$

since $P_{\pi_y}(\delta) \neq 0$ for every $y \in \Omega$. Hence, for every $n \in \mathbf{N}$, we can find

$$f_n \in \left(\int_Y^{\oplus} P_{\pi_y}(\delta) d\nu(y) \right) V_{\Omega_n} \quad \text{with } f_n \neq 0.$$

Since $\Omega_n \cap \Omega_m = \emptyset$ for $n \neq m$, it follows from (4) that the family $(f_n)_{n \in \mathbf{N}}$ is linearly independent. Therefore, on the one hand, the family $(S^{-1}f_n)_{n \in \mathbf{N}}$ of elements in $L_p(X, \mu)$ is linearly independent. On the other hand, it follows from (3) that

$$S^{-1}f_n \in P_\pi(\delta)(S^{-1}(V_{\Omega_n})) \subset P_\pi(\delta)(S^{-1}(V_\Omega)) \quad \text{for all } n \in \mathbf{N}.$$

This implies that the subspace $P_\pi(\delta)(S^{-1}(V_\Omega))$ of $L_p(X, \mu)$ is infinite dimensional. So, the projection $P_\pi(\delta)$ does not have finite-dimensional range. This is not possible since π is completely irreducible (see [God52, Theorem 2] or [HC53, Lemma 33]). This is a contradiction shows that ν is an atomic measure.

• *Fourth step.* We claim that there exists a point $x_0 \in X$ such that $\mu(Gx_0) = 1$. Indeed, it follows from the third step that there exists a point $x_0 \in X$ such that $\mu(Kx_0) > 0$ and therefore $\mu(Gx_0) > 0$. If we show that the G -action of (X, μ) is ergodic, then it will follow that $\mu(Gx_0) = 1$.

Assume, by contradiction, that there exists a partition $X = X_1 \coprod X_2$ into two G -invariant Borel subsets X_i with $\mu(X_i) > 0$ for $i = 1, 2$. Then

$$L_p(X, \mu) = V_{X_1} \oplus V_{X_2},$$

where

$$V_{X_i} = \{f \in L_p(X, \mu) : \mathbf{1}_{X_i} f = f\} \cong L_p(X_i, \mu|_{X_i});$$

moreover, Formula (1) shows that V_{X_i} is $\pi(G)$ -invariant. Since $V_{X_i} \neq \{0\}$ for $i = 1, 2$, this contradicts the fact that π is irreducible and the claim is proved.

The stabilizer of $x_0 \in X$ as above is a closed subgroup H . Hence, upon disregarding a set of measure zero, we may and will identify X , as *Borel G -space*, with the space G/H , equipped with its quotient Borel structure.

Next, we draw some consequences for the cocycle $c : G \times X \rightarrow \mathbf{S}^1$ over the transitive G -space X . First, upon passing to a cohomologous cocycle, we can assume that c is a strict cocycle, that is, Equation (2) holds for all $g_1, g_2 \in G$ and $x \in X$. Moreover, choose a Borel section $s : G/H \rightarrow G$ for the projection $G \rightarrow G/H$ with $s(H) = e$. Observe that $gs(x)s(gx)^{-1}$ belongs to H , for every

$g \in G$ and $x \in X$. Then there exists a continuous homomorphism $\chi : H \rightarrow \mathbf{S}^1$ such that c is cohomologous to the cocycle $c_\chi : G \times X \rightarrow \mathbf{S}^1$ given by

$$c_\chi(g, x) = \chi(gs(x)s(gx)^{-1}) \quad \text{for all } g \in G, x \in X;$$

for all this, see Theorem 5.27 in [Var85].

We may and will in the sequel assume that the cocycle c coincides with c_χ . Also, as is well-known (see e.g. [Mac52, Theorem 1.1]), all G -quasi-invariant σ -finite Borel measures on G/H are mutually equivalent; so, we may assume that the G -quasi-invariant μ on G/H is one of the standard G -invariant Borel measures on G/H constructed in [Mac52, §.1].

- *Fifth step.* We claim that K acts transitively on the space G/H .

Indeed, on the one hand, since G is a connected Lie group, $X = G/H$ is a C^∞ -smooth connected manifold; moreover, the measure μ on X is locally Lebesgue measure, that is, μ is given by a smooth density times Lebesgue measure in any local coordinates on X ; indeed, this follows from the fact that the maps $x \mapsto gx$ are diffeomorphisms of X for every $g \in G$.

On the other hand, it follows from the third step that there exists a K -orbit \mathcal{O} in X with $\mu(\mathcal{O}) > 0$. As is well-known, \mathcal{O} is a submanifold of X , diffeomorphic to K/L where L is the stabilizer in K of a point in K (see e.g. [Bou72, III, §1, 7.]). In view of what we said about μ , this is possible only if $\dim \mathcal{O} = \dim X$, that is, if \mathcal{O} is open in X . Since X is connected, it follows that $\mathcal{O} = X$.

- *Sixth step.* We claim that H is contained as a finite index subgroup in a parabolic subgroup of G , as defined in the Introduction. Indeed, it follows from the fifth step and Lemma 3 below, that there exists a parabolic subgroup Q which contains H and which has the following properties; let $Q = M_Q A_Q N_Q$ be the Langlands decomposition of Q . Write $Q^0 = C S A_Q N_Q$ for the connected identity component Q^0 of Q , where C is the maximal compact factor of M_Q^0 and S is the product of all the non-compact simple factors of M_Q^0 . Then

$$H^0 = C' S A_Q N_Q,$$

where C' is a connected closed subgroup of C .

We claim H has finite index in Q . Assume, by contradiction, that H has infinite index in Q . Observe that Q is a direct product $Q = ZQ^0$ for a finite subgroup Z of its center (see [Kna96, Theorem 7.53]). So, H is of the form $H = T S A_Q N_Q$ for a subgroup T of infinite index in the compact group $L := ZC$.

Let χ be the unitary character of H associated to c . Observe that χ is trivial on the semisimple Lie group S . Moreover, we have $[\mathfrak{a}_F, \mathfrak{n}_F] = \mathfrak{n}_F$ for the Lie algebras \mathfrak{a}_F and \mathfrak{n}_F of A_Q and N_Q , where F is the set of simple roots associated to Q as in the Introduction; it follows that the commutator subgroup of $A_Q N_Q$ coincides with N_Q and hence that χ is also trivial on N_Q .

Choose a Borel fundamental domain $\Omega_0 \subset G$ for G/Q and a fundamental domain $\Omega_1 \subset L$ for L/T . Then $\Omega_0 \Omega_1$ is a fundamental domain for G/H and every $g \in G$ has unique decompositions

$$g = \omega_0(g)q(g) = \omega_0(g)\omega_1(g)h(g) \quad \text{for } \omega_i(g) \in \Omega_i, q(g) \in Q, h(g) \in H.$$

Let ν_1 be the L -invariant probability measure on $\Omega_1 \cong L/T$. Identify $L_p(G/H, \mu)$ with $L_p(\Omega_0 \times \Omega_1, \nu_0 \otimes \nu_1)$ for a G -quasi-invariant probability measure ν_0 on $\Omega_0 \cong G/Q$. For $g \in G$ and $F \in L_p(\Omega_0 \times \Omega_1, \nu_0 \otimes \nu_1)$, we have

$$\pi(g)F(x) = \chi(h(g^{-1}x))F(\omega_0(g^{-1}x)\omega_1(g^{-1}x)) \quad \text{for all } x \in \Omega_0 \times \Omega_1.$$

Define a representation π_Q of Q on $L_p(\Omega_1, \nu_1)$ by

$$\pi_Q(q)f(x_1) = \chi(h(q^{-1}x_1))f(\omega_1(q^{-1}x_1))$$

for $f \in L_p(\Omega_1, \nu_1)$, $x_1 \in \Omega_1$, $q \in Q$.

Since $\Omega_1 \cong Q/H$ is infinite, $L_p(\Omega_1, \nu_1)$ is infinite dimensional; therefore, the restriction $\pi_Q|_L$ of π_Q to the compact group L is not irreducible (see [War72, 4.2.2.4]). Hence, there exists a closed $\pi_Q(L)$ -invariant subspace V of $L_p(\Omega_1, \nu_1)$ with $V \neq \{0\}$ and $V \neq L_p(\Omega_1, \nu_1)$. Observe that $\pi_Q(q)$ is a multiple of $\chi(q)$ for $q \in A_Q$ and $\pi_Q(q)$ is the identity for $q \in SN_Q$. Since $Q = LSA_QN_Q$, it follows that V is Q -invariant.

For $F \in L_p(\Omega_0 \times \Omega_1, \nu_0 \otimes \nu_1)$ and $x_0 \in \Omega_0$, let $F_{x_0} \in L_p(\Omega_1, \nu_1)$ be defined by

$$F_{x_0}(x_1) = F(x_0x_1).$$

Let \tilde{V} be subspace of $L_p(\Omega_0 \times \Omega_1, \nu_0 \otimes \nu_1)$ of all $F \in L_p(\Omega_0 \times \Omega_1, \nu_0 \otimes \nu_1)$ such that $F_{x_0} \in V$ for ν_0 -almost every $x_0 \in \Omega_0$. Then \tilde{V} is a proper non trivial closed subspace of $L_p(\Omega_0 \times \Omega_1, \nu_0 \otimes \nu_1)$.

We claim that \tilde{V} is $\pi(G)$ -invariant. Indeed, let $F \in \tilde{V}$, $g \in G$ and $F' = \pi(g)F$; for $x_0 \in \Omega_0$, $x_1 \in \Omega_1$, set $x = x_0x_1$ and let $q = q(x_0^{-1}g)$ be the Q -component of $x_0^{-1}g$. We have $\omega_0(g^{-1}x) = \omega_0(g^{-1}x_0)$ and

$$\begin{aligned} F'_{x_0}(x_1) &= F'(x_0x_1) = \chi(h(g^{-1}x))F(\omega_0(g^{-1}x_0)\omega_1(g^{-1}x)) \\ &= \chi(h((x_0^{-1}g)^{-1}x_1))F_{\omega_0(g^{-1}x_0)}(\omega_1((x_0^{-1}g)^{-1}x_1)) \\ &= \chi(h(q^{-1}x_1))F_{\omega_0(g^{-1}x_0)}(\omega_1(q^{-1}x_1)) \\ &= (\pi_Q(q)F_{\omega_0(g^{-1}x_0)})(x_1) \end{aligned}$$

Since $F_{\omega_0(g^{-1}x_0)} \in V$ for almost every x_0 and since V is Q -invariant, it follows that $F' \in \tilde{V}$ and the claim is proved.

Now, π is irreducible. So, we have obtained a contradiction which shows that H has finite index in Q .

- *Seventh step.* We claim that $H = Q$. Indeed, H contains Q^0 , since H has finite index in Q . As mentioned in the sixth step, Q is a direct product $Q = ZQ^0$ for a finite abelian subgroup Z . Hence, we have $Q = Z'H$ for a subgroup Z' of Z . We have to show that Z' is trivial.

Assume, by contradiction, that $Z' \neq \{e\}$. Then there exists a non trivial character δ of Z' . We proceed as in the sixth step. Let $\Omega_0 \subset G$ a fundamental domain for G/Q . We identify $L_p(G/H, \mu)$ with $L_p(\Omega_0 \times Z', \nu_0 \otimes \nu_1)$ for a convenient G -quasi-invariant probability measure ν_0 on $\Omega_0 \cong G/Q$ and a probability measure ν_1 on Z' .

For $F \in L_p(\Omega_0 \times Z', \nu_0 \otimes \nu_1)$ and $x_0 \in \Omega_0$, let $F_{x_0} \in L_p(\Omega_1, \nu_1)$ be defined by $F_{x_0}(x_1) = F(x_0x_1)$. Let \tilde{V}_δ be subspace of $L_p(\Omega_0 \times \Omega_1, \nu_0 \otimes \nu_1)$ of all F such that $F_{x_0}(zx_1) = \delta(z)F_{x_0}(x_1)$ for all $z \in Z'$ and ν_0 -almost every $x_0 \in \Omega_0$. Then \tilde{V}_δ is a proper non trivial closed subspace of $L_p(\Omega_0 \times \Omega_1, \nu_0 \otimes \nu_1)$. Moreover, one checks that \tilde{V}_δ is $\pi(G)$ -invariant; this contradicts the irreducibility of π .

- *Eighth step.* There exists a real linear form $\lambda \in \mathfrak{a}^*$ and a unitary character χ of M_Q such that π is equivalent to the representation $\pi(Q, \chi, \lambda, p)$ on $L_p(G/Q, \mu_Q)$ described in Theorem A.

Indeed, by the seventh step, we have $H = Q = M_Q A_Q N_Q$. The unitary character of Q associated to the cocycle c is trivial on N_Q and M_Q centralizes A_Q ; hence, this character is of the form $\chi \otimes \chi'$ for unitary characters χ of M_Q and χ' of A_Q . Let $F \subset \Sigma$ be the set of simple positive roots associated to Q . There exists $\lambda \in \mathfrak{a}^*$ such that

$$\chi'(\exp X) = e^{-i\lambda(X)} \quad \text{for all } X \in \mathfrak{a}_F = \log(A_Q).$$

Let μ be the unique quasi-invariant Borel probability measure on G/Q which is K -invariant. For $g \in G$, decompose g according to $G = KM_QA_QN_Q$ as

$$g = \kappa(g)\mu(g)\exp(H(g))n(g) \quad \text{for } \kappa(g) \in \Omega_Q, \mu(g) \in M_Q, H(g) \in \mathfrak{a}_F, n(g) \in N_Q.$$

The Radon-Nikodym derivative of $g_*\mu$ with respect to μ is given by (see e.g. [Kna96, Proposition 8.44])

$$\frac{dg_*(\mu)}{d\mu}(x) = e^{-\rho_F(H(g))}$$

for

$$\rho_F = \sum_{\alpha \in \Sigma^+ \setminus \text{span}(\mathbb{F})} \dim(\mathfrak{g}_\alpha)\alpha.$$

This finishes the proof of Theorem A.

The following lemma, which was used in the proof above, is a consequence of the description of cocompact subgroups of semisimple Lie groups from [Wit90] (see also [GW72]).

Lemma 3. *Let G be a noncompact connected semisimple linear Lie group with maximal compact subgroup K . Let H be a closed subgroup of G with the property that the natural action of K on G/H is transitive. Upon replacing H by one of its conjugate, there exists a standard parabolic subgroup Q which contains H and which has the following properties; denote by $Q = M_QA_QN_Q$ the Langlands decomposition of Q ; write $Q^0 = CSA_QN_Q$ for the identity component Q^0 of Q , where C is the maximal compact factor of M_Q^0 and S is the product of all the non-compact simple factors of M_Q^0 . Then there exists a connected closed subgroup C' of C such that*

$$H^0 = C'SA_QN_Q.$$

Proof. Since K acts transitively on G/H , we have $G = KH$. In particular, H is a cocompact subgroup of G . By [Wit90, Theorem 1.2] and upon conjugating H , there exists a standard parabolic subgroup Q of G with the following properties:

- $Q = M_QA_QN_Q$ contains H ;
- there exist a connected, normal subgroup X of S and a connected closed subgroup Y of CA_Q such that $H^0 = YXN_Q$, where S and C are as in the statement of the Lemma.

Since H/H^0 is at most countable, we have

$$G = \bigcup_{n \geq 1} KH_0g_n$$

for some sequence $(g_n)_{n \geq 1}$ of elements in G . As K is compact and H_0 is closed, KH_0g_n is a closed subset of G for every n . Hence, by Baire's category theorem, KH_0 has a non empty interior. By homogeneity, it follows that KH_0 is open in G . Since G is connected, we have therefore $G = KH_0$, that is,

$$(6) \quad G = KYXN_Q.$$

Recall that the multiplication map $K \times A \times N \rightarrow G$ from the Iwasawa decomposition $G = KAN$ is a diffeomorphism. The reductive Lie group M_Q has also an Iwasawa decomposition

$$M_Q = K_1A_1N_1,$$

for $K_1 := K \cap M_Q$; we have $A = A_1A_Q$ as a direct product and $N = N_1N_Q$ as semi-direct product (see [Kna96, Proposition 7.82]). In particular, we have $C \subset K_1$. Let $\Omega \subset K$ be a fundamental domain for K/K_1 . So,

$$(7) \quad \text{the product map } \Omega \times K_1 \times A_1 \times A_Q \times N_1 \times N_Q \rightarrow G \quad \text{is a bijection.}$$

The Iwasawa decomposition of the semisimple Lie group $X \subset S$ is $X = K_2 A_2 N_2$, where $K_2 := K_1 \cap X$ and where A_2 and N_2 are subgroups of A_1 and N_2 . By (6), we have therefore

$$G = KYXN_Q = KK_2YA_2N_2N_Q.$$

Since $Y \subset CA_Q$, it follows from (7) that

$$\dim A_2 = \dim A_1, \quad \dim N_2 = \dim N_1, \quad \text{and} \quad A_Q \subset Y.$$

So, we have $S = X$ and $Y = C'A_Q$ for a connected closed subgroup C' of C . \square

3. PROOF OF PROPOSITION 2

Let $Q = M_Q A_Q N_Q$ be a parabolic subgroup of G ; for a real linear form $\lambda \in \mathfrak{a}^*$ and a unitary character $\chi : M_Q \rightarrow \mathbf{S}^1$, consider the representation $\pi(Q, \lambda, \chi, p)$ of G on $L_p(G/Q, \mu)$ as in the Introduction.

We may realize $\pi(Q, \lambda, \chi, p)$ in the so-called compact picture as follows. Let $C(K, \chi)$ be the vector space of continuous functions $F : K \rightarrow \mathbf{C}$ such that

$$F(xk) = \overline{\chi(k)}F(x) \quad \text{for all } k \in K \cap M_Q, x \in K;$$

equip $C(K, \chi)$ with the norm

$$\|F\|_p = \left(\int_{K/(K \cap M_Q)} |F(x)|^p d\mu(x) \right)^{1/p},$$

where μ is the unique K -invariant probability measure on $K/(K \cap M_Q)$. Let $L_p(K, \chi)$ be the completion of $(C(K, \chi), \|\cdot\|_p)$. Let $\rho_{Q,p} \in \mathfrak{a}^*$ be defined as in the Introduction. Then $\pi(Q, \lambda, \chi, p)$ acts on $L_p(K, \chi)$ through

$$\pi(Q, \lambda, \chi, p)(g)F(x) = \overline{\chi(\mu(g^{-1}x))} e^{-(i\lambda + \rho_{Q,p})H(g^{-1}x)} F(\kappa(g^{-1}x))$$

for $F \in L_p(K, \chi), x \in K$ and $g \in G$ with decomposition

$$g = \kappa(g)\mu(g) \exp(H(g))n(g), \quad \kappa(g) \in K, \mu(g) \in M_Q, H(g) \in \log(A_Q), n(g) \in N_Q$$

Observe that, for $g \in K$, the operator $\pi(Q, \lambda, \chi, p)(g)$ is simply left translation on $L_p(K, \mu)$ by g^{-1} .

Set $E := L_p(K, \chi)$. Recall that the subspace E_K of K -finite vectors in E is the space of functions $F \in E$ such that

$$\pi(Q, \lambda, \chi, p)(K)F = \{\pi(Q, \lambda, \chi, p)(g)F : g \in K\}$$

spans a finite-dimensional subspace of E . The space E^∞ of C^∞ -vectors in E consists of smooth functions $F \in C(K, \chi)$. As a K -module, E^∞ is the representation of K differentiably induced by $\chi|_{K \cap M_Q}$ (in the sense of [War72, 5.3.1]). Therefore, by Frobenius reciprocity (which remains valid in this context), the multiplicity of $\delta \in \widehat{K}$ in E^∞ is equal to the multiplicity of χ in $\delta|_{K \cap M_Q}$ and is therefore finite, since δ is finite dimensional. Hence, the multiplicity of δ in $E_K \cap E^\infty$ is finite. Since $E_K \cap E^\infty$ is dense in E (see e.g. [War72, Theorem 4.4.5.16]), it follows that the multiplicity of δ in E is finite. In other words, $\pi(Q, \lambda, \chi, p)$ is an *admissible* representation of G (as defined [Kna01, p.207] and [Wal71, 3.3.5]) or is a *K-finite* representation (as defined in [War72, 4.5.1]).

As is well-known, E_K is a module over the universal enveloping algebra $\mathcal{U}(\mathfrak{g}_{\mathbf{C}})$ of the complexification of the Lie algebra \mathfrak{g} of G . Set

$$(8) \quad \nu := i\lambda + \rho_{Q,p} - \rho_{Q,2} = i\lambda + \left(\frac{2}{p} - 1 \right) \rho_{Q,2};$$

then E_K coincides with the (\mathfrak{g}, K) -module associated to the (nonunitary) induced representation $\text{Ind}_Q^G(\chi \otimes \nu \otimes 1_{N_Q})$. This (\mathfrak{g}, K) -module consists of the functions $f : G \rightarrow \mathbf{C}$ which are C^∞ and have the following properties:

- $f(gman) = \overline{\chi(m)} e^{-(i\lambda + \nu + \rho_{Q,2})(\log a)} f(g)$ for all $g \in G$ and $man \in Q$;
- the linear span of $K \cdot f$ is finite dimensional space, where $K \cdot f$ is the space of translates $g \mapsto f(k^{-1}g)$ for $k \in K$.

(for this, see [Wal71, 5.2.1]). So, the (\mathfrak{g}, K) module associated to $\pi(Q, \chi, \lambda, p)$ is the principal series representation denoted $I_{Q, \chi, \nu}$ in [Wal71] for ν as in (8).

Now, since $\pi(Q, \chi, \lambda, p)$ is admissible, $\pi(Q, \chi, \lambda, p)$ is completely irreducible if and only if its associated (\mathfrak{g}, K) -module $I_{Q, \chi, \nu}$ is algebraically irreducible. Moreover, two such representations $\pi(Q_1, \chi_1, \lambda_1, p_1)$ and $\pi(Q_2, \chi_2, \lambda_2, p_2)$ are Naimark equivalent if and only if the (\mathfrak{g}, K) -modules I_{Q_1, χ_1, ν_1} and I_{Q_2, χ_2, ν_2} are algebraically equivalent (for all this, see Theorems 4.5.5.4 and 4.5.5.2 in [War72]).

4. PROOF OF THEOREM B

We assume from now on that G is a simple Lie group with real rank one. In this case, we have $\dim \mathfrak{a} = 1$.

Choose $\alpha \in \Sigma^+$ and $H \in \mathfrak{a}$ such that $\frac{1}{2}\alpha \notin \Sigma$ and $\alpha(H) = 1$. We identify $\mathfrak{a}_{\mathbf{C}}^*$ with \mathbf{C} by means of the map $\lambda \mapsto \lambda(H)$.

There is, up to conjugation, only one parabolic subgroup, namely $P := MAN$. For $p \in (1 + \infty)$, we will write ρ_p instead of $\rho_{P,p}$. In view of Proposition 2, we have to study, given $\lambda \in \mathbf{R}$ and $\chi \in \widehat{M^{\text{Ab}}}$, the irreducibility of the (non unitary) principal series representation $I_{P, \chi, \nu_{\lambda, p}}$ for

$$\nu_{\lambda, p} = i\lambda + \left(\frac{2}{p} - 1\right) \rho_2$$

and to decide when two such representations are equivalent. We will apply several times the following result of Kostant about the case where $\nu = 1_M$. Set

$$t_\alpha := \begin{cases} (\dim \mathfrak{g}_\alpha)/2 & \text{if } 2\alpha \notin \Sigma \\ (\dim \mathfrak{g}_\alpha)/2 + 1 & \text{if } 2\alpha \in \Sigma, \end{cases}$$

where $\mathfrak{g}_\alpha \subset \mathfrak{g}$ is the usual root space corresponding to α ; set also

$$n_\alpha := \begin{cases} 1 & \text{if } 2\alpha \notin \Sigma \\ 2 & \text{if } 2\alpha \in \Sigma. \end{cases}$$

Recall that

$$\rho_2 = (\dim \mathfrak{g}_\alpha)/2 + \dim \mathfrak{g}_{2\alpha}.$$

For $\nu \in \mathbf{C}$, we have (see Theorem 2 in [Kos69] and Theorem 2.9.8 in [Kos75]):

$$(9) \quad I_{P, 1_M, \nu} \text{ is not irreducible} \iff \nu \notin (-t_\alpha, t_\alpha) \text{ and } \nu + \rho_2 \in n_\alpha \mathbf{Z}.$$

We will treat separately the different simple groups G which may occur; for the data we will use concerning these groups, we refer to [Hel78, Chap. X].

• Let $G = \text{SO}_0(n, 1)$ for $n \geq 2$. Here, $K \cong \text{SO}(n)$ and $M \cong \text{SO}(n-1)$. Moreover, we have $\dim \mathfrak{g}_\alpha = n-1$ and $2\alpha \notin \Sigma$. So,

$$t_\alpha = \frac{n-1}{2}, \quad n_\alpha = 1, \quad \text{and} \quad \rho_2 = \frac{n-1}{2};$$

hence, for $\lambda \in \mathbf{R}$, we have

$$\nu_{\lambda, p} = i\lambda + \left(\frac{2}{p} - 1\right) \frac{n-1}{2}.$$

Since $\left| \frac{2}{p} - 1 \right| < 1$, it follows from (9) that

$I_{P,1M,\nu_{\lambda,p}}$ is irreducible for all $\lambda \in \mathbf{R}$ and $p \in (1, +\infty)$ with $p \neq 2$.

The case where $n \neq 3$ is settled since the abelianization of M is then trivial and so $\widehat{M^{\text{Ab}}} = 1_M$.

Assume now that $n = 3$. Observe that in this case $G \cong \text{SL}_2(\mathbf{C})$ and $M \cong \text{U}(1)$. We use here the results from [Thi73, §. 13, B]: identifying $\widehat{M^{\text{Ab}}}$ with $\frac{1}{2}\mathbf{Z}$, we have, for $m \in \widehat{M^{\text{Ab}}}$ and $\nu \in \mathbf{C}$,

$$I_{P,m,\nu} \text{ is irreducible} \iff (\nu + l + 1)(\nu - l - 1) \neq 0$$

for all $l \in \{|m| + k : k = 0, 1, 2, \dots\}$. Since $(n - 1)/2 = 1$, this condition is clearly satisfied for $\nu_{\lambda,p}$ and so we obtain that $I_{P,\chi,\nu_{\lambda,p}}$ is irreducible for all $\lambda \in \mathbf{R}$, $\chi \in \widehat{M^{\text{Ab}}}$, and $p \in (1, +\infty)$ with $p \neq 2$.

• Let $G = \text{SU}(n, 1)$ for $n \geq 2$. Here, $K \cong \text{SU}(n)$ and $M \cong \text{U}(n - 1)$. Moreover, we have $\dim \mathfrak{g}_\alpha = 2(n - 1)$ and $\dim \mathfrak{g}_{2\alpha} = 1$. So,

$$t_\alpha = n, \quad n_\alpha = 2, \quad \text{and} \quad \rho_2 = n;$$

hence, for $\lambda \in \mathbf{R}$, we have

$$\nu_{\lambda,p} = i\lambda + \left(\frac{2}{p} - 1 \right) n.$$

Since $\left| \frac{2}{p} - 1 \right| < 1$, it follows from (9) that

$I_{P,1M,\nu_{\lambda,p}}$ is irreducible for all $\lambda \in \mathbf{R}$ and $p \in (1, +\infty)$ with $p \neq 2$.

In [Kra76, Proposition 1] precise necessary and sufficient conditions are given on $\sigma \in \widehat{M}$ and $\nu \in \mathbf{C}$ for $I_{P,\sigma,\nu}$ to be irreducible. In the special case of $\sigma \in \widehat{M^{\text{Ab}}}$, this criterion reads as follows. Identify $\widehat{M^{\text{Ab}}}$ with $\frac{1}{n+1}\mathbf{Z}$ as in [Kra76] and for $\sigma \in \widehat{M^{\text{Ab}}}$ and $\nu \in \mathbf{C}$, define $(s_1, \dots, s_{n+1}) \in \mathbf{C}^{n+1}$ by

$$s_i = \begin{cases} \frac{1}{2}(\nu - (n - 1)\sigma) & \text{if } i = 1 \\ \sigma + \frac{n}{2} - i + 1 & \text{if } 2 \leq i \leq n \\ -\frac{1}{2}(\nu + (n - 1)\sigma) & \text{if } i = n + 1. \end{cases}$$

Then $I_{P,m,\nu}$ is not irreducible if and only if either $s_1 - s_i \in \mathbf{Z} \setminus \{0\}$ for all $2 \leq i \leq n$ or $s_{n+1} - s_i \in \mathbf{Z} \setminus \{0\}$ for all $2 \leq i \leq n$.

It follows immediately that $I_{P,\sigma,\nu_{\lambda,p}}$ is irreducible if $\lambda \neq 0$. So, we may assume that $\lambda = 0$.

Writing $\sigma = \frac{m}{n+1}$ for $m \in \mathbf{Z}$, the criterion above implies that $I_{P,\sigma,\nu_{0,p}}$ is not irreducible if and only if either

$$\frac{1}{2} \left(\nu_{p,0} - (n - 1) \frac{m}{n+1} \right) - \frac{m}{n+1} + \frac{n}{2} \in \mathbf{Z} \setminus \{0\}$$

or

$$-\frac{1}{2} \left(\nu_{p,0} + (n - 1) \frac{m}{n+1} \right) - \frac{m}{n+1} + \frac{n}{2} \in \mathbf{Z} \setminus \{0\}$$

that is, if and only if

$$\text{either } \nu_{p,0} - (m + n) \in 2\mathbf{Z} \setminus \{0\} \quad \text{or} \quad \nu_{p,0} - (m + n) \in 2\mathbf{Z} \setminus \{0\}$$

As $\nu_{\lambda,0} = \left(\frac{2}{p} - 1\right)n$, we see that $I_{P,\sigma,\nu_0,p}$ is not irreducible if only if

$$(10) \quad \text{either } 2\left(\frac{1}{p} - 1\right)n - m \in 2\mathbf{Z} \setminus \{0\} \quad \text{or} \quad \frac{2n}{p} - m \in 2\mathbf{Z} \setminus \{0\}.$$

It is clear that (10) is equivalent to

$$p \in \left\{ \frac{2n}{k} : k \in \{1, \dots, 2n-1\}, k \equiv m \pmod{2}, k \neq m, k \neq 2n+m \right\}.$$

• Let $G = \mathrm{Sp}(n, 1)$ for $n \geq 2$. Here, $K \cong \mathrm{Sp}(n) \times \mathrm{Sp}(1)$ and $M \cong \mathrm{Sp}(n-1) \times \mathrm{Sp}(1)$. Moreover, we have $\dim \mathfrak{g}_\alpha = 4(n-1)$ and $\dim \mathfrak{g}_{2\alpha} = 3$. So,

$$t_\alpha = 2n-1, \quad n_\alpha = 2, \quad \text{and} \quad \rho_2 = 2n+1;$$

hence, for $\lambda \in \mathbf{R}$, we have

$$\nu_{\lambda,p} = i\lambda + \left(\frac{2}{p} - 1\right)(2n+1).$$

Observe that the abelianization of M is trivial and so $\widehat{M^{\mathrm{Ab}}} = \{1_M\}$. Moreover, we have

$$\left| \frac{2}{p} - 1 \right| (2n+1) \geq 2n-1 \iff p \in (1, \frac{2n+1}{2n}] \cup [2n+1, +\infty).$$

and

$$\left(\frac{2}{p} - 1\right)(2n+1) + 2n-1 \in 2\mathbf{Z} \iff p \in \left\{ \frac{2n+1}{k} : k \in \mathbf{N}^* \right\}.$$

It follows from (9) that

- $I_{P,1_M,\nu_{\lambda,p}}$ is irreducible for every $\lambda \in \mathfrak{a}^* \setminus \{0\}$
- $I_{P,1_M,\nu_0,p}$ is not irreducible if and only if $p = \frac{2n+1}{2n}$ or $p = 2n+1$.

• Let $G = \mathrm{F}_{4(-20)}$. Here, $K \cong \mathrm{Spin}(9)$ and $M \cong \mathrm{Spin}(7)$. Moreover, we have $\dim \mathfrak{g}_\alpha = 8$ and $\dim \mathfrak{g}_{2\alpha} = 7$. So,

$$t_\alpha = 5, \quad n_\alpha = 2, \quad \text{and} \quad \rho_2 = 11;$$

hence, for $\lambda \in \mathbf{R}$, we have

$$\nu_{\lambda,p} = i\lambda + 11\left(\frac{2}{p} - 1\right).$$

The abelianization of M is trivial and so $\widehat{M^{\mathrm{Ab}}} = \{1_M\}$. Moreover, we have

$$11\left| \frac{2}{p} - 1 \right| \geq 5 \iff p \in (1, \frac{11}{8}] \cup [\frac{11}{3}, +\infty).$$

and

$$11\left(\frac{2}{p} - 1\right) + 5 \in 2\mathbf{Z} \iff p \in \left\{ \frac{11}{k} : k \in \mathbf{N}^* \right\}.$$

It follows from (9) that

- $I_{P,1_M,\nu_{\lambda,p}}$ is irreducible for every $\lambda \in \mathfrak{a}^* \setminus \{0\}$
- $I_{P,1_M,\nu_0,p}$ is not irreducible if and only if $p \in \left\{ \frac{11}{10}, \frac{11}{9}, \frac{11}{8}, \frac{11}{3}, \frac{11}{2}, 11 \right\}$.

In all the cases above, if two representations $I_{P,\chi,\nu}$ and $I_{P,\chi',\nu'}$ are irreducible, then they are equivalent if and only if

$$(\chi', \nu) = (\bar{\chi}, \nu) \quad \text{or} \quad (\chi', \nu') = (\bar{\chi}, -\nu).$$

Now,

$$-\nu_{\lambda,p} = -(i\lambda + \left(\frac{2}{p} - 1\right)\rho_2) = (-i\lambda + \left(\frac{2}{q} - 1\right)\rho_2) = \nu_{\lambda,q}.$$

Therefore, $I_{P,\chi',\nu_{\lambda',p'}}$ and $I_{P,\chi,\nu_{\lambda,p}}$ are equivalent if and only if

$$(\chi', \lambda', p') = (\bar{\chi}, \lambda, p) \quad \text{or} \quad (\chi', \lambda', p') = (\bar{\chi}, -\lambda, q).$$

5. AN EXAMPLE: $SL_2(\mathbf{R})$

Let $G = SL_2(\mathbf{R})$, with maximal compact subgroup $K = SO(2)$. The standard minimal parabolic subgroup is

$$P = \left\{ \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} : a \in \mathbf{R}, a \neq 0, b \in \mathbf{R} \right\}.$$

We have $P = MAN$ for

$$A = \left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} : a \in \mathbf{R}, a > 0 \right\},$$

$$M = \left\{ \pm \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right\},$$

and

$$N = \left\{ \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} : b \in \mathbf{R} \right\}.$$

Identifying $\mathfrak{a}_{\mathbf{C}}^*$ with \mathbf{C} as above, we have

$$\rho_{P,p} = \frac{1}{p}$$

for every $p \in (1, \infty)$. We identify G/P as G -space with the real projective line $\mathbf{P}(\mathbf{R})$, with G acting by Möbius transformations on this latter space. The Lebesgue measure μ is the unique K -invariant probability Borel measure on $\mathbf{P}(\mathbf{R})$. The measurable space G/P can further be identified with $\mathbf{R} = \mathbf{P}(\mathbf{R}) \setminus \{\infty\}$.

The Radon-Nikodym derivative for $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ acting on \mathbf{R} is given by

$$\frac{dg_*(\mu)}{d\mu}(x) = \frac{1}{(cx + d)^2}.$$

Let ε denote the non trivial character of M and let $\lambda \in \mathbf{R}$. The representation $\pi(P, \chi, \lambda, p)$ of G by isometries on $L_p(\mathbf{R}, \mu)$ is defined by

$$\pi(P, 1_M, \lambda, p)(g)f(x) = |cx + d|^{-i\lambda - \frac{2}{p}} f\left(\frac{ax + b}{cx + d}\right) \quad \text{if} \quad g^{-1} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

and

$$\pi(P, \varepsilon, \lambda, p)(g)f(x) = \text{sgn}(cx + d)|cx + d|^{-i\lambda - \frac{2}{p}} f\left(\frac{ax + b}{cx + d}\right) \quad \text{if} \quad g^{-1} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

The Harish-Chandra (\mathfrak{g}, K) -module underlying $\pi(P, \varepsilon, \lambda, p)$ is $I_{P,\chi,\nu}$ for

$$\nu = i\lambda + \delta_p \rho_{P,2} = i\lambda + \frac{2}{p} - 1$$

Assume that $p \neq 2$. It is well-known that $I_{P,\chi,\nu}$ is irreducible (see e.g. Proposition 1.3.3 in [Vog81]); observe that the irreducibility of $I_{P,\varepsilon,\nu}$ depends on the fact that

$\frac{2}{p} - 1$ is not an integer). Moreover, the modules $I_{P,\chi,\nu}$ are pairwise non equivalent for fixed p .

In summary, the L_p -dual space of $G = \mathrm{SL}_2(\mathbf{R})$ for $p \in (1, +\infty)$ with $p \neq 2$ is

$$\widehat{G}_{L_p} = \{\pi(P, 1_M, \lambda, p) : \lambda \in \mathbf{R}\} \cup \{\pi(P, \varepsilon, \lambda, p) : \lambda \in \mathbf{R}\} \cup \{1_G\}.$$

REFERENCES

- [Ban93] Stefan Banach, *Théorie des opérations linéaires*, Éditions Jacques Gabay, Sceaux, 1993 (French). Reprint of the 1932 original. MR1357166 ↑5
- [Bou72] N. Bourbaki, *Éléments de mathématique. Fasc. XXXVII. Groupes et algèbres de Lie. Chapitre II: Algèbres de Lie libres. Chapitre III: Groupes de Lie*, Actualités Scientifiques et Industrielles [Current Scientific and Industrial Topics], No. 1349, Hermann, Paris, 1972. MR573068 ↑9
- [Bru56] François Bruhat, *Sur les représentations induites des groupes de Lie*, Bull. Soc. Math. France **84** (1956), 97–205 (French). MR0084713 ↑3
- [dJR17] Marcel de Jeu and Jan Rozendaal, *Disintegration of positive isometric group representations on L^p -spaces*, Positivity **21** (2017), no. 2, 673–710, DOI 10.1007/s11117-017-0499-4. MR3656017 ↑6, 7
- [EW11] Manfred Einsiedler and Thomas Ward, *Ergodic theory with a view towards number theory*, Graduate Texts in Mathematics, vol. 259, Springer-Verlag London, Ltd., London, 2011. MR2723325 ↑6
- [Fel65] J. M. G. Fell, *The dual spaces of Banach algebras*, Trans. Amer. Math. Soc. **114** (1965), 227–250, DOI 10.2307/1993999. MR0172131 ↑1
- [GN57] I. M. Gelfand and M. A. Neumark, *Unitäre Darstellungen der klassischen Gruppen*, Akademie-Verlag, Berlin, 1957 (German). MR0085262 ↑3
- [God52] Roger Godement, *A theory of spherical functions. I*, Trans. Amer. Math. Soc. **73** (1952), 496–556, DOI 10.2307/1990805. MR0052444 ↑1, 8
- [GW72] Morikuni Goto and Hsien-chung Wang, *Non-discrete uniform subgroups of semisimple Lie groups*, Math. Ann. **198** (1972), 259–286, DOI 10.1007/BF01419560. MR354934 ↑11
- [HC53] Harish-Chandra, *Representations of a semisimple Lie group on a Banach space. I*, Trans. Amer. Math. Soc. **75** (1953), 185–243, DOI 10.2307/1990730. MR0056610 ↑1, 8
- [Hel78] Sigurdur Helgason, *Differential geometry, Lie groups, and symmetric spaces*, Pure and Applied Mathematics, vol. 80, Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1978. MR514561 ↑13
- [Kna01] Anthony W. Knaapp, *Representation theory of semisimple groups*, Princeton Landmarks in Mathematics, Princeton University Press, Princeton, NJ, 2001. An overview based on examples; Reprint of the 1986 original. MR1880691 ↑2, 12
- [Kna96] ———, *Lie groups beyond an introduction*, Progress in Mathematics, vol. 140, Birkhäuser Boston, Inc., Boston, MA, 1996. MR1399083 ↑9, 11
- [Kos69] Bertram Kostant, *On the existence and irreducibility of certain series of representations*, Bull. Amer. Math. Soc. **75** (1969), 627–642, DOI 10.1090/S0002-9904-1969-12235-4. MR0245725 ↑3, 13
- [Kos75] ———, *On the existence and irreducibility of certain series of representations*, Lie groups and their representations (Proc. Summer School, Bolyai János Math. Soc., Budapest, 1971), Halsted Press, New York-Toronto, Ont., 1975, pp. 231–329. MR0399361 ↑13
- [Kra76] Hrvoje Kraljević, *On representations of the group $SU(n, 1)$* , Trans. Amer. Math. Soc. **221** (1976), no. 2, 433–448, DOI 10.2307/1997409. MR409725 ↑14
- [Lam58] John Lamperti, *On the isometries of certain function-spaces*, Pacific J. Math. **8** (1958), 459–466. MR0105017 ↑5
- [LW73] James Lepowsky and Nolan R. Wallach, *Finite- and infinite-dimensional representation of linear semisimple groups*, Trans. Amer. Math. Soc. **184** (1973), 223–246, DOI 10.2307/1996409. MR0327978 ↑3
- [Mac52] George W. Mackey, *Induced representations of locally compact groups. I*, Ann. of Math. (2) **55** (1952), 101–139, DOI 10.2307/1969423. MR0044536 ↑6, 9
- [PRRV67] K. R. Parthasarathy, R. Ranga Rao, and V. S. Varadarajan, *Representations of complex semi-simple Lie groups and Lie algebras*, Ann. of Math. (2) **85** (1967), 383–429, DOI 10.2307/1970351. MR0225936 ↑3
- [Thi73] Ernest Thieleker, *On the quasi-simple irreducible representations of the Lorentz groups*, Trans. Amer. Math. Soc. **179** (1973), 465–505, DOI 10.2307/1996515. MR0325856 ↑14
- [Var85] V. S. Varadarajan, *Geometry of quantum theory*, 2nd ed., Springer-Verlag, New York, 1985. MR0805158 ↑9

- [Vog81] David A. Vogan Jr., *Representations of real reductive Lie groups*, Progress in Mathematics, vol. 15, Birkhäuser, Boston, MA, 1981. MR0632407 ↑16
- [Wal71] Nolan R. Wallach, *Cyclic vectors and irreducibility for principal series representations*, Trans. Amer. Math. Soc. **158** (1971), 107–113, DOI 10.2307/1995774. MR0281844 ↑3, 12, 13
- [Wal88] ———, *Real reductive groups. I*, Pure and Applied Mathematics, vol. 132, Academic Press, Inc., Boston, MA, 1988. MR0929683 ↑3
- [War72] Garth Warner, *Harmonic analysis on semi-simple Lie groups. I*, Die Grundlehren der mathematischen Wissenschaften, vol. Band 188, Springer-Verlag, New York-Heidelberg, 1972. MR0498999 ↑1, 2, 10, 12, 13
- [Wit90] Dave Witte, *Cocompact subgroups of semisimple Lie groups*, Lie algebra and related topics (Madison, WI, 1988), Contemp. Math., vol. 110, Amer. Math. Soc., Providence, RI, 1990, pp. 309–313, DOI 10.1090/conm/110/1079114. MR1079114 ↑11
- [Zim84] Robert J. Zimmer, *Ergodic theory and semisimple groups*, Monographs in Mathematics, vol. 81, Birkhäuser Verlag, Basel, 1984. MR0776417 ↑5

BACHIR BEKKA, UNIV RENNES, CNRS, IRMAR-UMR 6625, CAMPUS BEAULIEU, F-35042
RENNES CEDEX, FRANCE

Email address: bachir.bekka@univ-rennes1.fr