

Supplementary material for: Sketched over-parametrized projected gradient descent for sparse spike estimation

Pierre-Jean Bénard^{1,*}, Yann Traonmilin¹, Jean-François Aujol¹

I. SUPPLEMENTARY MATERIAL

A. Proof of Lemma 1

Proof. Let $\theta^* \in \arg \min_{\theta \in \Theta} g_{\mathcal{S}_d}(\theta)$. We have, for any $\theta \in \Theta$

$$\sqrt{g_{\mathcal{S}}(\theta^*)} = \|\mathcal{S}A^{id}\phi(\theta^*) - \mathcal{S}A^{id}\phi(\theta_0)\|_2 \quad (1)$$

$$\begin{aligned} &= \|\mathcal{S}A^{id}\phi(\theta^*) - \mathcal{S}_dA\phi(\theta_0) \\ &\quad + \mathcal{S}_dA\phi(\theta_0) - \mathcal{S}A^{id}\phi(\theta_0)\|_2 \end{aligned} \quad (2)$$

With the triangle inequality, we obtain

$$\begin{aligned} \sqrt{g_{\mathcal{S}}(\theta^*)} &\leq \|\mathcal{S}A^{id}\phi(\theta^*) - \mathcal{S}_dA\phi(\theta_0)\|_2 \\ &\quad + \|\mathcal{S}_dA\phi(\theta_0) - \mathcal{S}A^{id}\phi(\theta_0)\|_2 \end{aligned} \quad (3)$$

$$= \sqrt{g_{\mathcal{S}_d}(\theta^*)} + \|(\mathcal{S} - \mathcal{S}_dT_{\Gamma})A^{id}\phi(\theta_0)\|_2 \quad (4)$$

$$\leq \sqrt{g_{\mathcal{S}_d}(\theta^*)} + \|\mathcal{S} - \mathcal{S}_dT_{\Gamma}\|_{\text{op}}^{A^{id}, \Sigma} \|y\|_2 \quad (5)$$

Let $\tilde{\theta} \in \arg \min_{\theta \in \Theta} g_{\mathcal{S}}(\theta)$, we have, similarly

$$\sqrt{g_{\mathcal{S}_d}(\theta^*)} \leq \sqrt{g_{\mathcal{S}_d}(\tilde{\theta})} \quad (6)$$

$$= \|\mathcal{S}A^{id}\phi(\tilde{\theta}) - \mathcal{S}_dA\phi(\theta_0)\|_2 \quad (7)$$

$$\leq \|\mathcal{S}A^{id}\phi(\tilde{\theta}) - \mathcal{S}A^{id}\phi(\theta_0)\|_2 \quad (8)$$

$$+ \|\mathcal{S}A^{id}\phi(\theta_0) - \mathcal{S}_dA\phi(\theta_0)\|_2 \quad (9)$$

$$\leq \sqrt{g_{\mathcal{S}}(\tilde{\theta})} + \|\mathcal{S} - \mathcal{S}_dT_{\Gamma}\|_{\text{op}}^{A^{id}, \Sigma} \|y\|_2 \quad (9)$$

We deduce from the two inequalities (5) and (9) that

$$\sqrt{g_{\mathcal{S}}(\theta^*)} \leq \sqrt{g_{\mathcal{S}}(\tilde{\theta})} + 2\|\mathcal{S} - \mathcal{S}_dT_{\Gamma}\|_{\text{op}}^{A^{id}, \Sigma} \|y\|_2 \quad (10)$$

□

B. Proof of Lemma 2

Proof. Let

$$\alpha_{i,k}(t) = \frac{f_k(z)}{2\pi\sigma_x\sigma_y} \times e^{-\left(\frac{(x_i-x)^2}{2\sigma_x^2} + \frac{(y_i-y)^2}{2\sigma_y^2}\right)} \quad (11)$$

the linear operator defined in our paper. Let us take the Fourier transform of (11),

$$[\mathcal{F}(A_k^{id}\delta_t)](w_l) = \int_{\mathbb{R}^2} \alpha_{i,k}(t) e^{-j\langle \omega_l, s_i \rangle} ds_i \quad (12)$$

*Contact: pierre-jean.benard@math.u-bordeaux.fr , ¹CNRS, Univ. Bordeaux, Bordeaux INP, CNRS, IMB, UMR 5251, F-33400 Talence, France. This work was supported by the French National Research Agency (ANR) under reference ANR-20-CE40-0001 (EFFIREG project). Experiments presented in this paper were carried out using the PlaFRIM experimental testbed (Inria, CNRS , Univ. Bordeaux, Bordeaux INP and Conseil Régional d'Aquitaine).

with $t = (x, y, z) \in \mathbb{R}^3$, $\omega_l = (\omega_{l,x}, \omega_{l,y}) \in \mathbb{R}^2$ and $s_i = (x_i, y_i) \in \mathbb{R}^2$.

$$\begin{aligned} [\mathcal{F}(A_k^{id}\delta_t)](w_l) &= \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{f_k(z)}{2\pi\sigma_x\sigma_y} \\ &\quad e^{-\left(\frac{(x_i-x)^2}{2\sigma_x^2} + \frac{(y_i-y)^2}{2\sigma_y^2}\right)} e^{-j\langle \omega_l, s_i \rangle} ds_i \end{aligned} \quad (13)$$

$$\begin{aligned} [\mathcal{F}(A_k^{id}\delta_t)](w_l) &= \frac{f_k(z)}{2\pi\sigma_x\sigma_y} \left(\int_{-\infty}^{+\infty} e^{-\frac{(x_i-x)^2}{2\sigma_x^2}} e^{-j\omega_{l,x}x_i} dx_i \right) \\ &\quad \left(\int_{-\infty}^{+\infty} e^{-\frac{(y_i-y)^2}{2\sigma_y^2}} e^{-j\omega_{l,y}y_i} dy_i \right) \end{aligned} \quad (14)$$

Let I be the first integral,

$$\begin{aligned} I &:= \int_{-\infty}^{+\infty} e^{-\frac{(x_i-x)^2}{2\sigma_x^2}} e^{-j\omega_{l,x}x_i} dx_i = \\ &\quad \int_{-\infty}^{+\infty} e^{-\frac{(x_i-x)^2+2j\omega_{l,x}x_i\sigma_x^2}{2\sigma_x^2}} dx_i \end{aligned} \quad (15)$$

We isolate x_i from the rest of the terms in the exponent,

$$\begin{aligned} (x_i - x)^2 + 2j\omega_{l,x}x_i\sigma_x^2 &= x_i^2 + 2x_i(j\omega_{l,x}\sigma_x^2 - x) + x^2 \\ &= (x_i + j\omega_{l,x}\sigma_x^2 - x)^2 - (j\omega_{l,x}\sigma_x^2 - x)^2 + x^2 \end{aligned} \quad (16)$$

$$= (x_i + j\omega_{l,x}\sigma_x^2 - x)^2 - (j\omega_{l,x}\sigma_x^2 - x)^2 + x^2 \quad (17)$$

We inject this result in our integral to get

$$I = \int_{-\infty}^{+\infty} e^{-\frac{(x_i+j\omega_{l,x}\sigma_x^2-x)^2-(j\omega_{l,x}\sigma_x^2-x)^2+x^2}{2\sigma_x^2}} dx_i \quad (18)$$

$$= e^{\frac{(j\omega_{l,x}\sigma_x^2-x)^2-x^2}{2\sigma_x^2}} \int_{-\infty}^{+\infty} e^{-\frac{(x_i+j\omega_{l,x}\sigma_x^2-x)^2}{2\sigma_x^2}} dx_i \quad (19)$$

(20)

By proceeding by substitution in this integral and using $x'_i := \frac{x_i+j\omega_{l,x}\sigma_x^2-x}{\sqrt{2}\sigma_x}$, we get that $dx'_i = \frac{1}{\sqrt{2}\sigma_x} dx_i \leftrightarrow dx_i = \sqrt{2}\sigma_x dx'_i$. We then obtain

$$I = \sqrt{2}\sigma_x e^{\frac{(j\omega_{l,x}\sigma_x^2-x)^2-x^2}{2\sigma_x^2}} \int_{-\infty}^{+\infty} e^{-(x'_i)^2} dx'_i \quad (21)$$

We recognize the integral of a one dimensional Gaussian which is equal to $\sqrt{\pi}$,

$$I = \sqrt{2\pi}\sigma_x e^{\frac{(j\omega_{l,x}\sigma_x^2-x)^2-x^2}{2\sigma_x^2}} \quad (22)$$

$$= \sqrt{2\pi}\sigma_x e^{-\frac{\omega_{l,x}^2\sigma_x^2}{2}-j\omega_{l,x}x} \quad (23)$$

We note that we obtain a similar result with the other integral in (14). By injecting our new result (23), we obtain,

$$[\mathcal{F}(A_k^{id}\delta_t)](w_l) = \frac{f_k(z)}{2\pi\sigma_x\sigma_y} \left(\sqrt{2\pi}\sigma_x e^{-\frac{\omega_{l,x}^2\sigma_x^2}{2}-j\omega_{l,x}x} \right) \\ \left(\sqrt{2\pi}\sigma_y e^{-\frac{\omega_{l,y}^2\sigma_y^2}{2}-j\omega_{l,y}y} \right) \quad (24)$$

By simplifying, we get

$$[\mathcal{F}(A_k^{id}\delta_t)](w_l) = f_k(z) e^{-\frac{\omega_{l,x}^2\sigma_x^2}{2}-j\omega_{l,x}x} e^{-\frac{\omega_{l,y}^2\sigma_y^2}{2}-j\omega_{l,y}y} \quad (25)$$

$$= f_k(z) e^{-\frac{\langle \omega_l^2, \sigma^2 \rangle}{2}-j\langle \omega_l, (x, y) \rangle}. \quad (26)$$

□

C. Gradients

We provide the gradients needed to perform OP-COMP with PGD from sketched measurements.

$$\frac{\partial}{\partial x} [\mathcal{F}(A_k^{id}\delta_t)](w_l) = -j\omega_x [\mathcal{F}(A_k^{id}\delta_t)](w_l) \quad (27)$$

$$\frac{\partial}{\partial y} [\mathcal{F}(A_k^{id}\delta_t)](w_l) = -j\omega_y [\mathcal{F}(A_k^{id}\delta_t)](w_l) \quad (28)$$

$$\frac{\partial}{\partial z} [\mathcal{F}(A_k^{id}\delta_t)](w_l) = f'_k(z) e^{-\frac{\langle \omega_l^2, \sigma^2 \rangle}{2}-j\langle \omega_l, (x, y) \rangle}. \quad (29)$$