



HAL
open science

Optical characterization of a single molecule complete spatial orientation using intra-molecular triplet-triplet absorption

Remigiusz Trojanowicz, Ludovic Douillard, Lydia Sosa Vargas, Fabrice Charra, Simon Vassant

► **To cite this version:**

Remigiusz Trojanowicz, Ludovic Douillard, Lydia Sosa Vargas, Fabrice Charra, Simon Vassant. Optical characterization of a single molecule complete spatial orientation using intra-molecular triplet-triplet absorption. *Physical Chemistry Chemical Physics*, inPress, 10.1039/D4CP00867G . hal-04585254

HAL Id: hal-04585254

<https://hal.science/hal-04585254>

Submitted on 23 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

PCCCP

Physical Chemistry Chemical Physics

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: R. Trojanowicz, L. Douillard, L. Sosa Vargas, F. Charra and S. Vassant, *Phys. Chem. Chem. Phys.*, 2024, DOI: 10.1039/D4CP00867G.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Cite this: DOI: 00.0000/xxxxxxxxxx

Optical characterization of a single molecule complete spatial orientation using intra-molecular triplet-triplet absorption[†]

Remigiusz Trojanowicz,^a Ludovic Douillard,^a Lydia Sosa Vargas^b, Fabrice Charra^a, and Simon Vassant^{*a}Received Date
Accepted Date

DOI: 00.0000/xxxxxxxxxx

Progress in single molecule fluorescence experiments have enabled an in-depth characterization of fluorophores, ranging from their photophysical rates to the orientation of their emission dipole moments in three dimensions. However, one crucial spatial information remains elusive: the molecule orientation relative to its emission dipole moment. One can retrieve the latter only by the use of another non-conlinear transition dipole moment. We experimentally demonstrate the optical retrieval of this information for single Terrylene (Tr) molecules in a 30 nm thin para-terphenyl matrix. We show, through second-order correlation measurements at varying excitation power and polarization, that Tr molecules experience an optically induced deshelling of their triplet states, mediated by two orthogonal intra-molecular triplet-triplet absorption dipole moments. We take advantage of these two transition dipole moments to retrieve the full orientation of the Tr molecule, employing a 3-level scheme for the molecule photophysics and analytical calculations for the exciting electric field distribution. This modelling approach enables us to accurately describe both varying power and polarization measurements, giving access to the molecule's photophysical rates and to its complete orientation in three dimensions. This includes the orientation of the singlet emission dipole moment in the laboratory frame, and the orientation of the molecule plane with respect to the singlet emission dipole moment.

1 Introduction

The groundbreaking work of Moerner and Kador¹ demonstrated the feasibility of reaching the single molecule limit with absorption optical measurements. Shortly thereafter, Orrit and Bernard² demonstrated that fluorescence provides a much better signal to noise ratio. The sensitivity achieved at the single-molecule level allowed for the observation of blinking, caused by quantum jump of the molecule from the bright singlet to the dark triplet manifold³. Progress in the field over the past decades enabled a profound understanding of phenomena at play in the photophysics of single emitters⁴. This has given rise to numerous applications, methods and experimental works in spectroscopy, super resolution microscopy and photonic quantum technologies^{5,6}.

An important property of a single-photon emitter is the orienta-

tion of its absorption and emission transition dipole moment. The interaction of both dipoles with their local environment strongly depends on their spatial orientation. This applies to the interaction of the absorption dipole with the excitation light, as well as the interaction of the emission dipole with its environment (such as interfaces or nearby objects) via the Local Density of Optical States (LDOS). Knowledge of the intrinsic and extrinsic properties of single emitters provides valuable information for their use as probes, spanning from fundamental physics experiments to practical applications.

Many experimental efforts have been devoted to measuring the orientation of the singlet emission dipole moment of emitters. Various techniques have been employed, including polarization analysis of fluorescence^{7,8}, back-focal plane imaging^{9,10}, defocused images¹¹ or excitation point-spread-function engineering¹². For more detailed information on these methods and additional references, we direct the reader to reviews from Moerner's¹³ and Brasselet's¹⁴ groups. However, to the best of our knowledge, there is currently no experimental optical technique capable of retrieving the orientation of the emitter around its emission dipole moment. To obtain this last information, another transition dipole moment is required. We will show that intra-

^a Université Paris-Saclay, CEA, CNRS, SPEC, 91191, Gif-sur-Yvette, France. Fax: +33 16908 8786; Tel: +33 16908 9597

^b Sorbonne Université, CNRS, Institut Parisien de Chimie Moléculaire (IPCM), F-75005, Paris, France.

* E-mail: simon.vassant@cea.fr

[†] Electronic Supplementary Information (ESI) available. See DOI: 10.1039/c4CP00000x/



molecular triplet-triplet transition dipole moments can serve this purpose and outline how we experimentally access this information, despite the fact that triplet states are usually dark.

To achieve this, we use Terrylene molecules (Tr, $C_{30}H_{16}$) embedded in a para-terphenyl matrix (pT, $C_{18}H_{14}$), a paradigmatic guest/host system in single molecule fluorescence experiments (See Fig. 1). We chose this particular system for the following reasons: i) Tr is highly photostable in a pT matrix, ii) the molecules are oriented preferentially perpendicular to the glass substrate, iii) the pT matrix has domains of low thicknesses (≈ 30 nm) and iv) it is a well-studied system. The earliest report on this dates back to 1994¹⁵. Since then, numerous studies have been conducted, spanning from liquid helium temperature^{3,16,17} to room temperature^{18–20}, and involving thick (several microns) pT crystals. In a notable study, Fleury et al.¹⁹ characterized all the photophysical rates of Tr, using a 3-level scheme (Fig. 1d). They observed an excitation power-dependent desheling of the triplet state (k_{32} in Fig. 1d). This observation was later confirmed by the group of Kozankiewicz²¹ who measured this desheling rate across a large range of temperatures (5 K to room temperature).




Fig. 1 Molecular system of the study. a) Guest: terrylene, b) Host: para-terphenyl, c) Coordinate system with angle definition for the terrylene molecule orientation. The black arrows (x, y, z) correspond to the laboratory frame. The red arrow correspond to the dipole moment transition of the absorption/emission of the singlet manifold. The smaller green arrow correspond to the dipole moment transition of the "orthogonal" triplet-triplet absorption (See text for details), d) 3-level system for terrylene and associated rates.

The Tr/pT guest/host system has also been employed in a thin film configuration, where Tr molecules showed an aligned emission dipole almost perpendicular to the glass substrate, accompanied by a remarkably high photo-stability²². This thin film configuration paved the way for pioneering nano-photonics experiments, enabling the quantitative measurement of the influence of nanoantennas on the very same single molecule, through scanning probe microscopy^{23,24}. Furthermore, the thin film configuration allowed the integration of single Tr molecules as tools to investigate the impact of more complex structures on their emission properties and photophysics, such as plasmonic nanoparticles^{23–26}, dielectric and metallic local probes²⁷, planar dielectric antennas²⁸, or hybrid nanostructures²⁹. In our specific case, the thin film configuration is particularly appealing as it provides a good knowledge of the excitation electric field distribution.

In this paper, we report fluorescence measurements of single Tr molecules in a sub-30 nm thick pT film under ambient conditions. We measure the second order auto-correlation function of the fluorescence signal with respect to excitation power and polarization on the same single molecule. Our findings reveal that two transition dipole moments are necessary to describe the optically induced desheling of the triplet state. This intricate process involves two intramolecular triplet-triplet transitions followed by reverse InterSystem Crossing (r-ISC) to the singlet manifold. Thanks to the simple excitation field distribution inside the thin film and within the framework of a 3-level system, we utilize these two triplet-triplet transitions to obtain the complete molecule orientation. This includes the three-dimensional orientation of the singlet transition dipole moment and the orientation of the molecule's plane around its singlet transition dipole moment. Furthermore, we extract all the rates involved in Tr fluorescence photophysics, identifying two populations of molecules, distinguished by the magnitude of the triplet state desheling rate as a function of the excitation polarization.

2 Results and Discussion

The electronic states involved in the fluorescence process of Tr molecules can be reduced to a 3-level system as illustrated in Fig. 1d^{18–20}. S_0 denotes the ground singlet state, S_1 is the first excited singlet state and T_1 is the lowest triplet state. Experimental findings have demonstrated that the spin degeneracy of the triplet state is only lifted at temperatures below 17.5 K³⁰. Since our work is conducted at room temperature, we consider only a single triplet level T_1 . The population dynamics of the various levels are governed by a system of three linear differential equations:

$$\begin{pmatrix} \dot{n}_{S_0} \\ \dot{n}_{S_1} \\ \dot{n}_{T_1} \end{pmatrix} = \begin{bmatrix} -k_{12} & k_{21} & k_{31} \\ k_{12} & -(k_{23} + k_{21}) & k_{32} \\ 0 & k_{23} & -(k_{31} + k_{32}) \end{bmatrix} \cdot \begin{pmatrix} n_{S_0} \\ n_{S_1} \\ n_{T_1} \end{pmatrix}, \quad (1)$$

where for example n_{S_0} represents the population of the S_0 level, and \dot{n}_{S_0} denotes its time derivative.

Referring to Fig. 1d, k_{12} represents the excitation rate from the singlet ground state S_0 to the bright excited state S_1 , while $1/k_{21}$ is the fluorescence lifetime, k_{23} the intersystem crossing (ISC) rate from S_1 to a dark triplet state T_1 , k_{31} the T_1 to ground singlet non-radiative depopulation rate, and k_{32} a power dependent triplet desheling rate¹⁹. The latter can be interpreted as a process in which the triplet level is pumped into higher triplet states ($T_1 - T_n$ transition, with $n=5$ or 6 ³¹), followed by a r-ISC to the singlet manifold. It's worth noting that such process is not unique to Tr but is observed in many other molecules³². A 4-level system model was tested in Ref.³³, using data from Ref.²⁸, and it showed no difference compared to the 3-level system beyond the experimental noise. In the end, in agreement with the literature^{18–20}, the electronic structure of the Tr molecule at the excitation wavelength used in this work (532 nm) is described by a 3-level system.

Solving Eq. 1 yields the time evolution of the level population, from which one can derive an analytical formula for the second order correlation function $g^{(2)}(\tau)$ for the steady state under a



continuous-wave excitation³⁴:

$$g^{(2)}(\tau) = \frac{\langle I(t) \rangle \langle I(t+\tau) \rangle}{\langle I(t) \rangle^2} = 1 - (1+A)e^{-\lambda_2\tau} + Ae^{-\lambda_3\tau}, \quad (2)$$

where $(-\lambda_2)$ and $(-\lambda_3)$ denotes the eigenvalues of the system in equation 1, and A is the bunching amplitude given by (see details in Supplementary Materials):

$$A = \frac{\lambda_2(k_{31} + k_{32} - \lambda_3)}{(k_{32} + k_{31})(\lambda_3 - \lambda_2)} \quad (3)$$

Assuming an emission quantum yield of 1 for Tr¹⁰, the total rate of photon emission R_t is given by :

$$R_t = \frac{k_{21}k_{12}k_{31}}{\lambda_2\lambda_3}. \quad (4)$$

These four quantities (λ_2 , λ_3 , A and R_t) are solely function of the rates of the system. The detected count rates R is then given by :

$$R = \eta R_t, \quad (5)$$

where η is the detection efficiency of our experimental system (around 4.5%).

The excitation rate k_{12} and the triplet state deshelving rate k_{32} both exhibit a linear dependence on excitation power^{19,20,28,33}. Therefore, by measuring $g^{(2)}(\tau)$ at different powers, one can extract all the rates involved in the fluorescence process, including those related to the dark triplet state, which is responsible for the molecule blinking. However, previous reports did not consider the respective orientation of the excitation electric field and the molecule. For instance, the power experienced by an absorption transition dipole moment varies with the input polarization as $|\mu \cdot \mathbf{E}|^2$, where μ is the transition dipole moment of the molecule and \mathbf{E} is the excitation field. In our study, we employ two complementary measurements : a variation of excitation power at fixed excitation polarization and a variation of the excitation polarization at fixed excitation power. Subsequently, we demonstrate that the latter type of measurement unveils a complex behavior: k_{32} involves two transition dipole moments.

Experimentally, we utilize a home-made confocal inverted microscope that is equipped with a spectrometer, a Hanbury-Brown-Twiss detection scheme and a home-made tuning-fork based atomic force microscope (TF-AFM). Terrylene molecules were synthesized following the procedure outlined in Ref.³⁵ and samples were prepared as in Ref.²². Using TF-AFM, we ensure that Tr molecules in pT films of thickness below 30 nm are studied (See Supplementary Materials). The excitation is provided by a 532 nm continuous-wave laser in total internal reflection via a high numerical aperture microscope objective. The laboratory reference frame is illustrated in Figure 1c, where (x,z) is the plane of incidence. We specifically select molecules exhibiting a maximum fluorescence intensity at 578 ± 1 nm. Employing a combination of half-wave plate and polarizer allows us to vary the excitation power in a continuous way. Another half-wave plate is used to rotate the excitation polarization before entering the microscope. The excitation polarization can thus be changed from transverse magnetic (or p, electric field in the (x,z) plane) to transverse elec-

tric (or s, electric field in the (y,z) plane). Fluorescence photons are collected through the same objective, filtered and recorded while continuously varying one of the half-wave plate (either for power or polarization). For clarity, we refer hereafter to a "power scan" for the measurement at different excitation power, and a "polarization scan" for the measurement at different excitation polarization angle. More comprehensive details on sample preparation, the optical setup and measurement procedures are provided in the Supplementary Materials.

The result of a typical power scan are depicted in Figure 2, where the measured count rates R , λ_2 , λ_3 and A are plotted as a function of excitation power (see Supplementary Materials for all molecules).




Fig. 2 Typical evolution of the measured count rates R , λ_2 , λ_3 and the bunching amplitude A as a function of excitation power (ramping up and down). The colored areas represent the 68% confidence intervals of the second order correlation function fit results.

Dashed black lines are result of a global fit function (see text for details).

The measurement includes data points for both increasing and decreasing excitation powers, which are almost indistinguishable in Figure 2. This shows the stability of the measurement apparatus, the pT matrix and the Tr molecule up to an excitation power density of 370 kW/cm^2 .

The polarization scan measurements are illustrated in Figure 3 and 4, presenting R , λ_2 , λ_3 and A as function of the excitation polarization angle (see Supplementary Materials for all molecules). When the detected count rates becomes very low, we lack sufficient photons to construct a meaningful $g^{(2)}(\tau)$ function, which accounts for the absence of experimental data points at excitation polarization angles with a low-emission rate.

Two distinct types of behavior are evident. The first one (Figure 3) where the count rates R , λ_2 , λ_3 have their maxima at the same excitation polarization angle. The bunching amplitude A shows a behavior that aligns with the power scan: it starts with a low value at low count rates, reaches a maximum when $R \approx 500$ kHz and then decreases at higher count rates. A slight asymmetry is observed for the value of A at 30° and 125° excitation polarization angles, which will be discussed later. We refer to such molecules as in-phase (ip) molecules.






Fig. 3 Evolution of R , λ_2 , λ_3 and A as a function of input polarization angle for an in-phase (ip) molecule. Colored markers are experimental data points. Dashed black lines are result of a global fit function (see text for details). The colored areas represent the 68% confidence intervals of the second order correlation function fit results.

Left and right panels are respectively linear and polar representation of the same data.

The second population (Figure 4) exhibits a similar behavior for R and λ_2 , but a distinctly different dependence for λ_3 and A : λ_3 is out of phase with R and λ_2 , with its lowest value being ≈ 10 kHz higher than the lowest values of ip molecules. The bunching amplitude shows a maximum, close to the count rates maximum, with a clear asymmetry around 100° . For these molecules, the evolution of λ_3 and A do not align with the power scans and cannot be easily explained. We refer to these molecules as out-of-phase (oop) molecules. It's worth noting that the values of λ_3 and A are very similar for both ip and oop molecules at the maximum count rates polarization angle, rendering them indistinguishable if one analyzes only power scans.

Theoretical calculations³⁶ show that the absorption dipole ($S_0 - S_1$ transition) and the emission dipole ($S_1 - S_0$ radiative transition) are oriented along the long axis of the molecule (red arrow in Figure 1c). Experimentally it was confirmed that the absorption and emission dipoles are aligned³⁷. Therefore, we assume colinear absorption and emission dipoles along the long-axis direction of the Terrylene molecule (red arrow in Figure 1a and c).

In our initial attempt to model the polarization scans, we considered a vertically oriented dipole above the interface, utilizing the rates deduced solely from power scans. We could reproduce qualitatively the main features of most of the ip molecules (except for the asymmetry in A , and an offset of the curves along the excitation polarization axis). However, it proved impossible to replicate the behavior of the oop molecules.

Subsequently, we incorporated the molecule's orientation with respect to the sample plane and to the excitation beam (i.e. the laboratory frame (x,y,z) in Figure 1c). We defined θ for the polar angle, ϕ for the azimuth angle (See Figure 1c for a schematic with angle definitions). For an accurate depiction of the molecule's interaction with the excitation field, we calculated the distribution of the excitation field both analytically³⁸ and numerically³⁹. We

emphasize here the critical role played by the 30 nm thickness of the para-terphenyl film. Although we do not precisely know the molecule's height within the film, the excitation field components remain relatively constant within the 30 nm pT thickness (see Supplementary Materials). This is not the case for thicker films.

However, considering the long-axis orientation of the molecule was not sufficient to describe the asymmetry in the bunching amplitude for ip molecule, and we couldn't even qualitatively reproduce the behavior of oop molecules. To develop a model capable of describing both ip and oop molecules, we must introduce two absorption transition dipole moments to characterize the triplet state deshelling process. These can be regarded as two distinct $T_1 - T_n$ transitions that later cross back to the singlet manifold by r-ISC. This hypothesis is consistent with a Density Functional Theory (DFT) study, predicting a $T_1 - T_5$ transition and a $T_1 - T_6$ transition respectively at 2.34 eV and 2.40 eV, with non-zero oscillator strength³¹. These energies are close to our excitation energy (2.33 eV). Additionally, this is in line with transient absorption experimental measurements of 2,5,10,13-tetra-*t*-butylterrylene (a Tr derivative with similar spectral properties) in benzene⁴⁰ revealing two triplet-triplet absorption broad peaks centered at 2.25 eV and 2.42 eV, overlapping at our excitation laser energy (2.33 eV). Consequently, we can excite these two intramolecular triplet-triplet transitions with our excitation laser. However, the orientation of the associated transition dipole moment in the molecule's reference frame is not reported to our knowledge. Symmetry consideration, as Tr belongs to the D_{2h} symmetry point group, suggests a contribution aligned with the singlet absorption dipole moment, i.e. along the long molecule axis, red arrow in Figure 1a, and a second one orthogonal to the singlet absorption dipole, i.e. (along the molecule short axis, green arrow in Figure 1a). We






Fig. 4 Evolution of R , λ_2 , λ_3 , and A as a function of input polarization angle for an out-of-phase (oop) molecule. Colored markers are experimental data points. Dashed black lines are result of a global fit function (see text for details). The colored areas represent the 68% confidence intervals of the second order correlation function fit results. Left and right panels are respectively linear and polar representation of the same data.

then express k_{32} as the sum of these two orthogonal transitions:

$$k_{32} = \sigma_{32a} |\mu_{32a} \cdot E|^2 + \sigma_{32o} |\mu_{32o} \cdot E|^2, \quad (6)$$

where we introduce two distinct triplet deshelling absorption cross-sections: σ_{32a} and σ_{32o} , respectively for aligned and orthogonal to the singlet transition dipole moment, μ_{32a} and μ_{32o} are the $T_1 - T_n$ transition dipole moment vectors and E the electric field of the excitation laser. The molecule orientation around the singlet dipole should then be specified. We define the angle α for the molecule rotation around its singlet dipole transition moment (See Figure 1c). It's important to note that σ_{32a} and σ_{32o} are not the direct $T_1 - T_n$ absorption cross sections, but the product of the $T_1 - T_n$ absorption cross sections and the r-ISC rate from the T_n levels back to the singlet manifold.

Considering all these elements, we define a global fit function to simultaneously adjust R , λ_2 , λ_3 and A , for both power and polarization scans. To obtain meaningful results in this fitting procedure, we fix the height of the molecule inside the pT film at 15 nm above the pT-glass interface. We also fix the singlet absorption cross-section of Tr at $\sigma_{12} = 1.2 \times 10^{-16} \text{ cm}^2$, which will be justified later. The free parameters include the molecular rates k_{21} , k_{23} and k_{31} , the molecule orientation angles θ , ϕ and α , and the two triplet deshelling absorption cross-sections σ_{32a} and σ_{32o} . This model allows for a perfect adjustment of all the measured molecules (both ip and oop), providing an excellent description of the small asymmetry in the bunching amplitude (A) of ip molecules polarization scans, and accurately describing the excitation polarization behavior of oop molecules. The power scans are also well reproduced, the global character of the fit function ensuring coherence between power scans and polarization scans. This is illustrated in Figures 3 and 4 where the fit result is represented by a black dashed line over the experimental datapoints.

The results obtained for 16 measured Tr molecules are presented in Fig.5 (see also Supplementary Materials). Except for one molecule, the Tr fluorescence lifetime ($1/k_{21}$) ranges within 17.5 to 23.2 ns, with a mean value of 19.78 ns, consistent with previous reports for Tr in pT thin films^{10,23}. For k_{23} , we obtain a mean value around 13 kHz. The dark triplet depopulation rate k_{31} has a mean value of 2.6 kHz. These results falls within the range of previously reported values^{18,19,21}. It's noteworthy that our results show much lower dispersion in k_{23} and k_{31} values than the previous reports. We believe that this is due to our illumination conditions being more constant from molecule to molecule in the thin film configuration, and to our spectral selection. Also, only stable molecules withstand our full measurement procedure and are thus selected in those statistics.




Fig. 5 Results for 16 Tr molecules. Triplet deshelling cross-sections power: σ_{32a} for the component aligned with the singlet transition dipole moment and σ_{32o} for the component orthogonal to the singlet transition dipole moment, the power independent rates (k_{23} , k_{31}) and the fluorescence lifetime ($1/k_{21}$).



Regarding the triplet deshelling absorption cross-sections (See Fig. 5), σ_{32a} shows lower dispersion for all molecules, with a mean value of $6.2 \times 10^{-20} \text{ cm}^2$, while σ_{32o} clearly shows two distinct populations: one around $3.7 \times 10^{-21} \text{ cm}^2$ and the other around $3.8 \times 10^{-20} \text{ cm}^2$. The low values of σ_{32o} correspond to ip molecules, while the higher values of σ_{32o} correspond to oop molecules. Setting σ_{32o} to zero for oop molecules completely changes the evolution of λ_3 and the bunching amplitude A in the polarization scan, making it similar to that of ip molecules, which contradicts experimental observations reported in Fig. 4. Concerning ip molecules, setting σ_{32o} to zero restores a perfect symmetry of the bunching amplitude, giving equal maximum values of A for an excitation polarization angle of 30° and 125° , also contradicting experimental results presented in Fig. 3. So, although σ_{32o} is one order of magnitude lower than σ_{32a} for ip molecules, it does have a noticeable influence on the polarization scan.

A single value for the singlet absorption cross-section σ_{12} is imposed for all molecules. The global fit procedure manages to find solutions for values of $\sigma_{12} \in [1 \times 10^{-16} - 1 \times 10^{-15}] \text{ cm}^2$. Outside this range, the fit quality starts to degrade (higher residuals), and some molecules cannot be fitted properly (unable to match both polarization and power scan). To assess σ_{12} with better accuracy, we compare θ and ϕ obtained from the global fit with results from detection polarization scans for 6 of the 16 molecules, using two analytical models^{7,8} (See Supplementary Material).

The global fit gives a very good match for ϕ , and very close values for θ with a singlet absorption cross-section fixed at $\sigma_{12} = 1.2 \times 10^{-16} \text{ cm}^2$. This value is 2 to 10 times higher than previously reported ones for single molecule measurements in thick crystals^{19,20}. Measurements in a CH_2Cl_2 solution gives $6.7 \times 10^{-16} \text{ cm}^2$ possibly measured at 568.2nm, where absorption is higher than at 532nm¹⁸. Another measurement of 2,5,10,13-tetra-*t*-butylterrylene (Tr derivative with similar spectral properties) in cyclohexane gives $7.8 \times 10^{-16} \text{ cm}^2$ at 557nm⁴⁰. We do not have any experimental means to check the angle α . The obtained values do not show any particular trend. The main uncertainty source in the molecule angle determination is the depth of the molecule inside the film because the weights of the electric field contributions (E_x and E_z) change. We did calculations assuming the extreme cases of a 1 nm and 29 nm depth and obtained uncertainties of $\pm 9^\circ$ for θ , $\pm 8^\circ$ for ϕ and $\pm 4^\circ$ for α . These uncertainties are much larger than the state-of-the-art single molecule orientation-localization microscopy methods^{14,41-43}, but we provide a supplementary information: the molecule orientation around its emission dipole moment.

It is tempting to discuss the Tr orientation in terms of possible insertion sites in the pT matrix. However, as observed by Werley and Moerner⁴⁴, thin pT crystal can show different orientation domains with cracks between them, which imposes special care in the analysis of ϕ and α as our 16 molecules are located in different areas in the pT film and thus likely to belong to different domains. Furthermore, we see no clear trends in angles values between ip and oop molecules.

We turn now to a discussion on the possible physical origin of populations of ip and oop molecules. As mentioned previously, our model is able to extract the optically induced deshelling

rate of the triplet state T_1 . This quantity is the product of the triplet-triplet (T_1-T_n) absorption cross-section and the subsequent r-ISC rate from the triplet to the singlet manifold. The difference between ip and oop molecules can thus come from either of these processes. We hypothesize that geometrical distortions of Tr molecules in the pT crystal could be at the origin of the existence of the two populations. At low temperature, pT has a complex crystalline structure with four non-equivalent pT sites resulting from the locking of the central phenyl ring of pT⁴⁵. Tr fluorescence measurements at 1.4 K indeed showed four different 0-0 phonon lines, suggesting a simple Tr substitution with a pT molecule of the crystal in these four non-equivalent sites¹⁵. However, at room temperature, the central phenyl rings or pT undergo large libration movements, making the molecule on average flat, leading to a mono-clinic P2₁a symmetry group with only 2 degenerate pT sites⁴⁵. Consequently we expect a single type of Tr insertion configuration in our room temperature experiments. The observation of two molecule populations is therefore surprising and suggests that a simple pT substitution might not be the only possible Tr configuration in the pT crystal.

Two different insertion schemes should thus lead to two distinct conformations and/or environments for the Tr molecule. Geometrical distortions can impact the triplet energy levels, resulting in different T_1-T_n oscillator strengths or absorption energies (and thus cross-sections at a fixed excitation energy). Additionally, these distortions can alter the energy and number of vibrational levels, as change in molecular symmetry affects possible vibrational eigenmodes. This, in turn may influence the $T_n \rightarrow S_n$ r-ISC rate. Surprisingly, the $S_1 \rightarrow T_1$ intersystem crossing rate (k_{23}) is similar for both ip and oop molecules. A more detailed discussion can be found in the Supplementary Materials. Based on the experimental absorption cross-sections of the T_1-T_n transitions⁴⁰, we estimate the r-ISC probability ($T_n \rightarrow S_n$) to be $\approx 5 \times 10^{-3}$. Verification of these hypotheses requires challenging experiments and involving DFT calculations which are beyond the scope of the present paper.

3 Conclusions

In conclusion, our investigation, combining excitation power and polarization scans on single fluorescent Tr molecules in a thin pT host matrix, revealed that the optically pumped deshelling rate of the T_1 state involves two orthogonal transition dipole moments. We attribute these dipole moments to $T_1 - T_5$ and $T_1 - T_6$ transitions with subsequent r-ISC to the singlet manifold. We derive values for all the rates at play in the molecule photophysics within a 3-level system framework. The presence of two orthogonal triplet-triplet transition dipole moments enables us to determine the full three-dimensional orientation of the molecule in the laboratory frame: the singlet dipole orientation and the orientation of the molecule plane in respect to the singlet transition dipole moment. The observation of two distinct molecule populations, exhibiting different T_1 deshelling rate along the Tr short-axis raises questions on the insertion geometry of Tr in pT at room-temperature and on the possible distortion of the Tr geometry. Nevertheless, our comprehensive photophysical characterization of Tr provides valuable insights and paves the way for



its enhanced utilization as a single photon source and a potent probe for various applications. The low dispersion of the obtained rates instills confidence for future experiments on the triplet state-associated rates at the single molecule level. In a more general perspective, we hope that our measurement method will enable progress where molecular orientation is crucial, such as the generation of pure spin state from single molecules where magnetic fields aligned with the molecule principal axis are needed⁴⁶, or the design of organic light emitting devices, where orientation plays an important role for both optical and electrical properties^{47,48}.

Author Contributions

Conceptualization, Data curation, Formal Analysis : R.T. and S.V., Funding acquisition and Visualization: S.V., Investigation and Writing - original draft: R.T., Methodology: R.T., S.V. and F.C., Project administration, Supervision and Validation: S.V. and F.C., Resources : L.S.V., L.D. and S.V., Software: R. T., L. D. and S. V., Writing Review and Editing : all authors.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The Acknowledgements come at the end of an article after Conflicts of interest and before the Notes and references.

Notes and references

- W. E. Moerner and L. Kador, *Phys. Rev. Lett.*, 1989, **62**, 2535–2538.
- M. Orrit and J. Bernard, *Phys. Rev. Lett.*, 1990, **65**, 2716–2719.
- T. Basché, S. Kummer and C. Bräuchle, *Nature*, 1995, **373**, 132–134.
- T. Basché, W. Moerner, M. Orrit and U. e. Wild, *Single-molecule optical detection, imaging and spectroscopy*, John Wiley & Sons, 2008.
- S. Adhikari and M. Orrit, *The Journal of Chemical Physics*, 2022, **156**, 160903.
- C. Toninelli, I. Gerhardt, A. S. Clark, A. Reserbat-Plantey, S. Götzinger, Z. Ristanović, M. Colautti, P. Lombardi, K. D. Major, I. Deperasińska, W. H. Pernice, F. H. L. Koppens, B. Kozankiewicz, A. Gourdon, V. Sandoghdar and M. Orrit, *Nature Materials*, 2021, **20**, 1615–1628.
- J. T. Fourkas, *Opt. Lett.*, 2001, **26**, 211–213.
- C. Lethiec, J. Laverdant, H. Vallon, C. Javaux, B. Dubertret, J.-M. Frigerio, C. Schwob, L. Coolen and A. Maître, *Phys. Rev. X*, 2014, **4**, 021037.
- M. A. Lieb, J. M. Zavislan and L. Novotny, *J. Opt. Soc. Am. B*, 2004, **21**, 1210–1215.
- B. C. Buchler, T. Kalkbrenner, C. Hettich and V. Sandoghdar, *Phys. Rev. Lett.*, 2005, **95**, 063003.
- M. Böhmer and J. Enderlein, *J. Opt. Soc. Am. B*, 2003, **20**, 554–559.
- M. P. Backlund, M. D. Lew, A. S. Backer, S. J. Sahl, G. Grover, A. Agrawal, R. Piestun and W. E. Moerner, *Proceedings of the National Academy of Sciences*, 2012, **109**, 19087–19092.
- M. P. Backlund, M. D. Lew, A. S. Backer, S. J. Sahl and W. E. Moerner, *ChemPhysChem*, 2014, **15**, 587–599.
- S. Brasselet and M. A. Alonso, *Optica*, 2023, **10**, 1486–1510.
- S. Kummer, T. Basché and C. Bräuchle, *Chemical Physics Letters*, 1994, **229**, 309–316.
- M. Vogel, A. Gruber, J. Wrachtrup and C. Von Borczyskowski, *Journal of Physical Chemistry*, 1995, **99**, 14915–14917.
- A. C. Brouwer, E. J. Groenen and J. Schmidt, *Physical Review Letters*, 1998, **80**, 3944–3947.
- F. Kulzer, F. Koberling, T. Christ, A. Mews and T. Basché, *Chemical Physics*, 1999, **247**, 23–34.
- L. Fleury, J. M. Segura, G. Zumofen, B. Hecht and U. P. Wild, *Physical Review Letters*, 2000, **84**, 1148–1151.
- L. Fleury, J.-M. Segura, G. Zumofen, B. Hecht and U. Wild, *Journal of Luminescence*, 2001, **94-95**, 805–809.
- M. Banasiewicz, O. Morawski, D. Wiacek and B. Kozankiewicz, *Chemical Physics Letters*, 2005, **414**, 374–377.
- R. J. Pfab, J. Zimmermann, C. Hettich, I. Gerhardt, A. Renn and V. Sandoghdar, *Chemical Physics Letters*, 2004, **387**, 490–495.
- S. Kühn, U. Håkanson, L. Rogobete and V. Sandoghdar, *Phys. Rev. Lett.*, 2006, **97**, 017402.
- H. Eghlidi, K. G. Lee, X.-W. Chen, S. Götzinger and V. Sandoghdar, *Nano Letters*, 2009, **9**, 4007–4011.
- K.-G. Lee, H. Eghlidi, X.-W. Chen, A. Renn, S. Götzinger and V. Sandoghdar, *Opt. Express*, 2012, **20**, 23331–23338.
- X. W. Chen, K. G. Lee, H. Eghlidi, S. Götzinger and V. Sandoghdar, *Opt. Express*, 2015, **23**, 32986–32992.
- S. Kühn, G. Mori, M. Agio and V. Sandoghdar, *Molecular Physics*, 2008, **106**, 893–908.
- K. G. Lee, X.-W. Chen, H. Eghlidi, P. Kukura, L. R., A. Renn, S. Götzinger and V. Sandoghdar, *Nature Photonics*, 2011, **5**, 166–169.
- X.-L. Chu, S. Götzinger and V. Sandoghdar, *Nature Photonics*, 2017, **11**, 58–62.
- M. Białkowska, A. Makarewicz, M. Banasiewicz and B. Kozankiewicz, *Chemical Physics Letters*, 2013, **555**, 131–134.
- T. Minami, S. Ito and M. Nakano, *The Journal of Physical Chemistry Letters*, 2012, **3**, 2719–2723.
- I. Carmichael and G. L. Hug, *Journal of Physical and Chemical Reference Data*, 1986, **15**, 1–250.
- K.-G. Lee, *Journal of the Korean Physical Society*, 2014, **64**, 1792–1796.
- R. Loudon, *The quantum theory of light*, OUP Oxford, 2000.
- Y. Avlasevich, C. Kohl and K. Müllen, *J. Mater. Chem.*, 2006, **16**, 1053–1057.
- T. M. Halasinski, J. L. Weisman, R. Ruitkamp, T. J. Lee, F. Salama and M. Head-Gordon, *The Journal of Physical Chemistry A*, 2003, **107**, 3660–3669.



- 37 J. Sepioł, J. Jasny, J. Keller and U. P. Wild, *Chemical Physics Letters*, 1997, **273**, 444–448.
- 38 D. Axelrod, *Journal of Microscopy*, 2012, **247**, 147–160.
- 39 J. Waxenegger, A. Trügler and U. Hohenester, *Computer Physics Communications*, 2015, **193**, 138–150.
- 40 Y. Meyer, P. Plaza and K. Müllen, *Chemical Physics Letters*, 1997, **264**, 643–648.
- 41 O. Zhang, W. Zhou, J. Lu, T. Wu and M. D. Lew, *Nano letters*, 2022, **22**, 1024–1031.
- 42 O. Zhang, Z. Guo, Y. He, T. Wu, M. D. Vahey and M. D. Lew, *Nature photonics*, 2023, **17**, 179–186.
- 43 P. Jouchet, A. R. Roy and W. Moerner, *Optics communications*, 2023, **542**, 129589.
- 44 C. A. Werley and W. E. Moerner, *The Journal of Physical Chemistry B*, 2006, **110**, 18939–18944.
- 45 P. Bordat and R. Brown, *Chemical Physics*, 1999, **246**, 323–334.
- 46 O. G. Reid, J. C. Johnson, J. D. Eaves, N. H. Damrauer and J. E. Anthony, *Accounts of Chemical Research*, 2023, **57**, 59–69.
- 47 T. D. Schmidt, T. Lampe, P. I. Djurovich, M. E. Thompson, W. Brütting *et al.*, *Physical Review Applied*, 2017, **8**, 037001.
- 48 A. Hofmann, M. Schmid and W. Brütting, *Advanced Optical Materials*, 2021, **9**, 2101004.

