



HAL
open science

Chronic exercise effects on overall depression severity and distinct depressive symptoms in older adults: A protocol of a systematic and meta-analytic review

Melanie Mack, Andreea Badache, Arzu Erden, Christoforos D Giannaki, Sandra Haider, Antonia Kaltsatou, Burcu Kömürcü Akik, Yaël Netz, Iuliia Pavlova, Pinelopi S Stavrinou, et al.

► **To cite this version:**

Melanie Mack, Andreea Badache, Arzu Erden, Christoforos D Giannaki, Sandra Haider, et al.. Chronic exercise effects on overall depression severity and distinct depressive symptoms in older adults: A protocol of a systematic and meta-analytic review. PLoS ONE, 2024, 19 (5), pp.e0297348. 10.1371/journal.pone.0297348 . hal-04587195

HAL Id: hal-04587195

<https://hal.science/hal-04587195>

Submitted on 24 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

STUDY PROTOCOL

Chronic exercise effects on overall depression severity and distinct depressive symptoms in older adults: A protocol of a systematic and meta-analytic review

Melanie Mack^{1,2*}, Andreea Badache³, Arzu Erden⁴, Christoforos D. Giannaki⁵, Sandra Haider⁶, Antonia Kaltsatou⁷, Burcu Kömürçü Akik⁸, Yaël Netz^{9,10}, Iuliia Pavlova¹¹, Pinelopi S. Stavrinou⁵, Claudia Voelcker-Rehage², Michel Audiffren¹², on behalf of the PhysAgeNet¹



OPEN ACCESS

Citation: Mack M, Badache A, Erden A, Giannaki CD, Haider S, Kaltsatou A, et al. (2024) Chronic exercise effects on overall depression severity and distinct depressive symptoms in older adults: A protocol of a systematic and meta-analytic review. *PLoS ONE* 19(5): e0297348. <https://doi.org/10.1371/journal.pone.0297348>

Editor: Claudio Damião Rosa, Universidade Estadual de Santa Cruz, BRAZIL

Received: June 9, 2023

Accepted: December 21, 2023

Published: May 23, 2024

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0297348>

Copyright: © 2024 Mack et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: No datasets were generated or analyzed during the current study. All relevant data from this study will be made available

1 Centre for the Interdisciplinary Study of Gerontology and Vulnerability, University of Geneva, Geneva, Switzerland, **2** Department of Neuromotor Behavior and Exercise, Institute of Sport and Exercise Sciences, University of Münster, Münster, Germany, **3** Department of Disability Research, National Research School on Ageing and Health, Örebro University, Örebro, Sweden, **4** Department of Physiotherapy and Rehabilitation, Faculty of Health Science, Karadeniz Technical University, Trabzon, Turkey, **5** Department of Life Sciences, University of Nicosia, Nicosia, Cyprus, **6** Center for Public Health, Department of Social and Preventive Medicine, Medical University of Vienna, Vienna, Austria, **7** Functional Architecture of Mammals in their Environment (FAME) Laboratory, Department of Physical Education and Sport Science, University of Thessaly, Volos, Greece, **8** Department of Psychology, Faculty of Languages and History-Geography, Ankara University, Ankara, Turkey, **9** The Levinsky-Wingate Academic Center, Tel Aviv, Israel, **10** Department of Health Promotion and Rehabilitation, Lithuanian Sports University, Kaunas, Lithuania, **11** Department of Theory and Methods of Physical Culture, Lviv State University of Physical Culture, Lviv, Ukraine, **12** Cognition and Learning Research Center, Maison des Sciences de l'Homme et de la Société, CNRS, University of Poitiers, Poitiers, France

¶ The complete membership list of the consortium is available here: <https://www.cost.eu/actions/CA20104/#tabs+Name:Working%20Groups%20and%20Membership>
* melanie.mack@unige.ch

Abstract

Introduction

There is high evidence that chronic exercise benefits overall depression severity in older adults. However, late-life depression is characterized by considerable heterogeneity in clinical manifestation emphasizing the need for more individualized exercise intervention programs. Therefore, the objective of the proposed review is to investigate the effects of chronic exercise on overall depression severity and on different symptoms of depression in randomized controlled trials (RCTs) including older adults with a mean age of at least 60 years, and by considering the moderating effects of intervention characteristics and individual characteristics.

Methods

This protocol is guided by the Preferred Reporting Items for Systematic Reviews and Meta-Analysis Protocols (PRISMA-P). We will use the Population-Intervention-Comparator-Outcomes-Study design (PICOS) criteria for study inclusion and will search the following

upon study completion on Open Science Framework, OSF (DOI: [10.17605/OSF.IO/ZNWE8](https://doi.org/10.17605/OSF.IO/ZNWE8)).

Funding: This publication is based upon work from EU COST Action CA20104 - Network on evidence-based physical activity in old age (PhysAgeNet), supported by COST (European Cooperation in Science and Technology). <https://www.cost.eu/>, <https://physagenet.eu/>. The funders did not and will not have a role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. There was no additional external funding received for this study.

Competing interests: All the authors of this manuscript are members of the COST network PhysAgeNet (CA 20104). The authors have declared that no competing interest exists. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

database sources for relevant RCTs: Web of Science, Academic Search Complete, CINAHL, APA Psycinfo, SPORTDiscuss, Cochrane. Two independent reviewers will conduct the study selection, data extraction, and quality assessment. Disagreement will be solved by a third reviewer. Primary outcome will be changes in overall depression severity and secondary outcomes will encompass changes in symptoms of depression as defined by the DSM-5, such as sleep quality, fatigue, anxiety, mood, apathy, changes in weight, information processing speed, and executive functions, from baseline until the end of the intervention and to any available intermediary measurement or follow up. Meta-analysis will be undertaken to synthesize the effects of chronic exercise on primary and secondary outcomes. Subgroup analysis will investigate the moderating effects of intervention characteristics (frequency, intensity, duration, type of exercise, cognitive demand, social interactions, exercise supervision, behavioral change techniques, compliance, study design, dropout-rate, type of control group) and individual characteristics (age, sex, education, functional capacity, global cognition, population) on primary and secondary outcomes. Additionally, we plan to assess quality of evidence and publication bias, and to carry out sensitivity analysis.

Conclusion

The results of the proposed review are anticipated to have a substantial impact on research and clinical practice. On the one hand, the review's conclusions could form the foundation for developing evidence-based recommendations for individualized exercise programs that alleviate depression in older adults. On the other hand, by revealing research gaps, the review results could encourage the formulation of research questions for further RCTs.

Protocol registration number

This protocol has been published in the Prospero repository (PROSPERO 2022 CRD42022361418, available from: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42022361418)

Introduction

Depression among older adults, also referred to as geriatric depression or late-life depression, presents a significant public health concern. The average global prevalence of geriatric depression, including estimates assessed between 1994 and 2020, is 31.74% [1], and rates of depressive disorders increase substantially in the highest age group 85 years and older, reaching up to 50% [2]. These high numbers may be attributed to various factors associated with aging, such as a higher proportion of women, more physical disability, higher cognitive impairment, and lower socioeconomic status [3, 4]. Given its high prevalence and potential negative consequences, such as increased physical illness, impaired functioning, suicide risk, and reduced quality of life, depression in older adults deserves special attention [5, 6].

Beneath the symptoms of depression described in the American Psychiatric Association's Diagnostic and Statistical Manual (DSM-5 [7]) and in the International Classification of Diseases (ICD-10 [8]), late-life depression is especially associated with cognitive and somatic changes [6, 9]. Cognitive changes may manifest as difficulties with concentration, speed of

information processing, and executive functions [10, 11], while somatic changes may include increased abdominal fat, type 2 diabetes, and hypertension [12]. In addition to medical and neurological comorbidity [13, 14], late-life depression is frequently associated with disability and psychosocial difficulties. These health domains often interact with one another, contributing to the development of geriatric depression [13, 15]. For instance, cerebrovascular disease may cause depression and cognitive impairment, which can lead to disability and social isolation, further worsening depressive symptoms [16, 17]. Once depression sets in, it may increase cognitive impairment, disability, psychosocial problems, and morbidity. Furthermore, older individuals with recurrent depression since younger age and those with late-life onset seems to have distinct clinical features, risk factors, and treatment responses [9, 18]. Compared with older patients with early-onset depression, patients with late-onset depression appear to have a higher prevalence of medical burden and neurological brain disorders [19], more impairment in neuropsychological tests [20] and dysfunctions of several brain regions [9, 20].

Given the complexity and heterogeneity of late-life depression, it is not surprising that many patients are not adequately treated [21]. The first line of treatment of late-life depression is commonly pharmacotherapy. However, antidepressants pose a considerable risk for side effects, especially in the presence of comorbidities and the administration of adjunctive medications, including risk of falls, osteoporosis, as well as cardiac and sexual dysfunction [21]. Thus, the majority of patients receive no treatment, an insufficient duration of treatment, or a lower than recommended dose [22, 23]. In addition, antidepressant efficacy in older adults is modest and declines with age, which may contribute to the higher rates of depression relapse in this population [21]. In recent years, regular physical exercise has gained attention in clinical practice and research as a promising noninvasive treatment option for late-life depression. By the majority of populations, exercise can be performed autonomously without additional costly equipment or external assistance, has no serious side effects, and is accessible to everyone.

Regular physical exercise, often termed as chronic or systematic exercise, is defined as planned, structured, and repetitive bodily movements done to improve or maintain one or more components of physical fitness [24]. Interventions including exercise aiming to reduce depression can be referred to as non-pharmacological interventions or behavioral medicine/therapy interventions and may fall within the scope of evidence-based behavioral medicine or behavioral therapy [25]. Evidence-based behavioral medicine is a systematic approach to use the best available scientific evidence from clinical research to help make decisions about the care of individual patients [26]. This approach has become particularly important in the development of clinical standards and guidelines to improve the quality of care. Systematic reviews of the available published evidence are required to identify interventions that result in improvements of behavior and health. The validity and explanatory power of reviews depends on the quality of the published research, but also on the compilation and descriptions of the aspects that are important for care [27–29]. The extension of the CONSORT statement for RCTs of non-pharmacological interventions specified a list of relevant aspects to be reported, such as the precise description of both the experimental treatment and the comparator, the implementation of the intervention, and how adherence was promoted and assessed [27]. In the following, we will discuss which aspects might be especially important regarding exercise interventions for late-life depression.

Intervention characteristics

Exercise interventions benefit depressive symptoms in older adults by promoting a variety of psychological and physiological mechanisms relevant to late-life depression. The former

includes psychosocial and cognitive factors, such as augmenting self-esteem [30], the latter includes anti-inflammatory effects or neuroplasticity mechanisms, beyond others [31, 32]. However, the heterogeneity of the effects is high, which might be caused by variations between exercise interventions. The induced mechanisms and their strength depend strongly on the specific training regime. One important aspect that impacts the effects of exercise interventions are the FITT principles of training, which stands for 'Frequency', 'Intensity', 'Time', and 'Type of exercise' [24]. 'Frequency' refers to the number of exercise sessions per week, while 'Intensity' refers to the level of exertion during exercise. 'Time' refers to the duration of each exercise session, and 'Type' refers to the specific mode of exercise, such as aerobic, resistance, or mind-body exercises. By manipulating the four components of the FITT principles, exercise programs can be designed to elicit specific adaptations in the body. For example, aerobic exercises or resistance exercises have a high metabolic demand that benefit depression by increasing brain volume (gray and white matter of primarily prefrontal and temporal regions) or peripheral biomarkers, such as BDNF [33]. Mind-body exercises, such as Yoga, Tai Chi, balance and other coordinative exercises, place high demands on the neuromuscular and neurocognitive system that might benefit depression by initiating functional and structural changes [34]. The effect size of exercise interventions is further modulated by frequency, intensity, and duration of exercise sessions [35]. Exercises with a higher intensity might induce higher acute neuroendocrine responses and long-term adaptations in the brain's neural architecture than exercises with lower intensity [36]. Meaningful responses may be only visible when the exercises are conducted with a certain frequency and duration [37].

Beneath the FITT principle, further important components of exercise interventions to improve late-life depression are psychosocial aspects, such as the degree of social interaction among participants or the use of behavioral change techniques. Social interaction differs between group-based and individual-based (including home-based) interventions but also within group-based interventions. Group-based training sessions, for example, may include exercises with partner(s) in cooperation and/or opposition or only verbal interactions [38]. The extent of social interaction can influence social connectedness and thus social identity and loneliness. Loneliness, defined as feelings of isolation, disconnectedness, and lack of belonging, is a strong and well-established risk factor for depression [39]. The social identity approach assumes that belonging to a group can protect against depression because of the associated access to resources (e.g., social support from other members), even in the absence of access to support from or contact with other group members [40, 41]. Behavioral change techniques, such as goal-setting, self-monitoring, and problem-solving, are strategies that are used to promote behavior change. In the context of exercise interventions, behavioral change techniques may be used to help individuals set and achieve realistic exercise goals, track their progress, and overcome barriers to exercise [42]. Parts of those techniques are also included in cognitive behavioral therapy programs that are known to have beneficial effects for older adults suffering from depression [43]. Although in exercise interventions, these techniques are applied to the context of exercise behavior, it is plausible that they can also be translated to other areas of life, such as dealing with depression. One might for example assume that participants who learn to make connections between their thoughts, feelings, and actions in relation to activity behavior can also transfer this strategy to depression and develop alternative ways of coping with the depressive symptoms.

RCTs are considered the gold standard for effectiveness research because they allow a cause-effect relationship to be established between the intervention and the outcome. However, for exercise interventions with psychosocial or behavioral components, it is difficult to identify appropriate control groups because the active/inactive components of the interventions are not as obvious as they are for medication interventions. This might lead to

methodological issues in the choice of the control group that might influence the results [44]. For example, an RCT including an active control group with social interactions to examine the effect of an individual-based exercise intervention (without social interaction) is not appropriate. Therefore, the choice of control group in RCTs on exercise interventions for late-life depression should be considered when interpreting results.

Individual characteristics

In attempting to generate reliable and clinically relevant evidence on the effects of exercise in late-life depression, the question of "what works for whom," also called precision medicine or personalized medicine is important to be considered. In this vein, the choice of the outcome but also the population to be treated might also be of importance. Late-life depression can be assessed with different validated depression scales, such as the Beck Depression Inventory-II (BDI-II [45]) or the Geriatric Depression Scale (GDS [46]). These standardized clinical instruments typically quantitatively measure the severity of depressive symptoms in a single score based on a multidimensional assessment of a range of depressive symptoms, including mood, appetite, sleep, energy, and concentration, as well as feelings of worthlessness, hopelessness, and guilt. By providing one overall depression score, depression rating scales can be helpful in clinical decision-making, tracking treatment progress, and determining the effectiveness of various interventions. Nevertheless, one single score provides only limited information on the severity of the specific symptoms that differ enormously in late-life depression, and which are linked to specific pathophysiological mechanisms as described above. Considering that exercise affects the brain and body by specific pathways, it is plausible that the effects of exercise on depression may vary depending on whether those pathways are impaired or vulnerable in the individual patient.

Older adults suffering from depression further differ concerning various individual characteristics, such as age, sex, physical fitness, cognitive functions, years of education and comorbidities. It is important to note that these individual characteristics might influence which exercise intervention is appropriate and leads to the most beneficial effects [47]. For instance, the physiological response to equivalent 'dosages' of exercise differs between males and females [48]. Therefore, gender might influence the intervention outcomes. Different comorbidities might also lead to differential effects of exercise. First, exercise is not without risk. Potential adverse events include cardiac events or hypoglycemia, and to ensure the safety of exercise certain precautions, such as reducing the intensity, or the duration of an exercise session, are necessary for certain populations. These precautions might imply that the impact of exercise is insufficient for reducing depression. However, it is also possible that exercise has a positive effect not only on depression but also on comorbid conditions. If these comorbidities are linked to depression (for instance, frailty leading to depression as a result of the loss of the ability to carry out daily activities), it could potentially result in even greater improvement in depression. Although these connections have not been investigated yet, they hold significance in the context of geriatric depression.

State of research

Several systematic reviews and meta-analyses investigated the effect of a large variety of chronic exercise interventions on depression in various populations of older adults (for an overview see umbrella reviews [49, 50]). Most of those overview articles on the effects of exercise on late-life depression target only overall depression severity as an outcome variable of interest, assessed with different validated depression scales [51–58]. Although there are quite many RCTs examining the effects of chronic exercise on specific symptoms of depression (to

name only a few [59–61]), to our knowledge, there is no published review or meta-analysis that attempts to uncover the effects of chronic exercise on the broadest possible range of symptoms as described in the DSM. However, given the substantial heterogeneity in clinical presentation of late-life depression, it underscores the need of customizing the exercise interventions to the unique symptom expressions of each patient.

Overall, already conducted reviews and meta-analysis suggest a moderate effect of chronic exercise on the severity of late-life depression, with the exercise group showing a higher decrease of depression severity compared to a control group with no exercise intervention [49]. A proportion of those overview articles also examined the effects of different moderator variables on this exercise-depression relationship. The studies addressed exercise intervention characteristics, such as intensity of exercise [47], type of exercise [47, 54], type of control group [53]; but also, individual characteristics, such as age [51, 56, 62], comorbidities [47, 62, 63], severity of depression [62, 64], or cognitive impairment [62]. The results show, on the one hand, a high degree of heterogeneity for the exercise characteristics and, on the other hand, partly different results for the individual characteristics. One example is the moderating effect of age. While one study found an increasingly beneficial effect of exercise with increasing age [51], another study found opposite results with decreasing positive effects with increasing age [62], and a third study found no moderating effect of age [56]. One reason for the varying and heterogeneous results could be that the studies included in the reviews differ in various aspects of individual and exercise characteristics. For example, the overview articles mentioned above differ in the type of exercises included: only mind-body exercises [51] versus only aerobic and resistance exercises [62] versus all three types of exercises [56]. It might be suggested that exercises, which vary in their characteristics, have different effects on depression due to the distinct neurobiological and psychological mechanisms they involve [54, 65, 66]. Thus, they may also have differential effects in different populations. For example, one might assume that moderate-intensity physical exercise with high cognitive demand is particularly beneficial for the elderly, as the likelihood of cognitive decline increases with age. Unfortunately, there is no review that comprehensively examines the influence of potentially relevant moderator variables on the relationship between chronic exercise interventions and depression in older adults.

In addition, there are numerous sources of bias that might threaten the validity of meta-analyses, such as low methodological quality or publication bias. In a recent umbrella review on the effects of chronic exercise on depression in older adults, only one of the twelve included reviews was considered to be of high methodological quality as defined by the rating of the ‘A MeaSurement Tool to Assess systematic Review– 2’ (AMSTAR-2 [67]) [49]. In terms of publication bias, there is evidence that previous meta-analyses on the effects of chronic exercise on depression have underestimated the benefits of exercise due to publication bias [57, 68]. This is different than for other treatments, such as psychotherapy whose effect sizes have consequently been overestimated [69], and would speak for an even higher beneficial effect of exercise on late-life depression as suggested by now. Since publication bias is seldom calculated, further evidence is needed to strengthen the evidence.

Rationale and objective

To sum up, available reviews provide evidence for the beneficial effect of chronic exercise on depression severity in older adults, but several issues remain unaddressed: (1) while most of the literature uses overall depression severity as an outcome, it is unclear whether chronic exercise differentially affects the various symptoms of depression; (2) the numerous reviews that have already examined the moderating effects of intervention characteristics and

individual characteristics have unfortunately found heterogeneous and partly differential results, probably because of their differential inclusion criteria with respect to those characteristics (e.g., only including mind-body interventions vs. only aerobic interventions or only including participants with neurodegenerative disease vs. only participants with major depressive disorder). Consequently, the moderating role of those characteristics and their interaction is still not fully understood; (3) low methodological quality of the included RCTs and publication bias seem to influence the validity and interpretation of the results but are only rarely considered. To derive evidence-based treatment decisions, reviews considering those aspects are necessary.

We hypothesize that chronic exercise interventions differ in their efficacy in reducing overall depression severity and the different symptoms of depression in older adults across participants with different individual's characteristics (i.e., age, sex, population, level of education at baseline, functional capacity at baseline, global cognition at baseline, depression severity at baseline) and across interventions with different exercise characteristics (i.e., frequency of exercise sessions, intensity of exercise, duration of exercise session, duration of intervention, type of exercise, cognitive demand of exercise, social interaction, use of behavioral change techniques, type of control group). Therefore, this protocol outlines a review aiming to investigate how the above-mentioned intervention characteristics and individual characteristics moderate the effects of chronic exercise on overall depression severity and the different symptoms of depression in older adults. The different symptoms of depression will be thereby defined using the criteria delineated by the DSM-5 [7] to make a diagnosis of depression: depressed mood, loss of interest/pleasure, weight loss or gain, insomnia or hypersomnia, psychomotor agitation or retardation, fatigue, feeling worthless or excessive/inappropriate guilt, decreased concentration, thoughts of death/suicide. Each of these symptoms will be used as secondary outcome in the present meta-analysis. In addition, we aim to assess quality of evidence and to consider publication bias [70].

The review will differ from previous reviews in several important ways: (1) by including a large sample of studies with a wide range of individual and exercise characteristics, the chronic effects of exercise on late-life depression will be examined as a function of a large variety of moderators related to these characteristics; (2) the effects of chronic exercise will be examined not only on the level of overall depression severity but on the level of the different symptoms of depression; (3) the results will be corrected for publication bias with different complementary techniques.

Method

This protocol has been published in the Prospero repository (PROSPERO 2022 CRD42022361418, available from: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42022361418) and follows the Preferred Reporting Items for Systematic Reviews and Meta-Analyses Protocol (PRISMA-P) statement (see [S1 Checklist](#)) [71].

Eligibility criteria

We will use the Population-Intervention-Comparator-Outcome-Study design (PICOS) criteria to determine primary study inclusion.

Population

To be considered in this review, studies must fulfill the following criteria: (1) age: the mean age of each arm of the RCT is 60 years or above and the minimal age of each arm of the RCT is 50 years (i.e., the mean age minus standard deviation of age must be 50 years or above); (2) level

of depression: the mean level of depression of each arm of the RCT is mild / moderate or higher as attested by any validated scale assessing depression (i.e., the mean score of depression + the standard deviation must be \geq cutoff value). For each depression scale we will therefore define the cutoff value based on the literature. The list of all validated scales and corresponding cutoffs will be made available on Open Science Framework (OSF). Our review further aims to encompass all populations of older adults independent of disease (e.g., dementia, cardiovascular disease, etc.), physical fitness and cognitive functioning or setting (e.g., clinical setting, community setting, etc.).

Intervention

To be eligible for this review, studies must include at least one intervention arm comprising a chronic exercise component defined as any kind of "planned, structured, and repetitive bodily movement done to improve or maintain one or more components of physical fitness" [24], such as aerobic exercise, resistance exercise, motor-coordinative exercise, mind-body exercise, exergames, hybrid exercise, multicomponent exercise (also including cognitive training) and concurrent training. The intervention must last at least one week and include more than one exercise session. Intervention arms comprising a chronic exercise component additionally to other intervention components, such as nutrition, cognitive behavioral therapy, or pharmacotherapy, are not eligible. However, this only applies if the cognitive behavioral therapy or pharmacotherapy component is specifically manipulated and tested in the study. If they are part of the usual treatment, the study will be included. Furthermore, we will only include studies in which frequency of exercise sessions, duration of exercise sessions, duration of the intervention and type of exercise are reported.

Comparator groups

Studies included in this review must include either a passive or an active control group. Eligible passive control groups include: (1) usual care / treatment as usual (TAU): the diseased participants continue to receive their usual treatment; (2) habitual activities: the healthy participants do not change their life habits; (3) waiting list: the participants do not change their life habits during the duration of the intervention but then follow the tested treatment. Eligible active control groups include: (1) health education: the participants follow a health education program; (2) non-physical leisure activities: the participants practice non-physical leisure activities, such as listening to music, reading newspapers, magazines or books, watching documentaries or movies, handicraft, playing games, etc.; (3) stretching / very light intensity exercise / relaxation: the participants practice exercises with a very light intensity, such as stretching, moving joints, relaxation; (4) social interactions: the participants communicate regularly with other participants.

Outcomes

Studies eligible for this review must include a measure of depression scores at baseline and at the end of the intervention, as measured by a validated scale (c.f., Population).

Study design

Eligible studies only include RCTs with either a parallel-group design (each participant is randomly assigned to one of multiple study arms) or a cluster design (pre-existing groups of participants, such as villages, nursing homes, are randomly assigned to one of multiple study arms).

Article type

For inclusion in this review, a study must be a peer-reviewed scientific report or a PhD thesis of an RCT written in English language. Abstracts from congresses and study protocols are not considered.

Information sources and search strategy

The following electronic databases will be searched from inception to shortly before publishing the meta-analysis: Web of Science (all bases); Academic Search Complete, MEDLINE with full text, CINAHL with full text, APA Psycinfo, SPORTDiscuss with full text through EBSCO host; Cochrane. The complete search strategy can be found in the [S1 File](#). In addition, reference lists of the included studies will be manually searched to identify additional relevant studies (e.g., from previous meta-analyses on the same topic). The search will be re-run prior to the final analysis for the months not covered by the initial search.

Study selection

Study selection will be performed with the reference management system Rayyan [72]. [Fig 1](#) illustrates the planned study selection process. The results of literature searches will be imported into Rayyan using RIS format, and duplicates will be removed. The selection of studies will be conducted in two steps. Pairs of two reviewers will be involved in the study selection process. In the first stage, the two reviewers will independently screen the title and abstract of




Fig 1. Planned study selection process.

<https://doi.org/10.1371/journal.pone.0297348.g001>

their assigned studies. If a disagreement occurs between the two reviewers, a third reviewer will independently examine the eligibility criteria. Selected studies at stage 1 will then be assigned to other pairs of reviewers for the screening of the full-text articles. It follows the same procedure as the screening of the title and abstract. To ensure compliance with the workflow and eligibility criteria, guidelines on the study selection process will be established for reviewers to follow and training sessions will be conducted. The studies selected in the different steps will be saved on Rayyan in respective directories. The guidelines for each stage of the selection process will be available on OSF.

Data extraction and management

Study data will be collected and managed using REDCap electronic data capture tools hosted at Yale University [73, 74]. Pairs of two reviewers will be involved in the data extraction process. The two reviewers will independently extract data from included studies. If a disagreement occurs between the two reviewers, a third reviewer will independently examine the extracted data and correct the wrong data. The following pieces of information will be extracted by each pair of reviewers and entered in the REDCap database: first author of the report, year of publication, characteristics of the RCTs (i.e., number of arms, type of control group, etc.), primary outcomes, secondary outcomes, moderator variables and risk of bias. Outcomes and moderators are detailed hereafter. If data is missing, incomplete or questionable, investigators will be contacted to obtain or confirm data. To ensure compliance with the workflow, guidelines on the study selection process will be established for reviewers to follow; and training sessions will be conducted. The guidelines and the questionnaire used for the data extraction process will be available on OSF.

Outcomes

All outcomes will be measured as the mean change of the values in the exercise group from baseline to the first post-intervention measurement time point, in comparison to the mean change of the control group. Therefore, we will extract the means and standard deviations from the exercise group and the control group at baseline and post-intervention measurement.

Primary outcome

The primary outcome will be depression severity assessed with a validated depression scale. If an author reported the results of two primary outcome measures meeting our criteria (i.e., mean change in depressive symptoms according to two different measures), we will use both.

Secondary outcomes

Secondary outcomes will include symptoms of depression delineated by the DSM-5 to make a diagnosis of depression. For the selection of the outcomes, we have adhered as closely as possible to each of the specific criteria, which are as follows: (1) depressed mood most of the day, nearly every day, as indicated by either subjective report (e.g., feelings of sadness, emptiness, hopelessness) or observation made by others (e.g., appears tearful); (2) markedly diminished interest or pleasure in all, or almost all, activities most of the day, nearly every day; (3) significant weight loss when not dieting or weight gain (e.g., a change of more than 5% of body weight in a month) or decrease or increase in appetite nearly every day; (4) psychomotor agitation or retardation nearly every day (observable by others, not merely subjective feelings of restlessness or being slowed down); (5) insomnia or hypersomnia nearly every day; (6) fatigue or loss of energy nearly every day; (7) feelings of worthlessness or excessive or inappropriate

guilt (which may be delusional) nearly every day (not merely self-reproach or guilt about being sick); (8) diminished ability to think or concentrate, or indecisiveness, nearly every day; (9) recurrent thoughts of death (not just fear of dying), recurrent suicidal ideation without a specific plan, or a suicide attempt or a specific plan for committing suicide. We have already completed a preliminary overview of RCTs on this topic to determine how many studies report each outcome of interest. If several instruments were available to assess a given symptom, we defined an order determined by the validity and frequency of use of the measurement instrument. An exception is the secondary outcome executive functions (see below). Here we will collect all reported measures. If not enough studies (less than three out of a total of approximately 100 studies) report a valid measure for a particular symptom, we will not assess that symptom as secondary outcome.

Sleep quality will be assessed with the following scales to measure aspects of depression related to the symptom “insomnia or hypersomnia nearly every day”: (1) Pittsburgh Sleep Quality index (PSQI) [75]; (2) Insomnia Severity Index (ISI) [76].

Fatigue will be assessed with the following scales to measure the symptom “fatigue or loss of energy nearly every day”: (1) Fatigue Severity Scale (FSS) [77]; (2) Parkinson’s Fatigue Scale (PFS-16) [78]; (3) Multidimensional Fatigue Inventory (MFI) [79]; and (4) Functional Assessment of Chronic Illness Therapy–Fatigue (FACIT-F) [80].

Anxiety will be assessed with the following scales to measure aspects of depression related to the symptom “depressed mood most of the day, nearly every day, as indicated by either subjective report (e.g., feels sad, empty, hopeless) or observation made by others (e.g., appears tearful)”: (1) Beck Anxiety Inventory (BAI) [81]; (2) Generalized Anxiety Disorder—7 items (GAD-7) [82]; (3) Hamilton Anxiety Rating Scale (HAM-A) [83]; (4) Hospital Anxiety and Depression Scale, subscale Anxiety (HAS) [84]; (5) State-Trait Anxiety Inventory, subscale Trait Anxiety (STAI) [85]; and (6) Symptom-Checklist, subscale Anxiety (SCL-90-R) [86].

Mood will be assessed with the following scales to measure aspects of depression related to the symptom “depressed mood most of the day, nearly every day, as indicated by either subjective report (e.g., feelings of sadness, emptiness, hopelessness) or observation made by others (e.g., appears tearful)”: (1) Alzheimer’s Mood Scale, subscale Negative Mood [87]; (2) Dementia Mood Scale, subscale Mood (DMAS) [88]; (3) Positive and Negative Affect Schedule, subscale Negative Affect (PANAS) [89]; (4) Profile of Mood States (POMS) [90]; and (5) State-Trait Anxiety Inventory, subscale State Anxiety (STAI) [85].

Apathy will be assessed with the following scales to measure aspects of depression related to the symptom “markedly diminished interest or pleasure in all, or almost all, activities most of the day, nearly every day (as indicated by either subjective account or observation)”: (1) Apathy Evaluation Scale (AES) [91]; (2) Neuropsychiatric Inventory Questionnaire, Apathy subscale (NIQ) [92]; and (3) Starkstein Apathy Scale (SAS) [93].

Changes of weight will be assessed with the Body Mass Index (BMI) to measure aspects of depression related to the symptom “significant weight loss when not dieting or weight gain (e.g., a change of more than 5% of body weight in a month) or decrease or increase in appetite nearly every day”. We are aware that regular exercise might influence BMI changes and that this variable carries the potential for misinterpretation, as weight changes could mistakenly imply loss of appetite or unintended weight loss. Nonetheless, given the robust U-shaped association between overall depression severity and obesity status (i.e., underweight, normal weight, overweight / obese) [94], we consider this symptom to be of significant importance. It might be plausible that exercise leads to a normalization of BMI, with individuals with excessively high BMIs experiencing weight loss and those with excessively low BMIs gaining weight, rather than resulting in an overall reduction in BMI.

Table 1. Overview of most used tests assessing executive functions in older adults.

Test or battery	Executive function	Outcome
Stroop task	Inhibitory control	Interference score, number of errors in incongruent trials
Random number generation task	Inhibitory control	Adjacency score
Hayling task	Inhibitory control	Inhibition score
Spatial Running Span task	Working memory	Number of correct responses
Verbal Running Span task	Working memory	Number of correct responses
2-back task	Working memory	Number of correct responses
Dimension-switching task	Cognitive flexibility	Local switch cost
Plus-Minus task	Cognitive flexibility	Shifting cost
Digit-Letter task	Cognitive flexibility	Shifting cost
Counting backward	Working memory	Number of correct responses
Digit Span Backward	Working memory	Number of correct responses
Stop Signal task	Inhibitory control	Stop Signal Reaction Time
Attention Network Test [96]	Inhibitory control	Error rate
Wisconsin Card-Sorting Test [97]	Cognitive flexibility	Number of shifts correctly performed, perseverance errors
Frontal Assessment Battery [98]	Executive functions	Global score
Go / No Go task	Inhibitory control	Number of failed inhibitions
Trail making test [95]	Cognitive flexibility	TMTB score–TMTA score, TMTB score, TMTB / TMTA

<https://doi.org/10.1371/journal.pone.0297348.t001>

Information processing speed will be assessed with the following cognitive tests to measure aspects of depression related to the symptom “psychomotor agitation or retardation nearly every day (observable by others, not merely subjective feelings of restlessness or being slowed down)”: (1) Trail Making Test, part A (TMT) [95]; and (2) reaction time (RT) tasks (simple RT task and two-choice RT task with compatible S-R mapping).

Executive functions will be assessed with the different cognitive tests to measure aspects of depression related to the symptom “diminished ability to think or concentrate, or indecisiveness, nearly every day” (see Table 1).

We will further consider changes in depression score and symptoms of depression from baseline to any available intermediary measurement or follow-up as secondary outcomes.

Moderators

We will extract information about moderator variables that are either commonly used in related previously published meta-analyses and / or are hypothesized to modulate the effect of chronic exercise on depression.

Intervention characteristics

Concerning exercise characteristics, we will extract information on the key components of the FITT principles:

- *Frequency of exercise sessions* assessed as mean number of sessions per week.
- *Intensity of exercise* categorized as follows: (1) light; (2) moderate; (3) vigorous / high [24]. The intensity of exercise will be determined by a dual approach utilizing both objective and subjective ratings. Objective ratings will be guided by cut-off points derived from literature for heart rate reserve (%HRR), oxygen uptake reserve (%VO₂R), maximal oxygen consumption (VO₂max), maximal heart rate (%HRmax), repetition maximum (%RM). Subjective ratings will be guided by predefined cut-off points for METs (metabolic equivalents of tasks),

ratings of perceived exertion (RPE), as well as specific exercise types classified by a Delphi survey by the COST action PhysAgeNet.

- *Duration of exercise sessions* assessed as mean duration of sessions in minutes.
- *Duration of the intervention* assessed as mean duration in weeks.
- *Type of exercise* categorized as follows: (1) aerobic exercise (e.g., Nordic Walking, swimming, running); (2) strength exercise / resistance exercise (e.g., weight-lifting, Pilates); (3) dance (Chinese scare dance, Turo, dance-based therapy, private ballroom dance, folk dance); (4) balance exercise / coordination exercise (postural reeducation, gentle gymnastics, range of motion exercises); (5) mind-body exercises (Tai Chi, Qigong, Baduanjin, Yoga, Alexander techniques, Feldenkrais); (6) exergames / video games / virtual reality (active game play, Wii games, Kinect games); and (7) multi-component exercises (combine two or more categories described above with or without a cognitive component).

We will extract information on intervention characteristics related to psychological / social / cognitive aspects:

- *Cognitive demand of exercise* categorized as either low or high and assessed by their affiliation to a specific type of exercise. Low cognitive demanding exercises include aerobic exercises and strength exercises / resistance exercises. High cognitively demanding exercises include balance exercises, coordination exercises, mind-body exercises, exergames, and cognitive tasks.
- *Social interaction* categorized by according to whether exercise sessions are conducted (1) in a group-based fashion, (2) an individual-based session containing only verbal interactions, and (3) an individual-based fashion containing exercising with partner(s) in cooperation and / or opposition [38].
- *Supervision during exercise practice* categorized as 1) home based, 2) supervised by a professional, and 3) not specified. Within home-based programs, participants engage in the program independently, relying on written or video materials for guidance but lacking real-time feedback during exercise. In supervised programs, a professional coach or instructor provides guidance and feedback to participants throughout the sessions.
- *Behavioral change techniques* categorized as follows: (1) no behavioral change technique; (2) goals and planning; (3) feedback and monitoring; (4) social support; (5) shaping knowledge, (6) natural consequences; (7) comparison of behavior; (8) associations; (9) repetition and substitution; (10) comparison of outcome; and (11) rewards and threat [42].
- *Compliance* assessed as percentage of sessions attended.

We will further extract information on study characteristics related to study design and statistical analysis aspects:

- *Study design* of the RCT categorized as either (1) parallel-group design or (2) cluster design.
- *Type of analysis* used by the authors to test the effect of the intervention categorized as follows: (1) Per-Protocol analysis (PP) where participants were excluded from final analysis due to low adherence rate and that may introduce a type 1 error; (2) Intention-To-Treat analysis (ITT) where missing data are replaced with different imputation methods and that increase the likelihood of type II errors; (3) analyses where all available data were considered, such as Complete-Case analysis (CC) where participants with missing data were excluded from final analysis, ITT where different sample sizes at baseline and at the end of the intervention were considered for final analysis, or analysis of RCTs with no missing data

- *Dropout rate* as the number of participants who were either excluded from the final analysis due to low adherence rates (PP) or dropped out during the study (CC and ITT) for each exercise and control group arm.
- Type of control group categorized as either (1) passive or (2) active. Passive control groups can be further subdivided to (1a) usual care / treatment as usual, (1b) habitual activities, (1c) waiting list. Active control groups can be further subdivided into (2a) health education, (2b) non-physical leisure activities, (2c) stretching / very light exercise / relaxation, (2d) social interaction.

Individual characteristics

Concerning individual characteristics, we will extract information on the following moderator variables:

- *Age* assessed in years.
- *Sex* assessed as percentage of women.
- *Level of education* at baseline assessed as years of education.
- *Functional capacity* at baseline assessed with the Hand Grip Strength Test and the 6-min walk test. We acknowledge the existence of additional suitable indicators for functional capacity, such as VO_2 max. However, in the process of selecting the individual characteristic variables, we followed a similar approach as for the secondary outcomes, selecting only those where enough studies (more than three out of a total of approximately 100 studies) reported a valid measure for a given characteristic.
- *Global cognition* at baseline assessed with the Mini Mental State Examination (MMSE) [99] or the Montreal Cognitive Assessment (MoCa) [100].
- *Population* categorized as follows: (1) neurodegenerative diseases (e.g., Alzheimer disease, Parkinson disease, dementia); (2) cardiovascular diseases (e.g., heart failures, strokes, hypertension); (3) other diseases (e.g., osteoarthritis, osteoporosis, diabetes, cancer, hemodialysis); (4) DSM diagnosis of depressive disorders (e.g., minor or major depression); (5) cognitive impairments (e.g., mild cognitive impairment, cognitive frailty, major neurocognitive disorder); (6) other disabilities (visual impairment, frailty, wheelchair); and (7) no specific disease / disorder / disability (e.g., healthy older adults, sleep complaints, pre-frailty, persons with high risk of falls).

Data analysis

Outcome measurement. For continuous data the standardized mean difference (SMD) as Cohen's *d* and Hedges' *g* together with the 95% confidence interval (CI) will be calculated. If dichotomous data is obtained, the risk ratio (RR) will be calculated with a 95% CI.

Dealing with missing data. If data or statistics (e.g., standard deviations) are missing, we will contact the authors. If we are unable to obtain the missing information, the missing statistics will be handled according to the Cochrane guidelines [101] by analyzing only the available data (i.e., ignoring the missing data). Missing data due to study dropout will be considered as part of the risk of bias assessment, noted in the summary table and considered in the discussion of the results.

Assessment of heterogeneity. We will visually inspect confidence intervals on forest plots for the degree of overlap to determine the potential direction and magnitude of heterogeneity.

We will quantify heterogeneity using Q (also known as χ^2 , or Chi^2) and I^2 ($[(Q - df) / Q] * 100\% = \text{variance}_{\text{between}} / \text{variance}_{\text{total}} * 100\%$) [101]. Q describes whether observed differences in results are compatible with chance alone (without considering the power of the specific studies). I^2 describes the percentage of the variability in effect estimates that is due to heterogeneity rather than chance and is a measure for the impact of the observed heterogeneity on the meta-analysis.

Assessment of reporting publication bias. To assess publication bias, we will visually inspect the funnel plot for asymmetry and apply statistical tests such as Egger's test [102]. If publication bias is detected, a sensitivity analysis will be performed to check whether the results are robust to this bias [70]. Other publication bias methods will be also used [70].

Assessment of evidence and study quality. Quality / certainty of evidence will be assessed with the GRADE framework (Grading of Recommendations, Assessment, Development, and Evaluations [103]) and the Cochrane Collaboration risk-of-bias tool for RCTs (RoB2) [104, 105]. Pairs of two reviewers will be involved in the quality assessment. The two reviewers will independently rate the quality of the included studies with RoB2. The GRADE framework will be used to assess the evidence accumulated for each outcome overall and for the specific domains: 1) risk of bias, 2) imprecision, 3) inconsistency, 4) indirectness, 5) publication bias. Disagreements will be resolved by a third reviewer.

Data synthesis. We aim to calculate summary effects for the outcomes. As it is expected that there will be considerable heterogeneity, a random-effects model will be used for the meta-analysis. Summary effects will be calculated as Hedges' g using random-effects weights. If data cannot be pooled between intervention studies (e.g., an insufficient number of RCTs investigating the respective outcome) findings will be reported in narrative syntheses for each outcome variable. All analyses will be performed using the software Comprehensive Meta-Analysis 4.0 [106] and R [107].

Subgroup analysis and meta-regression. One of the present review's hypothesis is that chronic exercise interventions differ in their effectiveness in reducing overall depression severity and the different symptoms of depression in older adults across participants with different individual's characteristics (i.e., age, sex, education, functional capacity, global cognition, population) and across interventions with different exercise characteristics (i.e., frequency, intensity, duration, type of exercise, cognitive demand, social interactions, exercise supervision, behavioral change techniques, compliance, study design, dropout-rate, type of control group). As such, to determine how different individual's and exercise characteristics / variables play a role, we will conduct subgroup analysis for categorical variables and meta-regression for continuous variables.

Sensitivity analysis. We plan to carry out sensitivity analysis for the primary outcomes to assess the extent to which the results are robust to methodological assumptions and decisions that were made when carrying out the synthesis, such as (1) the effect of study quality, (2) the effects of fixed vs. random effects models, (3) the effects of potential publication bias.

Ethics and dissemination

The present study will use published data and does not require ethics approval.

Discussion and conclusion

In this article, we have outlined a protocol for a systematic and meta-analytical review on the effects of chronic exercise on depression in older adults that is consistent with current best practice guidelines. Any deviations from this protocol will be indicated in the final manuscript. Dissemination will be through presentations at national and international conferences and through a robust peer-reviewed publication.




Fig 2. Working model of the role of intervention characteristics and individual characteristics in exercise interventions effects on late-life depression.

<https://doi.org/10.1371/journal.pone.0297348.g002>

Our meta-analytical review will be the first that examines the effects of chronic exercise on different symptoms of depression in older adults and as a function of a large variety of moderators depending on intervention characteristics and individual characteristics. In line with the already published meta-analyses, we expect to find a moderate effect of exercise on depression compared to active and passive control groups with no exercise components [49]. We further expect the results to be moderated by intervention characteristics and individual characteristics (Fig 2).

The present review aims to make significant contributions to both research and clinical practice. Firstly, the conclusions drawn from this review may serve as a basis for developing evidence-based recommendations for individualized exercise programs that can help alleviate depression in older adults. For instance, it can help determine the exercise regimen in terms of the FITT principles. Secondly, the review results may reveal research gaps that can stimulate the formulation of research questions for further RCTs. For instance, it can reveal whether certain moderator variables, such as comorbidities or physical fitness level still require further research to understand their influence. Finally, this review aims to enhance the understanding of the effects of chronic exercise on various health aspects in older adults within the "Network on evidence-based physical activity in old age" (<https://physagenet.eu/>). In this network, several reviews on the impact of chronic exercise on different health aspects in older age will be carried out and the findings will be compared and consolidated. By aligning research in this network on similar protocols and assessment tools, gaps in terms of generalizability in the large body of research on physical exercise in old age will be filled.

Besides the above-mentioned strengths, the study has some limitations as well. Treatment of depression requires a multidimensional approach including pharmacological treatment and psychotherapy. We plan to exclude RCTs with tested effects of pharmacological treatment and / or psychotherapy. Therefore, no conclusions can be drawn for exercise interventions as add-on treatment. We are aware that by that we might miss some clinically relevant insights. We chose this approach to extract the "pure" impact of chronic exercise on depression in old age and to obtain results comparable to reviews of other aspects of health in old age that will be

conducted in the “Network on evidence-based physical activity in old age” [108]. We further anticipate limitations in form of heterogeneous data and a wide variance in the reporting of the exercise characteristics (e.g., detailedness) and the individual characteristics. We aim to deal with those limitations by being cautious in interpreting the results.

In summary we believe that our review can, on the one hand, contribute to clinical praxis and research on chronic exercise in late-life depression and, on the other hand, contribute to the understanding of the influence of chronic exercise on various aspects of health in old age.

Supporting information

S1 Checklist. PRISMA-P 2015 checklist.
(DOCX)

S1 File. Search strategy.
(PDF)

Acknowledgments

We would like to acknowledge the support and resources provided by the all members of the EU COST Action CA20104—Network on evidence-based physical activity in old age (PhysAgeNet), whose collaboration is crucial for this study. The complete membership list of the consortium is available here: <https://www.cost.eu/actions/CA20104/#tabs+Name:Working%20Groups%20and%20Membership>.

Author Contributions

Conceptualization: Melanie Mack, Christoforos D. Giannaki, Antonia Kaltsatou, Claudia Voelcker-Rehage, Michel Audiffren.

Data curation: Melanie Mack, Andreea Badache, Arzu Erden, Christoforos D. Giannaki, Sandra Haider, Antonia Kaltsatou, Burcu Kömürçü Akik, Yaël Netz, Iuliia Pavlova, Pinelopi S. Stavrinou, Claudia Voelcker-Rehage, Michel Audiffren.

Funding acquisition: Yaël Netz.

Investigation: Melanie Mack, Andreea Badache, Arzu Erden, Christoforos D. Giannaki, Sandra Haider, Antonia Kaltsatou, Burcu Kömürçü Akik, Yaël Netz, Iuliia Pavlova, Pinelopi S. Stavrinou, Claudia Voelcker-Rehage, Michel Audiffren.

Methodology: Melanie Mack, Antonia Kaltsatou, Michel Audiffren.

Project administration: Melanie Mack, Christoforos D. Giannaki, Claudia Voelcker-Rehage, Michel Audiffren.

Software: Melanie Mack, Claudia Voelcker-Rehage, Michel Audiffren.

Supervision: Melanie Mack, Christoforos D. Giannaki, Claudia Voelcker-Rehage, Michel Audiffren.

Validation: Melanie Mack, Christoforos D. Giannaki, Claudia Voelcker-Rehage, Michel Audiffren.

Visualization: Melanie Mack, Michel Audiffren.

Writing – original draft: Melanie Mack.

Writing – review & editing: Andreea Badache, Arzu Erden, Christoforos D. Giannaki, Sandra Haider, Antonia Kaltsatou, Burcu Kömürçü Akik, Yaël Netz, Iuliia Pavlova, Pinelopi S. Stavrinou, Claudia Voelcker-Rehage, Michel Audiffren.

References

1. Zenebe Y, Akele B, Selassie M, Necho M. Prevalence and determinants of depression among old age: A systematic review and meta-analysis. *Ann Gen Psychiatry*. 2021; 20: 55. <https://doi.org/10.1186/s12991-021-00375-x> PMID: 34922595
2. Luppá M, Sikorski C, Luck T, Ehreke L, Konnopka A, Wiese B, et al. Age- and gender-specific prevalence of depression in latest-life—Systematic review and meta-analysis. *J Affect Disord*. 2012; 136: 212–221. <https://doi.org/10.1016/j.jad.2010.11.033> PMID: 21194754
3. Blazer DG. Depression in late life: review and commentary. *J Gerontol A Biol Sci Med Sci*. 2003; 58: 249–265. <https://doi.org/10.1093/gerona/58.3.m249> PMID: 12634292
4. Blazer DG, Hybels CF. Origins of depression in later life. *Psychol Med*. 2005; 35: 1241–1252. <https://doi.org/10.1017/S0033291705004411> PMID: 16168147
5. Hussenoeder FS, Jentzsch D, Matschinger H, Hinz A, Kilian R, Riedel-Heller SG, et al. Depression and quality of life in old age: A closer look. *Eur J Ageing*. 2021; 18: 75–83. <https://doi.org/10.1007/s10433-020-00573-8> PMID: 33746683
6. Fiske A, Wetherell JL, Gatz M. Depression in older adults. *Annu Rev Clin Psychol*. 2009/03/31. 2009; 5: 363–389. <https://doi.org/10.1146/annurev.clinpsy.032408.153621> PMID: 19327033
7. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. American Psychiatric Association Publishing; 2022. <https://doi.org/10.1176/appi.books.9780890425787> PMID: 38300502
8. World Health Organization(WHO). *The ICD-10 classification of mental and behavioural disorders*. Genève, Switzerland: World Health Organization; 1993.
9. Alexopoulos GS. Depression in the elderly. *The Lancet*. 2005; 365: 1961–1970. [https://doi.org/10.1016/S0140-6736\(05\)66665-2](https://doi.org/10.1016/S0140-6736(05)66665-2) PMID: 15936426
10. Lockwood KA, Alexopoulos GS, van Gorp WG. Executive dysfunction in geriatric depression. *Am J Psychiatry*. 2002; 159: 1119–26. <https://doi.org/10.1176/appi.ajp.159.7.1119> PMID: 12091189
11. Elderkin-Thompson V, Kumar A, Bilker WB, Dunkin JJ, Mintz J, Moberg PJ, et al. Neuropsychological deficits among patients with late-onset minor and major depression. *Archives of Clinical Neuropsychology*. 2003; 18: 529–549. [https://doi.org/10.1016/s0887-6177\(03\)00022-2](https://doi.org/10.1016/s0887-6177(03)00022-2) PMID: 14591448
12. Brown ES, Varghese FP, McEwen BS. Association of depression with medical illness: Does cortisol play a role? *Biol Psychiatry*. 2004; 55: 1–9. [https://doi.org/10.1016/s0006-3223\(03\)00473-6](https://doi.org/10.1016/s0006-3223(03)00473-6) PMID: 14706419
13. Alexopoulos GS, Buckwalter K, Olin J, Martinez R, Wainscott C, Krishnan KRR. Comorbidity of late life depression: An opportunity for research on mechanisms and treatment. *Biol Psychiatry*. 2002; 52: 543–558. [https://doi.org/10.1016/s0006-3223\(02\)01468-3](https://doi.org/10.1016/s0006-3223(02)01468-3) PMID: 12361668
14. Carney RM, Freedland KE. Depression, mortality, and medical morbidity in patients with coronary heart disease. *Biol Psychiatry*. 2003; 54: 241–247. [https://doi.org/10.1016/s0006-3223\(03\)00111-2](https://doi.org/10.1016/s0006-3223(03)00111-2) PMID: 12893100
15. Alexopoulos GS. Mechanisms and treatment of late-life depression. *Transl Psychiatry*. 2019; 9: 1–16. <https://doi.org/10.1038/s41398-019-0514-6> PMID: 31383842
16. Alexopoulos GS, Meyers BS, Young RC, Campbell S, Silbersweig D, Charlson M. “Vascular Depression” Hypothesis. *Arch Gen Psychiatry*. 1997; 54: 915–922. <https://doi.org/10.1001/archpsyc.1997.01830220033006> PMID: 9337771
17. Williamson W, Lewandowski AJ, Forkert ND, Griffanti L, Okell TW, Betts J, et al. Association of Cardiovascular Risk Factors With MRI Indices of Cerebrovascular Structure and Function and White Matter Hyperintensities in Young Adults. *JAMA*. 2018; 320: 665–673. <https://doi.org/10.1001/jama.2018.11498> PMID: 30140877
18. Potter GG, Steffens DC. Contribution of depression to cognitive impairment and dementia in older adults. *Neurologist*. 2007; 13: 105–117. <https://doi.org/10.1097/01.nrl.0000252947.15389.a9> PMID: 17495754
19. Alexopoulos GS, Young RC, Meyers BS. Geriatric depression: age of onset and dementia. *Biol Psychiatry*. 1993; 34: 141–145. [https://doi.org/10.1016/0006-3223\(93\)90383-o](https://doi.org/10.1016/0006-3223(93)90383-o) PMID: 8399805
20. Schweitzer I, Tuckwell V, O'Brien J, Ames D. Is late onset depression a prodrome to dementia? *Int J Geriatr Psychiatry*. 2002; 17: 997–1005. <https://doi.org/10.1002/gps.525> PMID: 12404648

21. Kok RM, Reynolds CF. Management of depression in older adults. *JAMA*. 2017; 317: 2114. <https://doi.org/10.1001/jama.2017.5706> PMID: 28535241
22. Barry LC, Abou JJ, Simen AA, Gill TM. Under-treatment of depression in older persons. *J Affect Disord*. 2012; 136: 789–796. <https://doi.org/10.1016/j.jad.2011.09.038> PMID: 22030136
23. Wang PS, Schneeweiss S, Brookhart MA, Glynn RJ, Mogun H, Patrick AR, et al. Suboptimal antidepressant use in the elderly. *J Clin Psychopharmacol*. 2005; 25: 118–126. <https://doi.org/10.1097/01.jcp.0000155819.67209.e5> PMID: 15738742
24. American College of Sports Medicine. ACSM's guidelines for exercise testing and prescription. Philadelphia: Lippincott Williams & Wilkins; 2013.
25. Davidson KW, Goldstein M, Kaplan RM, Kaufmann PG, Knatterud GL, Orleans CT, et al. Evidence-based behavioral medicine: What is it and how do we achieve it? *Annals of Behavioral Medicine*. 2003; 26: 161–171. https://doi.org/10.1207/S15324796ABM2603_01 PMID: 14644692
26. Sackett DL, Rosenberg WMC, Gray JAM, Haynes RB, Richardson WS. Evidence based medicine: What it is and what it isn't. *Br Med J*. 1996; 312: 71. <https://doi.org/10.1136/bmj.312.7023.71> PMID: 8555924
27. Boutron I, Moher D, Altman DG, Schulz KF, Ravaud P. Extending the CONSORT statement to randomized trials of nonpharmacologic treatment: Explanation and elaboration. *Ann Intern Med*. 2008; 148: 295. <https://doi.org/10.7326/0003-4819-148-4-200802190-00008> PMID: 18283207
28. Luan X, Tian X, Zhang H, Huang R, Li N, Chen P, et al. Exercise as a prescription for patients with various diseases. *J Sport Health Sci*. 2019; 8: 422–441. <https://doi.org/10.1016/j.jshs.2019.04.002> PMID: 31534817
29. Hecksteden A, Faude O, Meyer T, Donath L. How to construct, conduct and analyze an exercise training study? *Front Physiol*. 2018; 9. <https://doi.org/10.3389/fphys.2018.01007> PMID: 30140237
30. Park SH, Han KS, Kang CB. Effects of exercise programs on depressive symptoms, quality of life, and self-esteem in older people: a systematic review of randomized controlled trials. *Appl Nurs Res*. 2014; 27: 219–226. <https://doi.org/10.1016/j.apnr.2014.01.004> PMID: 24602398
31. De Sousa RAL, Rocha-Dias I, de Oliveira LRS, Improta-Caria AC, Monteiro-Junior RS, Cassilhas RC. Molecular mechanisms of physical exercise on depression in the elderly: A systematic review. *Mol Biol Rep*. 2021; 48: 3853–3862. <https://doi.org/10.1007/s11033-021-06330-z> PMID: 33864590
32. Zhao JL, Jiang WT, Wang X, Cai ZD, Liu ZH, Liu GR. Exercise, brain plasticity, and depression. *CNS Neurosci Ther*. 2020; 26: 885–895. <https://doi.org/10.1111/cns.13385> PMID: 32491278
33. Netz Y. Is there a preferred mode of exercise for cognition enhancement in older age?—A narrative review. *Front Med (Lausanne)*. 2019; 6: 57. <https://doi.org/10.3389/fmed.2019.00057> PMID: 30984760
34. Voelcker-Rehage C, Niemann C. Structural and functional brain changes related to different types of physical activity across the life span. *Neurosci Biobehav Rev*. 2013; 37: 2268–2295. <https://doi.org/10.1016/j.neubiorev.2013.01.028> PMID: 23399048
35. Stanton R, Reaburn P. Exercise and the treatment of depression: A review of the exercise program variables. *J Sci Med Sport*. 2014; 17: 177–182. <https://doi.org/10.1016/j.jsams.2013.03.010> PMID: 23602562
36. Nebiker L, Lichtenstein E, Minghetti A, Zahner L, Gerber M, Faude O, et al. Moderating effects of exercise duration and intensity in neuromuscular vs. endurance exercise interventions for the treatment of depression: A meta-analytical review. *Front Psychiatry*. 2018; 9: 305. <https://doi.org/10.3389/fpsy.2018.00305> PMID: 30072923
37. Chin EC, Yu AP, Leung CK, Bernal JD, Au WW, Fong DY, et al. Effects of exercise frequency and intensity on reducing depressive symptoms in older adults with insomnia: A pilot randomized controlled trial. *Front Physiol*. 2022; 13: 468. <https://doi.org/10.3389/fphys.2022.863457> PMID: 35450161
38. Parlebas P. Contribution à un lexique commenté en science de l'action motrice. Paris: INSEP; 1981.
39. Lee SL, Pearce E, Ajnakina O, Johnson S, Lewis G, Mann F, et al. The association between loneliness and depressive symptoms among adults aged 50 years and older: A 12-year population-based cohort study. *Lancet Psychiatry*. 2021; 8: 48–57. [https://doi.org/10.1016/S2215-0366\(20\)30383-7](https://doi.org/10.1016/S2215-0366(20)30383-7) PMID: 33181096
40. Cruwys T, South EI, Greenaway KH, Haslam SA. Social identity reduces depression by fostering positive attributions. *Soc Psychol Personal Sci*. 2015; 6: 65–74. <https://doi.org/10.1177/1948550614543309>
41. Stevens M, Lieschke J, Cruwys T, Cárdenas D, Platow MJ, Reynolds KJ. Better together: How group-based physical activity protects against depression. *Soc Sci Med*. 2021; 286: 114337. <https://doi.org/10.1016/j.socscimed.2021.114337> PMID: 34450391

42. Michie S, Richardson M, Johnston M, Abraham C, Francis J, Hardeman W, et al. The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: Building an international consensus for the reporting of behavior change interventions. *Annals of Behavioral Medicine*. 2013; 46: 81–95. <https://doi.org/10.1007/s12160-013-9486-6> PMID: 23512568
43. Jayasekara R, Procter N, Harrison J, Skelton K, Hampel S, Draper R, et al. Cognitive behavioural therapy for older adults with depression: A review. *Journal of Mental Health*. 2015; 24: 168–171. <https://doi.org/10.3109/09638237.2014.971143> PMID: 25358075
44. Kinser PA, Robins JL. Control group design: Enhancing rigor in research of mind-body therapies for depression. *Evid Based Complement Alternat Med*. 2013; 2013. <https://doi.org/10.1155/2013/140467> PMID: 23662111
45. Beck A, Steer R, Brown G. Beck Depression Inventory, 2nd edition. Manual. San Antonio: The Psychological Cooperation; 1996. <https://doi.org/10.1037/t00742-000>
46. Yesavage JA, Sheikh JI. 9/Geriatric Depression Scale (GDS). *Clin Gerontol*. 1986; 5: 165–173. https://doi.org/10.1300/J018v05n01_09
47. Schuch FB, Dunn AL, Kanitz AC, Delevatti RS, Fleck MP. Moderators of response in exercise treatment for depression: A systematic review. *J Affect Disord*. 2016; 195: 40–49. <https://doi.org/10.1016/j.jad.2016.01.014> PMID: 26854964
48. Ansdell P, Thomas K, Hicks KM, Hunter SK, Howatson G, Goodall S. Physiological sex differences affect the integrative response to exercise: acute and chronic implications. *Exp Physiol*. 2020; 105: 2007–2021. <https://doi.org/10.1113/EP088548> PMID: 33002256
49. Bigarella LG, Ballotin VR, Mazurkiewicz LF, Ballardini AC, Rech DL, Bigarella RL, et al. Exercise for depression and depressive symptoms in older adults: An umbrella review of systematic reviews and meta-analyses. *Aging Ment Health*. 2021; 1–11. <https://doi.org/10.1080/13607863.2021.1951660> PMID: 34328049
50. Catalan-Matamoros D, Gomez-Conesa A, Stubbs B, Vancampfort D. Exercise improves depressive symptoms in older adults: An umbrella review of systematic reviews and meta-analyses. *Psychiatry Res*. 2016; 244: 202–209. <https://doi.org/10.1016/j.psychres.2016.07.028> PMID: 27494042
51. Bo A, Mao W, Lindsey MA. Effects of mind-body interventions on depressive symptoms among older Chinese adults: A systematic review and meta-analysis. *Int J Geriatr Psychiatry*. 2017; 32: 509–521. <https://doi.org/10.1002/gps.4688> PMID: 28220964
52. Bridle C, Spanjers K, Patel S, Atherton NM, Lamb SE. Effect of exercise on depression severity in older people: Systematic review and meta-analysis of randomised controlled trials. *British Journal of Psychiatry*. 2018/01/02. 2012; 201: 180–185. <https://doi.org/10.1192/bjp.bp.111.095174> PMID: 22945926
53. Heinzel S, Lawrence JB, Kallies G, Rapp MA, Heissel A. Using exercise to fight depression in older adults. *The Journal of Gerontopsychology and Geriatric Psychiatry*. 2015; 28: 149–162. <https://doi.org/10.1024/1662-9647/a000133>
54. Miller KJ, Gonçalves-Bradley DC, Areerob P, Hennessy D, Mesagno C, Grace F. Comparative effectiveness of three exercise types to treat clinical depression in older adults: A systematic review and network meta-analysis of randomised controlled trials. *Ageing Res Rev*. 2020; 58: 100999. <https://doi.org/10.1016/j.arr.2019.100999> PMID: 31837462
55. Pérez-López FR, Martínez-Domínguez SJ, Lajusticia H, Chedraui P. Effects of programmed exercise on depressive symptoms in midlife and older women: A meta-analysis of randomized controlled trials. *Maturitas*. 2017; 106: 38–47. <https://doi.org/10.1016/j.maturitas.2017.09.001> PMID: 29150165
56. Rhyner KT, Watts A. Exercise and depressive symptoms in older adults: A systematic meta-analytic review. *J Aging Phys Act*. 2016; 24: 234–246. <https://doi.org/10.1123/japa.2015-0146> PMID: 26372020
57. Schuch FB, Vancampfort D, Rosenbaum S, Richards J, Ward PB, Veronese N, et al. Exercise for depression in older adults: A meta-analysis of randomized controlled trials adjusting for publication bias. *Brazilian Journal of Psychiatry*. 2016; 38: 247–254. <https://doi.org/10.1590/1516-4446-2016-1915> PMID: 27611903
58. Tu RH, Zeng ZY, Zhong GQ, Wu WF, Lu YJ, Bo ZD, et al. Effects of exercise training on depression in patients with heart failure: A systematic review and meta-analysis of randomized controlled trials. *Eur J Heart Fail*. 2014; 16: 749–757. <https://doi.org/10.1002/ehf.101> PMID: 24797230
59. Wu PL, Lee M, Wu SL, Ho HH, Chang MH, Lin HS, et al. Effects of home-based exercise on motor, non-motor symptoms and health-related quality of life in Parkinson's disease patients: A randomized controlled trial. *Japan Journal of Nursing Science*. 2021; 18. <https://doi.org/10.1111/jjns.12418> PMID: 33876562

60. Yi D, Yim J. Remote home-based exercise program to improve the mental state, balance, and physical function and prevent falls in adults aged 65 years and older during the COVID-19 pandemic in Seoul, Korea. *Medical Science Monitor*. 2021; 27. <https://doi.org/10.12659/MSM.935496> PMID: 34930888
61. Yu DSF, Lee DTF, Woo J, Hui E. Non-pharmacological interventions in older people with heart failure: Effects of exercise training and relaxation therapy. *Gerontology*. 2007; 53: 74–81. <https://doi.org/10.1159/000096427> PMID: 17057393
62. Klil-Drori S, Klil-Drori AJ, Pira S, Rej S. Exercise intervention for late-life depression: A meta-analysis. *J Clin Psychiatry*. 2020; 81: 6914. <https://doi.org/10.4088/JCP.19r12877> PMID: 31967748
63. Dauwan M, Begemann MJH, Slot MIE, Lee EHM, Scheltens P, Sommer IEC. Physical exercise improves quality of life, depressive symptoms, and cognition across chronic brain disorders: a transdiagnostic systematic review and meta-analysis of randomized controlled trials. *J Neurol*. 2021; 268: 1222–1246. <https://doi.org/10.1007/s00415-019-09493-9> PMID: 31414194
64. Li X, Guo R, Wei Z, Jia J, Wei C. Effectiveness of exercise programs on patients with dementia: A systematic review and meta-analysis of randomized controlled trials. *Biomed Res Int*. 2019;2308475. <https://doi.org/10.1155/2019/2308475> PMID: 31886182
65. Voelcker-Rehage C, Godde B, Staudinger UM. Physical and motor fitness are both related to cognition in old age. *European Journal of Neuroscience*. 2010; 31: 167–176. <https://doi.org/10.1111/j.1460-9568.2009.07014.x> PMID: 20092563
66. Fissler P, Küster O, Schlee W, Kolassa IT. Novelty interventions to enhance broad cognitive abilities and prevent dementia: Synergistic approaches for the facilitation of positive plastic change. *Prog Brain Res*. 2013; 207: 403–434. <https://doi.org/10.1016/B978-0-444-63327-9.00017-5> PMID: 24309264
67. Shea BJ, Reeves BC, Wells G, Thuku M, Hamel C, Moran J, et al. AMSTAR 2: A critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ*. 2017; 358. <https://doi.org/10.1136/bmj.j4008> PMID: 28935701
68. Netz Y. Is the comparison between exercise and pharmacologic treatment of depression in the clinical practice guideline of the American College of Physicians evidence-based? *Front Pharmacol*. 2017; 8. <https://doi.org/10.3389/fphar.2017.00257> PMID: 28555108
69. Cuijpers P, Smit F, Bohlmeijer E, Hollon SD, Andersson G. Efficacy of cognitive-behavioural therapy and other psychological treatments for adult depression: Meta-analytic study of publication bias. *The British Journal of Psychiatry*. 2010; 196: 173–178. <https://doi.org/10.1192/bjp.bp.109.066001> PMID: 20194536
70. Carter EC, Schönbrodt FD, Gervais WM, Hilgard J. Correcting for bias in psychology: A comparison of meta-analytic methods. *Adv Methods Pract Psychol Sci*. 2019; 2: 115–144. <https://doi.org/10.1177/2515245919847196>
71. Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: Elaboration and explanation. *BMJ*. 2015;349. <https://doi.org/10.1136/bmj.g7647> PMID: 25555855
72. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for systematic reviews. *Syst Rev*. 2016; 5: 210. <https://doi.org/10.1186/s13643-016-0384-4> PMID: 27919275
73. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—A metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009; 42: 377–381. <https://doi.org/10.1016/j.jbi.2008.08.010> PMID: 18929686
74. Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O'Neal L, et al. The REDCap consortium: Building an international community of software platform partners. *J Biomed Inform*. 2019; 95: 103208. <https://doi.org/10.1016/j.jbi.2019.103208> PMID: 31078660
75. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh sleep quality index: A new instrument for psychiatric practice and research. *Psychiatry Res*. 1989; 28: 193–213. [https://doi.org/10.1016/0165-1781\(89\)90047-4](https://doi.org/10.1016/0165-1781(89)90047-4) PMID: 2748771
76. Bastien CH, Vallières A, Morin CM. Validation of the Insomnia Severity Index as an outcome measure for insomnia research. *Sleep Med*. 2001; 2: 297–307. [https://doi.org/10.1016/s1389-9457\(00\)00065-4](https://doi.org/10.1016/s1389-9457(00)00065-4) PMID: 11438246
77. Krupp LB, Larocca NG, Muir Nash J, Steinberg AD. The Fatigue Severity Scale: Application to patients With multiple sclerosis and systemic lupus erythematosus. *Arch Neurol*. 1989; 46: 1121–1123. <https://doi.org/10.1001/ARCHNEUR.1989.00520460115022> PMID: 2803071
78. Nilsson MH, Bladh S, Hagell P. Fatigue in Parkinson's disease: Measurement properties of a generic and a condition-specific rating scale. *J Pain Symptom Manage*. 2013; 46: 737–746. <https://doi.org/10.1016/j.jpainsymman.2012.11.004> PMID: 23507131

79. Smets EMA, Garssen B, Bonke B, De Haes JCJM. The multidimensional Fatigue Inventory (MFI) psychometric qualities of an instrument to assess fatigue. *J Psychosom Res.* 1995; 39: 315–325. [https://doi.org/10.1016/0022-3999\(94\)00125-o](https://doi.org/10.1016/0022-3999(94)00125-o) PMID: 7636775
80. Cella D, Yount S, Sorensen M, Chartash E, Sengupta N, Grober J. Validation of the Functional Assessment of Chronic Illness Therapy Fatigue Scale relative to other instrumentation in patients with rheumatoid arthritis. *J Rheumatol.* 2005; 32: 811–9. Available: <http://www.ncbi.nlm.nih.gov/pubmed/15868614> PMID: 15868614
81. Beck AT, Epstein N, Brown G, Steer RA. An inventory for measuring clinical anxiety: Psychometric properties. *J Consult Clin Psychol.* 1988; 56: 893–897. <https://doi.org/10.1037//0022-006x.56.6.893> PMID: 3204199
82. Spitzer RL, Kroenke K, Williams JBW, Löwe B. A brief measure for assessing generalized anxiety disorder: The GAD-7. *Arch Intern Med.* 2006; 166: 1092–1097. <https://doi.org/10.1001/archinte.166.10.1092> PMID: 16717171
83. Hamilton M. The assessment of anxiety states by rating. *British Journal of Medical Psychology.* 1959; 32: 50–55. <https://doi.org/10.1111/j.2044-8341.1959.tb00467.x> PMID: 13638508
84. Zigmond AS, Snaith RP. The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand.* 1983; 67: 361–370. <https://doi.org/10.1111/j.1600-0447.1983.tb09716.x> PMID: 6880820
85. Spielberger CD, Gorsuch RL, Lushene R, Vagg P, Jacobs G. Test manual for the State-Trait Anxiety Inventory for adults. Redwood City, California. 1983.
86. Derogatis LR, Unger R. Symptom Checklist-90-Revised. The Corsini Encyclopedia of Psychology. Wiley; 2010. pp. 1–2. <https://doi.org/10.1002/9780470479216.corpsy0970>
87. Tappen RM, Williams CL. Development and testing of the Alzheimer's disease and related dementias mood scale. *Nurs Res.* 2008; 57: 426–435. <https://doi.org/10.1097/NNR.0b013e31818c3d3c> PMID: 19018217
88. Sunderland T, Alterman IS, Yount D, Hill JL, Tariot PN, Newhouse PA, et al. A new scale for the assessment of depressed mood in demented patients. *American Journal of Psychiatry.* 1988; 145: 955–959. <https://doi.org/10.1176/ajp.145.8.955> PMID: 3394879
89. Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive and negative affect: The PANAS scales. *J Pers Soc Psychol.* 1988; 54: 1063–1070. <https://doi.org/10.1037//0022-3514.54.6.1063> PMID: 3397865
90. McNair DM, Lorr M, Droppleman LF. Manual profile of mood states. San Diego, CA: Educational & Industrial testing service; 1971.
91. Marin RS, Biedrzycki RC, Firinciogullari S. Reliability and validity of the apathy evaluation scale. *Psychiatry Res.* 1991; 38: 143–162. [https://doi.org/10.1016/0165-1781\(91\)90040-v](https://doi.org/10.1016/0165-1781(91)90040-v) PMID: 1754629
92. Cummings JL, Mega M, Gray K, Rosenberg-Thompson S, Carusi DA, Gornbein J. The Neuropsychiatric Inventory. *Neurology.* 1994; 44: 2308–2308. <https://doi.org/10.1212/WNL.44.12.2308> PMID: 7991117
93. Starkstein SE, Jorge R, Mizrahi R, Carver LA. The prevalence, clinical correlates and treatment of apathy in Alzheimer's disease. *Eur J Psychiatry.* 2006; 20: 96–106.
94. Noh JW, Kwon YD, Park J, Kim J. Body mass index and depressive symptoms in middle aged and older adults. *BMC Public Health.* 2015; 15: 1–7. <https://doi.org/10.1186/S12889-015-1663-Z/TABLES/2>
95. Reitan RM. The relation of the Trail Making Test to organic brain damage. *J Consult Psychol.* 1955; 19: 393–394. <https://doi.org/10.1037/h0044509> PMID: 13263471
96. Sohlberg MM, Mateer CA. Effectiveness of an attention-training program. *J Clin Exp Neuropsychol.* 1987; 9: 117–130. <https://doi.org/10.1080/01688638708405352> PMID: 3558744
97. Grant DA, Berg E. A behavioral analysis of degree of reinforcement and ease of shifting to new responses in a Weigl-type card-sorting problem. *J Exp Psychol.* 1948; 38: 404–411. <https://doi.org/10.1037/h0059831> PMID: 18874598
98. Dubois B, Slachevsky A, Litvan I, Pillon B. The FAB: A frontal assessment battery at bedside. *Neurology.* 2000; 55: 1621–1626. <https://doi.org/10.1212/wnl.55.11.1621> PMID: 11113214
99. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state": A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res.* 1975; 12: 189–198.
100. Nasreddine ZS, Phillips NA, Bédirian V, Charbonneau S, Whitehead V, Collin I, et al. The Montreal Cognitive Assessment, MoCA: A brief screening tool for mild cognitive impairment. *J Am Geriatr Soc.* 2005; 53: 695–699. <https://doi.org/10.1111/j.1532-5415.2005.53221.x> PMID: 15817019

101. Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al. Cochrane handbook for systematic reviews of interventions version 6.3. John Wiley & Sons; 2022. Available: www.training.cochrane.org/handbook.
102. Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997; 315: 629–634. <https://doi.org/10.1136/bmj.315.7109.629> PMID: 9310563
103. Guyatt G, Oxman AD, Akl EA, Kunz R, Vist G, Brozek J, et al. GRADE guidelines: 1. Introduction- GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol*. 2011; 64: 383–394. <https://doi.org/10.1016/j.jclinepi.2010.04.026> PMID: 21195583
104. Higgins JP, Sterne JA, Savovic J, Page MJ, Hróbjartsson A, Boutron I, et al. A revised tool for assessing risk of bias in randomized trials. *Cochrane Database of Systematic Reviews*. 2016; 10: 29–31. Available: <https://research.monash.edu/en/publications/a-revised-tool-for-assessing-risk-of-bias-in-randomized-trials>
105. Sterne JAC, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I, et al. RoB 2: A revised tool for assessing risk of bias in randomised trials. *BMJ*. 2019; 366. <https://doi.org/10.1136/bmj.l4898> PMID: 31462531
106. Borenstein M, Hedges L, Higgins J, Rothstein H. *Comprehensive Meta-Analysis Version 4*. Biostat, Englewood, NJ; 2022.
107. R Core Team. *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing; 2020. Available: <https://www.R-project.org/>
108. Brach M, de Bruin ED, Levin O, Hinrichs T, Zijlstra W, Netz Y. Evidence-based yet still challenging! Research on physical activity in old age. *European Review of Aging and Physical Activity*. 2023; 20: 7. <https://doi.org/10.1186/s11556-023-00318-3> PMID: 36932320