



**HAL**  
open science

# Explicit formulae for the mean value of products of values of Dirichlet L-functions at positive integers

Stéphane Louboutin

► **To cite this version:**

Stéphane Louboutin. Explicit formulae for the mean value of products of values of Dirichlet L-functions at positive integers. Proc. Amer. Math. Soc., 2024. hal-04590245

**HAL Id: hal-04590245**

**<https://hal.science/hal-04590245>**

Submitted on 28 May 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Explicit formulae for the mean value of products of values of Dirichlet $L$ -functions at positive integers

Stéphane R. LOUBOUTIN  
Aix Marseille Université, CNRS, Centrale Marseille, I2M,  
Marseille, France  
stephane.louboutin@univ-amu.fr

May 28, 2024

To appear in Proc. Amer. Math. Soc.

## Abstract

Let  $m \geq 1$  be a rational integer. We give an explicit formula for the mean value

$$\frac{2}{\phi(f)} \sum_{\chi(-1)=(-1)^m} |L(m, \chi)|^2,$$

where  $\chi$  ranges over the  $\phi(f)/2$  Dirichlet characters modulo  $f > 2$  with the same parity as  $m$ . We then adapt our proof to obtain explicit means values for products of the form  $L(m_1, \chi_1) \cdots L(m_{n-1}, \chi_{n-1}) \overline{L(m_n, \chi_1 \cdots \chi_{n-1})}$ .

## 1 Introduction

For  $m \geq 1$  and  $f > 2$ , let

$$M(m, f) = \frac{2}{\phi(f)} \sum_{\chi(-1)=(-1)^m} |L(m, \chi)|^2$$

denote the mean value of  $|L(m, \chi)|^2$ , where  $\chi$  ranges over the  $\phi(f)/2$  Dirichlet characters modulo  $f$  such that  $\chi(-1) = (-1)^m$ , i.e. over the Dirichlet characters modulo  $f$  of the same parity as  $m$ .

The first step towards the study of these  $M(m, f)$ 's was made in [Wal], where the formula for  $M(1, f)$  was obtained, but only for prime moduli  $f$ . Secondly, in [QiMG] and [Lou93] the formula for  $M(1, f)$  for non necessarily prime moduli  $f$  was obtained. Thirdly, in [Lou01] a method was developed for obtaining recursively a formula for  $M(m, f)$  for a given  $m$ . This method was applied to obtain the formulae below for  $M(m, f)$  for  $m = 1, 2, 3$  and 4. Fourthly, in [Lin] the author developed another way for obtaining recursively a formula for  $M(m, f)$  for a given  $m$ . S. Louboutin and X. Lin's methods are easy to use with any software for mathematical computation for obtaining a formula for  $M(m, f)$  for a given  $m$ . But they did not give an explicit formula for these  $M(m, f)$ 's. Fifthly, in [LZ] the authors gave a rather complicated but explicit formula for  $M(m, f)$  and the mean value of the product of the values of two Dirichlet  $L$ -functions at positive integers of the same parity.

---

<sup>0</sup>1991 Mathematics Subject Classification. Primary 11M06, 11M20, 33B10.

Key words and phrases. Dirichlet  $L$ -functions, characters, cotangent, mean value.

Here, inspired by [BY, Proof of Theorem 2.1], we obtain in Theorem 1 an explicit formula for these  $M(m, f)$ 's. Then, we extend our method developed to obtain Theorem 1 to get in Theorem 7 an explicit formula for the mean value of more general products of the form  $L(m_1, \chi_1) \cdots L(m_{n-1}, \chi_{n-1}) \overline{L(m_n, \chi_1 \cdots \chi_{n-1})}$ . In particular, for  $n = 2$  we obtain in Theorem 9 a much simpler formula than the one given in [LZ, Theorem 1.1] for the mean value of the product of the values of two Dirichlet  $L$ -functions at positive integers of the same parity.

**Theorem 1.** *For  $m = 1$  we have*

$$M(1, f) = \frac{\pi^2}{6} \left( \phi_2(f) - 3 \frac{\phi_1(f)}{f} \right) \quad (1)$$

and for  $m \geq 2$  we have

$$M(m, f) = \zeta(2m) \times \left( \phi_{2m}(f) + (-1)^{m-1} \binom{2m}{m} \sum_{k=1}^{\lfloor m/2 \rfloor} \frac{m}{m-k} \binom{m}{2k} \frac{B_{2k} B_{2(m-k)}}{B_{2m}} \frac{\phi_{2k}(f)}{f^{2m-2k}} \right),$$

where  $\phi_l(f) = \prod_{p|f} (1 - 1/p^l)$  and the Bernoulli rational numbers  $B_k$  are given by

$$\frac{t}{e^t - 1} = \sum_{k \geq 0} \frac{B_k}{k!} t^k.$$

In particular,  $M(m, f)$  is asymptotic to  $\zeta(2m)\phi_{2m}(f)$  as  $f$  goes to infinity.

For example, in accordance with [Lou93, Theorem 2] and [Lin, Page 371] we have:

$$\begin{aligned} M(2, f) &= \frac{\pi^4}{90} \left( \phi_4(f) + 10 \frac{\phi_2(f)}{f^2} \right), \\ M(3, f) &= \frac{\pi^6}{945} \left( \phi_6(f) - 21 \frac{\phi_2(f)}{f^4} \right), \\ M(4, f) &= \frac{\pi^8}{9450} \left( \phi_8(f) + \frac{14}{3} \frac{\phi_4(f)}{f^4} + \frac{200}{3} \frac{\phi_2(f)}{f^6} \right), \\ M(5, f) &= \frac{\pi^{10}}{93555} \left( \phi_{10}(f) - 22 \frac{\phi_4(f)}{f^6} - 231 \frac{\phi_2(f)}{f^8} \right). \end{aligned}$$

## 2 Proof of Theorem 1

Theorem 1 readily follows from the well known formula

$$\zeta(2m) = (-1)^{m-1} \frac{(2\pi)^{2m} B_{2m}}{2 \cdot (2m)!}. \quad (2)$$

and from following Lemmas 2, 3 and 4.

**Lemma 2.** (See [Lou01, Proposition 3 and Proof of Proposition 5]). *Let  $m \geq 1$  and  $f > 2$  be rational integers. Let  $\cot^{(k)}$  denote the  $k$ th derivative of  $\cot := \cos / \sin$ . We have*

$$M(m, f) = \frac{\pi^{2m}}{2 \cdot ((m-1)!)^2 f^{2m}} \sum_{\substack{d|f \\ d>1}} \mu(f/d) S_m(d),$$

where

$$S_m(d) := \sum_{k=1}^{d-1} \left( \cot^{(m-1)}(\pi k/d) \right)^2 \quad (d > 1).$$

**Lemma 3.** *If  $R(X) = \sum_{l=0}^k r_{k,l} X^l \in \mathbb{Q}[X]$  is such that  $R(1) = 0$ , then for  $f > 2$  we have*

$$\sum_{\substack{d|f \\ d>1}} \mu(f/d) R(d) = \sum_{d|f} \mu(f/d) R(d) = \sum_{l=1}^k r_{k,l} \phi_l(f) f^l.$$

*Proof.* Notice that  $\sum_{d|f} \mu(f/d) d^l = \sum_{d|f} \mu(d) (f/d)^l = f^l \phi_l(f)$ . □

**Lemma 4.** *Set  $R_1(X) = (X^2 - 3X + 2)/3$  and for  $m \geq 2$  set*

$$R_m(X) = 2^{2m} \cdot \left( (-1)^{m-1} \frac{B_{2m}}{m^2 \binom{2m}{m}} X^{2m} + \sum_{k=0}^{\lfloor m/2 \rfloor} \frac{B_{2k} B_{2(m-k)}}{m(m-k)} \binom{m}{2k} X^{2k} - \frac{1 + (-1)^m B_m^2}{2} \frac{B_m^2}{m^2} \right).$$

*Then  $R_m(1) = 0$  for  $m \geq 1$  and  $S_m(d) = R_m(d)$  for  $m \geq 1$  and  $d \geq 2$ .*

*Proof.* For  $d \geq 1$ , set

$$g_m(d, z) = d \cot(dz) \left( \cot^{(m-1)}(z) \right)^2 = \frac{1}{z^{2m+1}} (dz \cot(dz)) (z^m \cot^{(m-1)}(z))^2,$$

a  $\pi$ -periodic meromorphic function of the complex variable  $z$ , with a simple pole at each  $k\pi/d$ ,  $k \in \mathbb{Z} \setminus d\mathbb{Z}$ , of residue

$$\operatorname{Res}_{z=k\pi/d} (g_m(d, z)) = \left( \cot^{(m-1)}(k\pi/d) \right)^2 \quad (k \in \mathbb{Z} \setminus d\mathbb{Z}) \quad (3)$$

and poles of order  $2m + 1$  at each  $k\pi$ ,  $k \in \mathbb{Z}$ . For  $R > 0$ , let  $C_R$  be the positively oriented rectangle of width  $\pi$ , height  $2R$  and center of gravity at  $\pi/2 - \pi/(2d)$ .

(i). We claim that

$$\lim_{R \rightarrow +\infty} \frac{1}{2\pi i} \int_{C_R} g_m(d, z) dz = \begin{cases} -d & \text{if } m = 1 \text{ and } d \geq 1, \\ 0 & \text{if } m \geq 2 \text{ and } d \geq 1. \end{cases} \quad (4)$$

Indeed, since  $g$  is  $\pi$ -periodic the integrals over the vertical sides of  $C_R$  cancel. Moreover, since  $|\cot(t + ib) + i \operatorname{sign}(b)| \leq 2/(e^{|b|} - 1)$  we have  $\lim_{b \rightarrow +\infty} \cot(a + ib) = -i$  and  $\lim_{b \rightarrow +\infty} \cot(a - ib) = i$  uniformly in  $a \in I$ , where  $I$  is a given closed interval. Hence for  $m = 1$  we have  $\lim_{R \rightarrow +\infty} g_1(d, t + iR) = di$  and  $\lim_{R \rightarrow +\infty} g_1(d, t - iR) = -di$ , uniformly in  $t \in [-\pi/(2d), \pi - \pi/(2d)]$ . The first case of claim (4) follows. Now assume that  $m \geq 2$ . Then  $\cot^{(m-1)} = Q_m(\cot)$  where the polynomials  $Q_k(X)$ 's are defined inductively by  $Q_1(X) = X$  and  $Q_{k+1}(X) = -(X^2 + 1)Q_k'(X)$ . Consequently, we have  $\lim_{R \rightarrow +\infty} g_m(d, t + iR) = Q_m(-i) = 0$  and  $\lim_{R \rightarrow +\infty} g_m(d, t - iR) = Q_m(i) = 0$ , uniformly in  $t \in [-\pi/(2d), \pi - \pi/(2d)]$ . Hence the integrals over the horizontal sides tend to 0 as  $R$  goes to infinity for  $m \geq 2$  and the second case of claim (4) follows.

(ii). Now, we claim that

$$R_m(d) = \lim_{R \rightarrow +\infty} \frac{1}{2\pi i} \int_{C_R} g_m(d, z) dz - \operatorname{Res}_{z=0} (g_m(d, z)) \quad (m \geq 1 \text{ and } d \geq 1). \quad (5)$$

Indeed, if  $m = 1$  then  $g_1(d, z) = d \cot(dz) \cot^2 z$ ,

$$\operatorname{Res}_{z=0} (g_1(d, z)) = -(d^2 + 2)/3$$

and the claims follow by (4) and the definition of  $R_1(X)$ . Now assume that  $m \geq 2$ . Using the power series expansion

$$z \cot(z) = \sum_{k \geq 0} (-1)^k \frac{2^{2k} B_{2k}}{(2k)!} z^{2k} \quad (|z| < \pi), \quad (6)$$

we have

$$z^m \cot^{(m-1)}(z) = (-1)^{m-1} (m-1)! + \sum_{k \geq m/2} (-1)^k \frac{2^{2k} B_{2k}}{2k \cdot (2k-m)!} z^{2k} \quad (|z| < \pi). \quad (7)$$

Using (6) and (7) to compute the coefficient of the term  $z^{2m}$  in the power series expansion of the function  $(dz \cot(dz))(z^m \cot^{(m-1)}(z))^2$ , we readily obtain that

$$\text{Res}_{z=0}(g_m(d, z)) = -R_m(d) \text{ for } m \geq 2 \text{ and } d \geq 1.$$

Hence claim (5) follows from (4).

(iii). Finally, we use the residue theorem. For  $d = 1$ , noticing that  $z = 0$  is the only pole of  $g_m(1, z)$  inside  $C_R$  and using (5) we obtain  $R_m(1) = 0$  for  $m \geq 1$ . For  $d \geq 2$ , noticing that  $z = 0$  and  $z = k\pi/d$ ,  $1 \leq k \leq d-1$ , are the only poles of  $g_m(d, z)$  inside  $C_R$  and using (5) and (3) we obtain  $R_m(d) = S_m(d)$  for  $d \geq 2$  and  $m \geq 1$ , as desired.  $\square$

In [EL, Corollary 6.8] the authors gave a reformulation of [BY, Corollary 2.2]. It would be interesting to see if their method can also be applied to a reformulation of our Lemma 4.

### 3 A general explicit formula

For  $f > 2$ ,  $n \geq 2$  and  $m_1, \dots, m_n \geq 1$  such that  $m_1 + \dots + m_n = 2s$  is even, we set  $\vec{m} = (m_1, \dots, m_n)$  and

$$M(\vec{m}, f) = \left( \frac{2}{\phi(f)} \right)^{n-1} \times \sum_{\chi_1(-1)=(-1)^{m_1}} \cdots \sum_{\chi_{n-1}(-1)=(-1)^{m_{n-1}}} L(m_1, \chi_1) \cdots L(m_{n-1}, \chi_{n-1}) \overline{L(m_n, \chi_1 \cdots \chi_{n-1})}.$$

Hence,  $(\chi_1 \cdots \chi_{n-1})(-1) = (-1)^{m_n}$ . Using the formula

$$L(m, \chi) = \frac{(-1)^{m-1} \pi^m}{2 \cdot f^m \cdot (m-1)!} \sum_{k=1}^{f-1} \chi(k) \cot^{(m-1)}(\pi k/f) \quad (\text{whenever } \chi(-1) = (-1)^m).$$

proved in [Lou01, (1)], we have

$$M(\vec{m}, f) = \frac{(-1)^n \pi^{2s}}{2^n f^{2s} \prod_{l=1}^n (m_l - 1)!} \times \sum_{k_n=1}^{f-1} \sum_{k_1=1}^{f-1} \cdots \sum_{k_{n-1}=1}^{f-1} \left( \prod_{l=1}^n \cot^{(m_l-1)}(\pi k_l/f) \right) \left( \prod_{l=1}^{n-1} \frac{2}{\phi(f)} \sum_{\chi_l(-1)=(-1)^{m_l}} \chi_l(k_l) \overline{\chi_l(k_n)} \right).$$

We have the following orthogonality relations for the  $\phi(f)/2$  characters  $\chi_l$ 's modulo  $f > 2$  for which  $\chi_l(-1) = (-1)^{m_l}$ :

$$\frac{2}{\phi(f)} \sum_{\chi_l(-1)=(-1)^{m_l}} \chi_l(k_l) \overline{\chi_l(k_n)} = \begin{cases} 1 & \text{if } k_l \equiv k_n \pmod{f} \text{ and } \gcd(k_n, f) = 1, \\ (-1)^{m_l} & \text{if } k_l \equiv -k_n \pmod{f} \text{ and } \gcd(k_n, f) = 1, \\ 0 & \text{otherwise.} \end{cases}$$

Noticing that  $\chi_l(-k_n) \cot^{(m_l-1)}(-\pi k_n/f) = \chi_l(k_n) \cot^{(m_l-1)}(\pi k_n/f)$  whenever  $\chi_l(-1) = (-1)^{m_l}$ , we obtain

$$M(\vec{m}, f) = \frac{(-1)^n \pi^{2s}}{2 f^{2s} \prod_{l=1}^n (m_l - 1)!} \sum_{\substack{k_n=1 \\ \gcd(k_n, f)=1}}^{f-1} \prod_{l=1}^n \cot^{(m_l-1)}(\pi k_n/f).$$

The following generalization of Lemma 2 follows:

**Lemma 5.** For  $f > 2$ ,  $n \geq 2$  and  $m_1, \dots, m_n \geq 1$  such that  $m_1 + \dots + m_n = 2s$  is even we have

$$M(\vec{m}, f) = \frac{(-1)^n \pi^{2s}}{2 \cdot (m_1 - 1)! \cdots (m_n - 1)! \cdot f^{2s}} \sum_{\substack{d|f \\ d > 1}} \mu(f/d) S_{\vec{m}}(d),$$

where

$$S_{\vec{m}}(d) = \sum_{k=1}^{d-1} \prod_{l=1}^n \cot^{(m_l-1)}\left(\frac{k\pi}{d}\right) \quad (d > 1).$$

Now, for  $d \geq 1$  a rational integer, we set

$$g_{\vec{m}}(d, z) = d \cot(dz) \prod_{l=1}^n \cot^{(m_l-1)}(z) = \frac{(dz \cot(dz)) \prod_{l=1}^n z^{m_l} \cot^{(m_l-1)}(z)}{z^{2s+1}},$$

a  $\pi$ -periodic meromorphic function of the complex variable  $z$ , with a simple pole at each  $k\pi/d$ ,  $k \in \mathbb{Z} \setminus d\mathbb{Z}$ , of residue

$$\operatorname{Res}_{z=k\pi/d}(g_{\vec{m}}(d, z)) = \prod_{l=1}^n \cot^{(m_l-1)}(k\pi/d) \quad (k \in \mathbb{Z} \setminus d\mathbb{Z}) \quad (8)$$

and poles of order  $2s + 1$  at each  $k\pi$ ,  $k \in \mathbb{Z}$ . Let  $T_{\vec{m}}(X)$  denote the coefficient of  $z^{m_1 + \dots + m_n}$  in the formal power series expansion

$$\left( \sum_{k \geq 0} (-1)^k \frac{2^{2k} B_{2k}}{(2k)!} X^{2k} z^{2k} \right) \prod_{l=1}^n \left( (-1)^{m_l-1} (m_l - 1)! + \sum_{k \geq m_l/2} (-1)^k \frac{2^{2k} B_{2k}}{2k \cdot (2k - m_l)!} z^{2k} \right).$$

Hence,  $T_{\vec{m}}(X)$  is a polynomial of degree  $2s = m_1 + \dots + m_n$ , with rational coefficients and leading coefficient

$$(-1)^{s-n} \frac{2^{2s} B_{2s}}{(2s)!} \prod_{l=1}^n (m_l - 1)!$$

and

$$\operatorname{Res}_{z=0}(g_{\vec{m}}(d, z)) = T_{\vec{m}}(d) \text{ for } d \geq 1. \quad (9)$$

With the same proof, we have the following generalization of (4):

$$\lim_{R \rightarrow +\infty} \frac{1}{2\pi i} \int_{C_R} g_{\vec{m}}(d, z) dz = \begin{cases} (-1)^{n/2} d & (\text{if } m_1 = \dots = m_n = 1), \\ 0 & \text{otherwise.} \end{cases} \quad (10)$$

( $n$  must be even if  $m_1 = \dots = m_n = 1$ ).

We have the following generalization of Lemma 4:

**Lemma 6.** Set

$$R_{\vec{m}}(X) = -T_{\vec{m}}(X) + \begin{cases} (-1)^{n/2} X & (\text{if } m_1 = \dots = m_n = 1), \\ 0 & \text{otherwise.} \end{cases}$$

Then  $R_{\vec{m}}(1) = 0$  and  $S_{\vec{m}}(d) = R_{\vec{m}}(d)$  for  $d \geq 2$ .

*Proof.* By (9) and (10), for  $d \geq 1$  we have

$$R_{\vec{m}}(d) := \frac{1}{2\pi i} \int_{C_R} g_{\vec{m}}(d, z) dz - \operatorname{Res}_{z=0}(g_{\vec{m}}(d, z)).$$

We use the residue theorem. For  $d = 1$ , noticing that  $z = 0$  is the only pole of  $g_{\vec{m}}(1, z)$  inside  $C_R$ , here again we get

$$R_{\vec{m}}(1) = 0. \quad (11)$$

For  $d \geq 2$ , noticing that  $z = 0$  and  $z = k\pi/d$ ,  $1 \leq k \leq d - 1$ , are the only poles of  $g_{\vec{m}}(d, z)$  inside  $C_R$  and using (8), we obtain  $R_{\vec{m}}(d) = S_{\vec{m}}(d)$ .  $\square$

By Lemma 6, Lemma 3 and (2), and noticing that  $T_{\vec{m}}(X)$  is even we end up with the following generalization of Theorem 1:

**Theorem 7.** *Write*

$$T_{\vec{m}}(X) = (-1)^{s-n} \frac{2^{2s} B_{2s}}{(2s)!} \left( \prod_{l=1}^n (m_l - 1)! \right) \left( X^{2s} + \sum_{k=0}^{s-1} r_{\vec{m}, 2k} X^{2k} \right),$$

Then, for  $f > 2$  we have

$$M(\vec{m}, f) = \zeta(2s) \times \left( \phi_{2s}(f) + \sum_{k=1}^{s-1} r_{\vec{m}, 2k} \frac{\phi_{2k}(f)}{f^{2s-2k}} \right) + \begin{cases} (-1)^{n/2} \frac{\pi^n \phi_1(f)}{2f^{n-1}} & (\text{if } m_1 = \dots = m_n = 1), \\ 0 & \text{otherwise.} \end{cases}$$

In particular, as  $f$  goes to infinity  $M(\vec{m}, f)$  is asymptotic to

$$\zeta(2s) \phi_{2s}(f), \text{ where } 2s = m_1 + \dots + m_n.$$

## 4 Several explicit examples

### 4.1 First example.

By Lemma 6 we have  $S_{\overrightarrow{(1,1,2)}}(d) = R_{\overrightarrow{(1,1,2)}}(d) = -T_{\overrightarrow{(1,1,2)}}(d)$ . By (9), we obtain

$$S_{\overrightarrow{(1,1,2)}}(d) = -T_{\overrightarrow{(1,1,2)}}(d) = -\text{Res}_{z=0}(d \cot(dz) \cot^2(z) \cot'(z)) = -\frac{1}{45}d^4 + \frac{1}{9}d^2 - \frac{4}{45}.$$

Therefore

$$M(\overrightarrow{(1, 1, 2)}, f) = -\frac{\pi^4}{2f^4} \sum_{d|f} \mu(f/d) S_{\overrightarrow{(1,1,2)}}(d) = \frac{\pi^4}{90} \left( \phi_4(f) - 5 \frac{\phi_2(f)}{f^2} \right),$$

by Lemma 5 and Lemma 3. We have recovered [Alk, Theorem 1].

### 4.2 Second example.

**Theorem 8.** *Let  $n \geq 2$  be even. For  $f > 2$  set*

$$M_n(f) = \left( \frac{2}{\phi(f)} \right)^{n-1} \sum_{\chi_1(-1)=-1} \dots \sum_{\chi_{n-1}(-1)=-1} L(1, \chi_1) \cdots L(1, \chi_{n-1}) \overline{L(1, \chi_1 \cdots \chi_{n-1})}.$$

Then

$$M_n(f) = \zeta(n) \times \left( \phi_n(f) + \frac{n!}{B_n} \sum_{k_0=1}^{n/2-1} C_{n, n/2-k_0} \frac{B_{2k_0}}{(2k_0)!} \frac{\phi_{2k_0}(f)}{f^{n-2k_0}} \right) + (-1)^{n/2} \frac{\pi^n \phi_1(f)}{2f^{n-1}},$$

where

$$C_{n, N} = \sum_{\substack{e_1, \dots, e_N \geq 0 \\ e_1 + 2e_2 + \dots + Ne_N = N}} \frac{n!}{(n - (e_1 + \dots + e_N))!} \prod_{l=1}^N \frac{1}{e_l!} \left( \frac{B_{2l}}{(2l)!} \right)^{e_l} \quad (n \geq N \geq 1).$$

*Proof.* Here

$$T_{\vec{m}}(d) = (-1)^{n/2} \frac{2^n B_n}{n!} \sum_{\substack{k_0, k_1, \dots, k_n \geq 0 \\ k_0 + k_1 + \dots + k_n = n/2}} \frac{n!}{B_n} \left( \prod_{l=1}^n \frac{B_{2k_l}}{(2k_l)!} \right) \frac{B_{2k_0}}{(2k_0)!} d^{2k_0}.$$

Now, for  $n \geq N \geq 1$  and  $F : \mathbb{Z}_{\geq 0} \rightarrow \mathbb{C}^*$  such that  $F(0) = 1$ , e.g. for  $F(k) = B_{2k}/(2k)!$ , we have

$$\sum_{\substack{k_1, \dots, k_n \geq 0 \\ k_1 + \dots + k_n = N}} \prod_{l=1}^n F(k_l) = \sum_{\substack{e_1, \dots, e_N \geq 0 \\ e_1 + 2e_2 + \dots + Ne_N = N}} \frac{n!}{(n - (e_1 + \dots + e_N))!} \prod_{l=1}^N \frac{F(l)^{e_l}}{e_l!},$$

by letting  $e_l$  denote the number of  $k_i$ 's which are equal to  $l$  and by noticing that we have  $\binom{n}{e_0}$  possible choices of the  $k_i$ 's which are equal to 0, then  $\binom{n-e_0}{e_1}$  possible choices of the  $k_i$ 's which are equal to 1, ..., and finally by using  $e_0 + e_1 + \dots + e_N = n$  and

$$\binom{n}{e_0} \binom{n-e_0}{e_1} \dots \binom{n-(e_0+e_1+\dots+e_{N-1})}{e_N} = \frac{n!}{e_0! \times e_1! \times \dots \times e_N!}.$$

□

For example,

$$M_2(f) = \frac{\pi^2}{6} \left( \phi_2(f) - 3 \frac{\phi_1(f)}{f} \right),$$

in accordance with (1),

$$M_4(f) = \frac{\pi^4}{90} \left( \phi_4(f) - 20 \frac{\phi_2(f)}{f^2} + 45 \frac{\phi_1(f)}{f^3} \right)$$

and

$$M_6(f) = \frac{\pi^4}{945} \left( \phi_6(f) - 21 \frac{\phi_4(f)}{f^2} + \frac{483}{2} \frac{\phi_2(f)}{f^4} - \frac{945}{2} \frac{\phi_1(f)}{f^5} \right)$$

For  $f = 3$ , there is only one odd character  $\chi$  modulo 3, and  $L(1, \chi) = \frac{\pi}{3\sqrt{3}}$ . Hence,  $M_n(3) = \frac{\pi^n}{3^{3n/2}}$ . Our formulae do give  $M_2(3) = \frac{\pi^2}{3^3}$ ,  $M_4(3) = \frac{\pi^4}{3^6}$  and  $M_6(3) = \frac{\pi^6}{3^9}$ .

### 4.3 Third example.

We have the following generalization of Theorem 1:

**Theorem 9.** For  $m \geq 1$ ,  $n \geq 1$  of the same parity and  $(m, n) \neq (1, 1)$ , we have

$$\begin{aligned} M(\overrightarrow{(m, n)}, f) &= \frac{2}{\phi(f)} \sum_{\chi(-1)=(-1)^m=(-1)^n} L(m, \chi) \overline{L(n, \chi)} \\ &= \zeta(m+n) \times \left( \phi_{m+n}(f) \right. \\ &\quad + (-1)^{m-1} \binom{m+n}{m} \sum_{k=1}^{\lfloor m/2 \rfloor} \frac{n}{m+n-2k} \binom{m}{2k} \frac{B_{2k} B_{m+n-2k}}{B_{m+n}} \frac{\phi_{2k}(f)}{f^{m+n-2k}} \\ &\quad \left. + (-1)^{n-1} \binom{m+n}{n} \sum_{k=1}^{\lfloor n/2 \rfloor} \frac{m}{m+n-2k} \binom{n}{2k} \frac{B_{2k} B_{m+n-2k}}{B_{m+n}} \frac{\phi_{2k}(f)}{f^{m+n-2k}} \right). \end{aligned}$$

In particular,  $M(\overrightarrow{(m, n)}, f)$  is asymptotic to  $\zeta(m+n)\phi_{m+n}(f)$  as  $f$  goes to infinity.



*Proof.* Use Lemma 6, notice that

$$\begin{aligned} T_{\overrightarrow{(m,n)}}(X) &= 2^{m+n} \times \left( (-1)^{\frac{m+n}{2}} \frac{B_{m+n}}{mn \binom{m+n}{n}} X^{m+n} \right. \\ &\quad - (-1)^{\frac{m-n}{2}} \sum_{k=1}^{\lfloor m/2 \rfloor} \frac{B_{2k} B_{m+n-2k}}{m(m+n-2k)} \binom{m}{2k} X^{2k} \\ &\quad - (-1)^{\frac{n-m}{2}} \sum_{k=1}^{\lfloor n/2 \rfloor} \frac{B_{2k} B_{m+n-2k}}{n(m+n-2k)} \binom{n}{2k} X^{2k} \\ &\quad \left. + (-1)^{\frac{m+n}{2}} \frac{1 + (-1)^m}{2} \frac{1 + (-1)^n}{2} \frac{B_m B_n}{mn} \right). \end{aligned}$$

and finally use Lemma 5. □

Our explicit formula for  $M(\overrightarrow{(m,n)}, f)$  is much simpler than the ones given in [LZ, Theorem 1.1] and [OO, Theorem 1.2]. For example, the first term in the formula given [LZ, Theorem 1.1] is

$$(-1)^{(m-n)/2} \frac{(2\pi)^{m+n}}{2 \cdot m! \cdot n!} \phi_{m+n}(f) \sum_{a=0}^m \sum_{b=0}^n B_{m-a} B_{n-b} \frac{\binom{m}{a} \binom{n}{b}}{a+b+1}.$$

By (2), this first term being equal to the one  $\zeta(m+n)\phi_{m+n}(f)$  of Theorem 9 is tantamount to saying that  $B_{m+n} = (-1)^n \binom{m+n}{n} \sum_{a=0}^m \sum_{b=0}^n B_{m-a} B_{n-b}$ . This is probably correct and at least we checked it for several values of  $m$  and  $n$ .

Notice that for  $m \geq 2$  we have  $M(\overrightarrow{(m,m)}, f) = M(m, f)$  and we recover the formula given in Theorem 1.

#### 4.4 Fourth example.

For  $m, n \geq 1$ , Theorem 7 gives a formula for

$$M(\overrightarrow{(m,n,m+n)}, f) = \frac{4}{\phi(f)^2} \sum_{\chi_1(-1)=(-1)^m} \sum_{\chi_2(-1)=(-1)^n} L(m, \chi_1) L(n, \chi_2) \overline{L(m+n, \chi_1 \chi_2)}$$

much simpler than the one given in [OO, Theorem 1.1].

## References

- [Alk] E. Alkan. Averages of values of  $L$ -series. *Proc. Amer. Math. Soc.* **141** (2013), 1161–1175.
- [BY] Bruce C. Berndt and Boon Pin Yeap. Explicit evaluations and reciprocity theorems for finite trigonometric sums. *Adv. in Appl. Math.* **29** (2002), 358–385.
- [EL] W. Ejsmont and F. Lehner. The trace method for cotangent sums. *J. Combin. Theory Ser. A* **177** (2021), Paper No. 105324, 32 pp.
- [Lin] Xin Lin. On the mean square value of the Dirichlet  $L$ -function at positive integers. *Acta Arith.* **189** (2019), 367–379.
- [Lou93] S. Louboutin. Quelques formules exactes pour des moyennes de fonctions  $L$  de Dirichlet. *Canad. Math. Bull.* **36** (1993), 190–196.
- [Lou94] S. Louboutin. Corrections à : Quelques formules exactes pour des moyennes de fonctions  $L$  de Dirichlet. *Canad. Math. Bull.* **37** (1994), p. 89.
- [Lou01] S. Louboutin. The mean value of  $|L(k, \chi)|^2$  at positive rational integers  $k \geq 1$ . *Colloq. Math.* **90** (2001), 69–76.
- [LZ] H. Liu and W. Zhang. On the mean value of  $L(m, \chi)L(n, \bar{\chi})$  at positive integers  $m, n \geq 1$ . *Acta Arith.* **122** (2006), 51–56.
- [OO] T. Okamoto and T. Onozuka. On various mean values of Dirichlet  $L$ -functions. *Acta Arith.* **167** (2015), 101–115.
- [QiMG] Qi Ming Gao. A kind of mean square formula for  $L$ -functions. (in Chinese) *J. Tsinghua Univ.* **31** (1991), 34–41. See [MR1168609(93g:11090)].
- [Wal] H. Walum. An exact formula for an average of  $L$ -series. *Illinois J. of Math.* **26** (1982), 1–3.