



Mid-Holocene marine faunas from the Bangkok Clay deposits in Nakhon Nayok, the Central Plain of Thailand

Parin Jirapatrasilp, Gilles Cuny, László Kocsis, Chirasak Sutcharit, Nom Ngamnisai, Thasinee Charoentitirat, Satapat Kumpitak, Kantapon Suraprasit

► To cite this version:

Parin Jirapatrasilp, Gilles Cuny, László Kocsis, Chirasak Sutcharit, Nom Ngamnisai, et al.. Mid-Holocene marine faunas from the Bangkok Clay deposits in Nakhon Nayok, the Central Plain of Thailand. *Zookeys*, 2024, 1202, pp.1-110. 10.3897/zookeys.1202.119389 . hal-04590679v1

HAL Id: hal-04590679

<https://hal.science/hal-04590679v1>

Submitted on 28 May 2024 (v1), last revised 30 May 2024 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Mid-Holocene marine faunas from the Bangkok Clay deposits in Nakhon Nayok, the Central Plain of Thailand

Parin Jirapatrasilp^{1,2}, Gilles Cuny³, László Kocsis⁴, Chirasak Sutcharit¹, Nom Ngamnisai⁵, Thasinee Charoentitirat⁶, Satapart Kumpitak⁶, Kantapon Suraprasit⁷

1 Animal Systematics Research Unit, Department of Biology, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand

2 Leibniz-Institut zur Analyse des Biodiversitätswandels - Standort Hamburg, Martin-Luther-King-Platz 3, Hamburg 20146, Germany

3 Universite Claude Bernard Lyon 1, LEHNA UMR 5023, CNRS, ENTPE, F-69622, Villeurbanne, France

4 Institute of Earth Surface Dynamics, University of Lausanne, Rue de la Mouline, 1015 Lausanne, Switzerland

5 Department of Geography, Faculty of Social Sciences, Srinakharinwirot University, Bangkok 10110, Thailand

6 Department of Geology, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand

7 Center of Excellence in Morphology of Earth Surface and Advanced Geohazards in Southeast Asia (MESA CE), Department of Geology, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand

Corresponding author: Kantapon Suraprasit (Kantapon.S@chula.ac.th)

Abstract

The Lower Central Plain of Thailand is an important area where the topography and climate change have been under the influence of sea level rise after the end of the Pleistocene. The area has shown evidence of an ancient sea-level change reaching the mid-Holocene highstand. Based on several field investigations, many molluscan shells and chondrichthyan teeth, together with other invertebrate and actinopterygian remains were found from the marine clay deposits (so called “Bangkok Clay”) at the clay pit in Ongkharak District, Nakhon Nayok Province, at a depth of about 5–7 m below the topsoil surface. In this study, we identify animal macrofossils recovered from these Holocene marine deposits and provide their chronological context, to reconstruct the paleoenvironments and paleoecosystems of the area at that time. The majority of marine fossils recovered from the site consist of molluscs, where a total of 63 species were identified, including 35 species of gastropods from 20 families, 27 species of bivalves from 15 families, and one species of scaphopod. The most common gastropods are *Architectonica perdix*, *Natica stellata*, and *Indothais lacera*, whereas the most common bivalves are *Corbula fortisulcata*, *Magallana* cf. *gigas*, and *Joannisella oblonga*. Other invertebrate remains include one species of stony coral (*Oulangia* cf. *stokesiana*), one unidentified species of mud lobster (*Thalassina* sp.), two species of barnacles (*Fistulobalanus kondakovi* and *Megabalanus* cf. *tintinnabulum*), and one species of sea urchin (*Temnotrema siamense*). The vertebrates are represented by fish remains, most of which are shark teeth belonging to the family Carcharhinidae with at least nine species. Other chondrichthyan remains include stingray teeth of the genus *Pastinachus* (Dasyatidae). In addition, few bony fish remains were recovered such as trichiurid teeth and one sciaenid otolith (*Johnius*). The molluscan fauna indicates that the

paleoenvironments of the area corresponded to intertidal to sublittoral zones, where some areas were mangrove forests and intertidal mudflats. The fish fauna is dominated by the carcharhinid *Glyphis* (also known as “river shark”), indicating freshwater influences and possibly occasional brackish conditions. The chronological results obtained from carbon-14 analysis of mollusc remains and charcoal show that deposition of the marine sediment sequence began during the mid-Holocene, spanning approximately from 8,800 to 5,300 cal yr BP. This study provides in-depth insights into the diversity of fishes, marine molluscs and other invertebrates from the Bangkok Clay deposits, supporting the existence of a marine transgression onto the Lower Central Plain of Thailand during the mid-Holocene.

Keywords carbon-14 dating, Chao Phraya River Basin, Chondrichthyes, marine transgression, mollusc, paleoenvironment

Introduction

The Holocene began approximately 11,700 years ago and belongs to one of the interglacial periods. Prior to the Holocene, during the period of the Last Glacial Maximum (LGM), the global climate was much colder and dryer and was a period of significant landscape changes in many areas due to a lowered sea level, approximately 120 m below the present-day stands (Clark et al. 2009). During the Holocene, however, the climate became warmer, and the sea level rose and inundated a previously emerged landmass. Southeast Asia is one of the regions impacted by sea level inundations (Sathiamurthy and Voris 2006), due in part to an increase in precipitation level after the cold and dry period (Chabangborn 2017). Differences in climatic and environmental conditions between the Pleistocene and Holocene have therefore had impacts on the ecological adaptations of both terrestrial and marine organisms, leading to dispersal, migration, and extinction of species.

The Central Plain of Thailand is an important area where the topography and climate changes were under the influence of a sea level rise starting at around 12 ka. The region covers the plains between the mountain ranges along the lower northern part of Thailand, including the Yom and Nan River basins, and the low-lying plains of the Chao Phraya, Tha Chin, Mae Klong and Bang Pakong River basins near the Gulf of Thailand (Sinsakul 2000). During the Holocene, the Sing Buri Plain (5–15 m above mean sea level (amsl)) and the Bangkok Lowland (0–5 m amsl), forming the Central plain of Thailand, were inundated (Takaya 1969). Around 8–7 ka, the sea level was 2–4 m higher than present-day stands and inundated the whole lowland areas of Bangkok. Later on, the sea level subsided along with a new sedimentation regime and a new landmass started to develop, shaping the current coastal shorelines (Umitsu et al. 2002; Tanabe et al. 2003).

The extent of marine intrusion into the Central Plain of Thailand during the Holocene was largely interpreted based on various studies (e.g., geomorphological, sedimentological, paleontological, palynological, and stratigraphical data as well as carbon-14 dating of peat layers). These studies indicated that the ancient coastal shoreline of the Gulf of Thailand was located further north in Phra Nakhon Si Ayutthaya Province during 7.3–6.5 ka (Jarupongsakul and Thiramongkol 1992). The ancient coastal shoreline was also reconstructed based on a 4,500-year-old coastal archeological site in Chonburi Province (Boyd et al. 1996), open-pit mining of clay deposits at a depth of 7 m under the ground surface in many provinces, dated to around 8–7 ka

(Songtham et al. 2007, 2015), and geological studies of coastal berms in Chumporn and Prachuap Khiri Khan provinces, dated between 8.9–5.6 ka and 6.5–6 ka, respectively (Nimnate et al. 2015; Surakiatchai et al. 2018).

In addition to the inference of ancient coastal shorelines during the Holocene, studies on faunas and floras along the ancient coast of the Gulf of Thailand are crucial to understand paleoenvironmental and paleoecological changes in response to climatic oscillations. The ancient coastal habitats occupied by these animals are investigated based on the identification and dating of fossils (e.g., molluscs, arthropods, and vertebrates). According to accurate species-level identification, microhabitats can be investigated based on comparisons between fossils and living species or related taxa. The paleoenvironments of the area are also reconstructed based on the identification of various groups of animals because each of them has a different mode of life in a specific habitat (Negri 2009, 2012; Oliver and Terry 2019).

Marine faunas from the Holocene Bangkok Clay deposits of the Central Plain of Thailand have infrequently been analysed in detail, and we identify the animal macrofossils recovered from the clay pit of Ongkharak, Nakhon Nayok Province in central Thailand (Fig. 1). We also date the Bangkok Clay sequence deposited in the area, using a carbon-14 dating technique. Although the Bangkok Clay deposits have previously been described from some areas of Nakhon Nayok Province (Songtham et al. 2007; Royal Irrigation Department 2009), the fauna and its paleoenvironments have not yet been thoroughly studied and compared to other sites from different regions (e.g., Negri 2009, 2012). This study not only fulfils the gap regarding the paleoenvironments of the Gulf of Thailand during the Holocene, but also illustrates the ancient coastlines in the eastern part of the Central Plain of Thailand to some extent. The reconstruction of ancient coastlines and paleoenvironments is crucial to advance the knowledge of archaeological contexts in terms of prehistoric human settlements and cultures, and probably helpful in forecasting changes in coastal ecosystems and environments in response to sea level oscillations due to the present-day global warming situation.

Geological settings of the study area

The study area is located at the clay pit in Ban Lad Chang, Chumphon Subdistrict, Ongkharak District, Nakhon Nayok Province ($13^{\circ}59'30.9''\text{N}$, $100^{\circ}55'11.5''\text{E}$) in the Lower Central Plain of Thailand (Fig. 1), which is covered by the thick marine deposits of the Chao Phraya River delta, also known as the Bangkok Clay. These marine clay deposits have been dated between the Late Pleistocene and Holocene (Cox 1968; Moh et al. 1969; Sinsakul 2000). The Bangkok clay sequence comprises three main zones (Ohtsubo et al. 2000): 1) weathered clay with a thickness of 0–2 m in the uppermost part, 2) soft marine clay in the middle part, and 3) medium stiff clay at the bottom. The Holocene Bangkok Soft Clay in the middle part, with a thickness ranging from 10 to 20 m, is olive grey and medium to dark grey in colour, indicating reducing environments (Sinsakul 2000; Choowong 2002; Tanabe et al. 2003). This Bangkok Soft Clay layer is further divided into lower transgressive peaty (mangrove swamp) and upper regressive deltaic sediments (Tanabe et al. 2003). Some marine shells and remains of mangrove trees were found from this middle layer, indicating intertidal and shallow sublittoral inner bay conditions, with local

mangrove and freshwater influences (Robba et al. 2002, 2003, 2004, 2007; Songtham et al. 2007; Negri 2009, 2012). This evidence supports the occurrence of a marine transgression into the Central Plain of Thailand after the LGM (Songtham et al. 2015).

The rectangular mine is 1,000 m long x 400 m wide and 30 m deep. It was opened to produce clay material used in construction sites. The active mining operations of the clay pit allow us to have access to the stratigraphic sequence of the Quaternary deposits of the area (Fig. 2). The topsoil (0.5–1 m thick) consists of coarse-grained sand with FeO grains and black mud nodules, underlain by a 5 m thick layer of black clay and organic soil (dark layer) partially interbedded with thin layers of silt to very fine-grained sand. Marine invertebrate and vertebrate remains were only found from this organic-rich clay unit, also known as the Bangkok Clay, which overlies the layers of Pleistocene stiff clay (Rau and Nutalaya 1983). In this site, the Pleistocene stiff clay contains seven different lithological units from the top to bottom of the pit: 1) reddish brown lateritic soil (1 m thick), 2) yellowish brown clay to silt-sized sediments with root-trace burrows (4 m thick), 3) reddish brown clay to silt-sized sediments with shell fragments (4 m thick), 4) purplish clay to sand-sized sediments (1 m thick), 5) greyish clay to sand-sized sediments (2 m thick), 6) reddish brown lateritic soil with pale pink and white spots of weathering soil (3 m thick), and 7) yellowish grey clay to silt-sized sediments (5 m thick), respectively (Fig. 2).

The Bangkok clay layer at the clay pit of Ongkharak has yielded nearly *in situ* fossil deposits as indicated by preservation of complete shells with both valves attached (Fig. 2C) as well as the orientation of shell valves of *Pholas orientalis*, which has been recovered in natural position within its burrows during field collection (see also Songtham et al. 2007: fig. 3). Although remains of marine molluscs were found throughout the sequence of dark clay deposits, they were most abundant in the middle part of the layer (Fig. 2D–F), at a depth of about 2 m below the upper contact that separates it from the topsoil. Numerous wood fragments, plant remains, and giant oyster shells were found from the lower part of the dark clay unit below the shell-rich layer.

Material and methods

Species identification of specimens is mainly based on the existing literature specified under the treatment of each taxon. The specimens are kept at the Department of Geology, Faculty of Science, Chulalongkorn University, and coded as the name of the collection (CUF: Chulalongkorn fossil collection), followed by the locality (NKNY: Nakhon Nayok) and the catalogue number. Fish remains including teeth and otolith as well as some small shells were photographed with a scanning electron microscope (SEM; JEOL, JSM-5410 LV) or a Leica M205C stereo light microscope with fusion optics and the Leica Application Suite Image System. L and R designate left and right valves of bivalve shells, respectively.

We selected six samples (shells and charcoal) collected from the marine clay deposits at different depths in the clay pit for carbon-14 dating (Table 1). The samples were pretreated and analyzed at the DirectAMS (USA), using an accelerated mass spectrometer (AMS). According to the convention outlined by Stuiver and Polach (1977), the ^{14}C age \pm one standard deviation (SD) was calculated using the Libby half-life of 5,568 years and an isotopic fractionation correction

based on $\delta^{13}\text{C}$ measurements obtained from the AMS. The DirectAMS results were calibrated and corrected for the marine reservoir effect, using the Calib 8.2 program with IntCal20.14c for charcoal (Reimer et al. 2020) and with Marine20 (Heaton et al. 2020) for shells.

Results

The ages of the Bangkok Clay deposits from the clay pit range from 8,784 cal yr BP to 5,318 cal yr BP (mid-Holocene) based on the radiocarbon dating of several shell and charcoal fragments collected along the stratigraphic section of the layer (Table 1). The majority of marine faunas are molluscs, where a total of 63 species were identified, including 35 gastropod species from 20 families, 27 bivalve species from 15 families, and one scaphopod species. The most common gastropods are *Architectonica perdix*, *Natica stellata*, and *Indothais lacera*, whereas the most common bivalves are *Corbula fortisulcata*, *Magallana cf. gigas*, and *Joannisiella oblonga*. Other invertebrate remains include one species of stony coral (*Oulangia cf. stokesiana*), one unidentified mud lobster species (*Thalassina* sp.), two species of barnacles (*Fistulobalanus kondakovi* and *Megabalanus cf. tintinnabulum*), and one species of a sea urchin (*Temnotrema siamense*).

The vertebrate remains include at least nine species (two families) of chondrichthyan fishes and at least two species (two families) of actinopterygian taxa. Altogether, one hundred cartilaginous fish fossils were recovered, including ninety-seven remains identified as belonging to the family Carcharhinidae and three to the family Dasyatidae. The latter teeth belong to the stingray genus *Pastinachus*. Among the carcharhinids, the genus *Glyphis* dominates the shark fauna (61%). There are also at least six species of *Carcharhinus*, representing 26% of the shark fauna. The accurate identification of *Carcharhinus* species is often hindered by the preservation of teeth, or by the morphological similarities of tooth positions among these taxa. A few teeth among the carcharhinids belong to the genus *Scoliodon*. Few actinopterygian remains were also found, including three teeth that belong to the family Trichiuridae and one sciaenid otolith that represents the genus *Johnius*.

Systematic paleontology

Phylum Cnidaria

Class Anthozoa Ehrenberg, 1834

Subclass Hexacorallia Haeckel, 1896

Order Scleractinia Bourne, 1900

Family Rhizangiidae d'Orbigny, 1851

Oulangia Milne Edwards & Haime, 1848

Oulangia cf. stokesiana Milne Edwards & Haime, 1848

Figs 3A, 5A

Oulangia stokesiana Milne Edwards & Haime, 1848: pl. 7, fig. 4, 4a. Type locality: Philippines.

Milne Edwards and Haime 1849: 183. Lam et al. 2008: 742, fig. 4e.

Oulangia stokesiana stokesiana—Cairns et al. 1999: 39.

Referred material. CUF-NKNY-O07 (1 specimen; Figs 3A, 5A).

Habitat. Shallow waters as well as in submarine caves (Lam et al. 2008).

Distribution. Indian Ocean; Indo-West Pacific, from Japan to the Philippines, and central Pacific (Cairns et al. 1999; Lam et al. 2008).

Taxonomic remarks and comparisons. This specimen is classified into the genus *Oulangia* based on the descriptions and figures of Cairns and Kitahara (2012) and Baron-Szabo and Cairns (2016). We tentatively identify this specimen as belonging to *Oulangia stokesiana* according to Lam et al. (2008).

Phylum Mollusca

Class Gastropoda Cuvier, 1795

Subclass Vetigastropoda Salvini-Plawen, 1980

Order Trochida Cox & Knight, 1960

Superfamily Trochoidea Rafinesque, 1815

Family Colloniidae Cossmann, 1917

***Homalopoma* Carpenter, 1864**

***Homalopoma cf. sangarensense* (Schrenck, 1861)**

Figs 3F, 5B

Turbo sangarensis Schrenck, 1861: 409, 410. Type locality: the Sangaric strait near the shore of the island of Jesso [Sangar (Tsugaru) Strait, Hakodate Bay, Hokkaido, Japan].

Homalopoma sangarensense—Habe 1958: 5, pl. 2, fig. 1. Golikov and Kussakin 1978: 65, 66, fig. 40a, b. Egorov 2000: 66, with in-text fig. Kantor and Sysoev 2006: 42, pl. 18h. Lebedev 2014: 57.

Referred material. CUF-NKNY-G24 (1 shell; Figs 3F, 5B).

Habitat. On silty-sandy or rarely on rocky substrates at a depth from 2 to 22 m (Egorov 2000; Kantor and Sysoev 2006).

Distribution. Sea of Japan and southern Kuril Islands (Kantor and Sysoev 2006; Lebedev 2014).

Record in Thailand. This is the first record of this species in Thailand.

Taxonomic remarks and comparisons. Although there is only one incomplete specimen, it is tentatively identified as *Homalopoma sangarensense* based on shell size and shape, and several strong spiral cords on the surface following the descriptions and figures in Habe (1958), Golikov and Kussakin (1978), and Egorov (2000).

Subclass Neritimorpha Golikov & Starobogatov, 1975

Order Cycloneritida Frýda, 1998

Superfamily Neritoidea Rafinesque, 1815

Family Neritidae Rafinesque, 1815

***Neripteron* Lesson, 1831**

***Neripteron cf. violaceum* (Gmelin, 1791)**

Figs 3B, 5C

Nerita violacea Gmelin, 1791: 3686. Type locality: unknown.

Neritina violacea—Tantanasiriwong 1978: 6, fig. 52. Nateewathana et al. 1981: 56. Wilson 1993: 41, pl. 2, figs 18a, b, 19a–d. Hylleberg and Kilburn 2003: 31. Robba et al. 2004: 24, 25, pl. 2, fig. 5a, b. Robba et al. 2007: 89 (appendix). Printrakoon et al. 2008: table 1. Tan and Clements 2008: 490, 491, fig. 3-33, 3-34. Nabhitabhata 2009: 39. Sanpanich and Duangdee 2013: 53. Ng et al. 2016: fig. 1-28. Yang et al. 2017: 20, fig. 75. Tudu et al. 2018: table 1.

Dostia violacea—Way and Purchon 1981: 314. Subba Rao and Dey 2000: 35. Sri-aroon et al. 2004: table 1, fig. 2-1. Dechruksa et al. 2014: fig. 2k.

Neripteron violacea—Swennen et al. 2001: 52, 108, text-fig. 275, fig. 275.

Neritina (Dostia) violacea—Gemert 2003: 103. Sri-aroon et al. 2005: tables 2, 3, 5, 6. Dey 2006: 22, figs 11, 12. Ramakrishna et al. 2007: 5, 37. Kesavan et al. 2009: 382, with in-text fig. Eichhorst 2016: 392–395, pl. 102, figs 1–15.

Neritina (Neripteron) violacea—Thach 2005: 38, pl. 6, figs 8, 9.

Neripteron violaceus—Eichhorst 2008: 268, pl. 79, fig. 7a, b.

Neripteron violaceum—Baharuddin et al. 2017: fig. 3o. BEDO 2017b: 97, with in-text fig. Kantharajan et al. 2017: table 1, fig. 4-34. Yadav et al. 2019: table 2, fig. 1h. Mustapha et al. 2021: 52, 53, figs 3e, f, 4i. Wells et al. 2021: 154, 155.

Referred material. CUF-NKNY-G26 (28 shells; Figs 3B, 5C).

Habitat. On Nipa palms, muddy surfaces, old wood and rocks in brackish habitats and mangrove forests (Swennen et al. 2001; Mustapha et al. 2021) as well as on gravels in high tidal zones (Yang et al. 2017).

Distribution. India (Kantharajan et al. 2017; Yadav et al. 2019); Indo-West Pacific, from Japan to Australia (Wilson 1993; Yang et al. 2017).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. The shell of *Neripteron violaceum* is highly similar to that of *N. cornucopia* (Benson, 1836) but differs in having (1) an orangish-red ventral side and (2) denticulations along the columellar edge often only present in the central part (Huang 1997; Tan and Clements 2008). Because the columella edge of the specimens is corroded and the colour of the ventral side cannot be accurately determined, these specimens are tentatively identified as belonging to *N. violaceum*.

Nerita Linnaeus, 1758

Nerita articulata Gould, 1847

Figs 3C, 5D

Nerita lineata Gmelin, 1791: 3684 [junior homonym of *Nerita lineata* Müller, 1774]. Type locality: Strait of Malacca. Way and Purchon 1981: 314. Hylleberg and Kilburn 2003: 29, 30. Yang et al. 2017: 20, fig. 67.

Nerita (Ritena) lineata—Cernohorsky 1972: 51, pl. 11, fig. 5.

Nerita articulata Gould, 1847: 220. Type locality: Tavoy. Wium-Andersen 1977: 5, fig. 8. Tantanasiriwong 1978: 6, fig. 42. Nateewathana et al. 1981: 56. Tan and Clements 2008:

483–485, fig. 2-3, 2-4. Nabhitabhata 2009: 36. Hamli et al. 2013: tables 2, 3, fig. 2g. Tudu et al. 2018: table 1.

Nerita balteata Reeve, 1855: pl. 6, fig. 28. Type locality: unknown. Wilson 1993: 40, pl. 2, fig. 9a, b. Nateewathana 1995: 95, with in-text fig. Nabhitabhata 2009: 36. Sanpanich and Duangdee 2013: 53. Chen et al. 2015: 75, fig. 1a–c. BEDO 2017b: 99, with in-text fig. Kantharajan et al. 2017: table 1, fig. 4-36. Tudu et al. 2018: table 1. Raven 2019: figs 4–6. Yadav et al. 2019: table 2, fig. 1i. Mustapha et al. 2021: 48, figs 2m, n, 4g. Wells et al. 2021: 155.

Nerita (Ritena) balteata—Cernohorsky 1978: 42, pl. 11, fig. 1. Dharma 2005: 68, pl. 9, fig. 3a, b.

Nerita (Amphinerita) articulata—Subba Rao and Dey 2000: 31. Dey 2006: 20, figs 7, 8.

Nerita (Nerita) articulata—Sri-aroon et al. 2004: table 1.

Nerita (Theliostyla) balteata—Thach 2005: 37, pl. 6, figs 36, 37.

Nerita balteata forma articulata—Eichhorst 2008: 280, pl. 85, fig. 1a, b.

Nerita (Cymostyla) balteata—Eichhorst 2016: 455–457, pl. 113, figs 1–9.

Referred material. CUF-NKNY-G61 (3 shells; Figs 3C, 5D).

Habitat. On tree trunks, branches, roots and on muddy banks and rocky areas or near mangrove forests (Tan and Clements 2008; Mustapha et al. 2021) as well as on rocks in intertidal zones (Yang et al. 2017).

Distribution. India (Kantharajan et al. 2017; Yadav et al. 2019); Indo-West Pacific, from China to Australia (Wilson 1993; Yang et al. 2017). Also reported from the Hawaiian Islands (Cernohorsky 1978).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized by having (1) numerous raised spiral cords and (2) a crenulated outer lip (Tan and Clements 2008). Although this species is widely known as *N. balteata* Reeve, 1855, the name *N. articulata* was made available earlier by Gould (1847). Thus, it has priority over *N. balteata*.

Subclass Caenogastropoda Cox, 1960

Cohort Sorbeoconcha Ponder & Lindberg, 1997

Subcohort Cerithiimorpha Golikov & Starobogatov, 1975

Superfamily Cerithioidea Fleming, 1822

Family Potamididae H. Adams & A. Adams, 1854

***Cerithidea* Swainson, 1840**

***Cerithidea obtusa* (Lamarck, 1822)**

Figs 3M, 5E

Cerithium obtusum Lamarck, 1822: 71. Type locality: Timor.

Cerithidea obtusa—Cernohorsky 1972: 60 (in part). Tantanasiriwong 1978: 7, fig. 70.

Nateewathana et al. 1981: 57. Way and Purchon 1981: 315. Nateewathana 1995: 97, with in-text fig. Poutiers 1998b: 450, with in-text figs. Gemert 2003: 104. Hylleberg and Kilburn 2003: 37, 38. Robba et al. 2004: 31, 32, pl. 3, fig. 4a, b. Dharma 2005: 92, pl. 21, fig. 8; 308, pl. 119, fig. 8a, b. Thach 2005: 49, pl. 10, fig. 4. Dey 2006: 35, 36, figs 38–40. Robba et al. 2007: 89 (appendix). Printrakoon et al. 2008: table 1. Kesavan et al. 2009: 382, with

in-text fig. Hamli et al. 2013: tables 2, 3, fig. 2a. Dechruksa et al. 2014: fig. 2f. Reid 2014: 27–29, 31, figs 2b, 8, 9. BEDO 2017b: 26, with in-text fig. Tudu et al. 2018: table 1. Wells et al. 2021: 125, 126.

Cerithidea (Cerithidea) obtusa—Brandt 1974: 192, 193, pl. 14, fig. 52. Houbrick 1984: 15, 16, fig. 5c. Sri-aroon et al. 2004: table 1, fig. 3-1. Sri-aroon et al. 2005: tables 2–6. Ramakrishna et al. 2007: 6, 47. Nabhitabhata 2009: 95.

Referred material. CUF-NKNY-G14 (23 shells; Figs 3M, 5E).

Habitat. On firm mud and on trunks and stilt roots in mangrove forests as well as in fully marine and estuarine areas (Reid 2014).

Distribution. Bay of Bengal; Indo-West Pacific, from southern Vietnam to Java and East Borneo (Reid 2014). Records of fossils from the Late Pliocene in Central and West Java, Indonesia, and from the Holocene in Thailand (Robba et al. 2004; Dharma 2005).

Record in Thailand. Chanthaburi River, Gulf of Thailand and Andaman Sea (Nabhitabhata 2009; Wells et al. 2021).

Taxonomic remarks and comparisons. This species has a more thickened and flared aperture, compared to two similar species, *Cerithidea anticipata* Iredale, 1929 and *Cerithidea reidi* Houbrick, 1986. These two species are also common in Sahul (New Guinea and Australia), whereas *C. obtusa* is present in Sunda (Reid 2014). See also comprehensive taxonomic remarks in Reid (2014).

Pirenella Gray, 1847

Pirenella incisa (Hombron & Jacquinot, 1848)

Figs 3I, 5F

Cerithium incisum Hombron & Jacquinot, 1848: 97, pl. 23, figs 8, 9. Type locality: Borneo.

Cerithidea (Cerithideopsilla) djadjariensis [non Martin]—Sri-aroon et al. 2004: table 1, fig. 2-7.

Cerithideopsilla djadjariensis [non Martin]—Lozouet 2008: 284, pl. 87, fig. 9; pl. 88, fig. 2.

Pirenella incisa—Reid and Ozawa 2016: 30–32, figs 1, 2g, h, 4i, 10, 11a. Zvonareva and Kantor 2016: 415, figs 5a–d. BEDO 2017b: 28, with in-text fig. Wells et al. 2021: 126.

Referred material. CUF-NKNY-G10 (47 shells; Figs 3I, 5F).

Habitat. On damp mud in mangrove forests. Also found on open surface areas of intertidal mudflats and on muddy shores of shrimp ponds (Reid and Ozawa 2016; Zvonareva and Kantor 2016).

Distribution. East India; Indo-West Pacific, from southern China, the Phillipines to West Sulawesi, and Flores Island (Reid and Ozawa 2016).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from the other two similar species, *P. pupiformis* Ozawa & Reid, 2016 and *P. caiyingyai* (Qian, Fang & He, 2013), by (1) the presence of a nearly columnar appearance, (2) a more oblique and more flared aperture with the projection of lips next to a deep anterior canal, and (3) three equal cords on each spire whorl with strong square nodules on axial ribs (Reid and Ozawa 2016). See comprehensive taxonomic remarks in Reid and Ozawa (2016).

***Telescopium* Montfort, 1810**

***Telescopium telescopium* (Linnaeus, 1758)**

Figs 3K, 5G

Trochus telescopium Linnaeus, 1758: 760. Type locality: unknown.

Telescopium telescopium—Tesch 1920: 58, 59, pl. 132, fig. 191. Cernohorsky 1972: 61, pl. 13, fig. 6. Brandt 1974: 196, pl. 15, fig. 61. Tantanasiwong 1978: 7, fig. 69. Nateewathana et al. 1981: 57. Way and Purchon 1981: 315. Houbrick 1991: 291–304, figs 1–6. Wilson 1993: 133, pl. 15, fig. 21. Bosch et al. 1995: 56, 57, fig. 186. Nateewathana 1995: 97, 98, with in-text fig. Poutiers 1998b: 451, with in-text figs. Subba Rao and Dey 2000: 53. Swennen et al. 2001: 52, 111, 112, with in-text fig., fig. 295. Hylleberg and Kilburn 2003: 38. Sri-aroon et al. 2004: table 1, fig. 3–3. Dharma 2005: 92, pl. 21, fig. 1; 308, pl. 119, fig. 4. Thach 2005: 45, pl. 9, figs 4, 7. Dey 2006: 31, 32, figs 29–32. Ramakrishna et al. 2007: 48, 49. Lozouet 2008: 284, pl. 87, figs 1,3. Printrakoon et al. 2008: table 1. Kesavan et al. 2009: 382, with in-text fig. Nabhitabhata 2009: 97. Hamli et al. 2013: tables 2, 3, fig. 2e. Sanpanich and Duangdee 2013: 54. Dechruksa et al. 2014: fig. 2h. BEDO 2017b: 29, with in-text fig. Kantharajan et al. 2017: table 1, fig. 3–18. Okutani 2017: 797, pl. 63, fig. 11. Yang et al. 2017: 28, fig. 109. Tudu et al. 2018: table 1. Yadav et al. 2019: table 2, fig. 1o. Palanisamy et al. 2020: 2, fig. 1a–d. George et al. 2021: 1532, fig. 2a, b. Wells et al. 2021: 126.

Referred material. CUF-NKNY-G64 (2 shells; Figs 3K, 5G).

Habitat. On muddy floors of mangrove forests and intertidal mud flats (Poutiers 1998b; Okutani 2017).

Distribution. Indian Ocean; Indo-West Pacific, from Japan to India (Poutiers 1998b; Okutani 2017; Palanisamy et al. 2020). Records of fossils from the Miocene to Quaternary in Indonesia (Tesch 1920; Dharma 2005).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized by (1) its high conical spire and broad, rather flat base, and (2) a twisted columella with a strong and central spiral ridge (Houbrick 1991; Poutiers 1998b).

Subcohort Hypsogastropoda Ponder & Lindberg, 1997

Superfamily Naticoidea Guilding, 1834

Family Naticidae Guilding, 1834

***Eunaticina* Fischer, 1885**

***Eunaticina papilla* (Gmelin, 1791)**

Figs 3G, 6A

Nerita papilla Gmelin, 1791: 3675. Type locality: Tranquebar [Tharangambadi, Tamil Nadu, India].

Eunaticina papilla—Cernohorsky 1971: 201, 202, fig. 69. Cernohorsky 1972: 102, pl. 27, fig. 5. Tantanasiwong 1978: 11, fig. 151. Nateewathana et al. 1981: 58. Way and Purchon 1981:

317. Majima 1989: 68, 69, pl. 10, fig. 16, text-figs 15.36, 23.1a–d. Wilson 1993: 223, pl. 36, fig. 7. Bosch et al. 1995: 87, fig. 326. Swennen et al. 2001: 54, 122, fig. 378. Gemert 2003: 105. Hylleberg and Kilburn 2003: 56. Beu et al. 2004: 206, 208–211, fig. 22d–f, h. Robba et al. 2004: 75, 76, pl. 9, fig. 7a, b. Dharma 2005: 176, pl. 63, fig. 25a–d; 342, pl. 136, fig. 12a, b. Ramakrishna et al. 2007: 9, 69, 70. Robba et al. 2007: 91 (appendix). Hollmann 2008: 500, pl. 195, figs 1, 2. Nabhitabhata 2009: 123. Beu 2010: 152, fig. 21a. Torigoe and Inaba 2011: 55, 56, pl. 2, fig. 17. Gopalakrishnan et al. 2012: 73. Öztürk and Bitlis 2013: 7, with in-text fig. Sanpanich and Duangdee 2013: 56. BEDO 2017b: 159, with in-text fig. Okutani 2017: 862, pl. 148, fig. 6. Yang et al. 2017: 52, fig. 215. Tudu et al. 2018: table 1. Albano et al. 2021: 12, fig. 5. Wells et al. 2021: 82.

Eunaticina papilla papilla—Thach 2005: 90, pl. 26, figs 3, 5.

Referred material. CUF-NKNY-G19 (5 shells; Figs 3G, 6A).

Habitat. Fine sandy bottoms in intertidal zones down to 30 m deep (Robba et al. 2004; Okutani 2017).

Distribution. Red Sea to Indian Ocean; Indo-West Pacific, from Japan to Australia and Fiji Islands (Robba et al. 2004; Okutani 2017), as well as the Israeli Mediterranean shelf (Albano et al. 2021). Several records of fossils during the Late Miocene to Holocene in Indonesia, Japan, Taiwan, and Thailand (Beu et al. 2004; Robba et al. 2004; Dharma 2005).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized by having (1) a regularly oval shell with a small spire and (2) a large body whorl with low and wide spiral cords, which are broader than their interspaces and crossed by prominent axial ridges (Öztürk and Bitlis 2013). See also comprehensive taxonomic remarks in Beu et al. (2004).

Natica Scopoli, 1777

Natica stellata Hedley, 1913

Figs 3D, 6B

Natica stellata Hedley, 1913: 299, 300. Type locality: unknown. Cernohorsky 1972: 94, 95, pl. 24, fig. 6. Wilson 1993: 217, pl. 36, fig. 24. Poutiers 1998b: 513, with in-text figs. Robba et al. 2004: 71, pl. 9, fig. 2a, b. Dharma 2005: 176, pl. 63, fig. 18. Robba et al. 2007: 91 (appendix). Hollmann 2008: 492, pl. 191, figs 9, 10a, b. Nabhitabhata 2009: 125. Torigoe and Inaba 2011: 76, 77. Mukhopadhyay et al. 2013: 152, 153, fig. 6. Sanpanich and Duangdee 2013: 57. Okutani 2017: 862, pl. 149, fig. 7. Wells et al. 2021: 83.

Natica (Natica) stellata—Cernohorsky 1971: 176, 177, figs 6, 8–13.

Referred material. CUF-NKNY-G05 (126 shells; Figs 3D, 6B).

Habitat. Sandy gravel bottoms in sublittoral zones down to 20 m deep (Hollmann 2008; Okutani 2017).

Distribution. Indo-West Pacific, from Japan to Australia (Cernohorsky 1972; Wilson 1993; Okutani 2017). Probably present in the Indian Ocean (Poutiers 1998b). Records of fossils from the Holocene in central Thailand (Robba et al. 2003, 2004).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from its similar species, *Natica vitellus*, in having a parietal callus forming a tongue-shaped extension over the posterior part of the umbilicus (Cernohorsky 1971). See also comprehensive taxonomic remarks in Cernohorsky (1971).

***Natica vitellus* (Linnaeus, 1758)**

Figs 3E, 6C

Nerita vitellus Linnaeus, 1758: 776. Type locality: Asiatic Ocean.

Natica vitellus—Tesch 1920: 70, 71, pl. 132, fig. 207a, b. Cernohorsky 1972: 94, pl. 24, fig. 5. Tantanasiwong 1978: 11, fig. 141. Nateewathana et al. 1981: 58. Way and Purchon 1981: 317. Majima 1989: 74–76, pl. 10, figs 1–12, text-figs 4.1, 15.38. Wilson 1993: 217, pl. 36, fig. 28. Bosch et al. 1995: 87, fig. 322. Poutiers 1998b: 515, with in-text figs. Subba Rao and Dey 2000: 78. Swennen et al. 2001: 54, 121, fig. 371. Hylleberg and Kilburn 2003: 57. Robba et al. 2003: table 5. Robba et al. 2004: 71–73, pl. 9, fig. 3a, b. Dharma 2005: 176, pl. 63, fig. 17; 342, pl. 136, fig. 5a, b. Ramakrishna et al. 2007: 8, 67. Robba et al. 2007: 91 (appendix). Hollmann 2008: 492, pl. 191, fig. 7. Nabhitabhata 2009: 125. Torigoe and Inaba 2011: 69, 70. Sanpanich and Duangdee 2013: 57. BEDO 2017b: 162, with in-text fig. Okutani 2017: 862, pl. 149, fig. 4. Yang et al. 2017: 56, fig. 230. Harzhauser et al. 2018: 7, 8, pl. 1, figs 7–15. Tudu et al. 2018: table 1. Wells et al. 2021: 83.

Natica (Natica) vitellus—Cernohorsky 1971: 173, 174, 176, figs 2–5. Ladd 1982: 39, pl. 6, fig. 7.

Natica vitellus vitellus—Thach 2005: 86, pl. 26, fig. 8.

Referred material. CUF-NKNY-G04 (5 shells; Figs 3E, 6C).

Habitat. Fine sandy or muddy bottoms in intertidal zones down to a depth of about 120 m (Robba et al. 2004); in clean coral-sand pockets and weedy-sand lagoons (Cernohorsky 1971).

Distribution. Persian Gulf to Indian Ocean; Indo-West Pacific, from Japan to Australia (Bosch et al. 1995; Poutiers 1998b; Okutani 2017). Several records of fossils from the Early Miocene to Holocene in India, Indonesia, Japan, Malaysia, Myanmar, the Philippines, Polynesia, Taiwan, Vanuatu, and Thailand (Ladd 1982; Robba et al. 2004; Dharma 2005).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from its similar species, *Natica stellata*, in lacking a tongue-shaped extension of a parietal callus over the posterior part of the umbilicus (Cernohorsky 1971). See also comprehensive taxonomic remarks in Cernohorsky (1971).

***Paratectonatica* Azuma, 1961**

***Paratectonatica tigrina* (Röding, 1798)**

Figs 3Q, 6D

Cochlis tigrina Röding, 1798: 147. Type locality: unknown.

Natica tigrina—Tantanasiwong 1978: 11, fig. 142. Nateewathana et al. 1981: 58. Dheeradilok et al. 1984: pl. 3, figs 9, 10. Wilson 1993: 217, pl. 36, fig. 10. Nateewathana 1995: 98, with in-text fig. Poutiers 1998b: 514, with in-text figs. Swennen et al. 2001: 54, 121, fig. 369.

Hylleberg and Kilburn 2003: 57. Dharma 2005: 176, pl. 63, fig. 14. Ramakrishna et al. 2007: 8, 66, 67, pl. 4, figs 45, 46. Nabhitabhata 2009: 125. Okutani 2017: 864, pl. 150, fig. 10. Yang et al. 2017: 56, fig. 229. Surakiatchai et al. 2018: table 5, pl. 1, fig. 6a, b.

Paratectonatica tigrina—Majima 1989: pl. 14, fig. 18. Robba et al. 2003: tables 2–5. Robba et al. 2004: 73, pl. 9, fig. 4a, b. Robba et al. 2007: 91 (appendix). Torigoe and Inaba 2011: 101, 102, pl. 3, fig. 30. Kang et al. 2018: 357, 358, 360, fig. 4.

Tectonatica tigrina—Thach 2005: 87, pl. 26, fig. 24.

Notocochlis tigrina—Gopalakrishnan et al. 2012: 73. Sanpanich and Duangdee 2013: 57. BEDO 2017b: 165, with in-text fig. Tudu et al. 2018: table 1. Yadav et al. 2019: table 2, fig. 1a. Wells et al. 2021: 84.

Referred material. CUF-NKNY-G03 (70 shells; Figs 3Q, 6D).

Habitat. Sandy mud in intertidal zones down to 30 m deep (Thach 2005; Okutani 2017).

Distribution. India (Yadav et al. 2019); Indo-West Pacific, from Japan to Australia (Wilson 1993; Poutiers 1998b; Okutani 2017). Records of fossils from the Late Miocene to Quaternary in Indonesia and from the Quaternary in Thailand (Robba et al. 2003, 2004; Surakiatchai et al. 2018).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized by (1) its tall-spired shell with deeply impressed sutures and (2) spiral rows of reddish-brown spots on a white background (Poutiers 1998b; Robba et al. 2004).

Superorder Latrogastropoda Riedel, 2000

Superfamily Calyptraeoidea Lamarck, 1809

Family Calyptraeidae Lamarck, 1809

***Ergaea* H. Adams & A. Adams, 1854**

***Ergaea walshi* (Reeve, 1859)**

Figs 3H, 6E

Crepidula walshi Reeve, 1859: *Crepidula*, pl. 3, sp. 17. Type locality: Singapore; Ceylon [Sri Lanka]. Tantanasiriwong 1978: 9. Bosch et al. 1995: 69, fig. 230. Subba Rao and Dey 2000: 64. Swennen et al. 2001: 54, 117, text-fig. 344, fig. 344. Gemert 2003: 104. Robba et al. 2003: tables 3–5. Robba et al. 2004: 66, 67, pl. 8, fig. 8a, b. Ramakrishna et al. 2007: 7, 55, 56. Robba et al. 2007: 91 (appendix).

Ergaea walshi—Nateewathana et al. 1981: 57. Nabhitabhata 2009: 109. Low and Tan 2014: 11–13, fig. 1. BEDO 2017b: 117, with in-text fig. Tudu et al. 2018: table 1. Wells et al. 2021: 75.

Crepidula walshii [sic]—Way and Purchon 1981: 316.

Siphopatella walshi—Hylleberg and Kilburn 2003: 47. Sanpanich and Duangdee 2013: 55.

Okutani 2017: 838, pl. 114, fig. 3. Yang et al. 2017: 32, fig. 117.

Crepidula (Siphopatella) walshi—Dharma 2005: 78, pl. 14, fig. 13a, b; 342, pl. 136, fig. 15a, b.

Crepidula (Ergaea) walshi—Thach 2005: 66, pl. 10, fig. 27.

Referred material. CUF-NKNY-G25 (10 shells; Figs 3H, 6E).

Habitat. Attached to the aperture of other shelled marine organisms, in intertidal zones down to 40 m deep (Robba et al. 2004; Thach 2005; Low and Tan 2014).

Distribution. Persian Gulf; Indo-West Pacific, from Japan to the Arafura Sea (Robba et al. 2004; Okutani 2017). Several records of fossils from the Late Miocene to Holocene in India, Indonesia, Japan, Taiwan, and Thailand (Robba et al. 2004; Dharma 2005).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized by having (1) an irregularly subrectangular and flattened shell, with a small apex close to the posterior margin, and (2) a shelf-like internal septum attached just inside the posterior margin (Robba et al. 2004). See also comprehensive taxonomic remarks in Low and Tan (2014).

Superfamily Tonnaidea Suter, 1913

Family Bursidae Thiele, 1925

Bufonaria Schumacher, 1817

Bufonaria rana (Linnaeus, 1758)

Figs 3N, 6F

Murex rana Linnaeus, 1758: 748. Type locality: Asiatic Ocean.

Bursa (Bufonaria) rana—Cernohorsky 1972: 119, pl. 32, fig. 8.

Bursa rana—Tantanasiriwong 1978: 12, fig. 170. Nateewathana et al. 1981: 59. Way and Purchon 1981: 317. Swennen et al. 2001: 55, 122, fig. 380.

Bufonaria rana—Wilson 1993: 226, pl. 43, fig. 1a, b. Aungtonya and Hylleberg 1998: 319.

Poutiers 1998b: 551, with in-text figs. Subba Rao and Dey 2000: 92. Hylleberg and Kilburn 2003: 59. Robba et al. 2004: 76, 77, pl. 9, fig. 9a, b. Dharma 2005: 194, pl. 72, fig. 1a–e; 352, pl. 141, fig. 7a–c. Ramakrishna et al. 2007: 10, 81, pl. 6, figs 69, 70. Robba et al. 2007: 91 (appendix). Beu 2008: 618, pl. 254, figs 6, 7a, b. Nabhitabhata 2009: 137.

Sanpanich and Duangdee 2013: 58. BEDO 2017b: 112, with in-text fig. Okutani 2017: 867, 868, pl. 154, fig. 5. Yang et al. 2017: 64, fig. 266. Harzhauser et al. 2018: 13, 14, pl. 3, figs 6–9. Tudu et al. 2018: table 1. Yadav et al. 2019: table 2, fig. 1b. Wells et al. 2021: 89.

Bufonaria (Bufonaria) rana—Cossignani 1994: 33–35, with in-text figs. Bosch et al. 1995: 102, fig. 372. Gemert 2003: 105. Thach 2005: 96, pl. 28, fig. 8.

Referred material. CUF-NKNY-G06, G07 (18 shells; Figs 3N, 6F).

Habitat. Muddy or sandy bottoms at a depth from 20 to 100 m (Okutani 2017; Yang et al. 2017).

Distribution. Red Sea to Indian Ocean; Indo-West Pacific, from Japan to Australia (Robba et al. 2004; Okutani 2017; Yadav et al. 2019). Several records of fossils from the Miocene to Quaternary in Indonesia, Japan, the Philippines, and Taiwan (Robba et al. 2004; Dharma 2005; Harzhauser et al. 2018).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Robba et al. (2004) and Harzhauser et al. (2018), specifically in having two prominent varices placed at either periphery, running vertically uninterrupted, or slightly staggered, from apex to base, and bearing three short spines.

Order Neogastropoda Wenz, 1938

Superfamily Volutoidea Rafinesque, 1815

Family Cancellariidae Forbes & Hanley, 1851

***Merica* H. Adams & A. Adams, 1854**

***Merica elegans* (Sowerby I, 1822)**

Figs 3L, 6G

Cancellaria elegans Sowerby I, 1822: *Cancellaria*, pl. 218, fig. 3. Type locality: unknown.

Tantanasiwong 1978: 17. Nateewathana et al. 1981: 61. Robba et al. 2007: 51, 52, fig. 19d, e; 93 (appendix).

Cancellaria asperella [sic]—Cernohorsky 1972: 179, pl. 50, fig. 3 right.

Merica elegans—Verhecken 1986: 40, 41, fig. 9. Verhecken 1997: 307, 308, fig. 36. Swennen et al. 2001: 57, 133, fig. 461. Hylleberg and Kilburn 2003: 100. Dharma 2005: 134, pl. 42, fig. 1. Thach 2005: 188, pl. 58, figs 6, 10. Hemmen 2007: 128, 129, with in-text fig. Verhecken 2008: 818, pl. 704, fig. 5a, b. BEDO 2017b: 221, with in-text fig. Tudu et al. 2018: table 1. Wells et al. 2021: 121.

Referred material. CUF-NKNY-G18 (3 shells; Figs 3L, 6G).

Habitat. Sandy and muddy bottoms at a depth from 10 to 30 m (Robba et al. 2004; Thach 2005).

Distribution. Indo-West Pacific, from Japan to the Philippines. Records of fossils from the Miocene in Indonesia, and from the Quaternary in Japan (Robba et al. 2007).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from its similar species, *Merica asperella* (Lamarck, 1822), in having (1) a slenderer and more fusiform shell with a narrower aperture and (2) a sculptured shell with finer and more numerous ribs crossed by spiral ridges nearly of the same strength, showing a rectangular reticulated pattern (Robba et al. 2007).

***Scalptia* Jousseaume, 1887**

***Scalptia scalariformis* (Lamarck, 1822)**

Figs 3J, 6H

Cancellaria scalariformis Lamarck, 1822: 115. Type locality: unknown.

Cancellaria (Trigonostoma) scalariformis—Tesch 1915: 39, pl. 79, fig. 81a, b.

Trigonostoma scalariformis—Cernohorsky 1972: 180, pl. 50, fig. 2, 2a. Garrard 1975: 27–29, figs 4 (3, 4). Way and Purchon 1981: 320. Wilson 1994: 178, pl. 37, fig. 7a, b.

Scalptia scalariformis—Verhecken 1986: 53–55, figs 28–35. Swennen et al. 2001: 57, 133, fig. 462. Hylleberg and Kilburn 2003: 101. Robba et al. 2004: 129, 130, pl. 17, fig. 6. Dharma 2005: 134, pl. 42, fig. 12a, b. Hemmen 2007: 277, 278, with in-text figs. Robba et al. 2007: 93 (appendix). Okutani 2017: 1054, pl. 344, fig. 4. Yang et al. 2017: 102, fig. 415. Tudu et al. 2018: table 1. Wells et al. 2021: 121. Chan and Lau 2022: 1, 2, figs 1–3.

Scalptia (Scalptia) scalariformis—Thach 2005: 188, pl. 58, figs 25, 27, 29.

Referred material. CUF-NKNY-G64 (1 shell; Figs 3J, 6H).

Habitat. Sand and muddy bottoms at a depth from 20 to 40 m in sublittoral zones (Thach 2005; Okutani 2017; Yang et al. 2017).

Distribution. Indian Ocean; Indo-West Pacific, from Japan to Australia (Robba et al. 2004; Okutani 2017). Several records of fossils from the Pliocene to Holocene in the Indo-Pacific area, including Indonesia and Thailand (Robba et al. 2004).

Record in Thailand. Gulf of Thailand (Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from its similar species, *Scalptia bicolor* (Hinds, 1843), by having (1) a higher and narrower body whorl and (2) a less widely open umbilicus (Robba et al. 2007).

Superfamily Buccinoidea Rafinesque, 1815

Family Buccinidae Rafinesque, 1815

***Pseudoneptunea* Kobelt, 1882**

***Pseudoneptunea varicosa* (Röding, 1798)**

Figs 4K, 7A

Neptunea varicosa Röding, 1798: 116. Type locality: unknown.

Siphonalia (Pseudoneptunea) aff. varicosa—Tesch 1915: 53, pl. 80, fig. 114a, b.

Pseudoneptunea varicosa—Cernohorsky 1975: 217, 218, figs 10–12. Way and Purchon 1981: 318. Swennen et al. 2001: 56, 128, fig. 428. Robba et al. 2004: 106, 107, pl. 14, fig. 5a, b. Dharma 2005: 100, pl. 25, fig. 7a, b; 312, pl. 121, fig. 9a, b. Thach 2005: 135, pl. 39, figs 12, 15. Robba et al. 2007: 93 (appendix). Nabhitabhata 2009: 148. Sanpanich and Duangdee 2013: 63. Wells et al. 2021: 95.

Referred material. CUF-NKNY-G13, G16, G17 (25 shells; Figs 4K, 7A).

Habitat. Shallow water at a depth from 10 to 15 m (Thach 2005).

Distribution. Indo-West Pacific, from Vietnam to Indonesia (Robba et al. 2004; Dharma 2005; Thach 2005). Records of fossils from the Pliocene and Quaternary in Indonesia and from the Holocene in Thailand (Robba et al. 2004).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Cernohorsky (1975) and Robba et al. (2004), specifically in having (1) subangulate whorls and (2) a sculpture of crisp main spirals with an occasional intermediate spiral thread.

Family Melongenidae Gill, 1871

***Brunneifusus* Dekkers, 2018**

***Brunneifusus ternatanus* (Gmelin, 1791)**

Figs 4A, 7B

Murex ternatanus Gmelin, 1791: 3554. Type locality: Ternate Island, North Maluku, Indonesia.

Pugilina ternatana—Tantanasiriwong 1978: 15. Nateewathana et al. 1981: 60. Poutiers 1998b: 586, with in-text figs.

Hemifusus ternatanus—Yokoyama 1927: 393. Way and Purchon 1981: 318. Swennen et al. 2001: 56, 130, fig. 445. Gemert 2003: 106, with in-text fig. Hylleberg and Kilburn 2003: 88. Robba et al. 2004: 122, 123, pl. 16, fig. 7a, b. Dharma 2005: 108, pl. 29, fig. 2a, b; 316, pl. 123, fig. 7a, b. Robba et al. 2007: 93 (appendix). Nabhitabhata 2009: 162. Sanpanich and Duangdee 2013: 61. BEDO 2017b: 288, with in-text fig. Tudu et al. 2018: table 1.

Brunneifusus ternatanus—Dekkers 2018: 41, 42, pl. 2, figs 5, 6. Alf and Thach 2021: pl. 1, figs 1–9, pl. 2, figs 1–3. Wells et al. 2021: 97.

Referred material. CUF-NKNY-G21, G22 (71 shells; Figs 4A, 7B).

Habitat. Mud and muddy sand bottoms from the low tide mark to about 10–50 m deep (Poutiers 1998b; Swennen et al. 2001; Dekkers 2018).

Distribution. Eastern Indian Ocean; Indo-West Pacific, from Taiwan to Indonesia and the Philippines (Poutiers 1998b; Robba et al. 2004). Records of fossils from the Pliocene and Quaternary in Indonesia, Japan, and Taiwan (Robba et al. 2004; Dharma 2005).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is the sole member of the genus *Brunneifusus*, characterized by a determinate outer lip, compared to the indeterminate growth of the outer lip in *Hemifusus* Swainson, 1840 (Alf and Thach 2021). See also comprehensive taxonomic remarks in Alf and Thach (2021).

Family Nassariidae Iredale, 1916

Nassarius Duméril, 1805

Nassarius micans (A. Adams, 1852)

Figs 3O, 7C

Nassa micans A. Adams, 1852: 106. Type locality: Cagayan, Misamis, Mindanao.

Nassarius (Zeuxis) micans—Cernohorsky 1984: 147, 148, pl. 29, figs 8–11. Robba et al. 2003: table 5. Robba et al. 2004: 120, pl. 16, fig. 4a, b. Dharma 2005: 106, pl. 28, fig. 16a, b.

Robba et al. 2007: 93 (appendix). Okutani 2017: 914, pl. 202, fig. 11

Nassarius micans—Gopalakrishnan et al. 2012: 77. Wells et al. 2021: 99.

Referred material. CUF-NKNY-G27 (53 shells; Figs 3O, 7C).

Habitat. Sandy bottoms in subtidal zones down to 50 m deep (Robba et al. 2004; Okutani 2017).

Distribution. Gulf of Oman to India; Indo-West Pacific, from Japan to Papua New Guinea (Cernohorsky 1984; Robba et al. 2004; Okutani 2017). Records of fossils from the Holocene in Thailand (Robba et al. 2004).

Record in Thailand. Gulf of Thailand (Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from its similar species, *Nassarius comptus* (A. Adams, 1852), by having (1) a longer body whorl with narrowly subcanaliculate sutures, (2) a more prominent varix, and (3) a narrower and more elongate aperture (Cernohorsky 1984). See also comprehensive taxonomic remarks in Cernohorsky (1984).

***Nassarius siquijorensis* (A. Adams, 1852)**
Figs 3P, 7D

Nassa siquijorensis A. Adams, 1852: 97. Type locality: Island of Siquijor, Philippines.
Nassa (Zeuxis) siquijorensis—Tesch 1915: 59, pl. 81, figs 128a, b, 129a, b.
Nassarius (Zeuxis) siquijorensis—Cernohorsky 1978: 88, pl. 27, fig. 8. Cernohorsky 1984: 134–136, pl. 25, figs 12–14, pl. 26, figs 1–5. Robba et al. 2003: table 5. Robba et al. 2004: 120, 121, pl. 16, fig. 5a, b. Dharma 2005: 106, pl. 28, fig. 14a–c. Robba et al. 2007: 93 (appendix).
Nassarius siquijorensis—Nobuhara 1993: fig. 8-1a, 8-1b. Swennen et al. 2001: 56, 130, fig. 442. Hylleberg and Kilburn 2003: 87. Kool 2007: figs 17–20. Martin 2008: 120, pl. 355, fig. 13; 128, pl. 359, fig. 13. Surakiatchai et al. 2018: table 5, pl. 1, fig. 9a, b. Wells et al. 2021: 99.
Zeuxis siquijorensis—Thach 2005: 148, pl. 44, fig. 25.
Nassarius siquinjorensis [sic]—Yang et al. 2017: 90, fig. 371.

Referred material. CUF-NKNY-G01 (10 shells; Figs 3P, 7D).

Habitat. Sandy and muddy bottoms from subtidal zones to 450 m deep (Cernohorsky 1984; Thach 2005; Yang et al. 2017).

Distribution. Red Sea to India; Indo-West Pacific, from Japan to Indonesia and New Caledonia (Cernohorsky 1984; Dharma 2005; Yang et al. 2017). Records of fossils from the Pliocene in Indonesia and from the Holocene in Thailand (Cernohorsky 1984; Surakiatchai et al. 2018).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from its similar species, *Nassarius castus* (Gould, 1850), in having (1) a more tapering spire, (2) canaliculate sutures, and (3) considerably more numerous axial ribs (Cernohorsky 1984). See also comprehensive taxonomic remarks in Cernohorsky (1984) and Kool (2007).

Superfamily Muricoidea Rafinesque, 1815

Family Muricidae Rafinesque, 1815

***Chicoreus* Montfort, 1810**

***Chicoreus capucinus* (Lamarck, 1822)**
Figs 4E, 7E

Murex capucinus Lamarck, 1822: 164. Type locality: unknown.
Naquetia capucina—Tantanasiwong 1978: 13. Nateewathana et al. 1981: 59. Way and Purchon 1981: 318. Bussarawit 1991: 31.
Chicoreus (Rhizophorimurex) capucinus—Houart 1992: 106–109, fig. 217. Tan 2000: 504. Dharma 2005: 164, pl. 57, fig. 20; 336, pl. 133, fig. 9a–c. Thach 2005: 114, pl. 36, fig. 2. Merle et al. 2011: 99, text-fig. 39.
Chicoreus capucinus—Wilson 1994: 27, pl. 3, fig. 12a, b. Middelfart 1997: 358, pl. 2, fig. 7. Hylleberg and Kilburn 2003: 69. Sri-aroon et al. 2005: tables 2, 3, 5, 6. Houart 2008: 154, pl. 372, fig. 1. Printrakoon et al. 2008: table 1. Nabhitabhata 2009: 141. BEDO 2017b: 300, with in-text fig. Sanpanich and Duangdee 2013: 62. Dechruksa et al. 2014: fig. 4d. Wells et al. 2021: 113.

Chicoreus (Naquetia) capucinus—Subba Rao and Dey 2000: 100.
Chicoreus copucinus [sic]—Sri-aroon et al. 2004: table 1.

Referred material. CUF-NKNY-G11, G12 (30 shells; Figs 4E, 7E).

Habitat. Muddy sand and rocks in mangrove forests (Thach 2005).

Distribution. Indo-West Pacific, from the Philippines to Australia to Fiji Islands (Houart 1992).

Records of fossils from the Middle Miocene to Late Pliocene in Indonesia (Dharma 2005).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Houart (1992) and Merle et al. (2011), specifically in having (1) a rounded last whorl usually associated with three spineless varices, each with a short and webbed expansion on adapical section, (2) an aperture with a characteristically small callus with straight, adherent columellar lips, and (3) a broad siphonal canal. See also comprehensive taxonomic remarks in Houart (1992).

Indothais Claremont et al., 2013

Indothais gradata (Jonas, 1846)

Figs 4F, 7F

Purpura gradata Jonas, 1846: 14, 15. Type locality: Indian Ocean, near Singapore.

Thais gradata—Bussarawit 1991: 33. Middelfart 1997: 367, 368, pl. 4, fig. 1. Tan 2000: 499.

Hylleberg and Kilburn 2003: 76. Dharma 2005: 170, pl. 60, fig. 11a, b; 340, pl. 135, fig. 1a–e. Printrakoon et al. 2008: table 1. Yang et al. 2017: 76, fig. 316.

Stramonita gradata—Wilson 1994: 47, pl. 4, fig. 5a, b. Thach 2005: 128, pl. 37, figs 33, 37.

Indothais gradata—Okutani 2017: 963, pl. 255, fig. 9. Wells et al. 2021: 114.

Referred material. CUF-NKNY-G29 (1 shell; Figs 4F, 7F).

Habitat. Rocky bottom at a depth down to 5 m (Thach 2005; Okutani 2017).

Distribution. Indo-West Pacific, from Japan to Australia (Okutani 2017; Yang et al. 2017).

Records of fossils from the Middle to Late Pliocene in Indonesia (Dharma 2005).

Record in Thailand. Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Middelfart (1997), specifically in having (1) an erect and crenulate outer lip, which has a pronounced indentation in the subsutural ramp area, and (2) an aperture with four denticles with proceeding lirae present inside.

Indothais lacera (Born, 1778)

Figs 4C, 8A

Murex lacerus Born, 1778: 107, 108. Type locality: unknown.

Cymia lacera—Bussarawit 1991: 32. Poutiers 1998b: 561, with in-text figs. Thach 2005: 123, pl. 36, fig. 1.

Thais lacera—Bosch et al. 1995: 123, fig. 491. Middelfart 1997: 369, pl. 4, fig. 4. Subba Rao and Dey 2000: 113. Tan 2000: 499. Swennen et al. 2001: 56, 127, fig. 422. Hylleberg and Kilburn 2003: 76, 77. Robba et al. 2003: table 2. Robba et al. 2004: 102, pl. 13, fig. 6a, b.

Dharma 2005: 170, pl. 60, fig. 12a–c; 338, pl. 134, fig. 15a–h. Dey 2006: 40, 41, figs 45, 46. Ramakrishna et al. 2007: 11, 93, pl. 7, figs 79, 80. Robba et al. 2007: 92 (appendix). Nolf 2009: 9, 10, figs 1–14. Kumar et al. 2017: 1101, fig. 1a, b.
Cuma lacera—Nabhitabhata 2009: 143. Yang et al. 2017: 78, fig. 324.
Indothais lacera—Sanpanich and Duangdee 2013: 63. BEDO 2017b: 305, with in-text fig. Kantharajan et al. 2017: table 1, fig. 3–11. Tudu et al. 2018: table 1. Wells et al. 2021: 114.

Referred material. CUF-NKNY-G08 (106 shells; Figs 4C, 8A).

Habitat. Mangrove forests (Kantharajan et al. 2017) and rocky bottom at a depth down to 25 m deep (Thach 2005).

Distribution. South Africa to India; Indo-West Pacific, from Japan to New Caledonia (Poutiers 1998b; Robba et al. 2004; Kantharajan et al. 2017; Yang et al. 2017) as well as from western Mediterranean Sea (Nolf 2009). Records of fossils from the Middle Miocene to Quaternary in Indonesia and from the Holocene in Thailand (Robba et al. 2004; Dharma 2005).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Middlefart (1997) and Poutiers (1998b), specifically in having (1) a body whorl bearing one to two spiral ridges, in which the prominent one bears eight spines, and (2) an elaborate crenulate outer lip with a strong incision where the spiral ridge(s) and the anal sulcus are positioned.

Murex Linnaeus, 1758

***Murex trapa* Röding, 1798**

Figs 4B, 8B

Murex trapa Röding, 1798: 145. Type locality: Tranquebar, India. Tantanasiriwong 1978: 13, fig. 174. Nateewathana et al. 1981: 59. Way and Purchon 1981: 318. Ponder and Vokes 1988: 41–45, figs 17–19, 67g, h; 71b, c, 73d, 83g, h. Bussarawit 1991: 31. Middelfart 1997: 352, pl. 1, fig. 1. Poutiers 1998b: 565, with in-text figs. Subba Rao and Dey 2000: 102. Tan 2000: 503. Swennen et al. 2001: 55, 126, fig. 410. Hylleberg and Kilburn 2003: 72, 73. Robba et al. 2003: table 5. Robba et al. 2004: 96, 97, pl. 12, fig. 9a, b. Dharma 2005: 158, pl. 54, fig. 4a, b; 334, pl. 132, fig. 3a–c. Ramakrishna et al. 2007: 10, 85, 86. Robba et al. 2007: 92 (appendix). Houart 2008: 136, pl. 636, fig. 2. Nabhitabhata 2009: 147. BEDO 2017b: 312, with in-text fig. Yang et al. 2017: 70, fig. 288. Surakiatchai et al. 2018: table 5, pl. 1, fig. 7a, b. Tudu et al. 2018: table 1. Wells et al. 2021: 116.

Murex cf. trapa—Aungtonya and Hylleberg 1998: 319.

Murex (Murex) trapa—Gemert 2003: 105. Thach 2005: 119, pl. 34, fig. 9. Merle et al. 2011: 59, text-fig. 27e; pl. 4, fig. 7; pl. 5, figs 8–10. Gopalakrishnan et al. 2012: 77. Sanpanich and Duangdee 2013: 63. Okutani 2017: 947, pl. 238, fig. 6.

Referred material. CUF-NKNY-G09 (33 shells; Figs 4B, 8B).

Habitat. Fine sand and muddy sand bottoms at a depth of 5 to 60 m (Swennen et al. 2001; Robba et al. 2004).

Distribution. Indian Ocean; Indo-West Pacific, from Japan to Fiji Islands (Dharma 2005; Okutani 2017). Records of fossils from the Middle Miocene to Holocene in Indonesia, Malaysia,

the Philippines, Taiwan, and Thailand (Ponder and Vokes 1988; Robba et al. 2004; Dharma 2005; Surakiatchai et al. 2018).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from other related species in having (1) higher spires, (2) angulated whorls, and (3) shorter spines (Ponder and Vokes 1988). See also comprehensive taxonomic remarks in Ponder and Vokes (1988).

Superfamily Conoidea Fleming, 1822

Family Borsoniidae Bellardi, 1875

***Maoritomella* Powell, 1942**

***Maoritomella vallata* (Gould, 1860)**

Figs 4S, 9A

Drillia vallata Gould, 1860: 336. Type locality: Hong Kong. Johnson 1964: 164, pl. 7, fig. 6.

Asthenotoma vallata—Yen 1944: 575, pl. 51, figs 6, 7.

Maoritomella vallata—Robba et al. 2003: table 5. Robba et al. 2004: 134, 135, pl. 18, fig. 5a, b.
Robba et al. 2007: 94 (appendix). Wells et al. 2021: 100.

Referred material. CUF-NKNY-G30 (5 shells; Figs 4S, 9A).

Habitat. Probably present in shallow sublittoral areas (Robba et al. 2004).

Distribution. Indo-West Pacific. Records of fossils from the Holocene in Thailand (Robba et al. 2004).

Record in Thailand. Gulf of Thailand (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Yen (1944), Johnson (1964), and Robba et al. (2004), specifically in having (1) a high-spired and claviform shell, with (2) the sculpture of one rather strong and peripheral spiral cord, (3) a weaker one abapical to it, and (4) one or two spiral threads on the shoulder slope with a total of 12–14 spirals occurring along the body whorl. This species is poorly known and has been assigned to *Maoritomella* due to a blunt, smooth, paucispiral protoconch (1.5 whorls in this species), which is the diagnostic character of the genus (Powell 1942).

Family Clathurellidae H. Adams & A. Adams, 1858

***Pseudoetrema* Oyama, 1953**

***Pseudoetrema fortilirata* (Smith, 1879)**

Figs 4M, 9B

Drillia fortilirata Smith, 1879b: 194, pl. 19, fig. 22. Type locality: Station 14, east of Goto Islands, 32° 48½' N. lat., 129° 6' E. long; 47 fathoms, and station 21, between southwestern extremity of Nippon and the island of Shikoku, 33° 45' N. lat., 132° 30' E. long., 30 fathoms. Yokoyama 1927: 393, 410, pl. 46, fig. 20.

Pseudoetrema fortilirata—Oyama 1953: 154. Robba et al. 2004: 141, pl. 19, fig. 6. Robba et al. 2007: 94 (appendix). Okutani 2017: 1017, pl. 311, fig. 7. Wells et al. 2021: 101.

Referred material. CUF-NKNY-G38 (5 shells; Figs 4M, 9B).

Habitat. Sand and sandy mud in sublittoral areas to a depth of 50 m (Robba et al. 2004; Okutani 2017).

Distribution. Japan and Thailand. Records of fossils from the Pliocene to Holocene in Japan, Taiwan, and Thailand (Robba et al. 2004).

Record in Thailand. Gulf of Thailand (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Smith (1879b), Yokoyama (1927), and Robba et al. (2004), specifically in having (1) a high-spired, claviform shell with (2) the sculpture of nine broad and low collabral ribs fading away shortly before the subsutural cord and overriding spirals, (3) three closely set spiral cords: adapical, peripheral, and abapical forms, and (4) a weaker more adapical one on the penultimate whorl with a total of 14 spirals occurring along the body whorl. This species is the type species of *Pseudoetrema* (Oyama 1953).

Family Clavatulidae Gray, 1853

Turridula Schumacher, 1817

Turridula javana (Linnaeus, 1767)

Figs 4I, 8C

Murex javanus Linnaeus, 1767: 1221. Type locality: Java.

Turridula (Vulpecula) javana—Tesch 1915: 48, 49, pl. 80, fig. 104a, b.

Turridula javana—Powell 1969: 235–237, pl. 192, figs 10, 11; pl. 201, fig. 11. Tantanasiriwong 1978: 17. Nateewathana et al. 1981: 61. Way and Purchon 1981: 320. Poutiers 1998b: 630, with in-text figs. Swennen et al. 2001: 57, 133, fig. 463. Hylleberg and Kilburn 2003: 104. Robba et al. 2003: tables 3, 5. Robba et al. 2004: 136, pl. 18, fig. 8a, b. Dharma 2005: 130, pl. 40, fig. 8a, b. Thach 2005: 214, pl. 59, fig. 16; pl. 60, fig. 24. Ramakrishna et al. 2007: 15, 133, 134. Robba et al. 2007: 93 (appendix). Nabhitabhata 2009: 196. Sanpanich and Duangdee 2013: 60. BEDO 2017b: 354, with in-text fig. Yang et al. 2017: 114, fig. 472. Surakiatchai et al. 2018: table 5, pl. 1, fig. 11a, b. Tudu et al. 2018: table 1. Wells et al. 2021: 101.

Referred material. CUF-NKNY-G15, 20 (53 shells; Figs 4I, 8C).

Habitat. Intertidal sand and rocks and muddy bottoms at a depth from 20 to 80 m (Robba et al. 2004; Thach 2005).

Distribution. Indian Ocean; Indo-West Pacific, from Japan to Australia (Robba et al. 2004; Yang et al. 2017). Records of fossils from the Late Miocene to Holocene in India, Indonesia, and Thailand (Robba et al. 2004; Surakiatchai et al. 2018).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Powell (1969) and Robba et al. (2004), specifically in having (1) an elongate-fusiform shell with (2) the sculpture of oblique nodes on the peripheral angulation and finer spiral cords over the shoulder slope, and (3) a twin cord forming a weak subsutural margin. See also comprehensive taxonomic remarks in Powell (1969).

Family Horaiclavidae Bouchet et al., 2011

***Paradrillia* Makiyama, 1940**

***Paradrillia melvilli* Powell, 1969**

Figs 4N, 9C

Paradrillia melvilli Powell, 1969: 314, 315, pl. 242, fig. 2; pl. 245, figs 1, 2. Type locality: Persian Gulf. Robba et al. 2004: 137, 139, pl. 19, fig. 2. Robba et al. 2007: 95 (appendix). Wells et al. 2021: 107.

Referred material. CUF-NKNY-G52 (7 shells; Figs 4N, 9C).

Habitat. Sublittoral and upper bathyal zones (Robba et al. 2004).

Distribution. Persian Gulf and Indian Ocean; Gulf of Thailand. Records of fossils from the Holocene in Thailand (Robba et al. 2004).

Record in Thailand. Gulf of Thailand (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Powell (1969) and Robba et al. (2004), specifically in having (1) a light built claviform shell and (2) with a medially situated blunt peripheral keel bearing rather weak cog-like axial nodes, approximately 18 or 19 per whorl.

Family Pseudomelatomidae Morrison, 1966

***Inquisitor* Hedley, 1918**

***Inquisitor vulpionis* Kuroda & Oyama, 1971**

Figs 4H, 8D

Inquisitor vulpionis Kuroda & Oyama in Kuroda et al., 1971: 215, pl. 56, fig. 4; pl. 110, fig. 15.

Type locality: Sagami Bay, Japan. Robba et al. 2004: 133, pl. 18, fig. 2a–c. Robba et al. 2007: 93 (appendix). Wells et al. 2021: 108.

Referred material. CUF-NKNY-G28 (1 shell; Figs 4H, 8D).

Habitat. Sandy bottoms at a depth from 10 to 100 m (Robba et al. 2004).

Distribution. Japan and the Philippines. Records of fossils from the Holocene in Thailand (Robba et al. 2004).

Record in Thailand. Gulf of Thailand (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Robba et al. (2004), specifically in having (1) a claviform shell with (2) the sculpture of nine collabral ribs resulting in a nodose appearance at the periphery and gradually fading away on the shoulder slope, and (3) very faint spiral sculpture all over the shell except for the base where more prominent cords are developed.

***Comitas* Finlay, 1926**

***Comitas ilariae* Bozzetti, 1991**

Figs 4Q, 9D

Comitas ilariae Bozzetti, 1991: 26–28, figs 1–3. Type locality: island of Bohol, central Philippines. Olivera and Sysoev 2008: 786, pl. 688, figs 9–10.

Referred material. CUF-NKNY-G55 (1 shell; Figs 4Q, 9D).

Habitat. Sandy bottoms at a depth from 100 to 150 m (Bozzetti 1991).

Distribution. Known only from the type locality (Bozzetti 1991).

Record in Thailand. New record in Thailand from this study.

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Bozzetti (1991) and Olivera and Sysoev (2008), specifically in having (1) a slight, fusiform, high-spired shell with (2) the axial sculpture made of 7–9 very pronounced, long, and thick tubercles positioned obliquely on each whorl, and (3) numerous dense spiral cords.

Family Terebridae Mörch, 1852

***Duplicaria* Dall, 1908**

***Duplicaria tricincta* (Smith, 1877)**

Figs 4O, 10A

Terebra tricincta Smith, 1877: 225. Type locality: Persian Gulf. Bratcher and Cernohorsky 1987: 76, pl. 17, fig. 58a, b. Wilson 1994: 229, with in-text fig. Hylleberg and Kilburn 2003: 106. Robba et al. 2004: 157, pl. 21, fig. 8. Robba et al. 2007: 95 (appendix).

Granuliterebra tricincta—BEDO 2017b: 360, with in-text fig. Okutani 2017: 1050, pl. 339, fig. 14. Tudu et al. 2018: table 1. Wells et al. 2021: 109.

Duplicaria tricincta—Fedosov et al. 2019: 365, fig. 3b. Aubry et al. 2021: pl. 36. Bibi et al. 2021: 51, pl. 1, figs 9–11.

Referred material. CUF-NKNY-G33 (5 shells; Figs 4O, 10A).

Habitat. Sandy mud bottoms at a depth from 10 to 50 m (Robba et al. 2004; Okutani 2017).

Distribution. Persian Gulf; Indo-West Pacific, from Japan to Australia (Robba et al. 2004; Okutani 2017). Records of fossils from the Holocene in Thailand (Robba et al. 2004).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Bratcher and Cernohorsky (1987) and Aubry et al. (2021), and differs from its similar species, *Granuliterebra bathyrhaphis* (Smith, 1875), in having (1) a smaller shell and (2) two additional minutely beaded spiral cords abapical to the subsutural margin and on the lower base.

***Granuliterebra* Oyama, 1961**

***Granuliterebra bathyrhaphis* (Smith, 1875)**

Figs 4P, 8F

Terebra (*Myurella*) *bathyraphae* Smith, 1875: 415. Type locality: Gulf of Yedo [Edo Bay, Honshu, Japan].

Terebra sp.—Dheeradilok et al. 1984: pl. 3, fig. 11.

Terebra bathyraphae—Bratcher and Cernohorsky 1987: 75, pl. 17, fig. 57a–e. Wilson 1994: 223, with in-text fig. Bosch et al. 1995: 171, fig. 765. Robba et al. 2004: 155, pl. 21, fig. 5. Robba et al. 2007: 95 (appendix).

Granuliterebra bathyraphae—Okutani 2017: 1049, 1050, pl. 339, fig. 13. Fedosov et al. 2019: 379, fig. 10a. Aubry et al. 2021: pl. 73. Bibi et al. 2021: 50, pl. 1, figs 14–17. Wells et al. 2021: 109.

Referred material. CUF-NKNY-G66 (1 shell; Figs 4P, 8F).

Habitat. Muddy and sandy bottoms from intertidal zones down to 200 m deep (Robba et al. 2004; Aubry et al. 2021).

Distribution. Persian Gulf and Indian Ocean; Indo-West Pacific, from Japan to Australia (Robba et al. 2004; Okutani 2017; Aubry et al. 2021). Records of fossils from the Quaternary in Japan and Thailand (Robba et al. 2004).

Record in Thailand. Gulf of Thailand (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Bratcher and Cernohorsky (1987) and Aubry et al. (2021), and differs from its similar species, *Duplicaria tricincta* (Smith, 1877), by (1) lacking the band anterior to the subsutural band and by (2) having the other two bands that are connected by well-developed ribs rather than by thin cords.

***Pristiterebra* Oyama, 1961**

***Pristiterebra miranda* (Smith, 1873)**

Figs 4D, 8E

Myurella miranda Smith, 1873: 267, 268. Type locality: Malacca.

Terebra miranda—Bratcher and Cernohorsky 1987: 78, pl. 18, fig. 63a, b; color pl. D, fig. 13.

Swennen et al. 2001: 57, 134, fig. 471. Robba et al. 2004: 155, 156, pl. 21, fig. 6a, b.

Dharma 2005: 114, pl. 32, fig. 5a, b. Robba et al. 2007: 95 (appendix).

Pristiterebra miranda—Aubry et al. 2021: pl. 296. Wells et al. 2021: 109.

Referred material. CUF-NKNY-G23 (17 shells; Figs 4D, 8E).

Habitat. At sea depth of 6–10 m (Bratcher and Cernohorsky 1987; Robba et al. 2004).

Distribution. Indo-West Pacific, from Thailand to Indonesia (Bratcher and Cernohorsky 1987; Robba et al. 2004). Records of fossils from the Holocene in Thailand (Robba et al. 2004).

Record in Thailand. Gulf of Thailand (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Bratcher and Cernohorsky (1987) and Aubry et al. (2021), specifically in having (1) a turreted and slightly cyrtoconoid shell with (2) the sculpture completely cancellate throughout and (3) with numerous and unevenly spaced axial cords crossed by about 6 spiral cords forming little bead-like nodes at intersections.

Subclass Heterobranchia Burmeister, 1837

Grade “Lower Heterobranchia”

Superfamily Architectonicae Gray, 1850

Family Architectonicidae Gray, 1850

***Architectonica* Röding, 1798**

***Architectonica perdix* (Hinds, 1844)**

Figs 4J, 11A

Solarium perdix Hinds, 1844: 22, 23. Type locality: Ceylon; north-west coast of Australia.

Architectonica perdix—Bieler 1993: 48–52, figs 35–38. Swennen et al. 2001: 57, 135, fig. 477.

Hylleberg and Kilburn 2003: 113. Robba et al. 2004: 158, pl. 21, fig. 10a, b. Dharma 2005: 204, pl. 77, fig. 2a, b; 358, pl. 144, fig. 7a, b. Thach 2005: 224. Robba et al. 2007: 95 (appendix). Nabhitabhata 2009: 200. BEDO 2017b: 73, with in-text fig. Yang et al. 2017: 118, 120, fig. 493. Surakiatchai et al. 2018: table 5, pl. 1, fig. 12a, b. Tudu and Balakrishnan 2018: 199, 200, fig. 1a–f. Wells et al. 2021: 131.

Referred material. CUF-NKNY-G02, G31 (155 shells; Figs 4J, 11A).

Habitat. Sandy and muddy bottoms at depth from 10 to 60 m (Bieler 1993; Thach 2005; Yang et al. 2017).

Distribution. Indian Ocean; Indo-West to Central Pacific, from China to Australia and Polynesia (Robba et al. 2004; Tudu and Balakrishnan 2018). Records of fossils from the Middle Pliocene in Indonesia, and from the Holocene in Thailand (Dharma 2005; Surakiatchai et al. 2018).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from its similar species, *Adelphotectonica reevei* (Hanley, 1862), by having (1) a much smaller protoconch and (2) a much higher upper point of attachment of the whorls (Bieler 1993). See also comprehensive taxonomic remarks in Bieler (1993).

Cohort Tectipleura Schrödl et al., 2011

Subcohort Euopisthobranchia Jörger et al., 2010

Order Cephalaspidea Fischer, 1883

Superfamily Cylichnoidea H. Adams & A. Adams, 1854

Family Cylichnidae H. Adams & A. Adams, 1854

***Cylichna* Lovén, 1846**

***Cylichna modesta* Thiele, 1925**

Figs 4R, 10B

Cylichna modesta Thiele, 1925: 241, 242 [275, 246], pl. 44 [32], fig. 7. Type locality: “Neu-Amsterdam”, Station 167 ($37^{\circ} 47'$ südl. Br., $77^{\circ} 33.7'$ östl. L., 496 m). Swennen et al. 2001: 137, fig. 497. Wells et al. 2021: 144.

Adamnestia modesta—Robba et al. 2004: 230, pl. 34, fig. 5a, b. Robba et al. 2007: 97 (appendix). Negri et al. 2014: table 3.

Referred material. CUF-NKNY-G39 (11 shells; Figs 4R, 10B).

Habitat. Sublittoral in muddy or fine sandy bottoms (Robba et al. 2004).

Distribution. Southwest Pacific. Records of fossils from the Holocene in Thailand (Robba et al. 2004).

Record in Thailand. Gulf of Thailand (Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from its similar species, *Cylichna sibogae* Schepman, 1913, by having (1) a less slender shell and (2) the adapical umbilicus unbounded by sharp angulation (Robba et al. 2004). The generic assignment to either *Adamnestia* Iredale, 1936 or *Cylichna* requires further investigation.

Subcohort Panpulmonata Jörger et al., 2010

Superorder Eupulmonata Haszprunar & Huber, 1990

Order Ellobiida Van Mol, 1967

Superfamily Ellobioidea Pfeiffer, 1854

Family Ellobiidae Pfeiffer, 1854

***Ellobium* Röding, 1798**

***Ellobium aurisjudaе* (Linnaeus, 1758)**

Figs 4G, 11B

Bulla aurisjudaе Linnaeus, 1758: 728. Type locality: unknown.

Ellobium aurisjudaе—Cernohorsky 1972: 211, pl. 60, fig. 7. Brandt 1974: 227, 228, pl. 16, fig.

94. Tantanasiwong 1978: 20, fig. 256. Nateewathana et al. 1981: 62. Way and Purchon 1981: 321. Bosch et al. 1995: 183, fig. 852. Nateewathana 1995: 100, with in-text fig. Poutiers 1998b: 643, with in-text figs. Subba Rao and Dey 2000: 187. Swennen et al. 2001: 60, 143, fig. 524. Gemert 2003: 107, with in-text fig. Hylleberg and Kilburn 2003: 133. Robba et al. 2003: tables 1, 2. Robba et al. 2004: 241, 242, pl. 35, fig. 5. Sri-aroon et al. 2004: table 1. Dharma 2005: 206, pl. 78, fig. 18; 360, pl. 145, fig. 11. Sri-aroon et al. 2005: tables 2, 3, 5, 6. Thach 2005: 232, pl. 71, figs 11, 14. Dey 2006: 56, figs 75, 76. Raven and Vermeulen 2007: 37, 39, pl. 2, figs 10–13. Robba et al. 2007: 98 (appendix). Printrakoon et al. 2008: table 1. Kesavan et al. 2009: 382, with in-text fig. Nabhitabhata 2009: 227. Groh 2010: 446, pl. 914, figs 8, 9. Hamli et al. 2013: tables 2, 3, fig. 2t. Hylleberg and Aungtonya 2013: 97. Sanpanich and Duangdee 2013: 64. Dechruksa et al. 2014: fig. 3e. BEDO 2017b: 81, with in-text fig. Tudu et al. 2018: table 1. Wells et al. 2021: 146.

Ellobium (*Ellobium*) cf. *aurisjudaе*—Matsubara and Komori 2007: 326–328, fig. 2a–d.

Ellobium (*Ellobium*) *aurisjudaе*—Matsubara and Komori 2007: fig. 2e–h.

Referred material. CUF-NKNY-G65 (25 shells; Figs 4G, 11B).

Habitat. In estuaries, mangrove and coastal forests, on salt marshes above the normal high tide line in or under rotting wood, and on sandy soil bordering a low, sandy beach (Raven and Vermeulen 2007).

Distribution. Arabian Gulf and Indian Ocean; Indo-West Pacific, from South China Sea to Australia (Poutiers 1998b; Robba et al. 2004). Records of fossils from the Early Miocene to Holocene in Indonesia, Japan, Malaysia, and Thailand (Robba et al. 2004; Dharma 2005; Matsubara and Komori 2007; Raven and Vermeulen 2007).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Brandt (1974), Poutiers (1998b), and Robba et al. (2004), specifically in having (1) an elongate-oval and unshouldered shell and (2) with the sculpture of numerous axial grooves and fine spiral lines.

Cassidula Féruſſac, 1821

Cassidula nucleus (Gmelin, 1791)

Figs 4L, 11C

Helix nucleus Gmelin, 1791: 3193. Type locality: “Tahiti” [Cooktown, Queensland, Australia].

Auricula mustelina Deshayes, 1830: 92. Type locality: “New Zealand”.

Cassidula nucleus—Cernohorsky 1972: 212, pl. 60, fig. 8. Bosch et al. 1995: 183, fig. 851.

Subba Rao and Dey 2000: 188. Swennen et al. 2001: 60, 143, fig. 526. Hylleberg and Kilburn 2003: 133. Robba et al. 2004: 242, 243, pl. 35, fig. 6. Dharma 2005: 208, pl. 79, fig. 1a, b. Thach 2005: 233, pl. 72, fig. 8. Dey 2006: 55, 56, figs 73, 74. Raven and Vermeulen 2007: 50, 51, pl. 4, figs 35, 36. Robba et al. 2007: 98 (appendix). Kesavan et al. 2009: 382, with in-text fig. Nabhitabhata 2009: 227. Groh 2010: 448, pl. 915, figs 9–11. Zvonareva and Kantor 2016: 429, 430, fig. 8s, t. BEDO 2017b: 80, with in-text fig.

Kanthalrajan et al. 2017: table 1, fig. 4-32. Yang et al. 2017: 134, fig. 545. Tudu et al. 2018: table 1. Wells et al. 2021: 145.

Cassidula mustelina—Brandt 1974: 221, pl. 16, fig. 88. Tantanasiriwong 1978: 20, fig. 259.

Nateewathana et al. 1981: 62. Way and Purchon 1981: 321. Hylleberg and Kilburn 2003: 133. Sri-aroon et al. 2004: table 1, fig. 2-9. Sri-aroon et al. 2005: tables 2–6. Dechruksa et al. 2014: fig. 3c. BEDO 2017b: 80, with in-text fig. Okutani 2017: 1128, pl. 427, fig. 4.

Referred material. CUF-NKNY-G63 (1 shell; Figs 4L, 11C).

Habitat. On mud in shaded areas and on tree trunks in mangrove and nipa palm forests, sometimes present in salt marsh or on muddy tidal flats (Brandt 1974; Raven and Vermeulen 2007; Zvonareva and Kantor 2016).

Distribution. Arabian Gulf and Indian Ocean; Indo-West Pacific, from Japan to Australia (Brandt 1974; Bosch et al. 1995; Subba Rao and Dey 2000; Okutani 2017). Records of fossils from the Holocene in Thailand (Robba et al. 2004).

Record in Thailand. Gulf of Thailand and Andaman Sea (Brandt 1974; Swennen et al. 2001; Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from its similar species, *Cassidula aurisfelis* (Bruguière, 1789), by having (1) a normal and unbifurcated columellar fold, (2) a less rounded outline with a less convex shoulder, (3) a straighter aperture, and (4) a thinner palatal ridge (Raven and Vermeulen 2007). See also comprehensive taxonomic remarks in Raven and Vermeulen (2007).

Class Bivalvia Linnaeus, 1758

Subclass Protobranchia Pelseneer, 1889

Order Nuculanida Carter et al., 2000

Superfamily Nuculanoidea H. Adams & A. Adams, 1858

Family Nuculanidae H. Adams & A. Adams, 1858

***Jupiteria Bellardi*, 1875**

***Jupiteria puellata* (Hinds, 1843)**

Figs 10C, 12O

Nucula puellata Hinds, 1843: 100. Type locality: Malacca; from ten to seventeen fathoms, coarse sand.

Nuculana puellata—Lyngé 1909: 105. Hylleberg and Kilburn 2003: 139.

Nuculana (Jupiteria) puellata—Robba et al. 2002: 53, pl. 1, fig. 4a, b. Robba et al. 2003: tables 1, 2, 4, 5. Robba et al. 2007: 83 (appendix). Nabhitabhata 2009: 279. Negri et al. 2014: table 3. Huber 2015: C666.

Jupiteria puellata—Wells et al. 2021: 50.

Referred material. CUF-NKNY-B25 (37 shells; Figs 10C, 12O).

Habitat. Subtidal on coarse sand, soft clay, and mud bottoms at a depth from 5 down to 40 m (Huber 2015).

Distribution. Indo-West Pacific, from southern China to New Guinea (Huber 2015). Records of fossils from the Holocene in Thailand (Robba et al. 2002).

Record in Thailand. Gulf of Thailand (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Robba et al. (2002), specifically in having an unsculptured surface bearing only faint growth markings.

Saccella Woodring, 1925

***Saccella mauritiana* (Sowerby I, 1833)**

Figs 10D, 12P

Nucula mauritiana Sowerby I, 1833: 15, fig. 7. Type locality: Mauritius.

Nuculana mauritiana—Lyngé 1909: 105. Swennen et al. 2001: 43, 62, fig. 4. Gopalakrishnan et al. 2012: 61.

Nuculana (Scaeoleda) caspidata [non Gould]—Robba et al. 2002: 53, pl. 1, fig. 3a, b.

Nuculana (Scaeoleda) mauritiana—Robba et al. 2003: tables 1, 3, 5. Robba et al. 2004: 245. Robba et al. 2007: 83 (appendix). Negri et al. 2014: table 3.

Saccella mauritiana [sic]—Nabhitabhata 2009: 278.

Nuculana (Saccella) mauritiana—Huber 2010: 97, with in-text fig. Huber 2015: C612.

Saccella mauritiana—Wells et al. 2021: 50.

Referred material. CUF-NKNY-B24 (20 shells; Figs 10D, 12P).

Habitat. Sublittoral on mainly mud, and clay with sand and shells at a depth from 11 down to 92 m (Huber 2015).

Distribution. Indian Ocean; Indo-West Pacific, from China to Indonesia (Huber 2015). Records of fossils from the Holocene in Thailand (Robba et al. 2002).

Record in Thailand. Gulf of Thailand and Andaman Sea (Swennen et al. 2001; Nabhitabhata 2009; Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Robba et al. (2002) and Huber (2010), specifically in having strong commarginal ridges in the umbonal area then changing into flat and somewhat imbricate low rugae which are more raised on crossing the postero-dorsal keel. This species often co-occurs with *Jupiteria puellata*, which has an unsculptured shell (Robba et al. 2002).

Superorder Pteriomorphia Beurlen, 1944

Order Arcida Stoliczka, 1871

Superfamily Arcoidea Lamarck, 1809

Family Arcidae Lamarck, 1809

***Anadara* Gray, 1847**

***Anadara inaequivalvis* (Bruguière, 1789)**

Figs 11D, 13D

Arca inaequivalvis Bruguière, 1789: 106, 107. Type locality: East India. Reeve 1844: *Arca*, pl. 8, sp. 54.

Scapharca inaequivalvis—Tantanasiriwong 1979: 4. Morris and Purchon 1981: 322.

Nateewathana et al. 1981: 63. Poutiers 1998a: 152, with in-text figs. Subba Rao and Dey 2000: 205. Robba et al. 2002: 58, 59, pl. 2, fig. 8. Robba et al. 2003: tables 4, 5. Robba et al. 2007: 83 (appendix). Nabhitabhata 2009: 294. Negri et al. 2014: table 3. Okutani 2017: 1168, pl. 468, fig. 3. Yang et al. 2017: 148, fig. 580.

Anadara (Scapharca) inaequivalvis—Vongpanich 1996: 182, figs 21–23. Aungtonya et al. 1999: 372. Dharma 2005: 242, pl. 96, fig. 10. Dey and Ramakrishna 2007: 151, 174. Huber 2010: 137, with in-text fig. Huber 2015: C3218.

Anadara cf. *inaequivalvis*—Swennen et al. 2001: 44, 65, 66, fig. 29.

Anadara inaequivalvis—Hylleberg and Kilburn 2003: 148. Poppe 2010a: 474, pl. 928, figs 5, 6. Sanpanich 2011: table 2. Gopalakrishnan et al. 2012: 59. Surakiatchai et al. 2018: table 6, pl. 3, fig. 1a, b. Tudu et al. 2018: table 1. Tudu et al. 2019: 38–40, fig. 3c–e. Wells et al. 2021: 52.

Referred material. CUF-NKNY-B03, B21 (21L+30R shells; Figs 11D, 13D).

Habitat. Sandy and muddy bottoms in intertidal and upper sublittoral zones (Robba et al. 2002; Huber 2015).

Distribution. Red Sea to India Ocean; Indo-West Pacific, from Japan to Australia as well as from Mediterranean and Black Sea (Robba et al. 2002; Tudu et al. 2019). Records of fossils from the Late Miocene and Pliocene in Indonesia and from the Holocene in Thailand (Robba et al. 2002; Surakiatchai et al. 2018).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Poutiers (1998a) and Tudu et al. (2019), specifically in having (1) an inflated, inequilateral, and roughly quadrate shell shape and (2) with 30–37 radial ribs.

***Anadara indica* (Gmelin, 1791)**

Figs 11E, 12G

Arca indica Gmelin, 1791: 3312. Type locality: the Indian Ocean. Reeve 1844: *Arca*, pl. 9, sp. 56.

Anadara (Scapharca) indica—Lynge 1909: 126, 127, pl. 2, figs 5–12.

Scapharca indica—Morris and Purchon 1981: 322. Bosch et al. 1995: 211, fig. 928. Poutiers 1998a: 153, with in-text figs. Robba et al. 2002: 59, pl. 2, fig. 9. Gemert 2003: 107. Robba et al. 2003: tables 1–5. Thach 2005: 243, pl. 75, fig. 17. Robba et al. 2007: 83 (appendix). Nabhitabhata 2009: 294. Yang et al. 2017: 148, fig. 581. Surakiatchai et al. 2018: pl. 3, fig. 3a, b.

Anadara indica—Vongpanich 1996: 183, figs 26, 27. Aungtonya et al. 1999: 372. Tudu et al. 2018: table 1. Wells et al. 2021: 52.

Anadara (Anadara) indica—Huber 2010: 136, with in-text fig. Huber 2015: C3220.

Referred material. CUF-NKNY-B06 (16L+24R shells; Figs 11E, 12G).

Habitat. In soft sand or sandy mud bottoms at bays, river mouths, and intertidal zones down to 15 m deep (Huber 2015).

Distribution. Persian Gulf to India Ocean; Indo-West Pacific, from Japan to Australia (Poutiers 1998a; Robba et al. 2002). Records of fossils from the Holocene in Thailand (Robba et al. 2002; Surakiatchai et al. 2018).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Poutiers (1998a), Robba et al. (2002), and Huber (2010), specifically in having (1) an inequilateral, laterally compressed, and subrectangular shell shape (2) with 30–36 flat radial ribs.

***Tegillarca* Iredale, 1939**

***Tegillarca granosa* (Linnaeus, 1758)**

Figs 11F, 12C

Arca granosa Linnaeus, 1758: 694. Type locality: southern European ocean. Reeve 1844: *Arca*, pl. 3, sp. 15. Yokoyama 1927: 403. Lutaenko 2015: 135, 136, pl. 1, fig. a–d.

Arca (Anadara) granosa—Lynge 1909: 118. Tesch 1920: 92, 93, pl. 137, figs 248a, b, 249a–c.

Anadara granosa—Tantanasiriwong 1979: 4. Morris and Purchon 1981: 322. Nateewathana et al. 1981: 63. Nateewathana 1995: 101 with in-text fig. Poutiers 1998a: 147, with in-text figs. Aungtonya et al. 1999: 372. Subba Rao and Dey 2000: 204. Hylleberg and Kilburn 2003: 148, pl. 1, fig. 3. Robba et al. 2002: 57, 58, pl. 2, fig. 4. Robba et al. 2003: tables 1–5. Dey 2006: 61, figs 86, 87. Dey and Ramakrishna 2007: 151, 172. Robba et al. 2007: 83 (appendix). Printrakoon et al. 2008: table 1. Nabhitabhata 2009: 284. Sanpanich 2011: table 2. Hylleberg and Aungtonya 2013: 94. Negri et al. 2014: table 3. Surakiatchai et al. 2018: table 6, pl. 2, fig. 7a, b.

Anadara (Tegillarca) granosa—Vongpanich 1996: 187, 189, figs 47–50. Dharma 2005: 242, pl. 96, fig. 9; 362, pl. 146, fig. 10. Lutaenko et al. 2019: 174, pl. 3, fig. c, d.

Tegillarca granosa—Swennen et al. 2001: 44, 66, fig. 35. Gemert 2003: 107. Thach 2005: 244, pl. 75, fig. 8. Huber 2010: 141, with in-text fig. Poppe 2010a: 486, pl. 934, figs 9, 10. Huber 2015: C3352. BEDO 2017a: 27, with in-text fig. Kantharajan et al. 2017: table 1, fig. 2–10. Okutani 2017: 1168, pl. 468, fig. 9. Yang et al. 2017: 150, fig. 586. Tudu et al. 2018: table 1. Tudu et al. 2019: 42–44, fig. 5c, d. Wells et al. 2021: 54.

Referred material. CUF-NKNY-B20 (15L+15R shells; Figs 11F, 12C).

Habitat. Mud down to 10 m deep in mangrove forests and muddy estuaries (Swennen et al. 2001; Thach 2005; Kantharajan et al. 2017).

Distribution. East Africa to India; Indo-West Pacific, from Japan to Australia and Polynesia (Poutiers 1998a; Kantharajan et al. 2017; Tudu et al. 2019). Records of fossils from the Late Miocene to Holocene in Indonesia, Japan, the Philippines, Taiwan, and Thailand (Robba et al. 2002; Dharma 2005; Surakiatchai et al. 2018).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from its similar species, *Tegillarca nodifera*, by having (1) a less elongated shell with (2) a lower number of ribs (15–21) and (3) a lower number of nodules on ribs (Tudu et al. 2019).

***Tegillarca nodifera* (Martens, 1860)**

Figs 12A, 14A

Arca nodifera Martens, 1860: 17. Type locality: Bangkok [Bangkok, Thailand].

Anadara nodifera—Nateewathana et al. 1981: 63. Nateewathana 1995: 101, with in-text fig.

Poutiers 1998a: 148, with in-text figs. Aungtonya et al. 1999: 372. Hylleberg and Kilburn 2003: 148, 149, pl. 1, fig. 5. Nabhitabhata 2009: 284. Hylleberg and Aungtonya 2013: 94.

Anadara (*Tegillarca*) *nodifera*—Vongpanich 1996: 189, 190, figs 51–54.

Tegillarca nodifera—Swennen et al. 2001: 44, 67, text-fig. 26, fig. 36. Thach 2005: 244, pl. 75, fig. 18. Huber 2010: 141, with in-text fig. Huber 2015: C3366. BEDO 2017a: 28, with in-text fig. Yang et al. 2017: 150, fig. 587. Tudu et al. 2018: table 1. Tudu et al. 2019: 44, fig. 5e, f. Wells et al. 2021: 54.

Referred material. CUF-NKNY-B01, B02, B15 (26L+31R shells; Figs 12A, 14A).

Habitat. On mud and sand bottoms, in intertidal or near mangrove forests and shallow sublittoral waters down to a depth of 10 m (Poutiers 1998a; Huber 2015).

Distribution. Eastern Indian Ocean; Indo-West Pacific, from East China Sea to Indonesia (Poutiers 1998a; Huber 2015; Tudu et al. 2019).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from its similar species, *Tegillarca granosa*, by having (1) a more elongated shell with (2) a higher number of ribs (19–23) and (3) a higher number of nodules on ribs (Tudu et al. 2019).

Family Noetiidae Stewart, 1930

***Estellacar* Iredale, 1939**

***Estellacar olivacea* (Reeve, 1844)**

Figs 12F, 14B

Arca olivacea Reeve, 1844: *Arca*, pl. 16, sp. 113. Type locality: San Nicolas, island of Zebu [the Philippines] (found in sandy mud at the depth of four fathoms).

Estellacar olivacea—Poutiers 1998a: 160, with in-text figs. Swennen et al. 2001: 44, 68, fig. 42. Robba et al. 2002: 61, pl. 2, fig. 13. Hylleberg and Kilburn 2003: 153. Robba et al. 2003: table 2. Vongpanich and Matsukuma 2004: 41, pl. 2, figs. r, s. Robba et al. 2007: 84 (appendix). Nabhitabhata 2009: 297. Huber 2010: 146, with in-text fig. Negri et al. 2014: table 3. Huber 2015: C3594. BEDO 2017a: 42, with in-text fig. Wells et al. 2021: 55.

Striarca olivacea—Printrakoon et al. 2008: table 1.

Referred material. CUF-NKNY-B18 (16L+14R shells; Figs 12F, 14B).

Habitat. In mud and sand from the intertidal zone, often near mangrove forests, down to 20 m deep (Poutiers 1998a; Robba et al. 2002).

Distribution. India; Indo-West Pacific, from South China to the Philippines (Huber 2015).

Records of fossils from the Holocene in Thailand (Robba et al. 2002).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from its similar species, *Estellacar galactodes* (Benson, 1842), by having a higher number of coarser riblets (Robba et al. 2002).

Noetiella Thiele, 1931

Noetiella pectunculiformis (Dunker, 1866)

Figs 12D, 14C

Barbatia pectunculiformis Dunker, 1866: 88, 89, pl. 28, figs 4–6. Type locality: Borneo.

Arca (Fossularca) pectunculiformis—Lyngé 1909: 115.

Striarca pectunculiformis—Tantanasiriwong 1979: 4. Morris and Purchon 1981: 323.

Nateewathana et al. 1981: 63. Aungtonya et al. 1999: 373.

Scelidionarca pectunculiformis—Robba et al. 2002: 61, pl. 2, fig. 14q, b. Robba et al. 2007: 84 (appendix). Nabhitabhata 2009: 297.

Noetiella pectunculiformis—Huber 2010: 146, with in-text fig. Huber 2015: C3575. BEDO 2017a: 43, with in-text fig. Kantharajan et al. 2017: table 1, fig. 2-11. Wells et al. 2021: 55.

Referred material. CUF-NKNY-B27 (4L+4R shells; Figs 12D, 14C).

Habitat. In soft clay, mud or sand, in subtidal zones from a depth of 2 to 43 m, and in mangrove forests (Huber 2015; Kantharajan et al. 2017).

Distribution. India Ocean; Indo-West Pacific, from South China to Indonesia (Huber 2015; Kantharajan et al. 2017). Records of fossils from the Holocene in Thailand (Robba et al. 2002).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Robba et al. (2002) and Huber (2010), specifically in having (1) a roundly trapezoidal and inequilateral shell shape with (2) the sculpture of dense, faint radials, and coarse growth markings.

Order Ostreida Féussac, 1822

Superfamily Ostreoidea Rafinesque, 1815
Family Ostreidae Rafinesque, 1815

***Magallana* Salvi & Mariottini, 2016**

***Magallana cf. gigas* (Thunberg, 1793)**
Figs 12E, 14D

Ostrea gigas Thunberg, 1793: 140–142, pl. 6, figs 1–3. Type locality: Japan. Yokoyama 1927: 402.

Crassostrea gigas—Tantanasiriwong 1979: 7. Nateewathana et al. 1981: 65. Poutiers 1998a: 233, with in-text figs. Aungtonya et al. 1999: 375. Robba et al. 2002: 69, 71, pl. 5, fig. 3a, b. Helleberg and Kilburn 2003: 162. Beu et al. 2004: 154–156, fig. 9d. Thach 2005: 257. Robba et al. 2007: 84 (appendix). Printrakoon et al. 2008: table 1. Huber 2010: 180, with in-text fig. Sanpanich 2011: table 2. Huber 2015: C5155. Okutani 2017: 1183, pl. 483, fig. 7. Yang et al. 2017: 176, 178, fig. 685. Surakiatchai et al. 2018: table 6, pl. 3, fig. 6a, b. Tudu et al. 2018: table 1.

Crassostrea cf. gigas—Swennen et al. 2001: 45, 72, text-fig. 62, fig. 62.

Crassostrea (Magallana) gigas—Lutaenko et al. 2019: 187, 188, pl. 15, fig. a–f.

Magallana gigas—Wells et al. 2021: 60.

Referred material. CUF-NKNY-B13 (130L+16R shells; Figs 12E, 14D).

Habitat. Attached to rocks from intertidal zones down to 30 m deep, in mud, bays, and sheltered areas that are often brackish with low salinity water (Thach 2005; Huber 2015).

Distribution. Cosmopolitan (Huber 2015). Records of fossils from the Miocene to Holocene in Japan, New Zealand, Taiwan, and Thailand (Robba et al. 2002; Beu et al. 2004; Surakiatchai et al. 2018).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. The cupped oysters in the subfamily Crassostreinae could not be identified at species level based on shell characters or soft tissue alone (Salvi and Mariottini 2021). Therefore, we tentatively identify these specimens of giant cupped oysters as belonging to *Magallana gigas*, due to large and thick shells and the presence of this species in Thailand (Wells et al. 2021).

Order Pectinida Gray, 1854

Superfamily Anomioidea Rafinesque, 1815
Family Placunidae Rafinesque, 1815

***Placuna* Lightfoot, 1786**

***Placuna placenta* (Linnaeus, 1758)**
Figs 13A, 14E

Anomia placenta Linnaeus, 1758: 703. Type locality: unknown.

Placuna placenta—Lyngé 1909: 107, 108. Tantanasiriwong 1979: 6. Morris and Purchon 1981: 324. Nateewathana et al. 1981: 64. Oliver 1992: 86, pl. 17, fig. 6a, b. Bosch et al. 1995:

234, fig. 1018. Poutiers 1998a: 218, with in-text figs. Aungtonya et al. 1999: 376. Subba Rao and Dey 2000: 230. Swennen et al. 2001: 46, 76, text-fig. 80, fig. 80. Robba et al. 2002: 77, pl. 6, fig. 10a, b. Hylleberg and Kilburn 2003: 169, 170. Robba et al. 2003: table 5. Dharma 2005: 246, pl. 98, fig. 17. Dey and Ramakrishna 2007: 154, 197. Robba et al. 2007: 84 (appendix). Nabhitabhata 2009: 20, with in-text fig. 347. Huber 2010: 190, with in-text fig. Sanpanich 2011: table 2. Huber 2015: C5627. Yang et al. 2017: 176, fig. 681. Surakiatchai et al. 2018: table 6, pl. 3, fig. 8a, b. Tudu et al. 2018: table 1. Wells et al. 2021: 64.

Placuna (Placuna) placenta—Dheeradilok et al. 1984: pl. 2, fig. 14. Thach 2005: 266, pl. 81, fig. 4.

Referred material. CUF-NKNY-B22 (19L+18R shells; Figs 13A, 14E).

Habitat. Sandy mud in intertidal zones and shallow waters down to 35 m deep, in quiet waters of lagoons, protected bays, and mangrove areas, or near estuaries (Poutiers 1998a; Thach 2005; Yang et al. 2017).

Distribution. Indian Ocean; Indo-West Pacific, from Japan to Australia (Huber 2015). Records of fossils from the Late Miocene to Holocene in Indonesia, Japan, the Philippines, Taiwan, and Thailand (Robba et al. 2002; Surakiatchai et al. 2018).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from its similar species, *Placuna ephippium* (Philipsson, 1788), by having (1) an almost circular shell and (2) a right valve having a hinge plate with arrowhead-shaped lamellar teeth with the posterior ridge distinctly longer than the anterior one (Poutiers 1998a; Swennen et al. 2001).

Superfamily Pectinoidea Rafinesque, 1815

Family Pectinidae Rafinesque, 1815

Volachlamys Iredale, 1939

Volachlamys singaporina (Sowerby II, 1842)

Figs 12B, 14F

Pecten singaporinus Sowerby II, 1842: 74, pl. 13, fig. 55; pl. 14, fig. 71. Type locality: Singapore. Lyngé 1909: 155.

Volachlamys singaporina—Roussy 1991: 21. Poutiers 1998a: 210, with in-text figs. Aungtonya et al. 1999: 376. Swennen et al. 2001: 45, 74, fig. 70. Gemert 2003: 108. Hylleberg and Kilburn 2003: 169. Dharma 2005: 248, pl. 99, fig. 13a–e; 364, pl. 147, fig. 3a, b. Thach 2005: 263, pl. 80, fig. 13. Nabhitabhata 2009: 342. Huber 2010: 203, with in-text fig. Huber 2015: C6118. BEDO 2017a: 236, with in-text fig. Yang et al. 2017: 172, fig. 667. Dijkstra and Beu 2018: 296–298, figs 98, 100j, l, 101d, e. Wells et al. 2021: 67.

Referred material. CUF-NKNY-B07 (32L+32R shells; Figs 12B, 14F).

Habitat. Byssally attached under coral boulders and rocks, on sand and sandy mud in intertidal zones down to 22 m deep (Huber 2015).

Distribution. Indo-West Pacific, from China to Australia (Huber 2015; Dijkstra and Beu 2018). Records of fossils from the Middle to Late Pliocene in Indonesia (Dharma 2005).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from its similar species, *Volachlamys tranquebaria* (Gmelin, 1791) and *V. hirasei* (Bavay, 1904), by having square and well-defined 18–24 ribs with wider interspaces between the ribs (Dharma 2005; Dijkstra and Beu 2018). See also comprehensive taxonomic remarks in Dijkstra and Beu (2018)

Superorder Heteroconchia Gray, 1854

Order Lucinida Gray, 1854

Superfamily Lucinoidea J. Fleming, 1828

Family Lucinidae J. Fleming, 1828

***Pegophysema* Stewart, 1930**

***Pegophysema bialata* (Pilsbry, 1895)**

Figs 13E, 15A

Loripes bialata Pilsbry, 1895: 133, 134, pl. 3, figs 13, 14. Type locality: Inland Sea, Japan.

Anodontia (?*Pegophysema*) *bialata*—Taylor and Glover 2005: 305, 306, figs 11a, 12a, 22, 23.

Pegophysema bialata—Huber 2015: 102, 455, 456, with in-text figs; C10244. Wells et al. 2021: 35.

Referred material. CUF-NKNY-B11 (3L+3R shells; Figs 13E, 15A).

Habitat. Muddy bottoms and sandy mud among weed in the intertidal zone down to 60 m deep (Huber 2015).

Distribution. Indian Ocean; Indo-West Pacific, from Japan to Malaysia (Taylor and Glover 2005; Huber 2015).

Record in Thailand. Gulf of Thailand (Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from its similar species,

Pegophysema philippiana (Reeve, 1850), by having (1) a ligament that is shallowly inset and (2) absence of secondary pallial attachment scars (Taylor and Glover 2005). See also comprehensive taxonomic remarks in Taylor and Glover (2005).

Order Venerida Gray, 1854

Superfamily Cyrenoidea Gray, 1840

Family Cyrenidae Gray, 1840

***Geloina* Gray, 1842**

***Geloina bengalensis* (Lamarck, 1818)**

Figs 13C, 15B

Cyrena bengalensis Lamarck, 1818: 554. Type locality: Bengal.

Polymesoda (*Geloina*) *bengalensis*—Brandt 1974: 310, 311, pl. 28, fig. 83. Nateewathana 1995:

106, with in-text fig. Aungtonya et al. 1999: 380. Robba et al. 2002: 106, pl. 16, fig. 9.

Robba et al. 2003: tables 1, 2. Robba et al. 2007: 87 (appendix). Nabhitabhata 2009: 21, with in-text fig. 471.

Polymesoda bengalensis—Poutiers 1998a: 319, with in-text figs. Dharma 2005: 266, pl. 108, fig.

25. Dey 2006: 74, 75, figs 113, 114. Tudu et al. 2018: table 1.

Geloina bengalensis—Huber 2015: 319; C20908. BEDO 2017a: 252, with in-text fig. Wells et al. 2021: 39.

Referred material. CUF-NKNY-B31 (2L+3R shells; Figs 13C, 15B).

Habitat. In intertidal zones, river deltas, estuaries, mud flats, and in mangrove areas (Poutiers 1998a; Huber 2015).

Distribution. Bay of Bengal; Indo-West Pacific, from Taiwan to Indonesia (Huber 2015).

Records of fossils from the Holocene in Thailand (Robba et al. 2002).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Robba et al. (2002) and Huber (2015), specifically in having a solid, inflated, pronounced inequilateral, and subtriangular shell shape.

Superfamily Mactroidea Lamarck, 1809

Family Mactridae Lamarck, 1809

***Lutraria* Lamarck, 1799**

***Lutraria complanata* (Gmelin, 1791)**

Figs 13H, 15C

Mactra complanata Gmelin, 1791: 3261. Type locality: the Indian Ocean.

Lutraria (Lutrophora) cf. complanata—Dheeradilok et al. 1984: pl. 2, fig. 5.

Lutraria complanata—Swennen et al. 2001: 47, 82, fig. 112. Hylleberg and Kilburn 2003: 186. Yang et al. 2017: 194, fig. 742. Wells et al. 2021: 40.

Lutraria (Lutrophora) complanata—Robba et al. 2002: 91, pl. 12, fig. 4. Robba et al. 2007: 86 (appendix). Huber 2010: 451, with in-text figs. Huber 2015: C21984.

Lutraria (Goniomactra) complanata—Beu 2006: 242.

Lutraria (Lutropbora) complanata [sic]—Nabhitabhata 2009: 412.

Referred material. CUF-NKNY-B29 (1L+2R shells; Figs 13H, 15C).

Habitat. Sandy and muddy sand in subtidal zones from a depth of 8 down to 20 m (Huber 2015).

Distribution. Indian Ocean; Indo-West Pacific, from South China to Indonesia (Huber 2015; Yang et al. 2017). Records of fossils from the Holocene in Thailand (Robba et al. 2002).

Record in Thailand. Gulf of Thailand (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Robba et al. (2002) and Huber (2010), specifically in having (1) an elongate-elliptical and markedly inequilateral thin shell with arched anterior and posterior margins and (2) an outer surface sculptured with irregular crenulated growth lines.

***Standella* Gray, 1853**

***Standella pellucida* (Gmelin, 1791)**

Figs 13B, 15D

Mactra pellucida Gmelin, 1791: 3260, 3261. Type locality: “Guinea coast.” Sanpanich 2011: table 2.

Standella pellucida—Lyngé 1909: 224. Nabhitabhata 2009: 418. Huber 2010: 451, with in-text figs. Huber 2015: C22028. Tudu et al. 2018: table 1. Wells et al. 2021: 42.

Meropesta pellucida—Bosch et al. 1995: 248, fig. 1100. Poutiers 1998a: 281, with in-text figs. Swennen et al. 2001: 47, 83, text-fig. 116, fig. 116. Robba et al. 2002: 91, 92, pl. 12, fig. 6. Dey and Ramakrishna 2007: 211. Robba et al. 2007: 86 (appendix). Meyer et al. 2008: tables 2, 3. Nabhitabhata 2009: 417. Wong 2009: 289, figs 9a, b, 22. BEDO 2017a: 155. Yang et al. 2017: 196, fig. 752.

Referred material. CUF-NKNY-B30 (1L+1R shells; Figs 13B, 15D).

Habitat. Fine sand and mud in intertidal zones down to 4 m deep, in lagoons, and at the fringe of mangrove areas (Poutiers 1998a; Huber 2015).

Distribution. Arabian Gulf and Indian Ocean; Indo-West Pacific, from Japan to Australia (Bosch et al. 1995; Wong 2009; Huber 2015). Records of fossils from the Quaternary in Indonesia and from the Holocene in Thailand (Robba et al. 2002).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Robba et al. (2002) and Huber (2010), specifically in having (1) a longer than high, inequilateral, and elliptical shell with an oval-shaped anterior margin and a somewhat pointed posterior margin and (2) an outer surface sculptured with uneven, fine growth markings, and faint posterior radial striation. An exceedingly shallow radial depression is sometimes observed in the mid-anterior part. Huber (2015) indicated that the type locality in Guinea is erroneous.

Superfamily Tellinoidea Blainville, 1814

Family Tellinidae Blainville, 1814

***Tellinides* Lamarck, 1818**

***Tellinides conspicuus* (Hanley, 1846)**

Figs 13J, 15E

Tellina conspicua Hanley, 1846b: 293, pl. 58, fig. 100. Type locality: unknown. Coan and Kabat 2012: 309.

Tellina (*Tellinides*) sp.—Robba et al. 2002: 98, pl. 14, fig. 7a, b.

Tellinides conspicuus—Huber 2015: 220, 639, with in-text figs; C14877. Wells et al. 2021: 30.

Referred material. CUF-NKNY-B04 (33L+40R shells; Figs 13J, 15E).

Habitat. Shallow water (Huber 2015).

Distribution. Indo-West Pacific, from southern China to New Guinea (Huber 2015).

Record in Thailand. Gulf of Thailand (Wells et al. 2021).

Taxonomic remarks and comparisons. *Tellina* (*Tellinides*) sp. in Robba et al. (2002) was identified as belonging to this species by Huber (2015). This species differs from its similar species, *Tellinides timorensis* Lamarck, 1818, by having (1) a higher shell that lacks the posterior

truncation and (2) with oblique anterior striae fading posteriorly (Robba et al. 2002; Huber 2015).

Superfamily Ungulinoidea Gray, 1854

Family Ungulinidae Gray, 1854

***Joannisiella* Dall, 1895**

***Joannisiella oblonga* (Hanley, 1846)**

Figs 12S, 15F

Cyrenoidea oblonga Hanley, 1846a: 10 (plate explanation). Type locality: Philippines. Hanley 1856: 353, pl. 15, fig. 6. Coan and Kabat 2012: 320.

Diplodonta (Joannisiella) oblonga—Lynge 1909: 176.

Cycladicama oblonga—Dheeradilok et al. 1984: pl. 2, figs 6, 7. Robba et al. 2002: 79, pl. 7, fig. 9a, b. Hylleberg and Kilburn 2003: 174. Robba et al. 2003: tables 3, 4. Robba et al. 2007: 85 (appendix). Nabhitabhatta 2009: 389. Yang et al. 2017: 182, fig. 704. Surakiatchai et al. 2018: table 6, pl. 3, fig. 10a, b.

Joannisiella oblonga—Huber 2015: 341, 829, with in-text figs; C21436. Wells et al. 2021: 42.

Referred material. CUF-NKNY-B09, B17 (59L+66R shells; Figs 12S, 15F).

Habitat. Muddy bottoms with soft clay, and muddy sediments on mud flats, in the shallow waters from intertidal zones down to 36 m deep (Huber 2015; Yang et al. 2017).

Distribution. Indo-West Pacific, from Japan to Indonesia and the Philippines (Huber 2015; Yang et al. 2017). Records of fossils from the Middle Miocene to Holocene in Indonesia, Japan, the Philippines, Taiwan, and Thailand (Robba et al. 2002; Surakiatchai et al. 2018).

Record in Thailand. Gulf of Thailand (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is characterized by its elongate, ovate-triangular shaped, and posteriorly grooved shell (Robba et al. 2002; Huber 2015).

Superfamily Veneroidea Rafinesque, 1815

Family Veneridae Rafinesque, 1815

***Dosinia* Scopoli, 1777**

***Dosinia dilecta* A. Adams, 1856**

Figs 12R, 16A

Dosinia dilecta A. Adams, 1856: 224. Type locality: Malacca. Lynge 1909: 249, 250, pl. 5, figs 11–13. Morris and Purchon 1981: 326. Swennen et al. 2001: 49, 95, text-fig. 201, fig. 201. Robba et al. 2002: 115, pl. 19, fig. 4. Dharmia 2005: 266, pl. 108, fig. 30. Robba et al. 2007: 88 (appendix). Sartori et al. 2008: table 1. Nabhitabhatta 2009: 482. Poppe 2010c: 304, pl. 1147, figs 5, 6. Surakiatchai et al. 2018: table 6, pl. 5, fig. 8a, b. Wells et al. 2021: 44.

Dosinia (Dosinella) dilecta—Fischer-Piette and Delmas 1967: 77, pl. 14, figs 1–3. Dheeradilok et al. 1984: pl. 2, figs 9–12. Huber 2010: 414, with in-text fig. Huber 2015: C19500.

Referred material. CUF-NKNY-B10 (2L+5R shells; Figs 12R, 16A).

Habitat. Soft clay and mud bottoms in subtidal zones at a depth from 5 to 20 m (Huber 2015).

Distribution. Indo-West Pacific, from southern China to Indonesia (Huber 2015). Records of fossils from the Holocene in Thailand (Robba et al. 2002; Surakiatchai et al. 2018).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Fischer-Piette and Delmas (1967), Robba et al. (2002), and Huber (2010), specifically in having (1) deeply demarcated and exceedingly small lunulae and (2) a shell with low concentric ribs. See also comprehensive taxonomic remarks in Fischer-Piette and Delmas (1967).

Paratapes Stoliczka, 1870

Paratapes undulatus (Born, 1778)

Figs 12Q, 16B

Venus undulata Born, 1778: 54, 55. Type locality: unknown.

Tapes (Paratapes) undulatus—Lynge 1909: 237.

Paphia undulata—Tantanasiriwong 1979: 13. Morris and Purchon 1981: 326. Nateewathana et al. 1981: 67. Dheeradilok et al. 1984: pl. 2, fig. 13. Oliver 1992: 191, pl. 43, fig. 6a, b. Bosch et al. 1995: 273, fig. 1227. Nateewathana 1995: 111, with in-text fig. Poutiers 1998a: 339, with in-text figs. Aungtonya et al. 1999: 382. Subba Rao and Dey 2000: 283. Swennen et al. 2001: 49, 95, text-fig. 198, fig. 198. Hylleberg and Kilburn 2003: 217. Dhama 2005: 270, pl. 110, fig. 16; 368, pl. 149, fig. 13. Thach 2005: 300, pl. 90, fig. 3. Dey and Ramakrishna 2007: 160, 244, 245. Nabhitabhata 2009: 492. Poppe 2010c: 296, pl. 1143, figs 7–9. Sanpanich 2011: table 2. Okutani 2017: 1248, pl. 543, fig. 4. Yang et al. 2017: 232, fig. 871. Surakiatchai et al. 2018: table 6, pl. 5, fig. 4a, b.

Paphia cf. undulata—Aungtonya and Hylleberg 1998: 320.

Paphia (Paphia) undulata—Gemert 2003: 109. Robba et al. 2002: 113, pl. 18, fig. 8a, b. Robba et al. 2003: tables 1, 3–5. Robba et al. 2007: 88 (appendix). Negri et al. 2014: table 3.

Neotapes undulata—Sartori et al. 2008: 115, table 1, figs 4l, 5l.

Paphia (Neotapes) undulata—Huber 2010: 423, with in-text fig.

Paratapes undulatus—Huber 2015: C19932. Tudu et al. 2018: table 1. Wells et al. 2021: 47.

Paratapes undulata—BEDO 2017a: 281, with in-text fig.

Referred material. CUF-NKNY-B12 (2L shells; Figs 12Q, 16B).

Habitat. Fine sandy sediments and mud as well as silty clay bottoms in intertidal to subtidal zones down to a depth of 50 m (Huber 2015).

Distribution. Red Sea and Indian Ocean; Indo-West Pacific, from Japan to Australia (Robba et al. 2002; Huber 2015). Records of fossils from the Late Miocene to Holocene in Indonesia, Japan, the Philippines, Taiwan, and Thailand (Robba et al. 2002; Dhama 2005; Surakiatchai et al. 2018).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. Despite having no colour and patterns, the elongate-oval and inequilateral shell shape with growth lines as well as slightly oblique and somewhat

undulating commarginal grooves on outer surface of the specimens conform to the characters of this species (Robba et al. 2002; Huber 2010).

***Placamen* Iredale, 1925**

***Placamen lamellatum* (Röding, 1798)**

Figs 12L, 16C

Venus lamellata Röding, 1798: 183. Type locality: the East Indian Seas.

Venus tiara Dillwyn, 1817: 162. Type locality: the East Indian Seas.

Venus calophylla Philippi, 1836: 229, 230, pl. 8, fig. 2 Type locality: the Chinese Sea.

Chione (Circomphalus) calophylla—Lynge 1909: 150, 151.

Callanaitis calophylla—Tantanasiriwong 1979: 13. Nateewathana et al. 1981: 67.

Placamen tiara—Tantanasiriwong 1979: 13. Nateewathana et al. 1981: 67. Poutiers 1998a: 351, with in-text figs. Aungtonya et al. 1999: 382. Hylleberg and Kilburn 2003: 219. Dharma 2005: 268, pl. 109, fig. 24. Thach 2005: 292. Dey and Ramakrishna 2007: 159, 238. Nabhitabhata 2009: 495. Poppe 2010c: 274, pl. 1132, figs 1–3.

Bassina calophylla—Bosch et al. 1995: 266, fig. 1191. Subba Rao and Dey 2000: 284.

Placamen calophyllum—Kilburn and Hylleberg 1998: 315. Aungtonya et al. 1999: 382.

Swennen et al. 2001: 48, 92, fig. 180. Robba et al. 2002: 107, pl. 17, fig. 4. Gemert 2003: 15. Hylleberg and Kilburn 2003: 218, 219. Dharma 2005: 268, pl. 109, fig. 25. Thach 2005: 292, pl. 89, fig. 11. Robba et al. 2007: 87 (appendix). Meyer et al. 2008: tables 2, 4. Sartori et al. 2008: table 1. Nabhitabhata 2009: 494. Poppe 2010c: 272, pl. 1131, figs 4–7.

Sanpanich 2011: table 2.

Placamen lamellatum—Huber 2010: 369, with in-text figs. Huber 2015: C18043. BEDO 2017a: 287. Tudu et al. 2018: table 1. Wells et al. 2021: 48.

Clausinella calophylla—Yang et al. 2017: 222, 224, fig. 834.

Clausinella tiara—Yang et al. 2017: 224, fig. 836.

Referred material. CUF-NKNY-B28 (1R shell; Figs 12L, 16C).

Habitat. Muddy sand, sand, and shell gravels from intertidal zones down to 100 m deep (Bosch et al. 1995; Robba et al. 2002; Yang et al. 2017).

Distribution. Arabian Gulf and Indian Ocean; Indo-West Pacific, from Japan to Indonesia (Bosch et al. 1995; Huber 2015; Yang et al. 2017). Records of fossils from the Late Neogene to Holocene in Fiji, India, Indonesia, Japan, the Philippines and Taiwan (Robba et al. 2002).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from its similar species, *Placamen chloroticum* (Philippi, 1849), by having (1) a somewhat shallower and trigonal pallial sinus and (2) more widely spaced and more raised commarginal lamellae (Robba et al. 2002).

Order Myida Stoliczka, 1870

Superfamily Myoidea Lamarck, 1809

Family Corbulidae Lamarck, 1818

***Corbula* Bruguière, 1797**

***Corbula fortisulcata* Smith, 1879**

Figs 12J, 16D

Corbula fortisulcata Smith, 1879a: 819, 820, pl. 50, fig. 23, 23b. Type locality: the Andaman Islands. Subba Rao and Dey 2000: 286. Swennen et al. 2001: 49, 97, text-fig. 212, fig. 212. Hylleberg and Kilburn 2003: 222. Poppe 2010b: 388, pl. 1189, figs 6, 7a, b. Yang et al. 2017: 240, 242, fig. 899. Surakiatchai et al. 2018: table 6, pl. 5, fig. 10a, b. Wells et al. 2021: 36.

Corbula (Corbula) fortisulcata—Robba et al. 2002: 117, 118, pl. 20, fig. 5. Robba et al. 2003: table 3. Robba et al. 2007: 88 (appendix). Nabhitabhata 2009: 506.

Corbula (Notocorbula) cf. fortisulcata—Thach 2005: 303, pl. 91, fig. 4.

Corbula (Notocorbula) fortisulcata—Huber 2010: 467, with in-text fig. Huber 2015: C22606.

Referred material. CUF-NKNY-B19 (154 shells; Figs 12J, 16D).

Habitat. Fine sand, sandy mud and muddy bottoms at intertidal to sublittoral zones down to a depth of 70 m (Huber 2015).

Distribution. Indian Ocean; Indo-West Pacific, from Taiwan to Australia (Huber 2015). Records of fossils from the Late Miocene in the Philippines and from the Holocene in Thailand (Robba et al. 2002; Surakiatchai et al. 2018).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species differs from its similar species, *Corbula tunicata* Reeve, 1843, by having a weaker keel on the juvenile shell part (Swennen et al. 2001).

***Potamocorbula* Habe, 1955**

***Potamocorbula* sp.**

Figs 12N, 16E

Referred material. CUF-NKNY-B26 (112 shells; Figs 12N, 16E).

Habitat. Mostly present in intertidal zones, estuarine and brackish habitats (Huber 2015).

Distribution. This genus is mostly distributed in the Indian Ocean and Indo-West Pacific, except for the species *Potamocorbula adusta* (Reeve, 1844), which is recorded only from West Africa (Huber 2015).

Taxonomic remarks and comparisons. These specimens are assigned to *Potamocorbula* based on the descriptions in Coan (2002), specifically in having smooth shells. Currently, there are a total of nine species in this genus (Huber 2015; MolluscaBase 2023), in which two species, *P. fasciata* (Reeve, 1843) and *P. laevis* (Hinds, 1843), are recorded from Thailand (Wells et al. 2021). However, our specimens have a more quadrate shell shape, probably suggesting the presence of other different species.

Superfamily Pholadoidea Lamarck, 1809

Family Pholadidae Lamarck, 1809

***Martesia* Sowerby I, 1824**

***Martesia striata* (Linnaeus, 1758)**

Figs 12M, 16F, G

Pholas striatus Linnaeus, 1758: 669. Type locality: southern Europe.

Pholas (Martesia) striata—Lyngé 1909: 283, 284.

Martesia striata—Cernohorský 1978: 188, pl. 68, fig. 4. Morris and Purchon 1981: 327. Oliver 1992: 203, pl. 45, fig. 4a, b. Bosch et al. 1995: 279, fig. 1263. Yoosukh and Jitkaew 1997: 402, fig. 1a, b. Poutiers 1998a: 358, with in-text figs. Subba Rao and Dey 2000: 287. Swennen et al. 2001: 49, 99, fig. 223. Hylleberg and Kilburn 2003: 224. Dey and Ramakrishna 2007: 161, 252. Nabhitabhata 2009: 21, with in-text fig. 509. Haga 2010: 394, pl. 1192, figs 3, 4. Gopalakrishnan et al. 2012: 61. Okutani 2017: 1270, pl. 563, fig. 5. Yang et al. 2017: 246, fig. 914. Tudu et al. 2018: table 1. Wells et al. 2021: 37.

Martesia (Martesia) striata—Robba et al. 2002: 121, 122, pl. 21, fig. 7a. Thach 2005: 304, pl. 91, fig. 10. Robba et al. 2007: 23, fig. 9p; 88 (appendix). Huber 2010: 476, with in-text figs. Huber 2015: C22956.

Referred material. CUF-NKNY-B14 (8L+8R shells; Figs 12M, 16F, G).

Habitat. Bored in old wood pilings, waterlogged tree trunks, and in driftwood but rarely in astreid corals or in soft rocks or mangroves *Rhizophora stylosa*, from intertidal zones down to 20 m deep (Huber 2015).

Distribution. Cosmopolitan (Huber 2015). Records of fossils from the Middle Miocene to Holocene in Japan, Indonesia, New Hebrides, and Thailand (Robba et al. 2002).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Poutiers (1998a), Robba et al. (2002), and Huber (2010), specifically in having (1) an outer surface bearing a radial groove at the anterior one-third of the shell and (2) sculptured with commarginal ridges that are minutely nodose and anterior to the radial groove with nodes arranged into radial rows.

Pholas Linnaeus, 1758

Pholas orientalis Gmelin, 1791

Figs 12U, 16H

Pholas orientalis Gmelin, 1791: 3216. Type locality: Siam [Thailand] and Tranquebar. Morris and Purchon 1981: 327. Nateewathana 1995: 111, with in-text fig. Poutiers 1998a: 356, with in-text figs. Subba Rao and Dey 2000: 286. Swennen et al. 2001: 50, 98, fig. 219. Haga 2010: 396, pl. 1193, figs 2, 3. BEDO 2017a: 168, with in-text fig. Hylleberg and Kilburn 2003: 224. Dey and Ramakrishna 2007: 160, 251, 252. Nabhitabhata 2009: 510. Gopalakrishnan et al. 2012: 61. Yang et al. 2017: 246, fig. 916. Tudu et al. 2018: table 1. Wells et al. 2021: 37.

Pholas siamensis Spengler, 1792: 88, 89. Type locality: the Gulf of Thailand, the mouth of the river Qweda, where it goes up to Alastav. Knudsen and Jensen 2001: 547–556, figs 1–4.

Pholas (Monothyra) orientalis—Lyngé 1909: 282. Aungtonya et al. 1999: 383. Robba et al. 2002: 121, pl. 21, fig. 5. Robba et al. 2003: tables 3–5. Dharma 2005: 270, pl. 110, fig. 25; 368, pl. 149, fig. 18a, b. Thach 2005: 304, pl. 91, fig. 6. Robba et al. 2007: 88 (appendix). Huber 2010: 473, with in-text fig. Huber 2015: C22846.

Referred material. CUF-NKNY-B05 (26L+73R shells; Figs 12U, 16H).

Habitat. Boring down to 50 cm deep in peat, soft rocks, stiff clay, or sticky and soft sandy-mud bottoms rich in silt and detritus, often near river mouths, from intertidal and sublittoral zones down to 30 m deep (Poutiers 1998a; Huber 2015).

Distribution. Indian Ocean; Indo-West Pacific, from Taiwan to Australia (Huber 2015). Records of fossils from the Middle Miocene to Holocene in Indonesia, Myanmar, and Thailand (Robba et al. 2002; Dharma 2005).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Poutiers (1998a), Swennen et al. (2001), and Robba et al. (2002), specifically in having antero-dorsal reflection that has axial supports.

Famliy Teredinidae Rafinesque, 1815

Teredinidae indet.

Fig. 16I

Referred material. CUF-NKNY-B14-4 (83 tube pieces; Fig. 16I).

Habitat. Boring into submerged wood and other plant material (Poutiers 1998a).

Taxonomic remarks and comparisons. These calcareous burrows have a septum dividing its cavity into half, indicating that these burrows belong to the shipworm family Teredinidae (e.g. Chan and Lau 2021). Due to the absence of shells, it is impossible to assign the inhabitants to species or even genus level based on these trace fossils.

Order Adapedonta Cossmann & Peyrot, 1909

Superfamily Solenoidea Lamarck, 1809

Family Pharidae H. Adams & A. Adams, 1856

Cultellus Schumacher, 1817

***Cultellus maximus* (Gmelin, 1791)**

Figs 13G, 17A

Solen maximus Gmelin, 1791: 3227. Type locality: Nicobar.

Solen lacteus Spengler, 1794: 94, 95. Type locality: Nicobar.

Cultellus lacteus—Tantanasiriwong 1979: 11. Morris and Purchon 1981: 325. Nateewathana et al. 1981: 66. Aungtonya et al. 1999: 379. Robba et al. 2002: 92, pl. 12, fig. 10. Robba et al. 2007: 86 (appendix). Nabhitabhata 2009: 423.

Cultellus maximus—Subba Rao and Dey 2000: 251. Swennen et al. 2001: 47, 83, text-fig. 121, fig. 121. Huber 2010: 271, with in-text fig. Negri et al. 2014: table 3. Huber 2015: C12478. BEDO 2017a: 48, with in-text fig. Tudu et al. 2018: table 1. Wells et al. 2021: 18.

Referred material. CUF-NKNY-B23 (6L+3R shells; Figs 13G, 17A).

Habitat. Deeply (down to 40 cm) buried in soft mud at seaward fringes of mangrove forests or intertidal zones down to 3 m deep (Huber 2015).

Distribution. Indian Ocean; Indo-West Pacific, from Taiwan to Borneo Island (Huber 2015). Records of fossils from the Holocene in Thailand (Robba et al. 2002).

Record in Thailand. Gulf of Thailand and Andaman Sea (Wells et al. 2021).

Taxonomic remarks and comparisons. Due to the large size and oblong-elliptical shape of shells (Robba et al. 2002; Huber 2010), we identified these specimens as belonging to *Cultellus maximus*.

Siliqua Megerle von Mühlfeld, 1811

Siliqua minima (Gmelin, 1791)

Figs 12I, 17B

Solen minimus Gmelin, 1791: 3227. Type locality: Tranquebar.

Siliqua minima—Lynge 1909: 278. Swennen et al. 2001: 47, 84, text-fig. 126, fig. 126. Robba et al. 2002: 93, pl. 12, fig. 11a, b. Robba et al. 2003: tables 3–5. Robba et al. 2007: 86 (appendix). Nabhitabhata 2009: 424. Negri et al. 2014: table 3. Yang et al. 2017: 218, fig. 822. Wells et al. 2021: 19.

Siliqua cf. *minima*—Hylleberg and Kilburn 2003: 195, 196, pl. 8, fig. 10. Thach 2005: 281.

Siliqua (*Neosiliqua*) *minima*—Huber 2010: 273, with in-text fig. Huber 2015: C12603.

Referred material. CUF-NKNY-B16 (6L+6R shells; Figs 12I, 17B).

Habitat. Sand and mud bottoms from intertidal zones down to 30 m deep (Huber 2015).

Distribution. Indian Ocean; Indo-West Pacific, from Japan to the Philippines (Robba et al. 2002; Huber 2015). Records of fossils from the Holocene in Thailand (Robba et al. 2002).

Record in Thailand. Gulf of Thailand (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Robba et al. (2002) and Huber (2010), specifically in having (1) an ovate-rectangular and inequilateral shell with a widely rounded and deep pallial sinus, the lower border of which coincides with the pallial line, and (2) a thin and slightly oblique inner rib.

Class Scaphopoda Brönn, 1862

Order Dentaliida Starobogatov, 1974

Family Dentaliidae Children, 1834

Dentalium variabile Deshayes, 1826

Figs 12H, 17C

Dentalium variabile Deshayes, 1826: 352, 353, pl. 16, fig. 30. Type locality: possibly India.

Scarabino 1995: 200, 201, fig. 16d, e. Robba et al. 2003: tables 1, 3–5. Robba et al. 2004: 13, pl. 1, fig. 1. Steiner and Kabat 2004: 660. Chaiwathee et al. 2007: table 1. Robba et al. 2007: 89 (appendix). Sahlmann and Poppe 2010: 406, pl. 1198, fig. 8. Negri et al. 2014: tables 3, 4. Wells et al. 2021: 164.

Dentalium (*Lentigodentalium*) *variabile*—Dey and Ramakrishna 2007: 162, 257.

Referred material. CUF-NKNY-O01 (48 shells; Figs 12H, 17C).

Habitat. Mud bottoms from a depth of 10 down to 75 m (Robba et al. 2004).

Distribution. Indo-West Pacific, from Japan to Reunion Island and New Caledonia. Records of fossils from the Quaternary in Thailand (Robba et al. 2004).

Record in Thailand. Gulf of Thailand (Wells et al. 2021).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Scarabino (1995) and Robba et al. (2004), specifically in having (1) a polygonal cross-section of apex and (2) a circular aperture. See also comprehensive taxonomic remarks in Scarabino (1995).

Phylum Arthropoda

Subphylum Crustacea Brünnich, 1772

Class Malacostraca Latreille, 1802

Order Decapoda Latreille, 1802

Family Thalassinidae Latreille, 1831

***Thalassina* Latreille, 1806**

***Thalassina* sp.**

Figs 13I, 17D

Referred material. CUF-NKNY-O09 (130 pieces; Figs 13I, 17D).

Habitat. Littoral and sublittoral zones, mangrove swamps and forests, and edges of estuaries (Ngoc-Ho and de Saint Laurent 2009).

Distribution. Indo-West Pacific, from Japan to Australia (Hyžný and de Angeli 2022). Records of fossils from the Miocene to Holocene in Indo-West Pacific, and from the Oligocene in Italy (Hyžný and de Angeli 2022).

Taxonomic remarks and comparisons. These specimens are assigned to *Thalassina* based on descriptions in Ngoc-Ho and de Saint Laurent (2009), Sakai and Türkay (2012), and Hyžný and de Angeli (2022), specifically in having a subchelate pereiopod 1 with its propodus having dorsomesially tuberculated or spinuous carinae. There is a total of 13 species within this genus (Hyžný and de Angeli 2022), in which three species, *T. anomala* (Herbst, 1804), *T. gracilis* Dana, 1852, and *T. squamifera* De Man, 1915, are recorded from Thailand (Ngoc-Ho and de Saint Laurent 2009; Sakai and Türkay 2012) (but see the comment on the record of *T. squamifera* in Thailand in Moh et al. (2013)). However, the pereiopod 1 observed in our specimens is shorter and has a more quadrate shape. The dactylus that has a hooked tip is nearly as long as the fixed finger, suggesting a yet unidentified species.

Class Thecostraca Gruvel, 1905

Subclass Cirripedia Burmeister, 1834

Order Balanomorpha Pilsbry, 1916

Family Balanidae Leach, 1817

Subfamily Amphibalaninae Pitombo, 2004

***Fistulobalanus* Zullo, 1984**

***Fistulobalanus kondakovi* (Tarasov & Zevina, 1957)**

Figs 12K, 17E

Balanus amphitrite var. *kondakovi* Tarasov & Zevina, 1957: 179, 191, fig. 76a–d. Type locality: Japan.

Balanus kondakovi—Henry and McLaughlin 1975: 114–123, text-figs 21, 22b, c, f; pl. 2, figs a–m. Yamaguchi 1977a: pl. 19, figs 6, 7; pl. 20, figs 4, 8, 12; pl. 21, fig. 4; pl. 22, figs 13–18. Yamaguchi 1977b: 176, 178, text-fig. 18. Kim and Kim 1980: 176, pl. 6, figs 1–8. Yamaguchi 1980: tables 1, 2. Jones et al. 2000: 279. Puspasari et al. 2000: tables 1, 2. Kim 2011: 119–121, fig. 64.

Balanus amphitrite kondakovi—Rosell 1973: 88, fig. 8a–j. Newman and Ross 1976: 63.

Fistulobalanus kondakovi—Zullo 1984: 1330. Pitombo 2004: 275. Prabowo and Yamaguchi 2005: table 3, fig. 3f. Chan et al. 2009: 248–251, figs 214–217. Jones and Hosie 2016: 287, 288. Kepel' 2018: 50, 51, fig. 1. Kim 2020: 75–78, figs 48, 49, table 2. Karasawa and Kobayashi 2022: 71, pl. 5, fig. 10a, b.

Referred material. CUF-NKNY-O06-1 (217 individuals; Figs 12K, 17E).

Habitat. Attached to hard substrates, mainly shells of oysters and gastropod molluscs and stalks of seaweed, as well as wooden piles and supports, bamboo, various objects installed under water on muddy beds for oyster culture, and buoys, in freshened closed inner parts of bays and estuarine coastal areas, from intertidal to subtidal zones (Henry and McLaughlin 1975; Chan et al. 2009; Kepel' 2018).

Distribution. Indian Ocean; Indo-West Pacific, from Japan to New Zealand (Henry and McLaughlin 1975; Puspasari et al. 2000; Jones and Hosie 2016). Records of fossils from the Pleistocene and Holocene in Japan (Yamaguchi 1977b; Karasawa and Kobayashi 2022).

Record in Thailand. Gulf of Thailand (Henry and McLaughlin 1975; Puspasari et al. 2000).

Taxonomic remarks and comparisons. The subfamily Amphibalaninae was erected to incorporate most of the species under the *Balanus amphitrite* complex/group (Pitombo 2004). Currently, there are a total of six species from this subfamily recorded from Thailand:

Amphibalanus amphitrite (Darwin, 1854), *A. reticulatus* (Utinomi, 1967), *A. thailandicus* (Puspasari, Yamaguchi & Angsupanich, 2001), *A. variegatus* (Darwin, 1854), *Fistulobalanus kondakovi* (Tarasov & Zevina, 1957), and *F. patelliformis* (Bruguière, 1789) (Henry and McLaughlin 1975; Puspasari et al. 2000, 2001; Pochai et al. 2017). The presence of two or more rows of parietal tubes on shell walls assigns these specimens to the genus *Fistulobalanus* (Zullo 1984; Chan et al. 2009). We attribute these specimens to *F. kondakovi* because of the smooth outer surface of shells without any longitudinal ribs (Henry and McLaughlin 1975; Chan et al. 2009; Kim 2011; Kepel' 2018).

Subfamily Megabalaninae Leach, 1817

Megabalanus Hoek, 1913

Megabalanus cf. *tintinnabulum* (Linnaeus, 1758)

Figs 13F, 17F

Lepas tintinnabulum Linnaeus, 1758: 668. Type locality: Amboina, Indonesia [lectotype designation by Henry and McLaughlin (1986)].

Balanus (Megabalanus) tintinnabulum tintinnabulum—Pilsbry 1916: 55–57, pl. 10, fig. 1, 1e.

Hiro 1939: 258, fig. 7a–b.

Balanus tintinnabulum var. *tintinnabulum*—Oliveira 1941: 11–14, text-fig. 1, pl. 2, figs. 1, 2; pl. 4, fig. 1; pl. 5, fig. 3; pl. 8, fig. 6.

Balanus tintinnabulum—Holthuis and Heerebout 1972: 24–31, pl. 1, fig. a–e.

Megabalanus tintinnabulum—Newman and Ross 1976: 68. Henry and McLaughlin 1986: 17–21, figs 1e, 2a, g, h, 3a–c, 5a–l. Jones et al. 2000: 282. Pitombo 2004: 275. Lozano-Cortés and Londoño-Cruz 2013: 466, 467. Jones and Hosie 2016: 289, 290. Pochai et al. 2017: 28, 29, fig. 11

Referred material. CUF-NKNY-O06-2 (2 individuals; Figs 13F, 17F).

Habitat. Attached on low exposed rocky shores in littoral areas, ship bottoms, and floating pontoons (Chan et al. 2009; Jones and Hosie 2016)

Distribution. Cosmopolitan (Jones and Hosie 2016).

Record in Thailand. Andaman Sea (Pochai et al. 2017). This is the first record of this species from Gulf of Thailand.

Taxonomic remarks and comparisons. These specimens are tentatively identified as belonging to *Megabalanus tintinnabulum* based on the descriptions and figures in Henry and McLaughlin (1986), Chan et al. (2009), and Pochai et al. (2017), specifically in having large cylindrical shells with smooth parietal surfaces and without spines.

Phylum Echinodermata

Class Echinoidea Schumacher, 1817

Order Camarodontia Jackson, 1912

Family Temnopleuridae Agassiz, 1872

***Temnotrema* Agassiz, 1864**

***Temnotrema siamense* (Mortensen, 1904)**

Figs 12T, 17G

Pleurechinus siamensis Mortensen, 1904: 79–82, pl. 1, figs 2, 7, 11, 20; pl. 2, figs 2, 9, 14, 15, 22; pl. 6, figs 16, 36; pl. 7, figs 14, 44, 53. Type locality: Koh Mesan, 3–15 fathoms; Koh Chuen, 15–38 fathms.; Koh Kram, 20–30 fathms.; Koh Kahdat, 10 faths.

Temnotrema siamense—Clark 1925: 92. Mortensen 1940: 45. Mortensen and Gislén 1940: 103. Mortensen 1943: 259–262, fig. 139. Clark and Rowe 1971: 155. Price 1981: 8. Liao 1998: table 1. Lane et al. 2000: 485. Samyn 2003: table 2. Marsh and Morrison 2004: tables 1, 2, 5. Gopalakrishnan et al. 2012: 32. Filander and Griffiths 2014: 53. Schultz 2015: 258, 263, 264, fig. 4.169b, c. Arachchige et al. 2017: table 3. Putchakarn 2018: 57, table 1.

Arachchige et al. 2019: table 1. Köhler 2020: 9, 72, 73, pl. 14, figs 11, 14. Mucharin and Sumitrakij 2022: 6, pl. 4, fig. 1.

Temnotrema siamensis—Putchakarn and Sonchaeng 2004: table 1.

Referred material. CUF-NKNY-O02 (23 shells; Figs 12T, 17G).

Habitat. Coarse and high energy subtidal sand from a depth of 5 down to 350 m (Price 1981; Schultz 2015).

Distribution. Persian Gulf and Indian Ocean; Indo-West Pacific, from South China Sea to Australia (Mortensen 1904; Price 1981; Schultz 2015; Arachchige et al. 2019).

Record in Thailand. Gulf of Thailand and Andaman Sea (Putchakarn and Sonchaeng 2004).

Taxonomic remarks and comparisons. This species is recognized based on the descriptions and figures in Mortensen (1904), Clark and Rowe (1971) and Schultz (2015), specifically in (1) having deep horizontal grooves in ambulacral and interambulacral plates and in (2) lacking a horizontal series of tubercles.

Phylum Chordata

Subphylum Vertebrata Lamarck, 1801

Class Chondrichthyes Huxley, 1880

Subcohort Selachimorpha Nelson, 1984

Superorder Galeomorphii Compagno, 1973

Order Carcharhiniformes Compagno, 1977

Family Carcharhinidae Jordan & Evermann, 1896

Carcharhinus Blainville, 1816

Carcharhinus cf. amblyrhynchoides (Whitley, 1934)

Fig. 18A–E

Gillisqualus amblyrhynchoides Whitley, 1934 189–191, text-fig. 4. Type locality: Cape Bowling Green, Queensland.

Carcharhinus amblyrhynchoides—Compagno 1984 458, 459, with in-text figs. Krajangdara et al. 2022: 49, with in-text figs.

Referred material. CUF-NKNY-S3-2 (Fig. 18A–E) (1 upper tooth).

Description. The crown of CUF-NKNY-S3-2 is triangular and erect with fine serrations, and displays well-developed heels mesially and distally. Its lingual face is distinctly more convex than the labial one. Its base presents a damaged lingual face.

Habitat. Tropical, inshore and offshore, coastal-pelagic species, found over the continental and insular shelves (Compagno 1984).

Distribution. Gulf of Aden and Indian Ocean; Indo-West Pacific, from southern China to Australia (Compagno 1984).

Record in Thailand. Gulf of Thailand and Andaman Sea (Compagno 1984; Krajangdara et al. 2022).

Taxonomic remarks and comparisons. The specimen CUF-NKNY-S3-2 represents an anterior upper tooth. Several species of *Carcharhinus* have similar upper teeth in terms of morphology. Additional teeth and larger assemblages are needed for a more precise identification.

Nevertheless, the tooth most resembles the upper teeth of *Carcharhinus amblyrhynchoides* (Garrick 1982: fig. 20). Male individuals of *C. brachyurus* have somewhat similar upper teeth (Garrick 1982: fig. 51), but the mesial cutting edge of their crown is often more convex.

Carcharhinus limbatus also shows the same characters, but the serration on the cusps at the base of the crown is much finer (Bass et al. 1973: pl. 5; Garrick 1982: fig. 18). The pattern of serration also resembles that of the lower teeth of *C. sorrah*, although in the latter, the base of the root is

more concave and the heels of the crown are better developed, the teeth being longer mesio-distally than high baso-apically (Voigt and Weber 2011). Regarding the fossil record in Southeast Asia, similar teeth were reported from the Late Miocene deposits of Brunei in Borneo (Kocsis et al. 2019).

***Carcharhinus* cf. *amblyrhynchos* (Bleeker, 1856)**

Fig. 18F–J

Carcharias (Prionodon) amblyrhynchos Bleeker, 1856 467, 468. Type locality: Java Sea near Solombo Island.

Carcharhinus amblyrhynchos—Compagno 1984 459–461, with in-text figs. Krajangdara et al. 2022: 49, with in-text figs.

Referred material. CUF-NKNY-S3-10 (Fig. 18F–J) (1 upper tooth).

Description. The crown of CUF-NKNY-S3-10 displays a rather narrow main cusp with well-developed heels. The main cusp is regularly serrated, with an almost straight mesial edge and a distal one that is slightly concave. It is thus inclined distally. The labial side of the main cusp is nearly flat, whereas the lingual one is convex. Both the mesial and distal heels of the crown are poorly preserved, and their serration pattern cannot thus be observed. The base of the root is concave and the nutritive groove on the bulged lingual face is poorly-developed or heavily worn. The labial face of the root is nearly flat.

Habitat. Continental and insular shelves and adjacent oceanic waters (Compagno 1984).

Distribution. Indo-western to Central Pacific (Compagno 1984).

Record in Thailand. Gulf of Thailand and Andaman Sea (Compagno 1984; Krajangdara et al. 2022).

Taxonomic remarks and comparisons. The general shape of the crown, the regular serration of the main cusp, and the arched root are reminiscent of *Carcharhinus amblyrhynchos* (Bass et al. 1973; Garrick 1982; Kocsis et al. 2019; Malyshkina et al. 2023), but the poorly preserved heels of the teeth make a definite identification difficult to ascertain.

***Carcharhinus* cf. *leucas* (Valenciennes, 1839)**

Fig. 19

Carcharias (Prionodon) leucas Valenciennes in Müller & Henle, 1839 42, 43. Type locality: Antilles.

Carcharhinus leucas—Compagno 1984 478–481, with in-text figs. Krajangdara et al. 2022: 52, with in-text figs.

Referred material. CUF-NKNY-Q04 (Fig. 19A–I), CUF-NKNY-SA-1 (Fig. 19J–N), CUF-NKNY-SB-3 and CUF-NKNY-SC-5 (4 upper teeth).

Description. Crowns show well-developed serrations, becoming larger and more complex in the basal part of the distal side of the crown. In labial or lingual view, the mesial side is almost straight, whereas the distal one is concave in its lower third. The outline of the base of the crown

is more concave on the lingual face than on the labial one. The root is deep, slightly concave at its base, and does not display a nutritive groove in lingual view.

Habitat. Close to inshore in reef-associated marine habitats, mostly in water less than 30 m deep (Compagno 1984; Krajangdara et al. 2022).

Distribution. Cosmopolitan in tropical and subtropical seas, but also reported from estuaries and rivers, as well as tolerate freshwater conditions (Compagno 1984).

Record in Thailand. Gulf of Thailand and Andaman Sea (Compagno 1984; Krajangdara et al. 2022).

Taxonomic remarks and comparisons. Teeth of *Carcharhinus amboiensis* are very similar to those of *C. leucas* so that it is very difficult to differentiate them (Kocsis et al. 2019). As a result, ten incomplete teeth may belong either *C. leucas* or *C. amboiensis*: CUF-NKNY-S2-1, CUF-NKNY-S3-9, CUF-NKNY-S5-1, CUF-NKNY-S5-5, CUF-NKNY-SA-6, CUF-NKNY-SB-2, CUF-NKNY-SC-10, CUF-NKNY-SD-7, CUF-NKNY-SE-3 and CUF-NKNY-SE-9.

***Carcharhinus cf. sorrah* (Valenciennes, 1839)**

Fig. 20

Carcharias (Prionodon) sorrah Valenciennes in Müller & Henle, 1839 45, 46, pl. 16. Type locality: Java.

Carcharhinus sorrah—Compagno 1984 500, 501, with in-text figs. Krajangdara et al. 2022: 57, with in-text figs.

Referred material. CUF-NKNY-3.1 (Fig. 20A–G) and CUF-NKNY-3.2 (Fig. 20H–O) (2 upper teeth).

Description. The crown is compressed labio-lingually. The labial face is slightly flatter than the lingual one. The cusp is inclined distally and presents a notch on its distal part. The serrations do not reach the apex of the main cusp and are enlarged, becoming more complex basally, especially on the distal heel. The mesial edge of the crown is almost straight in labial or lingual view. The base of the root is slightly concave in lingual or labial view. There is a well-developed groove with a nutritive foramen at the base of the root in lingual view.

Habitat. Coastal, shallow-water zones of the continental and insular shelves, primarily around coral reefs at intertidal zones down to 73 m deep (Compagno 1984; Krajangdara et al. 2022).

Distribution. Red Sea and Indian Ocean; Indo-West Pacific, from China to Australia (Compagno 1984).

Record in Thailand. Gulf of Thailand and Andaman Sea (Compagno 1984; Krajangdara et al. 2022).

Taxonomic remarks and comparisons. Some upper teeth of *Carcharhinus sorrah* appear to display a mesial cutting edge slightly more convex than on our specimens, but this character is known to depend on ontogenetic stages and the position of the teeth in the jaw (e.g., Bass et al. 1973: pl. 13; Garrick 1982: fig. 76). Juveniles of *C. dussumieri* may have upper teeth with similar characteristics, but the distal cutting edge of the main cusp is not serrated. This latter character appears in larger specimens, but the mesial cutting edge of the main cusp becomes slightly concave. In addition, the distal heels and the distal root lobe of the fossil teeth are more elongated than the upper teeth of *C. dussumieri* (White 2012).

***Carcharhinus* spp.**

Figs 21, 22

Referred material. CUF-NKNY-16.1 (Fig. 21A–H), CUF-NKNY-S2 (Fig. 21I–M), CUF-NKNY-S3-4, CUF-NKNY-S3-8 and CUF-NKNY-S4-7 (5 lower teeth) and CUF-NKNY-S1 (Fig. 22A–H) and CUF-NKNY-S3-7 (two upper teeth).

Description. The specimens CUF-NKNY-16.1, CUF-NKNY-S3-4, CUF-NKNY-S3-8, and CUF-NKNY-S4-7 display a narrow main cusp that is inclined lingually. Their labial face is flatter than the lingual one. The serrations are well-developed in the upper two-third of the cusp. The serrations appear to be very fine on the mesial and distal heels in CUF-NKNY-16.1, CUF-NKNY-S3-4, and CUF-NKNY-S3-8, but the serrations are damaged in CUF-NKNY-S4-7. The root is well preserved only in CUF-NKNY-S3-4 and CUF-NKNY-S3-8, both of which display a slightly arched base, a moderately bulged lingual face with a nutritive foramen in its center in CUF-NKNY-S3-4 and a faint nutritive groove in CUF-NKNY-S3-4. The specimen CUF-NKNY-S2 is asymmetric, with a mesial heel almost twice as elongated as the distal one and displays a sigmoid main cusp in labial or lingual view as well as in mesial or distal view, separated from the distal heel by a notch. The heels are serrated, but the main cusp is smooth. This tooth appears quite bulbous in mesial or distal view, being less compressed labio-lingually than the specimens described above. There is a well-developed nutritive groove on the root in lingual view, forming a basal notch in labial view.

A heavily worn tooth (CUF-NKNY-S3-7) is better attributed to an upper tooth. The lingual and distal faces of the crown are strongly damaged and a part of the distal end of the root is missing. The serration is only preserved in the middle part of the mesial cutting edge and appears to be quite regular. The root has a basal face slightly arched with a well-developed nutritive groove on its lingual face, forming a notch at the base of the root in labial view.

Another upper tooth, CUF-NKNY-S1, displays a crown with a quite coarse, irregular serration pattern nearly reaching the apex of the main cusp. The latter is bent lingually and slightly inclined distally. Its root shows a well-developed nutritive groove in lingual view, forming a notch at the base of the root in labial view. Apart from this notch, the base is almost straight.

Taxonomic remarks and comparisons. Lower teeth of *C. amblyrhynchos* (*C. wheeleri* in Garrick (1982): fig 51) display features similar to the specimen CUF-NKNY-16.1. They are characterized by a narrow main cusp that displays serrations ending at mid-height. Some of the lower teeth of *C. longimanus* also have these morphological patterns, though their crown is often more robust and wider at their base (Bass et al. 1973: pl. 17; Garrick 1982: fig. 70). Nevertheless, teeth very similar to the specimens CUF-NKNY-16.1, CUF-NKNY-S3-4, CUF-NKNY-S3-8, and CUF-NKNY-S4 from the Pliocene of Italy were attributed to *C. longimanus* (Marsili 2007).

The sigmoid crown of the specimen CUF-NKNY-S2 is reminiscent of the posterior lower teeth of *C. borneensis* and *C. tjutjot* (Voigt and Weber 2011). The dentition attributed to *C. dussumieri* by Voigt and Weber (2011) belongs in fact to *C. tjutjot* (White 2012). A sigmoid cusp, when present, is less developed in *C. sealei* and *C. sorrah* (Voigt and Weber 2011), and the distal heel is coarsely serrated in *C. coatesi* (White 2012). The morphology of the crown recalls that of *Rhizoprionodon*, but in the latter genus, the distal heel is often more convex than in the

specimen CUF-NKNY-S2 (Cappetta 2012: fig. 283; Carrillo-Briceño et al. 2015). This set of lower teeth is therefore attributed to the genus *Carcharhinus* but represents more than a single species, whereas the specimen CUF-NKNY-S3-7 is poorly preserved. Moreover, the specimen CUF-NKNY-S1 preserves only its posterior part and does not allow us to reach the species-level identification.

There are 27 species of the genus *Carcharhinus* reported from the Southeast Asian region (Froese and Pauly 2024), and at least 20 taxa were recorded in Thai waters (Krajangdara et al. 2022) including all the ones mentioned above.

Glyphis Agassiz, 1843

Glyphis sp. Figs 23–28

Referred material. CUF-NKNY-8.2 (Fig. 23A–H), CUF-NKNY-Q03 (Fig. 24A–H), CUF-NKNY-S5-2, CUF-NKNY-S5-4, CUF-NKNY-S5-6, CUF-NKNY-S5-7, CUF-NKNY-SA-2 to 5, CUF-NKNY-SA-7, CUF-NKNY-SA-8, CUF-NKNY-SB-1, CUF-NKNY-SB-4 to 9, CUF-NKNY-SC-1, CUF-NKNY-SC-3 (Fig. 25A–E), CUF-NKNY-SC-4, CUF-NKNY-SC-6, CUF-NKNY-SC-8, CUF-NKNY-SC-9, CUF-NKNY-SD1 to 3 (Fig. 25F–J), CUF-NKNY-SD6, CUF-NKNY-SD-8 to 10, CUF-NKNY-SE1 to 2, CUF-NKNY-SE-4 to 7, CUF-NKNY-SE-10 to 15, CUF-NKNY-SE-17 to 18 (46 upper teeth), and CUF-NKNY-14 (Fig. 26A–H), CUF-NKNY-S1-1 to 4 (Fig. 26I–M), CUF-NKNY-S3-1 (Fig. 27A–E), CUF-NKNY-S3-5 to 6 (Fig. 27F–J), CUF-NKNY-S4-1 to 2 (Fig. 28A–E), CUF-NKNY-S4-4 to 6 (Fig. 28F–J) (13 lower teeth).

Description. Many teeth are poorly preserved. The crown of the upper teeth is triangular in outline, with a concavity at the base on both the mesial and distal sides in labial and lingual view, although the distal one is better marked, especially in anterior teeth (CUF-NKNY-S5-7). The serrations are quite fine and homogeneous in size along the cutting edges. The crown is strongly compressed labio-lingually and may be slightly curved lingually. When being broken, the crown displays a pulp cavity compressed labio-lingually, indicating an orthodont tooth histology (CUF-NKNY-SE-11). The base of the root is slightly concave in labial or lingual view. A well-developed nutritive groove and/or a nutritive foramen may be present on the lower half of the lingual side of the root.

The lower teeth may reach 40 mm in height. They are narrower and less compressed labio-lingually than the upper ones. The enameloid is smooth and the labial face of the crown is flatter than the lingual one. The main cusp is inclined lingually, sometimes with a sigmoid outline (CUF-NKNY-S4-2, CUF-NKNY-S4-5). Three different crown morphologies are noted.

(1) On the specimens CUF-NKNY-S1-1, CUF-NKNY-S1-1-2, and CUF-NKNY-S1-1-4, the crown is devoid of heels and the tip of the main cusp is spearhead-shaped, the latter being regularly serrated, whereas the rest of the crown is smooth. (2) The specimens CUF-NKNY-S4-1 and CUF-NKNY-S4-4 have a narrow and upright triangular main cusp devoid of serration, showing a tiny accessory cusplet on each side of the main cusp in CUF-NKNY-S4-4 or only lingually in CUF-NKNY-S4-1. (3) On the specimen CUF-NKNY-S3-6, the main cusp is also upright, narrow, triangular in shape, and serrated on their mesial and distal edges. Well-

developed heels are present on each side of the main cusp, separated from the latter by a notch. The heel is sometimes less marked on the mesial side (CUF-NKNY-S3-1). The heels display less developed serration than on the main cusp. Many teeth belonging to this morphotype display a broken apex of the main cusp (CUF-NKNY-14).

The root displays a bulged lingual face. It is more massive and more arched in spearhead-shaped teeth than in heeled ones. There is a nutritive groove in the lingual side of the root with a nutritive foramen in its middle part that can be observed in most of the teeth. The specimen CUF-NKNY-S1-1 displays a double groove.

Taxonomic remarks and comparisons. Upper anterior teeth (CUF-NKNY-Q03) are more erect and narrower than the lateral ones (CUF-NKNY-8.2). Lower teeth with a spearhead-shaped apex correspond to the larger teeth in our sample, in agreement with the fact that this morphology is known only in anterior teeth of adult specimens of *Glyphis glyphis* and *G. garricki*. However, lateral teeth of these species display serrated heels and teeth of juveniles can display a pair of tiny lateral cusplets (Compagno et al. 2008; White et al. 2015). As a result, the three lower tooth morphotypes described above could be encountered into a single species during its ontogeny. Otherwise, it could indicate the presence of more than one species, possibly three: *G. glyphis*, *G. garricki*, and *G. gangeticus*, as the latter species often possesses anterior lower teeth with tiny lateral cusplets but lacks the spearhead-shaped apex (Roberts 2006; Cappetta 2012). Consequently, the teeth described above are left in open nomenclature as *Glypis* sp. This taxon appears to be most common in the studied fauna, although the discrepancies between the number of upper and lower teeth recovered suggest some degree of collecting bias.

The genus is specific to the Indo-West Pacific tropical region and often referred as “river shark” due to its habitat in or nearby rivers and estuaries. They are quite rare, hence difficult to study. Five living species are known (Froese and Pauly 2024), but DNA analyses suggest that some of these are conspecific (Li et al. 2015). On the other hand, teeth of *Glypis* are often recorded from Southeast Asia (Shimada et al. 2016; Kocsis et al. 2019 and references therein).

***Scoliodon* Müller & Henle, 1837**

***Scoliodon cf. laticaudus* Müller & Henle, 1838**

Fig. 29

Carcharias (Scoliodon) laticaudus Müller & Henle, 1838 28, 29, pl. 8. Type locality: India.
Scoliodon laticaudus—Compagno 1984 534, 535, with in-text figs. Krajangdara et al. 2022: 62, with in-text figs.

Referred material. CUF-NKNY-S3–S5 (Fig. 29A–O) (3 teeth).

Description. The specimen CUF-NKNY-S3 is a small and elongated tooth with a low crown and a worn off root. The mesial cutting edge is concave, smooth, and continues along the mesial heel. The distal edge of the crown is convex and it is separated by a notch from a distinct distal heel.

The specimen CUF-NKNY-S4 is a well-preserved tooth, with a distally strongly curved crown. The mesial cutting edge is sinusoid, starting convex, then becoming concave towards the apex. The apex of the crown overhangs distally the basal part of the tooth. The distal cutting edge

is also sinusoid, though less so than the mesial one, and it is separated by a distinct notch from a well-developed distal heel. The enameloid of the crown extends more basally on the labial side of the tooth. The root is thick and the lobes run rather horizontally. It bears a well-developed and deep nutritive groove in the centre.

The cusp of the specimen CUF-NKNY-S5 is strongly sigmoid in labial and lingual view, and similar to that of CUF-NKNY-S4, but its cusp is narrower and less wide at the base, with more apparent mesial heels. The cutting edge is smooth and does not reach the apex of the cusp on its mesial side. The heels of the crown display faint serrations. The root is asymmetric and projected lingually and distally. There is a well-defined groove distally on the lingual side of the basal face, forming a deep notch in apical view.

Habitat. Tropical zones of continental and insular shelves close to inshore, frequently in rocky areas (Compagno 1984).

Distribution. Indian Ocean; Indo-West Pacific, from Japan to Indonesia (Compagno 1984).

Record in Thailand. Andaman Sea (Krajangdara et al. 2022).

Taxonomic remarks and comparisons. The three teeth best fit the dentition of the modern species *Scoliodon laticaudus* (spadenose shark). The specimen CUF-NKNY-S5 has a slender sigmoidal cusp and its mesial cutting edge does not reach the apex and it displays asymmetric root, corresponding perfectly to a parasymphyseal tooth of a male specimen (Herman et al. 1991). The specimen CUF-NKNY-S4 likely represents an upper anterior-anterolateral tooth, whereas the specimen CUF-NKNY-S3 is probably a lower anterolateral tooth (see Springer 1964: fig. 3a). White et al. (2010) recognized another species, *S. macrorhynchos*, in the Western Pacific, meanwhile a molecular study also indicates a possible third *Scoliodon* species from the region (Lim et al. 2022), hence these fossil teeth are described in open nomenclature.

It may be mentioned that specimen CUF-NKNY-S3 also resembles that of the lower anterolateral teeth of other common sharks in Southeast Asia (Krajangdara et al. 2022) such as *Loxodon macrorhinus* (sliteye shark) and two *Rhizoprionodon* species, *R. acutus* (milk shark) and *R. oligolepis* (grey sharpnose shark). For the teeth of *Loxodon*, the apex of the crown is higher and somewhat aligned in a more distal position than that of the specimen CUF-NKNY-S3 (Springer 1964: fig. 4; Bass et al. 1975: pl. 8). The teeth of *R. oligolepis* seem to have a less concave mesial cutting edge (Springer 1964: fig. 13), whereas for *R. acutus*, the distal heel appears less elongated and vertically higher, but a clear distinction is difficult here (Springer 1964: fig. 6; Bass et al. 1975: pl. 9).

Carcharhinidae indet.

Referred material. CUF-NKNY-S3-3, CUF-NKNY-S4-3, CUF-NKNY-S5-3, CUF-NKNY-SC-2, CUF-NKNY-SC-7, CUF-NKNY-SD-4, CUF-NKNY-SD-5, CUF-NKNY-SD-11, CUF-NKNY-SE-8, CUF-NKNY-SE-16 (10 teeth).

Description. These teeth are poorly preserved. The crown is generally triangular, more or less narrow, and symmetric with serration incompletely preserved. The root is broken away or does not display the details of its vascularization pattern.

Taxonomic remarks and comparisons. The presence of triangular serrated crowns suggests that most of the teeth belong to upper ones of *Carcharhinus* and/or *Glyptodon*, but their poor preservation does not allow a more precise identification. The specimen CUF-NKNY-SC-7 may

represent a lower tooth of *C. leucas/amboinensis*, but without the root and the base of the crown preserved, this is impossible to ascertain.

Superorder Batomorphii Capetta, 1980
Order Myliobatiformes Compagno, 1973
Family Dasyatidae Jordan & Gilbert, 1879

***Pastinachus* Rüppell, 1829**

***Pastinachus* sp.**

Fig. 30

Referred material. CUF-NKNY-7.1 (Fig. 30A–F), CUF-NKNY-7.2 (Fig. 30G–L) and CUF-NKNY-8.1 (Fig. 30M–R) (3 teeth).

Description. The crown is hexagonal to diamond-shaped in apical view, longer mesio-distally than labio-lingually. The crown surface is rather smooth to heavily pitted. The labial face of the crown displays a salient horizontal bulge. There is a well-developed horizontal groove in the basal part of the crown on the lingual face. The vascularization of the teeth is holaulacorhize. There is a row of small foramina positioned under the crown on the labial face, and between one and four foramina are present in the groove separating the two branches of the root in basal view. **Taxonomic remarks and comparisons.** Heavily pitted crowns probably belong to non-functional teeth (Adnet et al. 2019). Four species of cowtail rays (*Pastinachus ater*, *P. gracilicaudus*, *P. solocirostris*, and *P. stellurostris*) are known in Southeast Asia (Last et al. 2016), all of them having been recorded in Thai waters (Krajangdara et al. 2022). Regarding the nearby fossil record, teeth of *Pastinachus* were reported from India, Taiwan, and Borneo (see discussion in Kocsis et al. 2019).

Class Actinopterygii Klein, 1885
Infraclass Teleostei Müller, 1845
Order Scombriformes Rafinesque, 1810
Family Trichiuridae Rafinesque, 1810

Trichiuridae indet.

Fig. 31

Referred material. CUF-NKNY-2.1 (Fig. 31A–F), CUF-NKNY-2.2 (Fig. 31G–K) and CUF-NKNY-18.2 (Fig. 31L–P) (3 teeth).

Description. Labiolingually flattened teeth seem to thin out at their mesial edge. They are covered with elongated and fine striation. This striation becomes coarser basally and distally on the more elongated specimens (specimens CUF-NKNY-2.1 and CUF-NKNY-2.2). The latter two teeth are strongly curved distally with a concave distal edge, and the more completely preserved specimen CUF-NKNY-2.1 bears an apical barb. The specimen CUF-NKNY-18.2 has a curved mesial margin, whereas the distal one is vertical and straight.

Taxonomic remarks and comparisons. The two strongly curved teeth (specimens CUF-NKNY-2 and CUF-NKNY-2.1) represent fang-like features from the front of the jaw, while the specimen CUF-NKNY-18.2 comes from a rather distal position. This latter tooth resembles

somewhat the lateral teeth of *Sphyranea* (barracuda, e.g., Gottfried et al. 2017), but those are more symmetrical. Fishbase (www.fishbase.org) reports twenty-seven cutlassfishes from the wider region of South and Southeast Asia (Froese and Pauly 2024). Two common genera, *Trichurus* and *Lepuracanthus*, contain some species that bear such barbed fang-like teeth (Nakamura and Parin 1993; 1998).

Order *incertae sedis* in Eupercaria Betancur-R. et al., 2014

Family Sciaenidae Cuvier, 1829

***Johnius* Bloch, 1793**

***Johnius* sp.**

Fig. 32

Referred material. CUF-NKNY-16.2 (Fig. 32A–D) (1 right sagitta otolith).

Description. Thick otolith with a smooth external surface, whereas the internal surface bears a sulcus characteristic of the genus. The ostium is shallow, vertically positioned, higher than long and it widens ventrally. The anterior and horizontal portion of the cauda is short and shallow, then opening posteriorly into a deep caudal funnel with an external circular shape.

Taxonomic remarks and comparisons. Otoliths of *Johnius* are unique and have distinct characteristics that make them easily distinguishable from those of other sciaenids or teleosts (Schwarzans 1993). There are 21 species of *Johnius* in Southeast Asia (Froese and Pauly 2024), and most of them inhabit shallow coastal waters and estuaries but some species are capable of entering rivers. They are also known from the fossil records in Japan (Ohe 1976) and Taiwan (Lin et al. 2022).

Discussion

Based on the survey of mid-Holocene marine faunas from the clay pit of Nakhon Nayok Province in Thailand, many fossils of molluscs together with other invertebrates and vertebrates have been found in the Bangkok clay layer at a depth of about 5–7 m below the surface. The faunal assemblages, especially molluscs (Table 2), depict the paleoecosystem of the area at that time as belonging to the intertidal to sublittoral zones. Some molluscan taxa, i.e., *Cerithidea obtusa*, *Pirenella incisa*, *Telescopium telescopium*, *Ellobium aurisjudae*, *Cassidula nucleus*, and *Geloina bengalensis*, also indicate that some parts of the area corresponded to mangrove forests and intertidal mudflats. The occurrence of mangrove forests has previously been corroborated by evidence of peat and pollens from the same, as well as nearby, sites (Songtham et al. 2007, 2015). In this study, several described mollusc taxa correspond to those reported from the Lower Central Plain of Bangkok and Samut Prakan, and inland of Phetchaburi coast (Somboon 1988; Somboon and Thiramongkol 1992; Robba et al. 2007; Negri 2009, 2012; Surakiatchai et al. 2018), suggesting similar paleoenvironments of these sites.

Most of the marine and mangrove shells were found at a depth of 2.2 m below the uppermost part of a marine clay layer, and the carbon-14 dating analysis indicates the age of this level as being approximately 5,900–5,300 cal yr BP, whereas charcoal material was found at a greater depth of 2.2–4.6 m, spanning approximately an age from 8,800 to 5,700 cal yr BP (Table 1). However, the abundance of tree trunks and carbonized woods as well as the accumulation of peat at greater depth than most shells indicate the predominance of mangrove forests, suggesting

an early stage of marine incursion further inland (Songtham et al. 2007). The ages of the Bangkok Clay deposits determined by this study are also in good agreement with those retrieved by Songtham et al. (2007). This duration corresponds to the depositional stage 2 (7,000–3,000 cal yr BP), a marine transgression time proposed by Tanabe et al. (2003). During this stage, a large mud shoal was formed around Samut Prakan Province, and the Chao Phraya delta prograded toward the head of the paleo-Gulf of Ayutthaya southward (Fig. 33; Tanabe et al. 2003). However, our chronological results suggest that the area of Ongkharak, Nakhon Nayok in central Thailand between 7,900 and 5,300 cal yr BP was possibly an ancient coastal shoreline, which was located further southward compared to the paleogeographic scheme proposed by Tanabe et al. (2003).

The vertebrate fauna is dominated by cartilaginous fishes among which the family Carcharhinidae is most common. The genus *Glypis*, also known as “river shark”, is dentally most abundant in the area due to its habitat in or nearby rivers and estuaries. It is a cryptic shark that is difficult to study, but it has been reported that these sharks also live in coastal, shallow marine region, while the young grow up in a river habitat where the predation pressure is reduced (Li et al. 2015). The dominant presence of this taxon is indicative of the vicinity of freshwater input and/or habitats close to rivers, which fits well with the paleogeographic context of the region (Fig. 33) and the results of other studies from the Bangkok Clay (Robba et al. 2002, 2003, 2004, 2007; Songtham et al. 2007; Negri 2009, 2012). Several *Carcharhinus* species were also found and the most likely species are *C. amblyrhynchoides*, *C. amblyrhinchos*, *C. leucas*, *C. amboinensis*, and *C. sorrah*, but other taxa could not be ruled out based on some similarities in tooth morphology, although the position of the teeth in the jaws cannot be ascertained. Nevertheless, the bull shark (*C. leucas*) appears to be also abundant and is known to live in freshwater environments as well. Moreover, the genus *Scoliodon*, represented in our records, was reported from brackish environments (Riede 2004). The other recovered taxa, the stingray (*Pastinachus*), cutlassfishes (Trichiuridae), and the sciaenid *Johnius* are all compatible with shallow marine coastal environments, as supported by the presence of molluscan faunas, although some of the species are reported from brackish environments. Due to the fact that the fish remains are almost entirely represented by isolated teeth and only one single otolith has been found so far (*Johnius*), the rarity of this type of remains could be the consequence of sampling biases and/or taphonomic processes. Therefore, targeting otoliths in future studies could shed more light on the bony fish fauna of the Bangkok Clay.

Acknowledgments

We thank the following colleagues from the Department of Biology, Faculty of Science, Chulalongkorn University): Wichase Khonsue who led us to the locality and provided information on the sites, Somsak Panha who facilitated access to collection storage and working area, Arthit Polyotha for providing SEM images, and Kittipum Chansri and Buntika A. Butcher for providing access and taking photographs with a Leica M205C stereo light microscope. We also thank Fa-Is Jindewha, Sutan Saehan, Jirapan Sumpaothong, and Suwijk Pata for their assistance in the field, Santi Pailoplee (Department of Geology, Faculty of Science, Chulalongkorn University) for his help in generating the map, Mana Rubbumrung (Department of Mineral Resources) for his/her (?) technical assistance and supports, and Thierry Backeljau (Royal Belgian Institute of Natural Sciences, Brussels) and Bernhard Hausdorf (Leibniz Institute

for the Analysis of Biodiversity Change, Hamburg) for the access to the relevant literature. Finally, we thank Jürgen Kriwet and an anonymous reviewer for their comments on the first version of this manuscript. This research was supported by the Sci-Super VII grant, Faculty of Science, Chulalongkorn University (Grant No. Sci-Super VII_64_001, to KS and CS).

References

- Adams A (1852) Catalogue of the species of *Nassa*, a genus of gasteropodous Mollusca belonging to the family Buccinidae, in the collection of Hugh Cuming, Esq., with the description of some new species. *Proceedings of the Zoological Society of London* 19: 94–114.
- Adams A (1856) Descriptions of twenty-five new species of shells from the collection of Hugh Cuming, Esq. *Proceedings of the Zoological Society of London* 23: 221–226.
- Adnet S, Mouana M, Charrault A-L, Essid EM, Ammar HK, Marzougui W, Merzeraud G, Tabuce R, Vianey-Liaud M, Marivaux L (2019) Teeth, fossil record and evolutionary history of the cowtail stingray *Pastinachus* Rüppell, 1829. *Historical Biology* 31: 1213–1222.
- Albano PG, Steger J, Bakker PAJ, Bogi C, Bošnjak M, Guy-Haim T, Huseyinoglu MF, LaFollette PI, Lubinevsky H, Mulas M, Stockinger M, Azzarone M, Sabelli B (2021) Numerous new records of tropical non-indigenous species in the Eastern Mediterranean highlight the challenges of their recognition and identification. *ZooKeys* 1010: 1–95.
- Alf A, Thach NN (2021) Some remarks on *Lenifusus*, *Brunneifusus* and *Hemifusus* and description of two new species of *Hemifusus* (Buccinoidea, Melongenidae). *Conchylia* 52: 95–108.
- Arachchige GM, Jayakody S, Mooi R, Kroh A (2017) A review of previous studies on the Sri Lankan echinoid fauna, with an updated species list. *Zootaxa* 4231: 151–168.
- Arachchige GM, Jayakody S, Mooi R, Kroh A (2019) An annotated species list of regular echinoids from Sri Lanka with notes on some rarely seen temnopleurids. *Zootaxa* 4571: 35–57.
- Aubry U, Gargiulo R, Russo S (2021) Terebridae: Descriptive and Iconographic Review from 1713 to 2021. L'Informatore Piceno, Ancona, Italy, 515 pp.
- Aungtonya C, Hylleberg J (1998) Check list of sublittoral molluscs, with nine new records for the Andaman Sea. *Phuket Marine Biological Center Special Publication* 18: 317–322.
- Aungtonya C, Thaiklang N, Srisawan D (1999) Recent records of Bivalvia in the reference collection of Phuket Marine Biological Center, Thailand. *Phuket Marine Biological Center Special Publication* 19: 371–383.
- Baharuddin N, Martin MB, Nagappan T, Borkhanuddin MH (2017) Checklist of mangrove snails (Mollusca: Gastropoda) in Setiu Wetlands. In: Abdullah DM, Nasir MZ, Mazlan NM, Jusoh WFAW (Eds) Prosiding Seminar Ekspedisi Saintifik Tanah Bencah Setiu 2016. WWF-Malaysia, pp.
- Baron-Szabo RC, Cairns SD (2016) Part F, Revised, Volume 2, Chapter 10: Systematic descriptions of the Scleractinia family Rhizangiidae. *Treatise Online* 80: 1–10.
- Bass AJ, D'Aubrey JD, Kistnasamy N (1973) Sharks of the East Coast of southern Africa. I. The genus *Carcharhinus* (Carcharhinidae). *Investigational Report – Issue 33*. Oceanographic Research Institute, Durban, South Africa, 168 pp.
- Bass AJ, D'Aubrey JD, Kistnasamy N (1975) Sharks of the East Coast of Southern Africa. III. The Families Carcharhinidae (excluding Mustelus and Carcharhinus) and Sphyrnidae. *Investigational Report Oceanographic Research Institute*, Durban, South Africa, 100 pp.
- BEDO (2017a) Marine Bivalves: Checklist of Molluscan Biodiversity in Thailand. BEDO, Bangkok [in Thai], 296 pp.
- BEDO (2017b) Marine Gastropods: Checklist of Molluscan Biodiversity in Thailand. BEDO, Bangkok [in Thai], 376 pp.

- Beu AG (2006) Marine Mollusca of oxygen isotope stages of the last 2 million years in New Zealand. Part 2. Biostratigraphically useful and new Pliocene to recent bivalves. *Journal of the Royal Society of New Zealand* 36: 151–338.
- Beu AG (2008) Bursidae. In: Poppe GT (Ed) *Philippine Marine Mollusks Volume I*. ConchBooks, Hackenheim, Germany, pp. 612–627.
- Beu AG (2010) Marine Mollusca of isotope stages of the last 2 million years in New Zealand. Part 3. Gastropoda (Vetigastropoda - Littorinimorpha). *Journal of the Royal Society of New Zealand* 40: 59–180.
- Beu AG, Alloway BV, Pillans BJ, Naish TR, Westgate JA (2004) Marine Mollusca of oxygen isotope stages of the last 2 million years in New Zealand. Part 1: Revised generic positions and recognition of warm-water and cool-water migrants. *Journal of the Royal Society of New Zealand* 34: 111–265.
- Bibi S, Terryn Y, Siddiqui PJA, Iqbal P (2021) In the wake of Captain Townsend – Poorly known and cryptic Terebridae (Gastropoda: Conoidea) of Pakistan. *Gloria Maris* 60: 43–59.
- Bieler R (1993) Architectonicidae of the Indo-Pacific (Mollusca, Gastropoda). *Abhandlungen des Naturwissenschaftlichen Vereins in Hamburg* 30: 1–376.
- Bleeker P (1856) *Carcharias (Prionodon) amblyrhynchos*, eene nieuwe haaisoort, gevangen nabij het eiland Solombo. *Natuurkundig Tijdschrift voor Nederlandsch Indië* 10: 467–468.
- Born I von (1778) *Index rerum naturalium Musei Cæsarei Vindobonensis*. Pars I.ma. Testacea. Verzeichniß der natürlichen Seltenheiten des k. k. Naturalien Cabinets zu Wien. Erster Theil. Schalthiere. [1-40], 1-458, [1-82]. Vindobonae [Vienna].
- Bosch DT, Dance SP, Moolenbeek RG, Oliver PG (1995) *Seashells of Eastern Arabia*. Montivate Publishing, Dubai, 296 pp.
- Boyd WE, Higham CFW, Thosarat R (1996) The Holocene palaeogeography of the southeast margin of the Bangkok plain, Thailand, and its archaeological implications. *Asian Perspectives* 35: 193–207.
- Bozzetti L (1991) *Comitas ilariae* new species from the Philippines. *La Conchiglia* 22: 26–28.
- Brandt RAM (1974) The non-marine aquatic Mollusca of Thailand. *Archiv für Molluskenkunde*, 105: 1–423.
- Bratcher T, Cernohorsky WO (1987) Living terebras of the world. A monograph of the recent Terebridae of the world. American Malacologists, Melbourne, Florida & Burlington, Massachusetts, 240 pp.
- Bruguière JG (1789) Encyclopédie méthodique ou par ordre de matières. Histoire naturelle des vers, volume 1. Pancoucke, Paris, xviii, 344 [Dates after Evenhuis, (2003)] pp.
- Bussarawit S (1991) The family Muricidae in the PMBC reference collection: Application of a database system. *Phuket Marine Biological Center Special Publication* 9: 29–34.
- Cairns S, Kitahara MV (2012) An illustrated key to the genera and subgenera of the Recent azooxanthellate Scleractinia (Cnidaria, Anthozoa), with an attached glossary. *ZooKeys* 227: 1–47.
- Cairns SD, Hoeksema BW, Van der Land J (1999) Appendix: List of extant stony corals. *Atoll Research Bulletin* 459: 13–46.
- Cappetta H (2012) Chondrichthyes. Mesozoic and Cenozoic Elasmobranchii: teeth. *Handbook of paleoichthyology*. Vol. 3E. Verlag Dr. Friedrich Pfeil, Munich, 512 pp.
- Carrillo-Briceño JD, Maxwell E, Aguilera OA, Sánchez R, Sánchez-Villagra MR (2015) Sawfishes and other elasmobranch assemblages from the Mio-Pliocene of the South Caribbean (Urumaco Sequence, Northwestern Venezuela). *PLoS ONE* 10: 1–30.
- Cernohorsky WO (1971) The family Naticidae (Mollusca: Gastropoda) in the Fiji Islands. *Records of the Auckland Institute and Museum* 8: 169–208.
- Cernohorsky WO (1972) *Marine Shells of the Pacific*: v. 2. Pacific Publications, Sydney, 411 pp.
- Cernohorsky WO (1975) The taxonomy of some Indo-Pacific Mollusca part 3. With descriptions of new taxa and remarks on an Ecuadorian fossil species of Turridae. *Records of the Auckland Institute and Museum* 12: 212–234.

- Cernohorsky WO (1978) Tropical Pacific Marine Shells. Pacific Publications, Sydney, 352 pp.
- Cernohorsky WO (1984) Systematics of the family Nassariidae (Mollusca: Gastropoda). Bulletin of the Auckland Institute and Museum 14: i–iv, 1–356.
- Chabangborn A (2017) Review of paleoclimatic reconstruction in Thailand between the last glacial maximum and the mid Holocene. Burapha Science Journal 22: 61–77 [in Thai]
- Chaiwathee S, Dumrongjowattana P, Matchacheep S, Chalermwat C (2007) Pre-list of some marine microsnails from Bangsaen beach, Chonburi province (Mollusca: Scaphopoda and Gastropoda). In: Proceedings of 45th Kasetsart University Annual Conference: Science. Kasetsart University, Bangkok, 573–576 pp.
- Chan BK-K, Prabowo RE, Lee K-S (2009) Crustacean Fauna of Taiwan: Barnacles, Volume 1. Cirripedia: Thoracica Excluding the Pyrgomatidae and Acastinae. National Taiwan Ocean University, Taiwan, 297 pp.
- Chan S-Y, Lau WL (2021) Biodiversity Record: The edible shipworm, *Bactrionophorus thoracites*. Nature in Singapore 14: e2021112.
- Chan S-Y, Lau WL (2022) Biodiversity Record: The scalariform nutmeg snail, *Scalptia scalariformis*, in Singapore. Nature in Singapore 15: e2022103.
- Chen Z-y, Tan Y-h, Lian X-p (2015) Taxonomic note on the Lineate Nerite: *Nerita balteata* Reeve, 1855 (Gastropoda, Neritidae, *Nerita*). Journal of Tropical Oceanography 34: 74–76.
- Choowong M (2002) The geomorphology and assessment of indicators of sea-level change to study coastal evolution from the gulf of Thailand. International Symposium on “Geology of Thailand”. Department of Mineral Resource, Bangkok, Thailand, pp. 207–220.
- Clark AM, Rowe FWE (1971) Monograph of shallow-water indo-west Pacific Echinoderms. Trustees of the British Museum (Natural History), London, x + 238 pp.
- Clark HL (1925) A catalogue of the recent sea-urchins (Echinoidea) in the collection of the British Museum (Natural History). Trustees of the Museum, London, 250 pp.
- Clark PU, Dyke AS, Shakun JD, Carlson AE, Clark J, Wohlfarth B, Mitrovica JX, Hostetler SW, McCabe AM (2009) The Last Glacial Maximum. Science 325: 710–714.
- Coan EV (2002) The eastern Pacific recent species of the Corbulidae (Bivalvia). Malacologia 44: 47–105.
- Coan EV, Kabat AR (2012) The malacological works and taxa of Sylvanus Hanley (1819–1899). Malacologia 55: 285–359.
- Compagno LJV (1984) FAO Species Catalogue. Vol. 4. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Part 2 - Carcharhiniformes. FAO Fisheries Synopsis No 125 4: 251–655.
- Compagno LJV, White WT, Last PR (2008) *Glyphis garricki* sp. nov., a new species of river shark (Carcharhiniformes: Carcharhinidae) from northern Australia and Papua New Guinea, with a redescription of *Glyphis glyphis* (Müller & Henle, 1839). CSIRO Marine and Atmospheric Research Paper 22: 203–225.
- Cossignani T (1994) Bursidae of the world. L'Informatore Piceno, Ancona, 119 pp.
- Cox JB (1968) A review of the engineering characteristics of therecent marine clays in southeast Asia. Asian Institute of Technology Research Report 6, Bangkok.
- Dechruksa W, Sritongtae S, Namchote S, Krailas D (2014) Species diversity of molluscs obtained from mangrove forest and estuary of river in the Eastern coast of Thailand. Tokyo International Conference on Life Science and Biological Engineering, Tokyo, Japan, 931–937 pp.
- Dekkers A (2018) Two new genera in the family Melongenidae from the Indo-Pacific and comments on the identity of *Hemifusus zhangyii* Kosuge, 2008 and *Pyrula elongata* Lamarck, 1822 (Gastropoda, Neogastropoda: Buccinoidea). Gloria Maris 57: 40–50.
- Deshayes GP (1826) Anatomie et monographie du genre Dentale. Mémoires de la Société d'Histoire Naturelle de Paris (2) 2: 321–378.
- Deshayes GP (1830) Encyclopédie méthodique ou par ordre de matières. Histoire naturelle des Vers et Mollusques. Vol. 2: viii+1-594 [part 1: i-vi, 1-256, Livraison 101, 1 Feb 1830; part 2: 1-144, Livraison 101, 1 Feb. 1830].

- Dey A (2006) Handbook on mangrove associate molluscs of Sundarbans. Zoological Survey of India, Kolkata, xi + 96 pp.
- Dey A, Ramakrishna (2007) Marine molluscs: Bivalvia, Scaphopoda and Cephalopoda. Fauna of Andhra Pradesh, State Fauna Series Vol 5 Part 7. Zoological Survey of India, Kolkata, pp. 149–260.
- Dharma B (2005) Recent and fossil Indonesian shells. ConchBooks, Hackenheim, 424 pp.
- Dheeradilok P, Chaimanee N, Piccoli G, Robba E (1984) On the Quaternary stratigraphy and fossils of Senanivate Housing Project area, Bangkok Metropolis. *Memorie di Scienze Geologiche* 36: 413–426.
- Dijkstra HH, Beu AG (2018) Living scallops of Australia and adjacent waters (Mollusca: Bivalvia: Pectinoidea: Propeamussiidae, Cycloclamyidae and Pectinidae). *Records of the Australian Museum* 70: 113–330.
- Dillwyn LW (1817) A descriptive catalogue of Recent shells, arranged according to the Linnean method; with particular attention to the synonymy, Vol. 1: 1–580 pp. John and Arthur Arch, London.
- Dunker W (1866) Novitates Conchologicae. Mollusca Marina. Beschreibung und Abbildung neuer oder wenig gekannter Meeres-Conchylien. Abt. II: Meeres Conchylien. Theodor Fischer, Cassel, iv + 144 pp., 145 pls pp.
- Egorov R (2000) Treasure of Russian Shells Vol. 4: Trochiformes: Trochidae, Calliostomatidae, Liotiidae, Turbinidae, Tricoliidae. Colus-Doverie, Moscow, 84 pp.
- Eichhorst TE (2008) Neritidae. In: Poppe GT (Ed) Philippine Marine Mollusks Volume I. ConchBooks, Hackenheim, Germany, pp. 266–281.
- Eichhorst TE (2016) Neritidae of the world. Volume 1. ConchBooks, Hackenheim, 695 pp.
- Fedosov AE, Malcolm G, Terryn Y, Gorson J, Modica MV, Holford M, Puillandre N (2019) Phylogenetic classification of the family Terebridae (Neogastropoda: Conoidea). *Journal of Molluscan Studies* 85: 359–388.
- Filander Z, Griffiths CL (2014) Additions to and revision of the South African echinoid fauna (Echinodermata: Echinoidea). *African Natural History* 10: 47–56.
- Fischer-Piette E, Delmas D (1967) Révision des Mollusques Lamellibranches du genre *Dosinia* Scopoli. Mémoires du Muséum national d'Histoire naturelle, Sér A – Zoologie 47: 1–91.
- Froese R, Pauly D (2024) FishBase. World Wide Web electronic publication. <https://www.fishbase.org>, version (02/2024). [Verified 18 February 2024]
- Garrard TA (1975) A revision of Australian Cancellariidae (Gastropoda: Mollusca). *Records of the Australian Museum* 30: 1–62.
- Garrick JAF (1982) Sharks of the genus *Carcharhinus*. NOAA Technical Report NMFS Circular vol. 445. NMFS Scientific Publications Office, Seattle, USA, 194 pp.
- Gemert L van (2003) Marine schelpen verzamelen op Ko Samui (Golf van Thailand). *De Kreukel* 39: 95–109.
- George N, Siddiqui G, Muhammad F, George Z (2021) DNA barcoding of gastropod *Telescopium telescopium* (Linnaeus, 1758) found at the Karachi coast, Pakistan. *Journal of Animal and Plant Sciences* 31: 1530–1536.
- Gmelin JF (1791) Vermes. In: Gmelin JF (Ed) Caroli a Linnaei Systema Naturae per Regna Tria Naturae, Ed 13 Tome 1(6). G.E. Beer, Lipsiae [Leipzig], pp. 3021–3910.
- Golikov AN, Kussakin OG (1978) Shell-bearing gastropods of the intertidal zone of the seas of the USSR. Opredeliteli po faune SSSR, izdavayemye Zoologicheskim Institutom AN SSSR 116: 1–292.
- Gopalakrishnan A, Divya PR, Basheer VS, Swaminathan TR, Kathirvelpandian A, Bineesh KK, Kumar RG, Jena JK (2012) Macro Flora & Fauna of the Gulf of Mannar: a checklist. National Bureau of Fish Genetic Resources, Lucknow, 127 pp.
- Gottfried MD, Samonds KE, Ostrowski SA, Andrianavalona TH, Ramihangihajason TN (2017) New evidence indicates the presence of barracuda (Sphyraenidae) and supports a tropical marine environment in the Miocene of Madagascar. *PLoS ONE* 12: e0176553.

- Gould AA (1847) Descriptions of new shells, received from Rev. Mr. Mason, of Burmah. Proceedings of the Boston Society of Natural History 2: 218–221.
- Gould AA (1860) Descriptions of new shells collected by the United States North Pacific Exploring Expedition. Proceedings of the Boston Society of Natural History 7: 323–340, 382–384.
- Groh K (2010) Ellobiidae. In: Poppe GT (Ed) Philippine Marine Mollusks Volume III. ConchBooks, Hackenheim, Germany, pp. 446–457.
- Habe T (1958) The Fauna of Akkeshi Bay: XXV. Gastropoda. Publication of the Akkeshi Marine Biological Station 8: 1–39.
- Haga T (2010) Pholadidae. In: Poppe GT (Ed) Philippine Marine Mollusks Volume IV. ConchBooks, Hackenheim, Germany, pp. 392–397.
- Hamli H, Idris MH, Abu Hena MK, Wong SK, Arshad A (2013) Checklist and habitat descriptions of edible gastropods from Sarawak, Malaysia. Journal of Fisheries and Aquatic Science 8: 412–418.
- Hanley SCT (1846a) An illustrated and descriptive catalogue of Recent shells, by Sylvanus Hanley, B.A., F.L.S. The plates forming a third edition of the Index Testaceologicus, by William Wood. W. Wood, London, 272 + 218 pp.
- Hanley SCT (1846b) Monograph of the genus *Tellina*. In: Sowerby II GB (Ed) Thesaurus conchyliorum; or, monographs of the genera of shells. Sowerby II, G. B., London, pp. 221–336.
- Hanley SCT (1856) An illustrated and descriptive catalogue of Revent bivalve shells. Williams & Norgate, London, xviii + 392 + 324 pp.
- Harzhauser M, Raven H, Landau BM, Kocsis L, Adnan A, Zuschin M, Mandic O, Briguglio A (2018) Late Miocene gastropods from northern Borneo (Brunei Darussalam, Seria Formation). Palaeontographica, Abt A: Palaeozoology – Stratigraphy 313: 1–79.
- Heaton TJ, Köhler P, Butzin M, Bard E, Reimer RW, Austin WEN, Bronk Ramsey C, Grootes PM, Hughen KA, Kromer B, Reimer PJ, Adkins J, Burke A, Cook MS, Olsen J, Skinner LC (2020) Marine20—the marine radiocarbon age calibration curve (0–55,000 cal BP). Radiocarbon 62: 779–820.
- Hedley C (1913) Studies on Australian Mollusca, part XI. Proceedings of the Linnean Society of New South Wales, Series 2 38: 258–339.
- Hemmen J (2007) Recent Cancellariidae. Annotated and illustrated catalogue of Recent Cancellariidae. Privately published, Wiesbaden, 428 pp.
- Henry DP, McLaughlin PA (1975) The barnacles of the *Balanus amphitrite* complex (Cirripedia, Thoracica). Zoologische Verhandelingen 141: 1–254.
- Henry DP, McLaughlin PA (1986) The recent species of *Megabalanus* (Cirripedia: Balanomorpha) with special emphasis on *Balanus tintinnabulum* (Linnaeus) sensu lato. Zoologische Verhandelingen 235: 3–69.
- Herman J, Hovestadt-Euler M, Hovestadt DC (1991) Contributions to the study of the comparative morphology of teeth and other relevant ichthyodorulites in living supraspecific taxa of chondrichthyan fishes. Part A: Selachii. No 2c: Order: Carcharhiniformes. Families: Proscylliidae, Hemigaleidae, Pseudotriakidae, Leptochariidae and Carcharhinidae. Bulletin de l'Institut royal des Sciences naturelles de Belgique, Biologie 61: 73–120.
- Hinds RB (1843) Descriptions of new species of *Nucula*, from the collections of Sir Edward Belcher, C.B., and Hugh Cuming, Esq. Proceedings of the Zoological Society of London 11: 97–101.
- Hinds RB (1844) Description of new species of shells. Proceedings of the Zoological Society of London 12(1844): 21–26.
- Hiro F (1939) Studies on the cirripedian fauna of Japan IV. Cirripeds of Formosa (Taiwan), with some geographical and ecological remarks on the littoral forms. Memoirs of the College of Science, Kyoto Imperial University, Series B 15: 245–284.
- Hollmann M (2008) Naticidae. In: Poppe GT (Ed) Philippine Marine Mollusks Volume I. ConchBooks, Hackenheim, Germany, pp. 482–501.
- Holthuis LB, Heerebout GR (1972) Vondsten van de zeepok *Balanus tintinnabulum* (Linnaeus, 1758) in Nederland. Zoologische Bijdragen 13: 24–31.

- Hombron JB, Jacquinot H (1848) Atlas d'Histoire Naturelle Zoologie par MM. Hombron et Jacquinot, Chirurgiens de l'Expédition. In: Jacquinot H (Ed) Voyage au Pole Sud et dans l'Océanie sur les corvettes l'Astrolabe et la Zélée...Livre 26. Gide et Cie, Paris, pp.
- Houart R (1992) The genus *Chicoreus* and related genera (Gastropoda: Muricidae) in the Indo-West Pacific. Mémoires du Muséum national d'Histoire naturelle Série A, Zoologie 154: 1–188.
- Houart R (2008) Muricidae. In: Poppe GT (Ed) Philippine Marine Mollusks Volume II. ConchBooks, Hackenheim, Germany, pp. 132–221.
- Houbrick RS (1984) Revision of higher taxa in genus *Cerithidea* (Mesogastropoda: Potamididae) based on comparative morphology and biological data. American Malacological Bulletin 2: 1–20.
- Houbrick RS (1991) Systematic review and functional morphology of the mangrove snails *Terebralia* and *Telescopium* (Potamididae; Prosobranchia). Malacologia 33: 289–338.
- Huang Q (1997) Morphological, allozymic, and karyotypic distinctions between *Neritina* (*Dostia*) *violacea* and *N. (D.) cornucopia* (Gastropoda: Neritoidea). Journal of Zoology (London) 241: 343–369.
- Huber M (2010) Compendium of bivalves. A full-color guide to 3,300 of the world's marine bivalves. A status on Bivalvia after 250 years of research. ConchBooks, Hackenheim, 901 pp.
- Huber M (2015) Compendium of Bivalves 2. A Full-Color Guide to the Remaining Seven Families. A Systematic Listing of 8'500 Bivalve Species and 10'500 Synonyms. ConchBooks, Hackenheim.
- Hylleberg J, Aungtonya C (2013) Biodiversity survey in Chalong Bay and surrounding areas, east coast of Phuket Island. Phuket Marine Biological Center Special Publication 32: 81–112.
- Hylleberg J, Kilburn RN (2003) Marine molluscs of Vietnam: annotations, voucher material, and species in need of verification. Phuket Marine Biological Center Special Publication 28: 1–300.
- Hyžný M, de Angeli A (2022) Mud lobster *Thalassina* Latreille, 1806 (Decapoda: Gebiidea: Thalassinidae), its Cenozoic occurrences in Italy and palaeobiogeography. Geodiversitas 44: 417–425.
- Jarupongsakul S, Thiramongkol N (1992) Holocene highstand shoreline of the Chao Phraya delta, Thailand. Journal of Southeast Asian Earth Sciences 7: 53–60.
- Johnson RI (1964) The recent Mollusca of Augustus Addison Gould. Bulletin of the United States National Museum 239: 1–182.
- Jonas JH (1846) Ueber die Gattung Proserpina, und Beschreibungen neuer Conchylien. Zeitschrift für Malakozoologie 1846: 10–16.
- Jones DS, Hewitt MA, Sampey A (2000) A checklist of the Cirripedia of the South China Sea. The Raffles Bulletin of Zoology Supplement No. 8: 233–307.
- Jones DS, Hosie AM (2016) A checklist of the barnacles (Cirripedia: Thoracica) of Singapore and neighbouring waters. Raffles Bulletin of Zoology Supplement No. 34: 241–311.
- Kang D-R, Tan KS, Liu L-L (2018) Egg-collar morphology and identity of nine species of Naticidae (Gastropoda) in Taiwan, with an assessment of their phylogenetic relationships. Journal of Molluscan Studies 84: 354–378.
- Kanthalrajan G, Pandey PK, Krishnan P, Deepak Samuel V, Bharti VS, Purvaja R (2017) Molluscan diversity in the mangrove ecosystem of Mumbai, west coast of India. Regional Studies in Marine Science 14: 102–111.
- Kantor YI, Sysoev AV (2006) Marine and brackish water Gastropoda of Russia and adjacent countries: an illustrated catalogue. KMK Scientific Press, Moscow, 372 pp.
- Karasawa H, Kobayashi N (2022) Cirripedes from the middle Pleistocene Atsumi Group, Japan, with a reevaluation of the genus *Adna* Sowerby, 1823 (Balanoidea: Pyrgomatidae). Bulletin of the Mizunami Fossil Museum 49: 67–93.
- Kepel' AA (2018) The first finding of *Fistulobalanus kondakovi* (Tarasov & Zevina, 1957) (Cirripedia: Thoracica) in the Russian waters of the Sea of Japan. Russian Journal of Biological Invasions 9: 50–52.
- Kesavan K, Palpandi C, Shanmugam A (2009) A checklist of malacofauna of the Vellar Estuarine Mangroves, India. Journal of Threatened Taxa 1: 382–384.

- Kilburn R, Hylleberg J (1998) Gastropods and bivalves from Thailand; with taxonomic notes and new records. Phuket Marine Biological Center Special Publication 18: 311–316.
- Kim HK (2020) Taxonomic Study of the Thoracican Barnacles in Korean Waters-Diversity, Host specificity, Distribution-. Seoul National University.
- Kim IH (2011) Flora and Fauna of Korea, Invertebrate Fauna of Korea, vol. 21, no. 6. Arthropoda: Crustacea: Cirripedia. Barnacles. National Institute of Biological Resources, Incheon, 144 pp.
- Kim IH, Kim HS (1980) Systematic studies on the cirripeds (Crustacea) from Korea. I. Balanomorph barnacles (Cirripedia, Thoracica, Balanomorpha). The Korean Journal of Zoology 23: 161–185.
- Knudsen J, Jensen KR (2001) Translations into English of Lorenz Spengler's papers on Bivalvia (1783–1798). Part 3. 1788: *Pholas siamensis* with notes on the type specimen. The oldest marine bivalve with a type locality in the Gulf of Thailand. Phuket Marine Biological Center Special Publication 25: 547–556.
- Kocsis L, Razak H, Briguglio A, Szabo M (2019) First report on a diverse Neogene cartilaginous fish fauna from Borneo (Ambug Hill, Brunei Darussalam). Journal of Systematic Palaeontology 17: 791–819.
- Köhler R (2020) Echinoids of the Indian Museum at Calcutta. III. Regular Echinoids: A Translation of Échinides du Musée indien à Calcutta. III. Echinides réguliers (J. M. Lawrence, Trans.). Herizos Press, Tampa, 147 pp.
- Kool HH (2007) *Nassarius garuda* n. sp., a new deepwater species from the Indonesian Tanimbar and Kai Islands and a review of the species *N. crematus* (Hinds, 1844), *N. euglyptus* (Sowerby III, 1914) and *N. siquijorensis* (A. Adams, 1852) (Gastropoda: Buccinoidea: Nassariidae). Miscellanea Malacologica 2: 87–92.
- Krajangdara T, Vidthayanon C, Ali A, Lim A, Sumontha M, Rodpradit S, Chansue N, Haetrakul T (2022) The Cartilaginous Fishes of Thailand and Adjacent Waters. Department of Fisheries, Thailand, 233 pp.
- Kumar R, Jaiswar AK, Jahageerdar S, Chakraborty SK, Pavan Kumar A, Prasad L (2017) Comparative taxonomic evaluation of *Thais* species (Order: Gastropoda; Family: Muricidae) of Mollusca from Maharashtra coast of India. Indian Journal of Geo Marine Sciences 46: 1098–1104.
- Kuroda T, Habe T, Oyama K (1971) The sea shells of Sagami Bay. Maruzen Co., Tokyo, xix, 741 (Japanese), 489 (English), 751 (index) pp.
- Ladd HS (1982) Cenozoic fossil mollusks from western Pacific islands; gastropods (Eulimidae and Volutidae through Terebridae). Professional Paper. - ed, 1–100 pp.
- Lam K, Morton B, Hodgson P (2008) Ahermatypic corals (Scleractinia: Dendrophylliidae, Oculinidae and Rhizangiidae) recorded from submarine caves in Hong Kong. Journal of Natural History 42: 729–747.
- Lamarck J-BM de (1818) Histoire naturelle des animaux sans vertèbres. Tome 5. Deterville/Verdière, Paris, 612 pp.
- Lamarck J-BM de (1822) Histoire naturelle des animaux sans vertèbres. Tome septième. de Lamarck, J.-B. M., Paris, 711 pp.
- Lane DJW, Marsh LM, Vandenspiegel D, Rowe FWE (2000) Echinoderm fauna of the South China Sea: An inventory and analysis of distribution patterns. The Raffles Bulletin of Zoology Supplement No. 8: 459–493.
- Last PR, White WT, de Carvalho MR, Séret B, Stehmann MFW, Naylor GJP (2016) Rays of the World. CSIRO Publishing,, Clayton South, 790 pp.
- Lebedev EB (2014) Intertidal gastropod mollusks of the Far Eastern Marine Biosphere Natural Reserve. In: Sayenko EM, Lutaenko KA (Eds) Abstracts of the Conference Mollusks of the Eastern Asia and Adjacent Seas. Dalnauka, Vladivostok, Russia, 56–59 pp.
- Li C, Corrigan S, Yang L, Straube N, Harris M, Hofreiter M, White WT, Naylor GJP (2015) DNA capture reveals transoceanic gene flow in endangered river sharks. Proceedings of the National Academy of Sciences 112: 13302–13307.

- Liao Y (1998) The Echinoderm fauna of Hainan Island. In: Morton B (Ed) The Marine Biology of the South China Sea III: Proceedings of the Third International Conference on the Marine Biology of the South China Sea. Hong Kong University Press, Hong Kong, 75–82 pp.
- Lim KC, White WT, Then AYH, Naylor GJP, Arunrugstichai S, Loh K-H (2022) Integrated taxonomy revealed genetic differences in morphologically similar and non-sympatric *Scoliodon macrorhynchos* and *S. laticaudus*. *Animals* 12: 681. DOI: 10.3390/ani12060681.
- Lin C-H, Wang Y-C, Ribas-Deulofeu L, Chang C-W, Li K-T (2022) Changes in marine resource consumption over the past 5000 years in southwestern Taiwan revealed by fish otoliths. *Journal of Archaeological Science: Reports* 42: 103400.
- Linnaeus C (1758) *Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis*. Editio decima, reformata [10th revised edition], vol. 1. Laurentius Salvius, Holmiae, 824 pp.
- Linnaeus C (1767) *Systema naturae per regna tria naturae: secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis*. Ed. 12. 1., Regnum Animale. 1 & 2. Laurentii Salvii, Holmiae [Stockholm].
- Low MEY, Tan SK (2014) *Ergaea* H. Adams & A. Adams, 1854, the correct genus for *Crepidula walshi* Reeve, 1859, with nomenclatural notes on *Syphopatella* Lesson, 1831, *Siphonipatella* L. Agassiz, 1846, and the incorrect subsequent spelling “*Syphopatella*” of authors (Gastropoda: Calyptraeoidea: Calyptraeidae). *Occasional Molluscan Papers* 3: 11–14.
- Lozano-Cortés DF, Londoño-Cruz E (2013) Checklist of barnacles (Crustacea; Cirripedia: Thoracica) from the Colombian Pacific. *Marine Biodiversity* 43: 463–471.
- Lozouet P (2008) Batillariidae & Potamididae. In: Poppe GT (Ed) Philippine Marine Mollusks Volume I. ConchBooks, Hackenheim, Germany, pp. 284–285.
- Lutaenko K (2015) The arcid collection of Carl Emil Lischke in the Zoological Institute, St. Petersburg. *Archiv für Molluskenkunde International Journal of Malacology* 144: 125–138.
- Lutaenko K, Noseworthy RG, Choi K-S (2019) Marine bivalve mollusks of Jeju Island (Korea). Part 1. *The Korean Journal of Malacology* 35: 149–238.
- Lynge H (1909) Marine Lamellibranchia. The Danish Expedition to Siam 1899–1900. Det Kongelige Danske Videnskabernes Selskab Skrifter, Naturvidenskab og Matematik Afdeling, Ser 7 5: 97–299.
- Majima R (1989) Cenozoic fossil Naticidae (Mollusca: Gastropoda) in Japan. *Bulletins of American Paleontology* 96: 1–159.
- Malyshkina TP, Ward DJ, Nazarkin MV, Nam G-S, Kwon S-H, Lee J-H, Kim T-W, Kim D-K, Baek D-S (2023) Miocene Elasmobranchii from the Duho Formation, South Korea. *Historical Biology* 35: 1726–1741.
- Marsh LM, Morrison SM (2004) Echinoderms of the Dampier Archipelago, Western Australia. Records of the Western Australian Museum Supplement No. 66: 293–342.
- Marsili S (2007) Revision of the teeth of the genus *Carcharhinus* (Elasmobranchii; Carcharhinidae) from the Pliocene of Tuscany, Italy. *Rivista Italiana di Paleontologia e Stratigrafia* 113: 79–95.
- Martens E von (1860) On the Mollusca of Siam. *Proceedings of the Zoological Society of London* 28: 6–18.
- Martin JC (2008) Nassariidae. In: Poppe GT (Ed) Philippine Marine Mollusks Volume II. ConchBooks, Hackenheim, Germany, pp. 114–129.
- Matsubara T, Komori K (2007) The first record of fossil *Ellobium* (Gastropoda: Ellobiidae) from northeastern Japan. *Venus* 65: 325–331.
- Merle D, Garrigues B, Pointier J-P (2011) Fossil and Recent Muricidae of the world. Part Muricinae. Conchbooks, Hackenheim, 648 pp.
- Meyer E, Nilkerd B, Glover EA, Taylor JD (2008) Ecological importance of chemosautotrophic lucinid bivalves in a peri-mangrove community in eastern Thailand. *Raffles Bulletin of Zoology Supplement* 18: 41–55.

- Middlefart P (1997) An illustrated checklist of Muricidae (Gastropoda: Prosobranchia) from the Andaman Sea, Thailand. Phuket Marine Biological Center Special Publication 17: 348–388.
- Milne Edwards H, Haime J (1848) Recherches sur les polypiers. Mémoire 3. Monographie des eupsammides. Annales des Sciences Naturelles, Zoologie, Series 3 10: 65–114.
- Milne Edwards H, Haime J (1849) Recherches sur les polypiers. Quatrième mémoire. Monographie des astréides (1). Suite. Quatrième section. Astréens agglomérés. Astreinae aggregatae. Annales des sciences naturelles, série 3, Zoologie 12: 95–197.
- Moh HH, Chong VC, Lim PE, Tan J (2013) Verification of four species of the mud lobster genus *Thalassina* (Crustacea: Decapoda: Gebiidea: Thalassinidae) using molecular and morphological characters. Raffles Bulletin of Zoology 61: 579–588.
- Moh ZC, Nelson JD, Brand EW (1969) Strength and deformation behavior of Bangkok Clay. In: Proceedings of the Seventh International Conference on Soil Mechanics, Vol 1. Mexico City, 287 pp.
- MolluscaBase (2023) MolluscaBase taxon details: *Potamocorbula* Habe, 1955. Available at <https://www.marinespecies.org/aphia.php?p=taxdetails&id=397174> [Verified 3 January 2023].
- Morris S, Purchon RD (1981) The marine shelled Mollusca of West Malaysia and Singapore part 3, Bivalvia. Journal of Molluscan Studies 47: 322–327.
- Mortensen T (1904) The Danish Expedition to Siam 1899–1900. III. Echinoidea (1). Kongelige Danske Videnskabelige Selskabs, Skrifter, Serie 7 1: 1–124.
- Mortensen T (1940) Contributions to the biology of the Philippine Archipelago and adjacent regions: Report on the Echinoidea collected by the United States Fisheries steamer Albatross during the Philippine Expedition, 1907–1910. Part 2. The Echinothuridae, Saleniidae, Arbaciidae, Aspidodiadematidae, Microphygidae, Diadematidae, Pedinidae, Temnopleuridae, Toxopneustidae, and Echinometridae. Bulletin of the United States National Museum 14: i–vi, 1–52.
- Mortensen T (1943) Monograph of the Echinoidea. III. 2 Camarodontia. I. Orthopsidae, Glyphocyphidae, Temnopleuridae and Toxopneustidae. C. A. Reitzel, Copenhagen, 533 pp.
- Mucharin A, Sumitrapijj R (2022) Echinoderms (Phylum Echinodermata) of Ko Yao Noi-Ko Yao Yai and nearby islands, Pang-Nga Province, Thailand. Science, Technology, and Social Sciences Procedia 2022: rsgp002.
- Mukhopadhyay AK, Sharma K, Ramakrishna (2013) Two new records of the genus *Polinices* and one of the *Natica* from India. Records of the Zoological Survey of India 113: 151–154.
- Müller J, Henle FGJ (1838) Systematische Beschreibung der Plagiostomen. Veit und Comp., Berlin, xxii, 200 [Pp. 201–228: 1838] pp.
- Müller J, Henle FGJ (1839) Systematische Beschreibung der Plagiostomen. Veit und Comp., Berlin, xxii, 200 [reset pp. 227–228, 229–102: 1839] pp.
- Mustapha N, Baharuddin N, Tan SK, Marshall DJ (2021) The neritid snails of Brunei Darussalam: their geographical, ecological and conservation significance. Ecologica Montenegrina 42: 45–61.
- Nabhitabhata J (2009) Checklist of Mollusca Fauna in Thailand. Office of Natural Resources and Environmental Policy and Planning, Bangkok, 576 pp.
- Nakamura I, Parin NV (1993) FAO species catalogue. Vol. 15. Snake mackerels and cutlassfishes of the world (Families Gempylidae and Trichiuridae). An annotated and illustrated catalogue of the snake mackerels, snoeks, escolars, gemfishes, sackfishes, domine, oilfish, cutlassfishes, scabbardfishes, hairtails, and frostfishes known to date. FAO Fisheries Synopsis No 125 15: 1–136.
- Nakamura I, Parin NV (1998) Trichiuridae Cutlassfishes. In: Carpenter KE, Niem VH (Eds) FAO species identification guide for fishery purposes The living marine resources of the Western Central Pacific Volume 6 Bony fishes part 4 (Labridae to Latimeriidae), estuarine crocodiles, sea turtles, sea snakes and marine mammals. FAO, Rome, pp. 3709–3720.
- Nateewathana A (1995) Taxonomic account of commercial and edible molluscs, excluding cephalopods, of Thailand. Phuket Marine Biological Center Special Publication 15: 93–116.

- Nateewathana A, Tantichodok P, Bussarawich S, Sirivejabandhu R (1981) Marine organisms in the Reference Collection. Phuket Marine Biological Center Research Bulletin 28: 43–86.
- Negri MP (2009) Fossil mollusc-faunas: Their bearing on the Holocene evolution of the Lower Central Plain of Bangkok (Thailand). Journal of Asian Earth Sciences 35: 524–544.
- Negri MP (2012) A Late Pleistocene transgression in Thailand: A marine molluscan fauna from Ban Praksa (Samut Prakan Province). Journal of Asian Earth Sciences 43: 145–149.
- Negri MP, Sanfilippo R, Basso D, Rosso A, Di Geronimo SI (2014) Molluscan associations from the Pak Phanang Bay (SW Gulf of Thailand) as a record of natural and anthropogenic changes. Continental Shelf Research 84: 204–218.
- Newman WA, Ross A (1976) A revision of the balanomorph barnacles: including a catalog of the species. Memoirs of the San Diego Society of Natural History 9: 1–108.
- Ng TH, Tan SK, Wong WH, Meier R, Chan S-Y, Tan HH, Yeo DCJ (2016) Molluscs for sale: Assessment of freshwater gastropods and bivalves in the ornamental pet trade. PLoS ONE 11: e0161130.
- Ngoc-Ho N, de Saint Laurent M (2009) The genus *Thalassina* Latreille, 1806 (Crustacea: Thalassinidea: Thalassinidae). Raffles Bulletin of Zoology Supplement No. 20: 121–158.
- Nimnate P, Chutakositkanon V, Choowong M, Pailoplee S, Phantuwongraj S (2015) Evidence of Holocene sea level regression from Chumphon coast of the Gulf of Thailand. ScienceAsia 41: 55–63.
- Nobuhara T (1993) The relationship between bathymetric depth and climate change and its effect on molluscan faunas of the Kakegawa Group, central Japan. Transactions and Proceedings of the Paleontological Society of Japan New series 170: 159–185.
- Nolf F (2009) *Thais lacera* (Born, 1778) (Mollusca: Gastropoda: Muricidae: Muricidae): an early record of an exotic species from the western Mediterranean Sea. Neptunea 8: 9–15.
- Ohe F (1976) Fossil otoliths of Sciaenid fishes from the Tertiary and Quaternary formations, Tobai district, central Japan and their paleontological vicissitudes. Bulletin of the Mizunami Fossil Museum 3: 73–97.
- Ohtsubo M, Egashira K, Koumoto T, Bergado DT (2000) Mineralogy and chemistry, and their correlation with the geotechnical index properties of Bangkok clay: comparison with Ariake clay. Soils and Foundations 40: 11–21.
- Okutani T (2017) Marine mollusks in Japan, ed. 2. 2 vols. Tokai University Press, Tokyo, Japan, 1375 pp.
- Oliver GJH, Terry JP (2019) Relative sea-level highstands in Thailand since the Mid-Holocene based on ¹⁴C rock oyster chronology. Palaeogeography, Palaeoclimatology, Palaeoecology 517: 30–38.
- Oliver PG (1992) Bivalved seashells of the Red Sea. National Museum of Wales, Cardiff, 330 pp.
- Olivera B, Sysoev A (2008) Turridae. In: Poppe GT (Ed) Philippine Marine Mollusks Volume II. ConchBooks, Hackenheim, Germany, pp. 764–787.
- Oliviera LPH de (1941) Contribuicao ao conhecimento dos coustaceos do Rio de Janeiro, subordem ‘Balanomorpha’ (Cirripedia, Thoracica). Memoirs of the Institute Oswaldo Crus 36: 1–31.
- Oyama K (1953) Review of the known species of the Japanese Turridae (1). Venus 17: 151–160.
- Öztürk B, Bitlis BB (2013) *Eunaticina papilla* (Gmelin, 1791) (Naticidae, Gastropoda): A new alien mollusc in the eastern Mediterranean. Triton 28: 7–8.
- Palanisamy SK, Prasanna Kumar C, Paramasivam P, Sundaresan U (2020) DNA barcoding of horn snail *Telescopium telescopium* (Linnaeus C, 1758) using mt-COI gene sequences. Regional Studies in Marine Science 35: 101109.
- Philippi RA (1836) Beschreibung einiger neuen Conchylien-Arten und Bemerkungen über die Gattung Lacuna von Turton. Archiv für Naturgeschichte 2: 224–235.
- Pilsbry HA (1895) Catalogue of the marine molluscs of Japan, with descriptions of new species, and notes on others collected by F. Stearns. F. Stearns, Detroit, viii + 196 pp.
- Pilsbry HA (1916) The sessile barnacles (Cirripedia) collected in the collections of the U. S. National Museum: Including a monograph of the American species. Bulletin of the United States National Museum 93: 1–366.

- Pitombo FB (2004) Phylogenetic analysis of the Balanidae (Cirripedia, Balanomorpha). *Zoologica Scripta* 33: 261–276.
- Pochai A, Kingtong S, Sukparangsi W, Khachonpisitsak S (2017) The diversity of acorn barnacles (Cirripedia, Balanomorpha) across Thailand's coasts: The Andaman Sea and the Gulf of Thailand. *Zoosystematics and Evolution* 93: 13–34.
- Ponder WF, Vokes EH (1988) A revision of the Indo-West Pacific fossil and recent species of *Murex* s.s. and *Haustellum* (Mollusca: Gastropoda: Muricidae). *Records of the Australian Museum Supplement* 8: 1–160.
- Poppe GT (2010a) Arcidae. In: Poppe GT (Ed) Philippine Marine Mollusks Volume III. ConchBooks, Hackenheim, Germany, pp. 468–491.
- Poppe GT (2010b) Corbulidae. In: Poppe GT (Ed) Philippine Marine Mollusks Volume IV. ConchBooks, Hackenheim, Germany, pp. 386–391.
- Poppe GT (2010c) Veneridae. In: Poppe GT (Ed) Philippine Marine Mollusks Volume IV. ConchBooks, Hackenheim, Germany, pp. 258–309.
- Poutiers JM (1998a) Bivalve (Acephala, Lamellibranchia, Pelecypoda). In: Carpenter KE, Niem VH (Eds) FAO Species Identification Guide for Fishery Purposes The Living Marine Resources of the Western Central Pacific Volume 1 Seaweeds, corals, bivalves and gastropods. Food and Agriculture Organization of the United Nations, Rome, Italy, pp. 123–362.
- Poutiers JM (1998b) Gastropods. In: Carpenter KE, Niem VH (Eds) FAO Species Identification Guide for Fishery Purposes The Living Marine Resources of the Western Central Pacific Volume 1 Seaweeds, corals, bivalves and gastropods. Food and Agriculture Organization of the United Nations, Rome, Italy, pp. 363–648.
- Powell AWB (1942) The New Zealand Recent and fossil Mollusca of the family Turridae with general notes on turrid nomenclature and systematics. *Bulletin of the Auckland Institute and Museum* 2: 1–188.
- Powell AWB (1969) The family Turridae in the Indo-Pacific. Part 2. The subfamily Turriculinae. *Indo-Pacific Mollusca* 2: 215–416.
- Prabowo RE, Yamaguchi T (2005) A new mangrove barnacle of the genus *Fistulobalanus* (Cirripedia: Amphibalaninae) from Sumbawa Island, Indonesia. *Journal of the Marine Biological Association of the United Kingdom* 85: 929–936.
- Price ARG (1981) Studies on the echinoderm fauna of the western Arabian Gulf. *Journal of Natural History* 15: 1–15.
- Printrakoon C, Wells FE, Chitramvong Y (2008) Distribution of molluscs in mangroves at six sites in the upper Gulf of Thailand. *The Raffles Bulletin of Zoology* 18: 247–257.
- Puspasari IA, Yamaguchi T, Angsupanich S (2000) Reexamination of a little-known mangrove barnacle, *Balanus patelliformis* Bruguière (Cirripedia, Thoracica) from the Indo-West Pacific. *Sessile Organisms* 16: 1–13.
- Puspasari IA, Yamaguchi T, Angsupanich S (2001) *Balanus thailandicus* sp. nov., a new mangrove barnacle of the *Balanus amphitrite* complex (Cirripedia, Balanomorpha) from Satun, Southwest Thailand. *Sessile Organisms* 18: 27–33.
- Putchakarn S (2018) Biodiversity of sea urchins (Echinoidea, Echinodermata) in Thai Waters. Institute of Marine Science, Burapha University, 108 pp.
- Putchakarn S, Sonchaeng P (2004) Echinoderm fauna of Thailand: History and inventory reviews. *ScienceAsia* 30: 417–428.
- Ramakrishna, Dey A, Barua S, Mukhopadhy A (2007) Marine molluscs: Polyplacophora and Gastropoda. Fauna of Andhra Pradesh, State Fauna Series Vol 5 Part 7. Zoological Survey of India, Kolkata, pp. 1–148.
- Rau JL, Nutalaya P (1983) Geology of the Bangkok clay. *Geological Society of Malaysia Bulletin* 16: 99–116.
- Raven H, Vermeulen JJ (2007) Notes on molluscs from NW Borneo and Singapore. 2. A synopsis of the Ellobiidae (Gastropoda, Pulmonata). *Vita Malacologica* 4: 29–62.

- Raven JGM (2019) Notes on molluscs from NW Borneo - Dispersal of molluscs through nipa rafts. *The Festivus* 51: 3–10.
- Reeve L (1844) Monograph of the genus *Arca*. *Conchologia Iconica*, or, illustrations of the shells of molluscous animals, vol 2, pls 1–17, and unpaginated text. Lovell Reeve & Co. , London, pp. [without pagination] [stated dates: pl. 1–2: 1843; 1843–1817: 1844].
- Reeve LA (1855) Monograph of the genus *Nerita*. *Conchologia Iconica*, or, illustrations of the shells of molluscous animals, vol 9, pls 1–19, and unpaginated text. L. Reeve & Co., London, pp.
- Reeve LA (1859) Monograph of the genus *Crepidula*. *Conchologia Iconica*, or, illustrations of the shells of molluscous animals, vol 11, pl 1–5 and unpaginated text. L. Reeve & Co., London, pp.
- Reid DG (2014) The genus *Cerithidea* Swainson, 1840 (Gastropoda: Potamididae) in the Indo-West Pacific region. *Zootaxa* 3775: 1–65.
- Reid DG, Ozawa T (2016) The genus *Pirenella* Gray, 1847 (= *Cerithideopsis* Thiele, 1929) (Gastropoda: Potamididae) in the Indo-West Pacific region and Mediterranean Sea. *Zootaxa* 4076: 1–91.
- Reimer PJ, Austin WEN, Bard E, Bayliss A, Blackwell PG, Bronk Ramsey C, Butzin M, Cheng H, Edwards RL, Friedrich M, Grootes PM, Guilderson TP, Hajdas I, Heaton TJ, Hogg AG, Hughen KA, Kromer B, Manning SW, Muscheler R, Palmer JG, Pearson C, van der Plicht J, Reimer RW, Richards DA, Scott EM, Southon JR, Turney CSM, Wacker L, Adolphi F, Büntgen U, Capone M, Fahrni SM, Fogtmann-Schulz A, Friedrich R, Köhler P, Kudsk S, Miyake F, Olsen J, Reinig F, Sakamoto M, Sookdeo A, Talamo S (2020) The IntCal20 Northern Hemisphere radiocarbon age calibration curve (0–55 cal kBP). *Radiocarbon* 62: 725–757.
- Riede K (2004) Global register of migratory species from global to regional scales. Final Report of the R & D Project 808 05 081 (p. 329). Federal Agency for Nature Conservation, Bonn, Germany.
- Robba E, Di Geronimo I, Chaimanee N, Negri M, Sanfilippo R (2003) Mollusc associations of the Northern Gulf of Thailand and Holocene analogs from the Bangkok Clay. *Mahasarakham University Journal* 22: 191–213.
- Robba E, Di Geronimo I, Chaimanee N, Negri MP, Sanfilippo R (2002) Holocene and Recent shallow soft-bottom mollusks from the northern Gulf of Thailand area: Bivalvia. *Bollettino Malacologico* 38: 49–132.
- Robba E, Di Geronimo I, Chaimanee N, Negri MP, Sanfilippo R (2004) Holocene and recent shallow soft-bottom mollusks from the Northern Gulf of Thailand area: Scaphopoda, Gastropoda, additions to Bivalvia. *La Conchiglia, International Shell Magazine* 35: 1–290.
- Robba E, Di Geronimo I, Chaimanee N, Negri MP, Sanfilippo R (2007) Holocene and recent shallow soft-bottom mollusks from the Western Gulf of Thailand: Pak Phanang Bay and additions to Phetchaburi fauna. *Bollettino Malacologico* 43: 1–98.
- Roberts TR (2006) Rediscovery of *Glyptis gangeticus*: Debunking the mythology of the supposed "Gangetic freshwater shark". *Natural History Bulletin of the Siam Society* 54: 261–278.
- Röding PF (1798) *Museum Boltenianum sive Catalogus cimeliorum e tribus regnis naturæ quæ olim collegerat Joa. Fried Bolten, M. D. p. d. per XL. annos proto physicus Hamburgensis. Pars secunda continens Conchylia sive Testacea univalvia, bivalvia & multivalvia.* Trapp, Hamburg, viii, 199 pp.
- Rosell NC (1973) Some thoracic barnacles (Crustacea: Cirripedia) of Manila Bay. *Kalikasan, the Philippine Journal of Biology* 2: 69–95.
- Roussy H (1991) Family Pectinidae recorded – a checklist. *Phuket Marine Biological Center Special Publication* 9: 21.
- Royal Irrigation Department (2009) Soil and Geological Survey Report, Preparation of Cross Sections and Soil Classification in Irrigation Bureau Area No. 11. Royal Irrigation Department, Bangkok, 1–28 pp.
- Sahlmann B, Poppe GT (2010) Dentaliidae. In: Poppe GT (Ed) *Philippine Marine Mollusks Volume IV*. ConchBooks, Hackenheim, Germany, pp. 402–411.

- Sakai K, Türkay M (2012) A review of the species of the genus *Thalassina* (Thalassinidea, Thalassinidae). *Crustaceana* 85: 1339–1376.
- Salvi D, Mariottini P (2021) Revision shock in Pacific oysters taxonomy: the genus *Magallana* (formerly *Crassostrea* in part) is well-founded and necessary. *Zoological Journal of the Linnean Society* 192: 43–58.
- Samyn Y (2003) Shallow-water regular echinoids (Echinodermata : Echinoidea) from Kenya. *African Zoology* 38: 193–212.
- Sanpanich K (2011) Marine bivalves occurring on the east coast of the Gulf of Thailand. *ScienceAsia* 37: 195–204.
- Sanpanich K, Duangdee T (2013) The biodiversity of marine gastropods of Thailand in the late decade. *Malaysian Journal of Science* 32: 47–64.
- Sartori AF, Printrakoon C, Mikkelsen PA, Bieler R (2008) Siphonal structure in the Veneridae (Bivalvia: Heterodontia) with an assessment of its phylogenetic application and a review of venerids in the Gulf of Thailand. *Raffles Bulletin of Zoology Supplement* 18: 103–125.
- Sathiamurthy E, Voris H (2006) Maps of Holocene sea level transgression and submerged lakes on the Sunda shelf. *The Natural History Journal of Chulalongkorn University* 2: 1–43.
- Scarabino V (1995) Scaphopoda of the tropical Pacific and Indian Oceans, with description of 3 new genera and 42 new species. *Mémoires du Muséum national d'histoire naturelle* 167: 189–379.
- Schrenck LI (1861) Vorläufige Diagnosen einiger neuer Molluskenarten aus der Meerenge der Tartarei und dem nord-japanischen Meere. *Bulletin de l'Académie Impériale des Sciences de St Pétersbourg* 4: 408–413.
- Schultz H (2015) *Handbook of Zoology: Echinodermata, Volume 1: Echinoidea*. Walter de Gruyter GmbH, Berlin/Boston, 386 pp.
- Schwarzans W (1993) A comparative morphological treatise of recent and fossil otoliths of the family Sciaenidae (Perciformes). *Piscium catalogus: Part Otolithi Piscium*. Vol. 1. Verlag Dr. Friedrich Pfeil, München, 245. pp.
- Shimada K, Egi N, Tsubamoto T, Maung-Maung MM, Thaung-Htike TH, Zin-Maung-Maung-Thein ZM, Nishioka Y, Sonoda T, Takai M (2016) The extinct river shark *Glypis pagoda* from the Miocene of Myanmar and a review of the fossil record of the genus *Glypis* (Carcharhiniformes: Carcharhinidae). *Zootaxa* 4161: 237–251.
- Sinsakul S (2000) Late Quaternary geology of the lower central plain, Thailand. *Journal of Asian Earth Sciences* 18: 415–426.
- Smith EA (1873) Remarks on a few species belonging to the family Terebridae, and descriptions of several new forms in the collection of the British Museum. *Annals and Magazine of Natural History* ser 4 11: 262–271.
- Smith EA (1875) A list of the Gasteropoda collected in Japanese seas by Commander H. C. St. John, R. N. *Annals and Magazine of Natural History* ser 4 15: 414–427.
- Smith EA (1877) Descriptions of new species of Conidae and Terebridae. *Annals and Magazine of Natural History* ser 4 19: 222–231.
- Smith EA (1879a) On a collection of marine shells from the Andaman Islands. *Proceedings of the Zoological Society of London* 1878: 804–821.
- Smith EA (1879b) On a collection of Mollusca from Japan. *Proceedings of the Zoological Society of London* 1879: 181–218.
- Somboon JRP (1988) Paleontological study of the recent marine sediments in the lower central plain, Thailand. *Journal of Southeast Asian Earth Sciences* 2: 201–210.
- Somboon JRP, Thiramongkol N (1992) Holocene highstand shoreline of the Chao Phraya delta, Thailand. *Journal of Southeast Asian Earth Sciences* 7: 53–60.
- Songtham W, Musika S, Mildenhall DC, Cochran UA, Kojevnikova D (2015) Development of the lower central plain of Thailand with history of human settlements: evidence from pollen, spores and diatoms. *Journal of Geological Resource and Engineering* 2: 98–107.

- Songtham W, Phanwong P, Seelanan T (2007) Middle Holocene peat and mollusk shells from Ongkharak area, Nakhon Nayok, central Thailand: evidence of in situ deposits during a marine transgression period. International Conference on Geology of Thailand: Towards Sustainable Development and Sufficiency Economy: 177–179.
- Sowerby I GB (1822) The genera of recent and fossil shells, for the use of students, in conchology and geology. Published in 42 numbers. Vol. 1, pls 1-126 [1821-1825]; vol. 2, pls 127-262 + text (unpaginated) [1825-1834]. privately published, London.
- Sowerby I GB (1833[1832-1841]) The conchological illustrations or, Coloured figures of all the hitherto unfigured recent shells. privately published, London.
- Sowerby II GB (1842) Monograph of the genus *Pecten*. In: II GBS (Ed) Thesaurus Conchyliorum, or monographs of Genera of Shells. Sowerby II, G. B., London, pp. 45–78.
- Spengler L (1792) Betragtninger og Anmærkninger ved den Linneiske Slægt *Pholas* bland de mangeskallede Muskeler, med dens hidindtil bekendte gamle og nye Arter, samt den dermed i Forbindelse staaende Slægt *Teredo* Linn. Skrivter af Naturhistorie-Selskabet, Kiøbenhavn 2: 72–106.
- Spengler L (1794) Nøiere Bestemmelse og Udvidelse af det Linneiske Genus *Solen*. Skrivter af Naturhistorie-Selskabet, Kiøbenhavn 3: 81–114.
- Springer VG (1964) A revision of the carcharhinid shark genera *Scoliodon*, *Loxodon*, and *Rhizoprionodon*. Proceedings of the United States National Museum 115: 559–632.
- Sri-aroon P, Lohachit C, Harada M (2004) Survey of brackish-water snails in eastern Thailand. The Southeast Asian Journal of Tropical Medicine and Public Health 35: 150–155.
- Sri-aroon P, Lohachit C, Harada M (2005) Brackish-water mollusks of Surat Thani Province, southern Thailand. The Southeast Asian Journal of Tropical Medicine and Public Health 36: 180–188.
- Steiner G, Kabat AR (2014) Catalog of species-group names of Recent and fossil Scaphopoda (Mollusca). Zoosystema 26: 549–726.
- Stuiver M, Polach HA (1977) Discussion: reporting of ¹⁴C data. Radiocarbon 19: 355–363.
- Subba Rao NV, Dey A (2000) Catalogue of the marine molluscs of Andaman and Nicobar Islands. Records of the Zoological Survey of India, Occasional Paper 187: 1–323.
- Surakiatchai P, Choowong M, Charusiri P, Charoentitirat T, Chawchai S, Pailoplee S, Chabangborn A, Phantuwongraj S, Chutakositkanon V, Kongsen S, Nimnate P, Bissen R (2018) Paleogeographic reconstruction and history of the sea level change at Sam Roi Yot National Park, Gulf of Thailand. Tropical Natural History 18: 112–134.
- Swennen C, Moolenbeek RG, Ruttanadakul N, Hobbelink H, Dekker H, Hajisamae S (2001) The Molluscs of the Southern Gulf of Thailand. The Biodiversity Research and Training Program, Bangkok, 210 pp.
- Takaya Y (1969) Topographical analysis of the southern basin of the central plain, Thailand. Southeast Asian Studies 7: 293–300.
- Tan KS (2000) Species checklist of Muricidae (Mollusca: Gastropoda) in the South China Sea. The Raffles Bulletin of Zoology Supplement No. 8: 495–512.
- Tan SK, Clements R (2008) Taxonomy and distribution of the Neritidae (Mollusca: Gastropoda) in Singapore. Zoological Studies 47: 481–494.
- Tanabe S, Saito Y, Sato Y, Suzuki Y, Sinsakul S, Tiyanairach S, Chaimanee N (2003) Stratigraphy and Holocene evolution of the mud-dominated Chao Phraya delta, Thailand. Quaternary Science Reviews 22: 789–807.
- Tantanasiwong R (1978) An illustrated checklist of marine shelled gastropods from Phuket Island, adjacent mainland and offshore island, western peninsular Thailand. Phuket Marine Biological Center Research Bulletin 21: 1–63.
- Tantanasiwong R (1979) A checklist of marine bivalves from Phuket Island, adjacent mainland and offshore island, western peninsular Thailand. Phuket Marine Biological Center Research Bulletin 27: 1–15.

- Tarasov NI, Zevina GB (1957) Barnacles (Cirripedia, Thoracica) of the Soviet Seas. Fauna SSSR, Crustacea 6: 1–267.
- Taylor JD, Glover EA (2005) Cryptic diversity of chemosymbiotic bivalves: A systematic revision of worldwide *Anodontia* (Mollusca: Bivalvia: Lucinidae). Systematics and Biodiversity 3: 281–338.
- Tesch P (1915) Jungtertiäre und quartäre Mollusken von Timor. I. Teil. Palaontologie von Timor, Stuttgart 5: 1–70.
- Tesch P (1920) Jungtertiäre und quartäre Mollusken von Timor. II. Teil. Palaontologie von Timor, Stuttgart 8: 41–121.
- Thach NN (2005) Shells of Vietnam. ConchBooks, Hackenheim, 338 pp.
- Thiele J (1925) Gastropoden der Deutschen Tiefsee-Expedition. II Teil. Wissenschaftliche Ergebnisse der Deutschen Tiefsee-Expedition auf dem Dampfer "Valdivia" 1898–1899 17: 35–382, pls 313–346 [reprints paginated 381–348, pls 381–334].
- Thunberg CP (1793) Tekning och Beskrifning på en stor Ostronsort ifrån Japan. Kongliga Vetenskaps Akademiens Nya Handlingar 14: 140–142.
- Torigoe K, Inaba A (2011) Revision on the classification of recent Naticidae. Bulletin of the Nishinomiya Shell Museum 7: 1–133.
- Tudu PC, Balakrishnan S (2018) First report of Partridge Sundial Snail *Architeconica perdix* (Hinds, 1844) from Northern part of East Coast of India. Records of the Zoological Survey of India 118: 198–201.
- Tudu PC, Yennawar P, Ghorai N, Tripathy B, Mohapatra A (2018) An updated checklist of marine and estuarine mollusc of Odisha coast. Indian Journal of Geo Marine Sciences 47: 1537–1560.
- Tudu PC, Yennawar P, Mohapatra A (2019) First report of two ark shells, *Anadara consociata* (E.A. Smith, 1885) and *A. troscheli* (Dunker, 1882) (Arcidae: Anadarinae) from Indian waters with notes on morpho-taxonomy of some related species from east coast of India. Records of the Zoological Survey of India 119: 34–48.
- Umitsu M, Tiyanairach S, Chaimanee N, Kawase K (2002) Late Holocene sea-level change and evolution of the central plain, Thailand. The Symposium on Geology of Thailand, Bangkok, Thailand, 201–206 pp.
- Verhecken A (1986) The recent Cancellariidae of Indonesia (Neogastropoda: Cancellariacea). Gloria Maris 25: 29–66.
- Verhecken A (1997) Mollusca, Gastropoda: Arafura Sea Cancellariidae collected during the KARUBAR Cruise. In: Crosnier A, Bouchet P (Eds) Résultats des Campagnes MUSORSTOM 16 Campagne Franco-Indonésienne KARUBAR Mémoires du Muséum national d'Histoire naturelle Série A, Zoologie, 172. pp. 295–324.
- Verhecken A (2008) Cancellariidae. In: Poppe GT (Ed) Philippine Marine Mollusks Volume II. ConchBooks, Hackenheim, Germany, pp. 816–825.
- Voigt M, Weber D (2011) Field guide for sharks of the genus *Carcharhinus*. Verlag Dr. Friedrich Pfeil, München.
- Vongpanich V (1996) The Arcidae of Thailand. Phuket Marine Biological Center Special Publication 16: 177–192.
- Vongpanich V, Matsukuma A (2004) Family Noetiidae in Thailand. Phuket Marine Biological Center Research Bulletin 65: 31–44.
- Way K, Purchon RD (1981) The marine shelled Mollusca of West Malaysia and Singapore part 2. Polyplacophora and Gastropoda. Journal of Molluscan Studies 47: 313–321.
- Wells FE, Sanpanich K, Tan SK, Duangdee T (2021) The Marine and Estuarine Molluscs of Thailand. Lee Kong Chian Natural History Museum National University of Singapore, Singapore, 195 pp.
- White WT (2012) A redescription of *Carcharhinus dussumieri* and *C. sealii*, with resurrection of *C. coatesii* and *C. tjutjot* as valid species (Chondrichthyes: Carcharhinidae). Zootaxa 3241: 1–34.
- White WT, Appleyard SA, Sabub B, Kyne PM, Harris M, Lis R, Baje L, Usu T, Smart JJ, Corrigan S, Yang L, Naylor GJP (2015) Rediscovery of the Threatened River Sharks, *Glyptis garricki* and *G. glyphis*, in Papua New Guinea. PLoS ONE 10: e0140075.

- White WT, Last PR, Naylor GJP (2010) *Scoliodon macrorhynchos* (Bleeker, 1852), a second species of spadenose shark from the Western Pacific (Carcharhiniformes: Carcharhinidae). In: Last PR, White WT, Pogonoski JJ (Eds) Descriptions of New Australian Chondrichthyans, CSIRO Marine and Atmospheric Research Paper 32. CSIRO, Hobart, Australia, pp. 61–76.
- Whitley GP (1934) Notes on some Australian sharks. Memoirs of the Queensland Museum 10: 180–200.
- Wilson B (1993) Australian marine shells, Prosobranch Gastropods, Vol. 1. Odyssey Publishing, Kallaroo, Australia, 408 pp.
- Wilson B (1994) Australian marine shells, Prosobranch Gastropods, Vol. 2. Odyssey Publishing, Kallaroo, Australia, 370 pp.
- Wium-Andersen G (1977) Marine *Nerita* species from Phuket island and their chromosome numbers (Gastropoda : Neritidae). Phuket Marine Biological Center Research Bulletin 15: 1–9.
- Wong HW (2009) The Mactridae (Mollusca: Bivalvia) of East Coast Park, Singapore. Nature in Singapore 2: 283–296.
- Yadav R, Malla PK, Dash D, Bhoi G, Patro S, Mohapatra A (2019) Diversity of gastropods and bivalves in the mangrove ecosystem of Paradeep, east coast of India: a comparative study with other Indian mangrove ecosystems. Molluscan Research 39: 325–332.
- Yamaguchi T (1977a) Taxonomic studies on some fossil and recent Japanese Balanoidea (Part 1). Transactions and Proceedings of the Paleontological Society of Japan New series 1977: 135–160.
- Yamaguchi T (1977b) Taxonomic studies on some fossil and recent Japanese Balanoidea (Part 2). Transactions and Proceedings of the Paleontological Society of Japan New series 1977: 161–201.
- Yamaguchi T (1980) A new species belonging to the *Balanus* amphitrite Darwin group (Cirripedia, Balanomorpha) from the late Pleistocene of Japan; an example of peripheral speciation. Journal of Paleontology 54: 1084–1101.
- Yang W, Cai Y, Kuang X (2017) Color Atlas of Molluscs of the South China Sea. China Agriculture Press, Beijing, 275 pp.
- Yen T-C (1944) Notes on some unfigured type-specimens of Chinese mollusks from the North Pacific Expedition. Proceedings of the California Academy of Sciences 23: 561–586.
- Yokoyama M (1927) Mollusca from the Upper Musashino of Western Shimosa and Southern Musashi. Journal of the Faculty of Science, University of Tokyo, section 2 1: 439–457.
- Yoosukh W, Jitkaew M (1997) Marine wood-boring bivalves (Pholadacea) in the Gulf of Thailand. Phuket Marine Biological Center Special Publication 17: 401–405.
- Zullo V (1984) New genera and species of balanoid barnacles from the Oligocene and Miocene of North America. Journal of Paleontology 58: 1312–1338.
- Zvonareva S, Kantor Y (2016) Checklist of gastropod molluscs in mangroves of Khanh Hoa province, Vietnam. Zootaxa 4162: 401–437.

Table 1. AMS ^{14}C ages of shells and charcoal collected from the clay pit of Ongkharak in Nakhon Nayok, central Thailand.

Lab Code	Sample type	Depths (m) below the uppermost part of a marine clay layer	^{14}C age (yr BP)	Calibrated ^{14}C (2-sigma) age (cal yr BP)
D-AMS 044190	Shell (<i>Tellinides conspicuus</i>)	2.2	5303 ± 26	5882–5464
D-AMS 044191	Shell (<i>Pholas orientalis</i>)	2.2	5202 ± 28	5757–5318
D-AMS 044193	Charcoal	2.2	5047 ± 26	5899–5726
D-AMS 044195	Charcoal	3.4	6186 ± 29	7166–6983
D-AMS 044194	Shell (<i>Magallana cf. gigas</i>)	4	7241 ± 32	7901–7510
D-AMS 044192	Charcoal	4.6	7913 ± 26	8784–8600

Table 2. Molluscan species assemblages from the Bangkok Clay deposits in Nakhon Nayok recovered in the present work, and from other sites retrieved from previous literature (indicated by an asterisks). NA = data not available.

Class	Species in this study	Lower Central Plain and Chao Phraya delta (Somboon 1988; Somboon and Thiramongkol 1992)	Lower Central Plain of Bangkok and inland of Phetchaburi coast (Robba et al. 2007; Negri 2009)	Ban Praks a, Samut Prakan (Negri 2012)	Sam Roi Yot National Park, Phetchaburi (Surakiathai et al. 2018)	Habitat	Substrate preference
Gastropoda	1. <i>Homalopoma cf. sangarensis</i>	-	-	-	-	Sublittoral	Sand, hard
	2. <i>Neripteron cf. violaceum</i>	-	-	-	-	Intertidal, mangrove	Mud, hard
	3. <i>Nerita articulata</i>	*	-	-	-	Intertidal	Mud, hard

						mangrove	
	4. <i>Cerithidea obtusa</i>	-	*	-	-	Intertidal, mangrove	Mud, hard
	5. <i>Pirenella incisa</i>	-	-	-	-	Intertidal, mangrove	Mud
	6. <i>Telescopium telescopium</i>	-	-	-	-	Intertidal, mangrove	Mud
	7. <i>Eunaticina papilla</i>	-	*	-	-	Intertidal — sublittoral	Sand
	8. <i>Natica stellata</i>	-	*	-	-	Sublittoral	Sand
	9. <i>Natica vitellus</i>	-	*	-	-	Intertidal — sublittoral	Sand, mud
	10. <i>Paratectonica tigrina</i>	*	*	*	*	Intertidal — sublittoral	Sand, mud
	11. <i>Ergaea walshi</i>	-	*	-	-	Intertidal — sublittoral	Hard
	12. <i>Bufoaria rana</i>	-	-	-	-	Sublittoral	Sand, mud
	13. <i>Merica elegans</i>	-	-	-	-	Sublittoral	Sand, mud
	14. <i>Scalptia scalariformis</i>	-	*	-	-	Sublittoral	Sand, mud
	15. <i>Pseudoneptuna varicosa</i>	-	*	-	-	Sublittoral	NA
	16. <i>Brunneifusus ternatanus</i>	*	-	-	-	Sublittoral	Sand, mud

	17. <i>Nassarius micans</i>	-	*	-	-	Intertidal — sublittoral	Sand
	18. <i>Nassarius siquijorensis</i>	-	*	-	*	Intertidal — sublittoral	Sand, mud
	19. <i>Chicoreus capucinus</i>	*	-	-	-	Intertidal , mangrove	Sand, mud, hard
	20. <i>Indothais gradata</i>	*	-	-	-	Sublittoral	Hard
	21. <i>Indothais lacera</i>	-	*	-	-	Intertidal — sublittoral	Mud, hard
	22. <i>Murex trapa</i>	*	*	-	*	Sublittoral	Sand, mud
	23. <i>Maoritomella vallata</i>	-	*	-	-	Sublittoral	NA
	24. <i>Pseudoetrema fortilirata</i>	-	*	-	-	Sublittoral	Sand, mud
	25. <i>Turricula javana</i>	*	*	-	*	Intertidal — sublittoral	Sand, mud, hard
	26. <i>Paradrillia melvilli</i>	-	*	-	-	Sublittoral–upper bathyal	NA
	27. <i>Inquisitor vulpionis</i>	-	*	-	-	Sublittoral	Sand
	28. <i>Comitas ilariae</i>	-	-	-	-	Sublittoral	Sand
	29. <i>Duplicaria tricincta</i>	-	*	-	-	Sublittoral	Sand, mud
	30. <i>Granuliterebra bathyraphae</i>	-	*	*	-	Intertidal — sublittoral	Sand, mud

	31. <i>Pristiterebra miranda</i>	-	*	-	-	Sublittoral	NA
	32. <i>Architectonica perdix</i>	-	*	-	*	Sublittoral	Sand, mud
	33. <i>Cylichna modesta</i>	-	*	-	-	Sublittoral	Sand, mud
	34. <i>Ellobium aurisjudeae</i>	-	*	-	-	Intertidal, mangrove	Sand, mud, hard
	35. <i>Cassidula nucleus</i>	-	*	-	-	Intertidal, mangrove	Mud, hard
Bivalvia	1. <i>Jupiteria puellata</i>	-	*	*	-	Sublittoral	Sand, mud
	2. <i>Sacella mauritiana</i>	-	*	-	-	Sublittoral	Sand, mud, hard
	3. <i>Anadara inaequivalvis</i>	-	*	-	*	Intertidal – Sublittoral	Sand, mud
	4. <i>Anadara indica</i>	-	*	-	*	Intertidal – sublittoral	Sand, mud
	5. <i>Tegillarca granosa</i>	*	*	-	*	Intertidal, mangrove	Mud
	6. <i>Tegillarca nodifera</i>	-	-	-	-	Intertidal – sublittoral, mangrove	Sand, mud
	7. <i>Estellacar olivacea</i>	*	*	-	-	Intertidal – sublittoral, mangrove	Sand, mud

	8. <i>Noetiella pectunculifor mis</i>	-	*	-	-	Intertidal – sublittoral, mangrove	Sand, mud
	9. <i>Magallana cf. gigas</i>	-	*	-	*	Intertidal – sublittoral	Mud, hard
	10. <i>Placuna placenta</i>	-	*	*	*	Intertidal – sublittoral	Sand, mud
	11. <i>Volachlamys singaporina</i>	-	-	-	-	Intertidal – sublittoral	Sand, mud, hard
	12. <i>Pegophysem a bialata</i>	-	-	-	-	Intertidal – sublittoral	Sand, mud
	13. <i>Geloina bengalensis</i>	-	*	-	-	Intertidal , mangrove	Mud
	14. <i>Lutraria complanata</i>	-	*	-	-	Sublitoral	Sand, mud
	15. <i>Standella pellucida</i>	-	*	-	-	Intertidal – sublittoral, mangrove	Sand, mud
	16. <i>Tellinides conspicuus</i>	-	-	-	-	Sublitoral	NA
	17. <i>Joannisiella oblonga</i>	*	*	-	*	Intertidal – sublittoral	Mud
	18. <i>Dosinia dilecta</i>	-	*	-	*	Sublitoral	Mud
	19. <i>Paratapes undulatus</i>	*	*	*	*	Intertidal –	Sand, mud

						sublittora 1	
	20. <i>Placamen lamellatum</i>	-	*	-	-	Intertidal — sublittora 1	Sand, mud
	21. <i>Corbula fortisulcata</i>	-	*	-	*	Intertidal — sublittora 1	Sand, mud, hard
	22. <i>Potamocorb ula</i> sp.	--	NA	NA	NA	Intertidal	NA
	23. <i>Martesia striata</i>		*	-	-	Intertidal — sublittora 1	Hard
	24. <i>Pholas orientalis</i>	-	*	-	-	Intertidal — sublittora 1	Sand, mud, hard
	25. Teredinidae indet.	-	NA	NA	NA	Intertidal — sublittora 1	Hard
	26. <i>Cultellus maximus</i>	-	*	-	-	Intertidal — sublittora 1, mangrov e	Mud
	27. <i>Siliqua minima</i>	-	*	-	-	Intertidal — sublittora 1	Sand, mud
Scaphopo da	1. <i>Dentalium variabile</i>	-	*	*	-	Sublitor al	Mud

Table 3. Fish remains recovered from the Bangkok Clay deposits in Nakhon Nayok.

Class Chondrichthyes						
Order	Family	Species			No.	Material
Carcharhiniformes	Carcharhinidae	<i>Carcharhinus</i> cf. <i>amblyrhynchoides</i>			1	tooth
		<i>Carcharhinus</i> cf. <i>amblyrhynchos</i>			1	tooth

		<i>Carcharhinus</i> cf. <i>leucas</i>	4	teeth
		<i>C. leucas/amboinensis</i>	10	teeth
		<i>Carcharhinus</i> cf. <i>sorrah</i>	2	teeth
		<i>Carcharhinus</i> spp.	7	teeth
		<i>Glyphis</i> sp.	59	teeth
		<i>Scoliodon</i> cf. <i>laticaudus</i>	3	teeth
		Carcharhinidae indet.	10	teeth
Myliobatiformes	Dasyatidae	<i>Pastinachus</i> sp.	3	teeth
Class Actinopterygii				
Order	Family	Species	No.	Material
Scombriformes	Trichiuridae	Trichiuridae indet.	3	teeth
<i>incertae sedis</i> in Eupercaria	Sciaenidae	<i>Johnius</i> sp.	1	otolith




Fig. 1. Map of the lower central plain of Thailand (orange). A red star indicates the location of the study area.




Fig. 2. The study area and the sediment profile of the clay pit. **A** the clay pit of Ongkharak, Nakhon Nayok Province in central Thailand **B** the Bangkok clay layer bearing numerous *in situ* complete shells of *Placuna placenta* **C** Shells of a bivalve *Anadara inaequivalvis* buried *in situ* in the Bangkok clay deposits **D** lithological profile of the clay pit **E** the contact between two

successive layers: coarse-grained sand (upper) and black clay (lower) **F** the contact between three successive layers: black clay (upper), lateritic soil (middle), and yellowish brown clay to silt (lower) **G** the contact between two successive layers: purplish (upper) and greyish (lower) clay to sand **H** the contact between two successive layers: greyish clay to sand (upper) and reddish brown lateritic layer (lower).




Fig. 3. Size comparison of coral and gastropods found in this study. **A** *Oulangia cf. stokesiana* **B** *Neripteron cf. violaceum* **C** *Nerita articulata* **D** *Natica stellata* **E** *Natica vitellus* **F** *Homalopoma cf. sangarense* **G** *Eunaticina papilla* **H** *Ergaea walshi* **I** *Pirenella incisa* **J** *Scalptia scalariformis* **K** *Telescopium telescopium* **L** *Merica elegans* **M** *Cerithidea obtusa* **N** *Bufonaria rana* **O** *Nassarius micans* **P** *Nassarius siquijorensis* **Q** *Paratectonatica tigrina*.



Fig. 4. Size comparison of gastropods found in this study. **A** *Brunneifusus ternatanus* **B** *Murex trapa* **C** *Indothais lacera* **D** *Pristiterebra miranda* **E** *Chicoreus capucinus* **F** *Indothais gradata* **G** *Ellobium aurisjudeae* **H** *Inquisitor vulpionis* **I** *Turridula javana* **J** *Architectonica perdix* **K** *Pseudoneptunea varicosa* **L** *Cassidula nucleus* **M** *Pseudoetrema fortilirata* **N** *Paradrillia melvilli* **O** *Duplicaria tricincta* **P** *Granuliterebra bathyrhaphes* **Q** *Comitas ilariae* **R** *Cyllichna modesta* **S** *Maoritomella vallata*.




Fig. 5. Coral and gastropods. **A** *Oulangia* cf. *stokesiana* **B** *Homalopoma* cf. *sangarensense* **C** *Neripteron* cf. *violaceum* **D** *Nerita articulata* **E** *Cerithidea obtusa* **F** *Pirenella incisa* **G** *Telescopium telescopium*.




Fig. 6. Gastropods. **A** *Eunaticina papilla* **B** *Natica stellata* **C** *Natica vitellus* **D** *Paratectonatica tigrina* **E** *Ergaea walshi* **F** *Bufoaria rana* **G** *Merica elegans* **H** *Scalptia scalariformis*.




Fig. 7. Gastropods. **A** *Pseudoneptunea varicosa* **B** *Brunneifusus ternatanus* **C** *Nassarius micans* **D** *Nassarius siquijorensis* **E** *Chicoreus capucinus* **F** *Indothais gradata*.




Fig. 8. Gastropods. **A** *Indothais lacera* **B** *Murex trapa* **C** *Turricula javana* **D** *Inquisitor vulpionis* **E** *Pristiterebra miranda* **F** *Granuliterebra bathyrhaphes*.




Fig. 9. Gastropods. **A** *Maoritomella vallata* **B** *Pseudoetrema fortilirata* **C** *Paradrillia melvilli* **D** *Comitas ilariae*.




Fig. 10. Gastropods and bivalves. **A** *Duplicaria tricincta* **B** *Cylichna modesta* **C** *Jupiteria puellata* **D** *Saccella mauritiana*.




Fig. 11. Gastropods and bivalves. **A** *Architectonica perdix* **B** *Ellobium aurisjudeae* **C** *Cassidula nucleus* **D** *Anadara inaequivalvis* **E** *Anadara indica* **F** *Tegillarca granosa*.



Fig. 12. Size comparison of bivalves, scaphopod and other invertebrates found in this study. **A** *Tegillarca nodifera* **B** *Volachlamys singaporina* **C** *Tegillarca granosa* **D** *Noetiella pectunculiformis* **E** *Magallana cf. gigas* **F** *Estellacar olivacea* **G** *Anadara indica* **H** *Dentalium variabile* **I** *Siliqua minima* **J** *Corbula fortisulcata* **K** *Fistulobalanus kondakovi* **L** *Placamen lamellatum* **M** *Martesia striata* **N** *Potamocorbula* sp. **O** *Jupiteria puellata* **P** *Saccella mauritiana* **Q** *Paratapes undulatus* **R** *Dosinia dilecta* **S** *Joannisiella oblonga* **T** *Temnotrema siamense* **U** *Pholas orientalis*.



Fig. 13. Size comparison of bivalves and other invertebrates found in this study. **A** *Placuna placenta* **B** *Standella pellucida* **C** *Geloina bengalensis* **D** *Anadara inaequivalvis* **E** *Pegophysema bialata* **F** *Megabalanus cf. tintinnabulum* **G** *Cultellus maximus* **H** *Lutraria complanata* **I** *Thalassina* sp. **J** *Tellinides conspicuus*.




Fig. 14. Bivalves. **A** *Tegillarca nodifera* **B** *Estellacar olivacea* **C** *Noetiella pectunculiformis* **D** *Magallana* cf. *gigas* **E** *Placuna placenta* **F** *Volachlamys singaporina*.




Fig. 15. Bivalves. **A** *Pegophysema bialata* **B** *Geloina bengalensis* **C** *Lutraria complanata* **D** *Standella pellucida* **E** *Tellinides conspicuus* **F** *Joannisiella oblonga*.




Fig. 16. Bivalves. **A** *Dosinia dilecta* **B** *Paratapes undulatus* **C** *Placamen lamellatum* **D** *Corbula fortisulcata* **E** *Potamocorbula* sp. **F**, **G** *Martesia striata* **G** showing ... **H** *Pholas orientalis* **I** calcareous burrows of Teredinidae indet.




Fig. 17. Bivalves and other invertebrates. **A** *Cultellus maximus* **B** *Siliqua minima* **C** *Dentalium variabile* **D** *Thalassina* sp. **E** *Fistulobalanus kondakovi* **F** *Megabalanus* cf. *tintinnabulum* **G** *Temnotrema siamense*.




Fig. 18. Carcharhinid shark teeth. **A–E** *Carcharhinus* cf. *amblyrhynchoides*, specimen CUF-NKNY-S3-2 in **A** lingual **B** labial **C** mesial **D** distal and **E** apical views **F–J** *Carcharhinus* cf. *amblyrhynchos*, specimen CUF-NKNY-S3-10 in **F** lingual **G** labial **H** mesial **I** distal and **J** apical views.




Fig. 19. Carcharhinid shark teeth - *Carcharhinus* cf. *leucas* **A–I** specimen CUF-NKNY-Q04 in **A** lingual **B** labial and **C–F** different apical views **G–I** close-ups on the serrations of the distal cutting edge at the level of **G** distal heel and mesial cutting edge at the level of **H** upper part of the cusp and **I** heel. **J–N** specimen CUF-NKNY-SA-1 in **J** lingual **K** labial **L** mesial **M** distal and **N** apical views.




Fig. 20. Carcharhinid shark teeth - *Carcharhinus* cf. *sorrah* **A–G** specimen CUF-NKNY-3.1 **A** lingual **B** apical **C** labial **D** mesial and **E** distal views **F**, **G** close-ups of the serration **F** on the distal heel and **G** on the middle part of the cusp mesial cutting-edge **H–N** specimen CUF-NKNY-3.2 **H** lingual **I** apical **J** labial **K** mesial and **L** distal views **M** close-up of the serration on the lower part of the mesial cutting-edge **N** close-up of the serration between the heel and the base of the distal cutting edge.




Fig. 21. Carcharhinid shark teeth - *Carcharhinus* spp. **A–G** specimen CUF-NKNY-16.1 in **A** lingual **B** labial **C** mesial **D** distal and **E, F** apical views **G** close-ups of the serration on the upper part of the cusp mesial cutting edge in lingual view **H–L** specimen S2 in **H** labial **I** apical **J** lingual **K** mesial and **L** distal views.




Fig. 22. Carcharhinid shark teeth - *Carcharhinus* sp. **A–H** specimen CUF-NKNY-S1 in **A** lingual **B** labial **C** mesial **D** distal and **E, F** apical views **G, H** close-ups on the serration **G** at mid-height of the cusp mesial cutting-edge and **H** at the level of the distal heel and base of the cusp.




Fig. 23. Carcharhinid shark teeth - *Glyphis* sp. specimen CUF-NKNY-8.2, upper lateral tooth in **A** lingual **B** labial **C** mesial **D** distal and **E** apical views **F–H** close-ups of the serration **F** at the base of the cusp mesial cutting-edge, **G** at the tip of the cusp and **H** on the distal heel.




Fig. 24. Carcharhinid shark teeth - *Glyphis* sp. specimen CUF-NKNY-Q03, upper anterior tooth in **A** lingual **B** labial **C** mesial **D** distal and **E** apical views **F–H** close-ups of the serration **F** at the base, **G** in the middle and **H** in the upper part of the cusp mesial cutting-edge.




Fig. 25. Carcharhinid shark teeth - *Glyphis* sp. **A–E** specimen CUF-NKNY-SC-3, upper anterolateral tooth in **A** lingual **B** labial **C** mesial **D** distal and **E** apical views **F–J** specimen CUF-NKNY-SD-3, upper lateral tooth in **F** lingual **G** labial **H** mesial **I** distal and **J** apical views.




Fig. 26. Carcharhinid shark teeth - *Glyphis* sp. **A–H** specimen CUF-NKNY-14, lower tooth in **A** lingual **B** labial **C** mesial **D** distal and **E** apical views **F–H** close-ups on the serration **F** on the distal heel, **G** at the base of the cusp distal cutting-edge and **H** in the middle part of the cusp mesial cutting-edge **I–M** specimen CUF-NKNY-S1-2, lower anterior tooth in **I** lingual **J** labial **K** mesial **L** distal and **M** apical views.




Fig. 27. Carcharhinid shark teeth - *Glyphis* sp. **A–E** specimen CUF-NKNY-S3-1, lower tooth in **A** lingual **B** labial **C** mesial **D** distal and **E** apical views **F–J** specimen CUF-NKNY-S3-6, lower tooth in **F** lingual **G** labial **H** mesial **I** distal and **J** apical views.




Fig. 28. Carcharhinid shark teeth - *Glyphis* sp. **A–E** specimen CUF-NKNY-S4-1, lower tooth in **A** lingual **B** labial **C** mesial **D** distal and **E** apical views **F–J** specimen CUF-NKNY-S4-4, lower anterior tooth in **F** lingual **G** labial **H** mesial **I** distal and **J** apical views.




Fig. 29. Carcharhinid shark teeth - *Scoliodon* cf. *laticaudus* **A–E** specimen CUF-NKNY-S3, lower anterolateral tooth in **A** lingual **B** labial **C** mesial **D** distal and **E** apical views **F–J** specimen CUF-NKNY-S4, upper anterior-anterolateral tooth in **F** lingual **G** labial **H** mesial **I** distal and **J** apical views **K–O** specimen CUF-NKNY-S5, parasympphyseal tooth in **K** lingual **L** labial **M** mesial **N** distal and **O** apical views.




Fig. 30. Dasyatid ray teeth - *Pastinachus* sp. **A–F** specimen CUF-NKNY-7.1 in **A** lingual **B** labial **C**, **F** profile **D** occlusal and **E** basal views **G–L** specimen CUF-NKNY-7.2 in **G** lingual **H** labial **I**, **L** profile **J** occlusal and **K** basal views **M–R** specimen CUF-NKNY-8.1 in **M** lingual **N** labial **O**, **R** profile **P** occlusal and **Q** basal views.




Fig. 31. Trichiuridae indet. **A–F** specimen CUF-NKNY-2.1, anterior tooth in **A**, **B** lateral **C** anterior **D** posterior and **E** apical views **F** close-up of the tip of the tooth **G–K** specimen CUF-NKNY-2.2, anterior tooth in **G**, **H** lateral **I** anterior **J** posterior and **K** apical views **L–P** specimen CUF-NKNY-18.2, lateral tooth in **L**, **M** lateral **N** anterior **O** posterior and **P** apical views.



Fig. 32. Sciaenid otolith belonging to *Johnius* sp. CUF-NKNY-16.2, right sagitta otolith in **A** inner **B** external **C** anterior and **D** posterior views.




Fig. 33. Paleogeographic maps illustrating the evolution of the Chao Phraya delta modified from Tanabe et al. (2003), during **A** 8,000–7,000 cal yr BP and **B** 7,000–6,000 cal yr BP. Red stars indicate the location of the study area.