



HAL
open science

Poly(ADP-ribose) polymerase-1 and its ambiguous role in cellular life and death

Maria Castedo, Antoine Lafarge, Guido Kroemer

► **To cite this version:**

Maria Castedo, Antoine Lafarge, Guido Kroemer. Poly(ADP-ribose) polymerase-1 and its ambiguous role in cellular life and death. *Cell Stress*, 2023, 7 (1), pp.1-6. 10.15698/cst2023.01.275 . hal-04596744

HAL Id: hal-04596744

<https://hal.science/hal-04596744>

Submitted on 31 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Poly(ADP-ribose) polymerase-1 and its ambiguous role in cellular life and death

Maria Castedo^{1,2,*}, Antoine Lafarge^{1,2,3} and Guido Kroemer^{1,2,4,*}

¹ Equipe 11 labellisée par la Ligue contre le Cancer, Université de Paris Cité, Sorbonne Université, INSERM U1138, Centre de Recherche des Cordeliers, 75006 Paris, France.

² Metabolomics and Cell Biology Platforms, Gustave Roussy Cancer Campus, Villejuif, France.

³ Faculté de médecine, Université de Paris Saclay, Kremlin Bicêtre, France.

⁴ Institut du Cancer Paris CARPEM, Department of Biology, Hôpital Européen Georges Pompidou, Assistance Publique-Hôpitaux de Paris, Paris, France.

* Corresponding Authors:

Maria Castedo, Equipe 11 labellisée par la Ligue contre le Cancer, Université de Paris, Sorbonne Université, INSERM U1138, Centre de Recherche des Cordeliers, 75006 Paris, France; Tel: +33 1 44 27 76 61; Fax: +33 1 44 27 76 74; E-mail: maria.castedo-delrieu@sorbonne-universite.fr

Guido Kroemer, Equipe 11 labellisée par la Ligue contre le Cancer, Université de Paris, Sorbonne Université, INSERM U1138, Centre de Recherche des Cordeliers, 75006 Paris, France; Tel: +33 1 44 27 76 67; Fax: +33 1 44 27 76 74; E-mail: kroemer@orange.fr

ABSTRACT The deletion of the gene coding for poly(ADP-ribose) polymerase-1 (PARP1) or its pharmacological inhibition protects mice against cerebral ischemia and Parkinson's disease. In sharp contrast, PARP1 inhibitors are in clinical use for the eradication of vulnerable cancer cells. It appears that excessive PARP1 activation is involved in a specific cell death pathway called parthanatos, while inhibition of PARP1 in cancer cells amplifies DNA damage to a lethal level. Hence, PARP1 plays a context-dependent role in cell fate decisions. In addition, it appears that PARP1 plays an ambiguous role in organismal aging.

doi: 10.15698/cst2023.01.275

Received originally: 10.10.2022;

in revised form: 31.12.2022

Accepted 02.01.2023.

Published 23.01.2023

Keywords: PARP1 inhibition, DNA damage, parthanatos, cancer, neurological diseases, aging, cellular stress response.

Abbreviations:

PAR – poly(ADP-ribose), **NAD⁺** – nicotinamide adenine dinucleotide, **PARP1** – poly(ADP-ribose) polymerase-1, **AIF** – apoptosis inducing factor, **MOMP** – mitochondrial outer membrane permeabilization, **PAAN** – parthanatos-associated apoptosis-inducing factor nuclease, **MIF** – macrophage migration inhibitor factor, **MPTP** – 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine, **AIMP2** – aminoacyl-tRNA synthetase complex interacting multi-functional protein-2, **MCAO** – middle cerebral artery occlusion, **AD** – Alzheimer's disease, **HD** – Huntington's disease, **ALS** – amyotrophic lateral sclerosis, **NAMPT** – nicotinamide phosphoribosyltransferase, **MMP** – mitochondrial membrane permeabilization, **MIMP** – mitochondrial inner membrane permeabilization, **DBC1** – protein deleted in breast cancer 1.

In biology, the integration of intracellular circuitries is achieved by the multifunctionality of molecules, molecular complexes and organelles [1]. Poly(ADP-ribose) polymerase-1 (PARP1) exemplifies a multitasking protein that fulfills several signaling functions in the context of cellular stress response, senescence and aging, as well as pathological cell death. Here, we will briefly summarize the multiple functions of PARP1 (**Figure 1**).

PARP1 IN DNA REPAIR: A TARGET FOR ANTICANCER DRUGS

PARP1 is best known as is a DNA repair enzyme present in the nuclei of mammalian cells. In response to single-strand DNA breaks, PARP1 initiates the synthesis of poly(ADP-ribose) (PAR) chains from nicotinamide adenine dinucleotide (NAD⁺). This process results in the covalent attach-




FIGURE 1: PARP1 context-dependent functions in cellular life and death. Poly(ADP-ribose) polymerase-1 (PARP1) is a DNA repair enzyme activated in cancer cells treated with DNA damaging agents, and pharmacological PARP1 inhibitors can be used to improve chemotherapy and radiotherapy efficacy. PARP1 inhibition may also have neuroprotective effects by decreasing parthanatos-induced neuronal death. However, the ambivalent role of PARP1 in organismal aging and cell death modalities should be considered in view of the possible side effects of long-term treatments with PARP1 inhibitors. PARP1, Poly(ADP-ribose) polymerase-1; PAR: Poly(ADP-ribose); AIF: apoptosis inducing factor.

ment of PAR to multiple proteins. PAR then acts as a signal for the recruitment and the activation of other DNA repair enzymes [2]. Logically, cancer cells that are treated with DNA damaging agents (such as radiotherapy or chemotherapeutic cytotoxicants such as cisplatin) must activate PARP1 to survive. Indeed, cisplatin-resistant cells upregulate PARP1 activity and hence contain higher PAR levels in their nuclei [3]. This is clinically relevant because high PAR levels detected by immunohistochemistry have a poor prognostic impact in several cancer types, correlating with reduced immunosurveillance [3-6]. In addition, BRCA2- and other homologous recombination-defective cells have increased PARP1 activation meaning that cell survival relies on a constitutive activation of PARP1 [7].

Pharmacological PARP1 inhibitors can be used to sensitize cancer cells to chemotherapy and radiotherapy [8-10]. Moreover, cells relying on constitutive PARP1 activation become sensitive to monotherapy with PARP1 inhibitors [3,6,7]. Indeed, several PARP1 inhibitors such as niraparib,

olaparib, rucaparib and talazoparib have been clinically approved for the treatment of oncological patients with germline mutations in BRCA1/2, as well as for epithelial ovarian, fallopian tube and primary peritoneal cancer [9,10]. These small-molecule inhibitors interact with the binding site of NAD^+ and inhibit the synthesis of PAR chains. However, the cytotoxic effect of PARP1 inhibitors against cancer cells is mostly mediated by the trapping of PARP1 at sites of DNA damage that generates stalled replication forks during the S phase of the cell cycle [11-13]. Therefore, novel PARP1 inhibitors with increased trapping capacity are promising candidates to target cancer cell death [14]. Of note, inhibition of PARP1 does not only have cancer cell-autonomous effects but also stimulates T lymphocyte-mediated anticancer immune response through yet-to-be-elucidated mechanisms [6, 12, 15]. Thus, combining PARP1 inhibition with immune checkpoint blockade holds promise for the treatment of ovarian cancer patients [16].

PRO-DEATH ACTIVITY OF PARP1 IN PARTHANATOS

Ted and Valina Dawson described a cell death modality relying on the overactivation of PARP1 and PAR-induced mitochondrial outer membrane permeabilization (MOMP) that they coined “parthanatos” [17]. In this lethal pathway, oxidative or nitrosative stress results in the activation of PARP1. PAR accumulating at mitochondrial membranes then triggers the release of the intermembrane protein apoptosis inducing factor (AIF, official gene/protein name AIFM1) that subsequently interacts with parthanatos-associated apoptosis-inducing factor (AIF) nuclease (PAAN), also known as macrophage migration inhibitor factor (MIF). PAAN is a member of the PD-D/E(X)K nuclease family and acts as a final executioner in parthanatos. The genetic depletion or pharmacological inhibition of PARP1 and PAAN prevents neurodegeneration of dopaminergic neurons induced by stereotactic intrastriatal injection of α -synuclein preformed fibrils, systemic administration of the Parkinsonian neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) or overexpression of the parkin substrate aminoacyl-tRNA synthetase complex interacting multifunctional protein-2 (AIMP2) [18-20]. Parthanatos has been involved in rodent models of ischemic stroke in which PARP1 inhibitors given within 4 to 6 hours after middle cerebral artery occlusion (MCAO) confer neuroprotection [21]. Other disease models in which PARP1 inhibitors have cytoprotective effects include Alzheimer's disease (AD), Huntington's disease (HD) and amyotrophic lateral sclerosis (ALS) [22], experimental retinal detachment [23], as well as ischemia/reperfusion injury of the kidney or the liver [24, 25]. Parthanatos has also been involved in human psoriasis in which PARP1 is overactivated downstream of the NAD⁺ generating enzyme nicotinamide phosphoribosyltransferase (NAMPT) and the translocation of AIF from mitochondria to nuclei can be observed in skin lesions [26]. As an aside, cancer cells can die in response to specific drugs or drug combinations in a PARP1-dependent fashion [27-29]. Altogether, it appears that, outside of the realm of oncology, PARP1 inhibition might be useful for the treatment of diseases, many of which are caused by the acute, massive or chronic, insidious loss of neurons.

PRO-APOPTOTIC CLEAVAGE AND INACTIVATION OF PARP1

Apoptosis usually involves two major hierarchically related events, namely (i) mitochondrial membrane permeabilization (MMP) and (ii) activation of specific set of proteases, which are called caspases. MMP may occur to variable proportions as mitochondrial inner membrane permeabilization (MIMP) and MOMP, leading to the arrest of oxidative phosphorylation and other essential metabolic functions, hence usually sealing the cell's fate [30, 31]. MOMP causes the release of proteins that are found in the intermembrane space or are loosely attached to the outer surface of the inner membrane into the cytosol. One prominent example is cytochrome *c* which, once released from mitochondria, interacts with APAF1 to stimulate the formation of the apoptosome, which is a caspase-9 activation

complex [32]. Caspase-9 is an initiator caspase that then activates other caspases such as caspase-3 and caspase-7 that are effector caspases and destroy multiple intracellular proteins, hence dismantling the cell from inside [31]. When caspases are inhibited, MMP causes cell death, though without the morphological appearance of apoptosis with nuclear pyknosis, rounding of the cellular contours, shrinkage of the cytoplasm and formation of apoptotic blebs. Rather, MMP without subsequent caspase activation results in necrotic cell death with cellular oncosis, organelle swelling and early plasma membrane rupture [33]. In this context, it is important to note that PARP1 is a prominent caspase-3/7 substrate and that caspase-digested PARP1 loses its enzymatic function. Thus, when cells die in the absence of caspase activation, PARP1 tends to become activated by damaged DNA, hence futilely consuming ATP and NAD⁺, which contributes to the lethal bioenergetic crisis that culminates in necrosis. In this context, the genetic or pharmacologic inhibition of PARP1 favors the apoptotic rather than necrotic demise of dying cells [34, 35]. Thus, in certain situations, the activity of PARP1 does not determine the death/life decision itself but rather affects the propensity to succumb to one or the other lethal subroutine: apoptosis or necrosis.

PRO-AGING AND ANTI-AGING EFFECTS OF PARP1

PARP1 is necessary for genomic stability, suggesting that its inhibition should derive genomic instability, which is one of the principal hallmarks of aging as well as of cancer. Indeed, transgenic PARP1 overexpression in basal skin keratinocytes from mice suppresses skin papilloma formation in a two-stage skin carcinogenesis protocol [36]. In *Drosophila*, conditional overexpression of PARP1 in the imago increases median lifespan of females and the maximum lifespan of males [37]. In mice, knockout of PARP1 accelerates aging and causes the precocious manifestation of spontaneous carcinogenesis, as well as a shift to a higher frequency of epithelial cancers of the lung, liver and uterus [38]. In patients with solid tumors, treatment with PARP1 inhibitors, results in an increased incidence of myelofibroblastic syndrome (which is a typical age-associated condition) and acute myeloid leukemia (which often develops from myelofibrosis) [39]. Possible mechanisms accounting for an age-related decline in PARP1 activity include a progressive depletion of the NAD⁺ [40]. This reduces PARP1 activity not only because NAD⁺ is an essential PARP1 substrate, but also due to the reduced binding of NAD⁺ to the protein deleted in breast cancer 1 (DBC1), which then engages in an inhibitory interaction with PARP1 [41]. Altogether, the aforementioned findings suggest that PARP1 has an anti-aging function.

In sharp contrast, there are also arguments in favor of a pro-aging activity of PARP1. Thus, pharmacological PARP1 inhibition rescues the short lifespan of hyperglycemic *Caenorhabditis elegans* [42], improves neurovascular and cognitive parameters in aging mice [43], and ameliorates cardiac performance in aging rats in which it also enhances acetylcholine-induced, nitric oxide-mediated vascular re-

laxation [44]. Conversely, mice expressing a human PARP1 transgene exhibit reduced healthspan and lifespan, accompanied by reduced hair growth and premature manifestation of inflammation and age-associated pathologies, such as anemia, adiposity, cardiomyopathy, dermatitis, hepatitis, kyphosis, nephropathy, pneumonitis, as well as an increase in the incidence of carcinomas [45]. The mechanisms accounting for these pro-aging effects of human PARP1 in mice are elusive. However, they plead for an ambiguous, dose-dependent implication of PARP1 in the aging process.

CONCLUSION

Pharmacological inhibition of PARP1 may kill those cancer cells that are undergoing constant endogenous DNA damage due to defects in DNA repair enzymes as well as cells that are being exposed to DNA damaging irradiation or cytotoxicants. Paradoxically, PARP1 inhibition has potent cell death-inhibitory effects in other contexts, in particular against parthanatos, a non-apoptotic modality of cellular demise that relies on the enzymatic overactivation of PARP1 and that appears to be particularly prevalent in neurological diseases. In addition, PARP1 inhibition may accelerate or retard age-related pathologies, suggesting that PARP1 has an ambiguous role in aging as well. This consideration is important in view of the possible side effects of long-term treatments with PARP1 inhibitors.

ACKNOWLEDGMENTS

GK is supported by the Ligue contre le Cancer (équipe labellisée); Agence National de la Recherche (ANR) – Projets blancs; AMMICA US23/CNRS UMS3655; Association pour la recherche sur le cancer (ARC); Association “Ruban Rose”; Cancéropôle Ile-de-France; Fondation pour la Recherche Médicale (FRM); a donation by Elior; Equipex Onco-Pheno-Screen; European

Joint Programme on Rare Diseases (EJPRD); Gustave Roussy Odyssey, the European Union Horizon 2020 Projects Oncobiome and Crimson; Fondation Carrefour; Institut National du Cancer (INCa); Inserm (HTE); Institut Universitaire de France; LabEx Immuno-Oncology (ANR-18-IDEX-0001); the Leducq Foundation; a Cancer Research ASPIRE Award from the Mark Foundation; the RHU Torino Lumière; Seerave Foundation; SIRIC Stratified Oncology Cell DNA Repair and Tumor Immune Elimination (SOCRATE); and SIRIC Cancer Research and Personalized Medicine (CARPEM). This study contributes to the IdEx Université de Paris ANR-18-IDEX-0001. AL was supported by a fellowship from Institut Thématique Multi-Organisme Cancer (ITMO Cancer) du Plan Cancer 2014-2019.

CONFLICT OF INTEREST

GK has been holding research contracts with Daiichi Sankyo, Eleor, Kaleido, Lytix Pharma, PharmaMar, Samsara, Sanofi, Sotio, Vascage and Vasculox/Tioma. GK is on the Board of Directors of the Bristol Myers Squibb Foundation France. GK is a scientific co-founder of everImmune, Osasuna Therapeutics, Samsara Therapeutics and Therafast Bio.

COPYRIGHT

© 2023 Castedo *et al.* This is an open-access article released under the terms of the Creative Commons Attribution (CC BY) license, which allows the unrestricted use, distribution, and reproduction in any medium, provided the original author and source are acknowledged.

Please cite this article as: Maria Castedo, Antoine Lafarge and Guido Kroemer (2023). Poly(ADP-ribose) polymerase-1 and its ambiguous role in cellular life and death. *Cell Stress* 7(1): 1-6. doi: 10.15698/cst2023.01.275

REFERENCES

- López-Otín C, Guido Kroemer (2021). Hallmarks of Health. *Cell* 184(1):33-63. doi: 10.1016/j.cell.2020.11.034
- Chaudhuri AR, Nussenzweig A (2017). The multifaceted roles of PARP1 in DNA repair and chromatin remodelling. *Nat Rev Mol Cell Biol* 18(10): 610–621. doi: 10.1038/nrm.2017.53
- Michels J, Vitale I, Galluzzi L, Adam J, Olaussen KA, Kepp O, Senovilla L, Talhaoui I, Guegan J, Enot DP, Talbot M, Robin A, Girard P, Oréar C, Lissa D, Sukkurwala AQ, Garcia P, Behnam-Motlagh P, Kohno K, Wu GS, Brenner C, Dessen P, Saparbaev M, Soria JC, Castedo M, Kroemer G (2013). Cisplatin resistance associated with PARP hyperactivation. *Cancer Res* 73(7):2271-2280. doi: 10.1158/0008-5472.CAN-12-3000
- Michels J, Adam J, Goubar A, Obrist F, Damotte D, Robin A, Alifano M, Vitale I, Olaussen KA, Girard P, Cremer I, Castedo M, Soria JC, Kroemer G (2015). Negative prognostic value of high levels of intracellular poly(ADP-ribose) in non-small cell lung cancer. *Ann Oncol* 26(12):2470–2477. doi: 10.1093/annonc/mdv393
- Joseph A, Juncheng P, Mondini M, Labaied N, Loi M, Adam J, Lafarge A, Astesana V, Obrist F, Klein C, Bloy N, Stoll G, Signolle N, Genestie C, Damotte D, Alifano M, Leary A, Pautier P, Morice P, Gouy S, Deutsch E, Chargari C, Dieu-Nosjean MC, Cremer I, Michels J, Kroemer G, Castedo M (2021). Metabolic features of cancer cells impact immu-

nosurveillance. *J Immunother Cancer* 9(6):e002362. doi: 10.1136/jitc-2021-002362

6. Juncheng P, Joseph A, Lafarge A, Martins I, Obrist F, Pol J, Saavedra E, Li S, Sauvat A, Cerrato G, Lévesque S, Leduc M, Kepp O, Durand S, Aprahamian F, Nirmalathansan N, Michels J, Kroemer G, Castedo M (2022). Cancer cell-autonomous overactivation of PARP1 compromises immunosurveillance in non-small cell lung cancer. *J Immunother Cancer* 10(6):e004280. doi: 10.1136/jitc-2021-004280

7. Gottipati P, Vischioni B, Schultz N, Solomons J, Bryant HE, Djureinovic T, Issaeva N, Sleeth K, Sharma RA, Helleday T (2010). Poly(ADP-ribose) polymerase is hyperactivated in homologous recombination-defective cells. *Cancer Res* 70(13):5389-98. doi: 10.1158/0008-5472.CAN-09-4716

8. Michels J, Vitale I, Senovilla L, Enot DP, Garcia P, Lissa D, Olaussen KA, Brenner C, Soria JC, Castedo M, Kroemer G (2013). Synergistic interaction between cisplatin and PARP inhibitors in non-small cell lung cancer. *Cell Cycle* 12(6):877-83. doi: 10.4161/cc.24034

9. Curtin NJ, Szabo C (2020). Poly(ADP-ribose) polymerase inhibition: past, present and future. *Nat Rev Drug Discov* 19(10):711-736. doi: 10.1038/s41573-020-0076-6.

10. Lord CJ, Tutt AN, Ashworth A (2015). Synthetic lethality and cancer therapy: lessons learned from the development of PARP inhibitors. *Annu Rev Med* 66:455-70. doi: 10.1146/annurev-med-050913-022545
11. Murai J, Huang SY, Das BB, Renaud A, Zhang Y, Doroshow JH, Ji J, Takeda S, Pommier Y (2012). Trapping of PARP1 and PARP2 by clinical PARP inhibitors. *Cancer Res* 72(21):5588-99. doi: 10.1158/0008-5472.CAN-12-2753
12. Chabanon RM, Muirhead G, Krastev DB, Adam J, Morel D, Garrido M, Lamb A, Hénon C, Dorvault N, Rouanne M, Marlow R, Bajrami I, Cardeñosa ML, Konde A, Besse B, Ashworth A, Pettitt SJ, Haider S, Marabelle A, Tutt AN, Soria JC, Lord CJ, Postel-Vinay S (2019). PARP inhibition enhances tumor cell-intrinsic immunity in ERCC1-deficient non-small cell lung cancer. *J Clin Invest* 129(3):1211-1228. doi: 10.1172/JCI123319
13. Kim C, Chen C, Yu Y (2021). Avoid the trap: Targeting PARP1 beyond human malignancy. *Cell Chem Biol* Apr 15;28(4):456-462. doi: 10.1016/j.chembiol.2021.02.004
14. Zandarashvili L, Langelier MF, Velagapudi UK, Hancock MA, Steffen JD, Billur R, Hannan ZM, Wicks AJ, Krastev DB, Pettitt SJ, Lord CJ, Talele TT, Pascal JM, Black BE (2020). Structural basis for allosteric PARP-1 retention on DNA breaks. *Science* 3;368(6486):eaax6367. doi: 10.1126/science.aax6367
15. Juncheng P, Lafarge A, Kroemer G, Castedo M (2022). PARP1 inhibition elicits immune responses against non-small cell lung cancer. *Oncoimmunology* 11(1):2111915. doi: 10.1080/2162402X.2022.2111915
16. Musacchio L, Cicala CM, Camarda F, Ghizzoni V, Giudice E, Carbone MV, Ricci C, Perri MT, Tronconi F, Gentile M, Salutari V, Scambia G, Lorusso D (2022). Combining PARP inhibition and immune checkpoint blockade in ovarian cancer patients: a new perspective on the horizon? *ESMO Open* 7(4):100536. doi: 10.1016/j.esmoop.2022.100536
17. Andrabi SA, Dawson TM, Dawson VL (2008). Mitochondrial and nuclear cross talk in cell death: parthanatos. *Ann N Y Acad Sci* 1147:233-41. doi: 10.1196/annals.1427.014
18. Kam TI, Mao X, Park H, Chou SC, Karuppagounder SS, Umanah GE, Yun SP, Brahmachari S, Panicker N, Chen R, Andrabi SA, Qi C, Poirier GG, Pletnikova O, Troncoso JC, Bekris LM, Leverenz JB, Pantelyat A, Ko HS, Rosenthal LS, Dawson TM, Dawson VL (2018). Poly(ADP-ribose) drives pathologic α -synuclein neurodegeneration in Parkinson's disease. *Science* 2;362(6414):eaat8407. doi: 10.1126/science.aat8407
19. Park H, Kam TI, Peng H, Chou SC, Mehrabani-Tabari AA, Song JJ, Yin X, Karuppagounder SS, Umanah GK, Rao AVS, Choi Y, Aggarwal A, Chang S, Kim H, Byun J, Liu JO, Dawson TM, Dawson VL (2022). PAAN/MIF nuclease inhibition prevents neurodegeneration in Parkinson's disease. *Cell* 185(11):1943-1959.e21. doi: 10.1016/j.cell.2022.04.020
20. Lee Y, Karuppagounder SS, Shin JH, Lee YI, Ko HS, Swing D, Jiang H, Kang SU, Lee BD, Kang HC, Kim D, Tessarollo L, Dawson VL, Dawson TM (2013). Parthanatos mediates AIMP2-activated age-dependent dopaminergic neuronal loss. *Nat Neurosci* 16(10):1392-400. doi: 10.1038/nn.3500.
21. Koehler RC, Dawson VL, Dawson TM (2021). Targeting parthanatos in ischemic stroke. *Front Neurol* 5;12:662034. doi: 10.3389/fneur.2021.662034
22. Park H, Kam TI, Dawson TM, Dawson VL (2020). Poly (ADP-ribose) (PAR)-dependent cell death in neurodegenerative diseases. *Int Rev Cell Mol Biol* 353:1-29. doi: 10.1016/bs.ircmb.2019.12.009
23. Dong K, Yan Y, Lu L, Wang Y, Li J, Zhang M, Ding J (2021). PJ34 protects photoreceptors from cell death by inhibiting PARP-1 induced parthanatos after experimental detachment. *Curr Eye Res* 46(1):115-121. doi: 10.1080/02713683.2020.1776881
24. Jang HR, Lee K, Jeon J, Kim JR, Lee JE, Kwon GY, Kim YG, Kim DJ, Ko JW, Huh W (2020). Poly (ADP-Ribose) polymerase inhibitor treatment as a novel therapy attenuating renal ischemia-reperfusion injury. *Front Immunol* 11:564288. doi: 10.3389/fimmu.2020.564288
25. Haga S, Kanno A, Morita N, Jin S, Matoba K, Ozawa T, Ozaki M (2022). Poly(ADP-ribose) polymerase (PARP) is critically involved in liver ischemia/reperfusion-injury. *J Surg Res* 270:124-138. doi: 10.1016/j.jss.2021.09.008
26. Martínez-Morcillo FJ, Cantón-Sandoval J, Martínez-Navarro FJ, Cabas I, Martínez-Vicente I, Armistead J, Hatzold J, López-Muñoz A, Martínez-Menchón T, Corbalán-Vélez R, Lacal J, Hammerschmidt M, García-Borrón JC, García-Ayala A, Cayuela ML, Pérez-Oliva AB, García-Moreno D, Mulero V (2021). NAMPT-derived NAD⁺ fuels PARP1 to promote skin inflammation through parthanatos cell death. *PLoS Biol* 19(11):e3001455. doi: 10.1371/journal.pbio.3001455
27. Sica V, Bravo-San Pedro JM, Izzo V, Pol J, Pierredon S, Enot D, Durand S, Bossut N, Chery A, Souquere S, Pierron G, Vartholomaiou A, Zamzami N, Soussi T, Sauvat A, Mondragón L, Kepp O, Galluzzi L, Martinou JC, Hess-Stumpp H, Ziegelbauer K, Kroemer G, Maiuri MC (2019). Lethal poisoning of cancer cells by respiratory chain inhibition plus dimethyl α -ketoglutarate. *Cell Rep* 27(3):820-834.e9. doi: 10.1016/j.celrep.2019.03.058
28. Ribeiro DL, Tuttis K, Oliveira LCB, Serpeloni JM, Gomes INF, Lengert AVH, Rocha CQD, Reis RM, Cólus IMS, Antunes LMG (2022). The antitumoral/antimetastatic action of the flavonoid bradydin A in metastatic prostate tumor spheroids in vitro is mediated by (parthanatos) PARP-related cell death. *Pharmaceutics* 14(5):963. doi: 10.3390/pharmaceutics14050963
29. Santofimia-Castaño P, Huang C, Liu X, Xia Y, Audebert S, Camoin L, Peng L, Lomberk G, Urrutia R, Soubeyran P, Neira JL, Iovanna J (2022). NUPR1 protects against hyperPARylation-dependent cell death. *Commun Biol* 5(1):732. doi: 10.1038/s42003-022-03705-1
30. Tang D, Kang R, Berghe TV, Vandenberghe P, Kroemer G (2019). The molecular machinery of regulated cell death. *Cell Res* 29(5):347-364. doi: 10.1038/s41422-019-0164-5
31. Kesavardhana S, Malireddi RKS, Kanneganti TD (2020). Caspases in cell death, inflammation, and pyroptosis. *Annu Rev Immunol* 38:567-595. doi: 10.1146/annurev-immunol-073119-095439
32. Zou H, Henzel WJ, Liu X, Lutschg A, Wang X (1997). Apaf-1, a human protein homologous to C. elegans CED-4, participates in cytochrome c-dependent activation of caspase-3. *Cell* 90(3):405-13. doi: 10.1016/s0092-8674(00)80501-2
33. Hirsch T, Marchetti P, Susin SA, Dallaporta B, Zamzami N, Marzo I, Geuskens M, Kroemer G (1997). The apoptosis-necrosis paradox. Apoptogenic proteases activated after mitochondrial permeability transition determine the mode of cell death. *Oncogene* 15(13):1573-81. doi: 10.1038/sj.onc.1201324
34. Herceg Z, Wang ZQ (1999). Failure of poly(ADP-ribose) polymerase cleavage by caspases leads to induction of necrosis and enhanced apoptosis. *Mol Cell Biol* 19(7):5124-33. doi: 10.1128/MCB.19.7.5124
35. Los M, Mozoluk M, Ferrari D, Stepczynska A, Stroch C, Renz A, Herceg Z, Wang ZQ, Schulze-Osthoff K (2002). Activation and caspase-mediated inhibition of PARP: a molecular switch between fibroblast necrosis and apoptosis in death receptor signaling. *Mol Biol Cell* 13(3):978-88. doi: 10.1091/mbc.01-05-0272
36. Bürkle A, Beneke S, Muir ML (2004). Poly(ADP-ribosyl)ation and aging. *Exp Gerontol* 39(11-12):1599-601. doi: 10.1016/j.exger.2004.07.010
37. Shaposhnikov MV, Moskalev AA, Plyusnina EN (2011). Effect of PARP-1 overexpression and pharmacological inhibition of NF- κ B on

the lifespan of *Drosophila melanogaster*. **Adv Gerontol** 24(3):405-19 PMID: 22184968

38. Piskunova TS, Yurova MN, Ovsyannikov AI, Semenchko AV, Zabezhinski MA, Popovich IG, Wang ZQ, Anisimov VN (2008). Deficiency in poly(ADP-ribose) polymerase-1 (PARP-1) accelerates aging and spontaneous carcinogenesis in mice. **Curr Gerontol Geriatr Res** 2008:754190. doi: 10.1155/2008/754190

39. Csizmar CM, Saliba AN, Swisher EM, Kaufmann SH (2021). PARP inhibitors and myeloid neoplasms: a double-edged sword. **Cancers** 13(24):6385. doi: 10.3390/cancers13246385

40. Abdellatif M, Sedej S, Kroemer G (2021). NAD⁺ metabolism in cardiac health, aging, and disease. **Circulation** 144(22):1795-1817. doi: 10.1161/CIRCULATIONAHA.121.056589

41. Li J, Bonkowski MS, Moniot S, Zhang D, Hubbard BP, Ling AJ, Rajman LA, Qin B, Lou Z, Gorbunova V, Aravind L, Steegborn C, Sinclair DA (2017). A conserved NAD⁺ binding pocket that regulates protein-protein interactions during aging. **Science** Mar 24;355(6331):1312-1317. doi: 10.1126/science.aad8242

42. Xia Q, Lu S, Ostrovsky J, McCormack SE, Falk MJ, Grant SFA (2018). PARP-1 inhibition rescues short lifespan in hyperglycemic *C. Elegans*

and improves GLP-1 secretion in human cells. **Aging Dis** 9(1):17-30. doi: 10.14336/AD.2017.0230

43. Tarantini S, Yabluchanskiy A, Csipo T, Fulop G, Kiss T, Balasubramanian P, DelFavero J, Ahire C, Ungvari A, Nyúl-Tóth Á, Farkas E, Benyo Z, Tóth A, Csiszar A, Ungvari Z (2019). Treatment with the poly(ADP-ribose) polymerase inhibitor PJ-34 improves cerebrovascular endothelial function, neurovascular coupling responses and cognitive performance in aged mice, supporting the NAD⁺ depletion hypothesis of neurovascular aging. **Geroscience** 41(5):533-542. doi: 10.1007/s11357-019-00101-2

44. Pacher P, Vaslin A, Benko R, Mabley JG, Liaudet L, Haskó G, Marton A, Bátkai S, Kollai M, Szabó C (2004). J A new, potent poly(ADP-ribose) polymerase inhibitor improves cardiac and vascular dysfunction associated with advanced aging. **Pharmacol Exp Ther** 311(2):485-91. doi: 10.1124/jpet.104.069658

45. Mangerich A, Herbach N, Hanf B, Fischbach A, Popp O, Moreno-Villanueva M, Bruns OT, Bürkle A (2010). Inflammatory and age-related pathologies in mice with ectopic expression of human PARP-1. **Mech Ageing Dev** 131(6):389-404. doi: 10.1016/j.mad.2010.05.005