



**HAL**  
open science

## Ionization potential depression and dense plasma collisional properties

Annette Calisti, Sandrine Ferri, Jean-Christophe Pain, Djamel Benredjem

► **To cite this version:**

Annette Calisti, Sandrine Ferri, Jean-Christophe Pain, Djamel Benredjem. Ionization potential depression and dense plasma collisional properties. 21st meeting on Atomic Processes in Plasmas, IAEA, May 2023, Vienne, Austria. pp.4. hal-04598477

**HAL Id: hal-04598477**

**<https://hal.science/hal-04598477>**

Submitted on 4 Jun 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **Ionization potential depression and dense plasma collisional properties.**

**Annette Calisti<sup>1</sup>, Sandrine Ferri<sup>2</sup>, Jean-Christophe Pain<sup>3</sup>, Djamel Benredjem<sup>4</sup>**

<sup>1</sup>*CNRS*

<sup>2</sup>*Aix Marseille University, CNRS, PIIM*

<sup>3</sup>*CEA, DAM, DIF*

<sup>4</sup>*Université Paris-Saclay, CNRS, Laboratoire Aimé Cotton*

The radiative properties of an atom or an ion surrounded by a plasma, are modified through various mechanisms. Depending on plasma conditions the electrons supposedly occupying the upper quantum levels of radiators no longer exist as they belong to the plasma free electron population. All the charges present in the radiator environment, electrons and ions, contribute to the lowering of the energy required to free an electron in the fundamental state. This mechanism is known as ionization potential depression (IPD). The knowledge of IPD is fundamental as it affects both the radiative properties of the various ionic states and their populations. Its evaluation deals with highly complex n-body coupled systems, involving particles with different dynamics and attractive ion-electron forces. Two distinct models, namely the Stewart–Pyatt (SP) [1] and Ecker–Kröll (EK) [2] models, are widely used to estimate the IPD.

More recently, an approach based on classical molecular dynamics simulation has been developed providing an alternative way to calculate the IPD [3]. Ions and electrons are treated as classical particles and a minimum of quantum properties are taken into account through a regularized potential allowing to model collisional ionization and recombination processes. The related numerical code, BinGo-TCP, has been designed to describe neutral mixtures composed of ions of the same atom with different charge states, and electrons. Within the limits of classical mechanics, all charge-charge interactions are accounted for in the particle motion.

In this work, after a brief reminder of the modeling basis, the importance of the choice of the IPD modelling will be emphasized through a study of the influence of the IPD on the magnitude of the cross-section of ionization by free-electron impacts in the high-density domain [4]. Additionally, we will discuss the ionization energy distributions obtained with BinGo-TCP due to the fluctuating environment of the ions.

## References

1. Stewart J C and Pyatt K D 1966, *Astrophys. J.* 144, 1203.
2. Ecker G and Kröll W 1963, *Phys. Fluids* 6, 62.
3. Calisti A, Ferri S and Talin B 2015, *J. Phys. B: At. Mol. Opt. Phys.* 48, 224003.
4. Benredjem D, Pain J.-C, Calisti A and Ferri S 2022, *J. Phys. B: At. Mol. Opt. Phys.* 55, 105001.

Presenting Author Email Address: *annette.calisti@univ-amu.fr*