



First record of the large-scale loach *Misgurnus dabryanus* (Guichenot, 1872) (Teleostei, Cobitidae) in France

Alain Cuinet, Gaël P.J. Denys, Sylvain Besson, Yves Marchiset, Julien Bouchard

► To cite this version:

Alain Cuinet, Gaël P.J. Denys, Sylvain Besson, Yves Marchiset, Julien Bouchard. First record of the large-scale loach *Misgurnus dabryanus* (Guichenot, 1872) (Teleostei, Cobitidae) in France. BioInvasions Records, 2024, 13 (2), pp.541 - 550. 10.3391/bir.2024.13.2.18 . hal-04600517

HAL Id: hal-04600517

<https://hal.science/hal-04600517v1>

Submitted on 10 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Rapid Communication

First record of the large-scale loach *Misgurnus dabryanus* (Guichenot, 1872) (Teleostei, Cobitidae) in France

Alain Cuinet¹, Gaël P.J. Denys^{2,3}, Sylvain Besson⁴, Yves Marchiset⁵ and Julien Bouchard⁴

¹Eaux Continentales, 29 rue Principale, 25440 Chay, France

²UAR Patrimoine Naturel – Centre d'expertise et de données (2006 OFB – CNRS – MNHN – IRD), Muséum national d'Histoire naturelle, 36 rue Geoffroy-Saint-Hilaire CP 41, 75005 Paris, France

³UMR Biologie des organismes et écosystèmes aquatiques (BOREA 8067), MNHN, CNRS, IRD, SU, UCN, UA, 57 rue Cuvier CP26, 75005 Paris, France

⁴Office Français de la Biodiversité, Direction Régionale Bourgogne / Franche-Comté, 57 rue de Mulhouse, 21000 Dijon, France

⁵Syndicat Mixte d'Aménagement de la Basse et Moyenne Vallée de l'Ognon, 8 Rue Fred Lipmann, 70190 Boulot, France

Corresponding author: Gaël P.J. Denys (gael.denys@mnhn.fr)

Citation: Cuinet A, Denys GPJ, Besson S, Marchiset Y, Bouchard J (2024) First record of the large-scale loach *Misgurnus dabryanus* (Guichenot, 1872) (Teleostei, Cobitidae) in France. *BioInvasions Records* 13(2): 541–550, <https://doi.org/10.3391/bir.2024.13.2.18>

Received: 22 September 2023

Accepted: 19 December 2023

Published: 21 April 2024

Handling editor: Tatenda Dalu

Thematic editor: Kenneth Hayes

Copyright: © Cuinet et al.

This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International - CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

OPEN ACCESS

Abstract

A specimen of weatherloach was caught in the Lanterne stream at Chevigney-sur-l’Ognon (Rhône catchment, Doubs department, eastern France) on the 22nd of July 2020. It was photographed and initially identified as European weatherloach *Misgurnus fossilis* before being released. Upon further examination of the photographs, looking at the adipose crests on the caudal peduncle, the coloration pattern and morphometric data of the head, the specimen was then correctly identified as a non-native large-scale loach *Misgurnus dabryanus*, thus being the first record of this species in France. Monitoring must be put in place to know if there is an established population and to follow its possible propagation.

Key words: weatherloach, Rhône drainage, invasive freshwater fish species, morphology

Introduction

The last decades have seen a steadily increasing number of introductions—intentional or not—in natural habitats of alien, and potentially invasive, freshwater fish species (Leprieur et al. 2008; Thomaz et al. 2015; Turbelin et al. 2017). One of the causes is the ornamental fish trade, which has allowed to disperse many exotic fish species all over the world (Padilla and Williams 2004; Strecker et al. 2011; Maceda-Veiga et al. 2013; Chan et al. 2019), thus having numerous impacts on the native aquatic fauna (Gozlan et al. 2010; Cucherousset and Olden 2011).

Asian weatherloaches *Misgurnus* spp. are among the most frequently introduced ornamental fish species in Europe (Kottelat and Freyhof 2007; Franch et al. 2008; Belle et al. 2017; Clavero et al. 2023). In addition to the native European weatherloach *Misgurnus fossilis* (Linnaeus, 1758), a species of international conservation concern, listed in the Annex II of the European Habitat Directive (Council of the European Union 1992), we see three

introduced Asian weatherloach species: (i) the dojo weatherloach *Misgurnus anguillicaudatus* (Cantor, 1842) in Italy (Razzetti et al. 2001; Geiger et al. 2014), Germany (Freyhof and Korte 2005), Spain (Franch et al. 2008; Aparicio et al. 2016; Clavero et al. 2023), England (Zięba et al. 2010), France (Chauvigné 2011; Denys and Manné 2024), and Netherlands (van Kessel et al. 2013); (ii) the large-scale loach *Misgurnus dabryanus* (Guichenot, 1872) – also known as *Paramisgurnus dabryanus* Guichenot in Dabry de Thiersant, 1872 before the taxonomic revision of Zhang H et al. (2021) – in Germany (Riffel et al. 1994; Stoeckle et al. 2019), Switzerland (Marchesi 2010), and Catalonia (Clavero et al. 2023); (iii) *Misgurnus bipartitus* (Sauvage & Dabry de Thiersant, 1874) in Austria (Zangl et al. 2020), and Catalonia (Clavero et al. 2023). The alien invasive weatherloaches may hybridize with the native species and therefore have an impact on its populations (Wanzenböck et al. 2021).

During intense field sampling of the Lanterne stream at Chevigney-sur-l’Ognon (Rhône catchment, Doubs department, eastern France, about 10 km NW from Besançon), one weatherloach specimen was caught, photographed, and initially identified on-site as *M. fossilis* before being released. Given the risk of misidentification, the photographs were sent to the Muséum national d’Histoire naturelle (Paris, MNHN) for expert assessment. Without any preserved material nor tissue for molecular analysis, the aim of this study is to (i) identify which *Misgurnus* species was caught in the Lanterne stream, and (ii) to report the first record of the large-scale loach *Misgurnus dabryanus* in France, identified from photographs. The potential impact of the species on the native fauna will be discussed as well as some advices for riverine managers about its monitoring.

Materials and methods

The Lanterne stream (12.5 km long) is a tributary of the Ognon River in the middle valley sector of eastern France (Figure 1). Its source is immediately upstream of Pouilley-les-Vignes; it then crosses the villages of Pelousey and Chaucenne before its confluence with the Ognon river on the left bank, between Emagny and Chevigney-sur-l’Ognon. Both water and habitat qualities are negatively impacted by multiple human-induced alterations to this stream: the watercourse has historically undergone a general recalibration of its bed, as well as more localized suppressions of sinuosities and riparian forest. Its current state is characterized by very high (2 m in some places) and steep banks, and an abnormally high wetted width. Thus the flows are highly homogenous and the waters pretty shallow, with little to contact with riparian vegetation. Furthermore, the original alluvial mattress has almost completely disappeared so, leaving the river bed almost naked, with some muddy deposits instead of pebbles and gravel. This stream presents essentially lentic flows, flats and pits, and ends upstream on the apron of an agricultural bridge.




Figure 1. Location (red square) of the first occurrence in France of the large-scale loach *Misgurnus dabryanus* in the Lanterne stream at Chevigney-sur-l’Ognon, the 22nd July 2020, and a picture of the habitat where the specimen was caught.

This causes a small localized acceleration, so a minor oxygenation zone. Despite those high banks, there is also locally a connective riparian forest composed of willows. A natural berm has formed downstream of the station, at Chevigney-sur-l’Ognon (latitude: 47.30256N; longitude: 5.85356E) and is the subject of a restoration program for the Lantern stream by the “Syndicat Mixte d’Aménagement de la Moyenne et Basse Vallée de l’Ognon” (SMAMBVO).

On the 22nd of July 2020, the French consulting engineer firm Eaux Continentales inventoried the Lanterne stream at Chevigney-sur-l’Ognon (Rhône catchment, Doubs department, eastern France) for the French Biodiversity Agency (OFB). Fish were caught by complete electrofishing with a “Héron” device from Dream Electronique, using two anodes, following the protocol of Belliard et al. (2012). All individuals were measured in total lengths and weighted before being released, and a specimen of each species was photographed.

The weatherloach was measured, weighted and sex determined: the first and second pectoral fin rays are enlarged and include a *lamina circularis* in males (vs. slim first and second pectoral fin rays and no *lamina circularis* in females) (Kim 2009). The specimen was identified from photographs using external traits as well as morphometric character of the head. The general examination was made following Kottelat and Freyhof (2007) (coloration pattern, adipose crests of the caudal peduncle) and comparison with the holotype specimen of *Paramisgurnus dabryanus* Guichenot in Dabry de




Figure 2. Holotype specimen of *Paramisgurnus dabryanus* Guichenot in Dabry de Thiersant 1872 (MNHN IC A-5087) (A); photo credit: L. Randrihasipa/MNHN. Asian weather loch identified as *Misgurnus dabryanus* (Guichenot, 1872) caught in the Lanterne stream at Chevigney-sur-l’Ognon the 22 of July 2020: caught specimen in lateral view (B), head in lateral view (C) and in dorsal view (D); scale bars represent 1 cm; white arrows indicate high adipose crests on the caudal peduncle, typical of the species; photos credit: A. Cuinet/Eaux Continentales.

Thiersant, 1872 (MNHN IC A-5087; Figure 2A) according to Bertin and Estève (1948); the morphometric measurements (head depth, head width, snout length, postorbital length, eye diameter, interorbital width, inner rostral barbel length, outer rostral barbel length and maxillary barbel length) were done using pixels numbers counted on photographs (Figure 2C–D) with the GIMP software (<https://www.gimp.org/>); their ratio with the pixels number of the head length were then compared with the data on male specimens from Zhang H et al. (2021) (Table 1).

Results

One male Asian weatherloach (Figure 2B–D) (25 cm TL, 102 g) was caught with 19 other fish and crayfish species, including native rheophilic species *Alburnoides bipunctatus* (Bloch, 1782), *Leuciscus leuciscus* (Linnaeus, 1758), *Squalius cephalus* (Linnaeus, 1758), *Gobio gobio* (Linnaeus, 1758), *Barbus barbus* (Linnaeus, 1758), *Barbatula* sp., *Cottus gobio* Linnaeus, 1758, some limnophylic species *Rutilus rutilus* (Linnaeus, 1758), *Tinca tinca* (Linnaeus, 1758) and *Perca fluviatilis* Linnaeus, 1758, as well as introduced species

Table 1. Morphometric measurements in percentage of the head length taken on photographs (Figure 2C and 2D; in pixel numbers) of the male Asian weatherloach caught in the Lanterne Stream at Chevigney-sur-l’Ognon the 22nd July 2020 compared to data on male specimens from Zhang H et al. (2021) on the three *Misgurnus* species introduced in Europe; * and \$ mean that measurements were made respectively on the Figure 2C and D; diagnostic characters are in bold.

	<i>Misgurnus</i> sp. from the Lanterne stream	<i>Misgurnus dabryanus</i>	<i>Misgurnus anguillicaudatus</i>	<i>Misgurnus bipartitus</i>
Head depth *	63.8	57.7–70.8	52.1–57.7	47.3–54.9
Head width \$	53.8	43.3–55.1	42.6–49.9	39.6–46.6
Snout length *	33.0	26.4–55.8	30.1–37.6	33.4–41.8
Postorbital length *	58.3	11.3–61.1	45–58.3	46.4–57.2
Eye diameter *	11.9	8.8–25.6	8.3–17.5	7.4–12.7
Interorbital width \$	25.8	21.2–55.4	14.6–23.3	16.2–21
Inner rostral barbel length \$	55.8	44–56.6	30.5–39.8	15.8–29.9
Outer rostral barbel length \$	44.6	44.6–55.4	34–38.5	17.1–30.4
Maxillary barbel length \$	51.7	44.5–62.8	30.5–44.9	17.9–30.6

Rhodeus amarus (Linnaeus, 1758), *Carassius* sp., *Alburnus alburnus* (Linnaeus, 1758), *Chondrostoma nasus* (Linnaeus, 1758), *Silurus glanis* Linnaeus, 1758, *Gymnocephalus cernua* (Linnaeus, 1758) and *Faxonius limosus* (Rafinesque, 1817) and three invasive alien species *Pseudorasbora parva* (Temminck & Schlegel, 1846), *Ameiurus melas* (Rafinesque, 1820) and *Lepomis gibbosus* (Linnaeus, 1758).

The specimen was encountered on a flat-type facies 20 cm deep, with a stony substrate including multi-centimeter clogging of silt and clay, downstream of a small stony pit sheltering sculpins *C. gobio* (Figure 1).

The following morphological data support its identification as *Misgurnus dabryanus*: no distinct coat coloration pattern (vs. bands for *M. fossilis*, dark spotted coat for *M. anguillicaudatus* or marbled coat for the other species), a caudal peduncle with high adipose crests (vs. low adipose crests for the other *Misgurnus* species), and no black spot at the upper caudal base (vs. presence of a black spot for *M. anguillicaudatus* and *M. bipartitus*) (Figure 2B). Morphometric data on the head (head depth, head width, interorbital with, inner rostral barbel length, outer rostral barbel length and maxillary barbel length) (Table 1, indicated in bold) also support this identification.

Discussion

The large-scale loach *Misgurnus dabryanus* is native to the Yangtze and Pearl drainages (central and southern China) as well as inland waters of Taiwan (Kottelat 2001; Shedko and Vasil’eva 2022). However, it was also introduced in other Asian countries such as Japan (Kanou et al. 2007) and Russia (Dyldin et al. 2023), as well as California (Kirsch et al. 2018) and in Europe: Germany (Riffel et al. 1994; Stoeckle et al. 2019), Switzerland (Marchesi 2010), and Catalonia (Clavero et al. 2023). This species will be integrated into the French taxonomic register TAXREF (Gargominy et al. 2022) of the National Inventory of Natural Heritage (*Inventaire du Patrimoine Naturel* – INPN). This record is then the fourth in Europe and the second Asian weatherloach species encountered in France, following the earlier

records of the dojo weatherloach *M. anguillicaudatus* (Chauvigné 2011; Denys and Manné 2024).

This specimen was morphologically identified from photographs, using the following criteria: coloration pattern and the presence of high adipose crests on the caudal peduncle (Marchesi 2010; Kirsch et al. 2018). We demonstrated that morphometric data on the head (head depth and width, interorbital width, barbel lengths) were also diagnostic (see Zhang H et al. 2021). The use of the comparison with the holotype specimen of *Paramisgurnus dabryanus* Guichenot in Dabry de Thiersant, 1872 in the expertise also supports our morphometric identification (e.g., Sluys 2021). Unfortunately, the specimen was released and we were not able to support our species identification with molecular data from other authors (e.g., Riffel et al. 1994; Kirsch et al. 2018; Stoeckle et al. 2019; Clavero et al. 2023). The additional use of a nuclear marker could have also flagged a potential hybrid with *M. anguillicaudatus* (Belle et al. 2021). Nevertheless, morphological identification is still feasible, as the morphological diagnostic characters (morphometry, coloration pattern, and the adipose crests on the caudal peduncle) are easy to observe, even for inexperienced riverine managers, if appropriately trained. This should prevent future confusion with the native and endangered European weatherloach *M. fossilis*, a misidentification that could adversely impact the riverine management (Bortolus 2008).

Knowledge about this species' potential invasiveness and its impact on the native European ichthyofauna is scarce. Some of its biological features are known, though: it lives in lentic and lotic environments, with muddy substrate with low gradient, slack water and high turbidity (Gao et al. 2010; Sato et al. 2011; Huang et al. 2013); it can adapt to hypoxic environments, breathing through its skin and intestines (Zhang et al. 2012; Wang et al. 2020; Liu et al. 2021) that have anatomical and physiological adaptations for air-breathing (Liu and Wang 2021); it tolerates a high range of water temperature (10 °C to 30 °C), burrowing into the sediment at a depth of 20–30 cm if the temperatures were to be too high or too low (Wang and Li 2005; Zhang et al. 2012); it also has a rapid growth, reaching its maximal length of about 30 cm in a few years (Wang et al. 2001; Gao et al. 2017). Moreover, considering its high reproductive potential (Chu et al. 2012), and its large range of diet, from zooplankton to macroinvertebrates and algae (Wang and Li 2005; Kanou et al. 2007), *M. dabryanus* fulfils all criteria to be considered an invasive species (e.g., Keller et al. 2011). Furthermore, this species is able to reproduce by asexual gynogenesis (You et al. 2008), which may accentuate its invasive potential (Fuad et al. 2021). In addition to a possible risk of competition with native benthic fish species (e.g., Mills et al. 2004), *M. dabryanus* may have a sanitary impact, for it is the intermediate host of nematodes *Contracaecum* spp. that might infect native fish species, as well as piscivorous birds and humans (Zhang C et al. 2021). However, unlike the Bavarian

record (Stoeckle et al. 2019), there would be no threat to the population of *M. fossilis*. Its closest known population is located more than 100 km aerial away in another catchment system, the Savoureuse stream, belonging to the Doubs sub-drainage. However, *M. fossilis* is not native in the Rhône drainage and its presence is certainly due to accidental introduction with restocking events (Keith et al. 2020). The two occurrences of *M. anguillicaudatus* known in France are located in the Orge river (Seine drainage), in the south suburb of Paris at Sainte-Geneviève-des-Bois in 2010 with only one fish, and an established population in the Schadgraben stream at Geispolsheim, near Strasbourg (Rhine catchment, eastern France) in autumn 2021 (Chauvigné 2011; Denys and Manné 2024). Their presences—and that of our large-scale weatherloach in the Lantern stream—are probably due to aquarium releases (Chan et al. 2019). However, the presence of a lonely specimen in an isolated location from any urban environment lets us to suggest potential other specimens in a widespread area in the catchment.

Surveys must be set up in this region to assess the situation, especially to verify if our record indicates the presence of an established population, or if it was a lonely individual. Thus, regular electrofishing operations must be undertaken, as well as molecular sampling, like measuring eDNA (Beng and Corlett 2020; Feng et al. 2023), but see Jerde (2020) about the risk of results misinterpretation. Furthermore, initiatives at the national scale must also be initiated to detect eventual cryptic populations of the large scale-loach in the French hydrosystems.

Authors' contribution

AC conceptualized research, designed sample and methodology, investigated and collected data. Data analysis and interpretation were performed by GPJD, SB and JB. YM provided funding. AC and GPJD wrote the original draft. All authors reviewed and edited the manuscript.

Acknowledgements

This study is supported by the MNHN, the UAR PatriNat 2006, the UMR BOREA 8067 and the OFB. We thank M. Hauteceur for checking the English. We thank the two anonymous reviewers for their comments to improve our manuscript.

Funding declaration

This study was funded by the “Syndicat Mixte d’Aménagement de la Basse et Moyenne Vallée de l’Ognon” (SMAMBVO).

Ethics and permits

Electrofishing permit was provided according to the prefectoral decree of the Doubs department (DDT25-ERNF-2020-06-18-003). No ethics agreement was required.

References

- Aparicio E, Alcaraz C, Carmona-Catot G, García-Berthou E, Pou-Rovira Q, Rocaspana R, Vargas MJ, Vinyoles D (2016) Peixos continentals de Catalunya. Ecologia, conservació i guia d’identificació. Lynx Edicions, Barcelona, 251 pp [in Catalan]

- Belle CC, Stoeckle BC, Cerwenka AF, Kuehn R, Mueller M, Pander J, Geist J (2017) Genetic species identification in weatherfish and first molecular confirmation of Oriental Weatherfish *Misgurnus anguillicaudatus* (Cantor, 1842) in Central Europe. *Knowledge and Management of Aquatic Ecosystems* 418: 31, <http://doi.org/10.1051/kmae/2017025>
- Belle CC, Stoeckle BC, Cerwenka AF, Kuehn R, Pander J, Geist JP (2021) Taxonomic requirements for better documenting and understanding biological invasions – the example of genetic weather fish *Misgurnus/Paramisgurnus* sp. identification. *BioInvasions Records* 10: 506–520, <http://doi.org/10.3391/bir.2021.10.3.01>
- Belliard J, Ditché JM, Roset N, Dembski S (2012) Guide pratique de mise en œuvre des opérations de pêche à l'électricité dans le cadre des réseaux de suivi des peuplements de poissons. ONEMA report, 31 pp [In French]
- Beng KC, Corlett RT (2020) Applications of environmental DNA (eDNA) in ecology and conservation: opportunities, challenges and prospects. *Biodiversity and Conservation* 29: 2089–2121, <https://doi.org/10.1007/s10531-020-01980-0>
- Bertin L, Estève R (1948) Catalogue des types de poissons du Muséum National d'Histoire Naturelle, 4e partie, Ostariophysaires (Cypriniformes). Imprimerie Nationale, Paris, 117 pp [In French]
- Bortolus A (2008) Error cascades in the biological sciences: the unwanted consequences of using bad taxonomy in ecology. *AMBIO A Journal of the Human Environment* 37: 114–118, [http://doi.org/10.1579/0044-7447\(2008\)37\[114:ECITBS\]2.0.CO;2](http://doi.org/10.1579/0044-7447(2008)37[114:ECITBS]2.0.CO;2)
- Chan F, Beatty S, Gilles A, Hill J, Kozic S, Luo D, Morgan D, Pavia R, Therriault T, Verreycken H, Vilizzi L, Wei H, Yeo D, Zeng Y, Zięba G, Copp G (2019) Leaving the fish bowl: the ornamental trade as a global vector for freshwater fish invasions. *Aquatic Ecosystems Health and Management* 22: 417–439, <http://doi.org/10.1080/14634988.2019.1685849>
- Chauvigné S (2011) Recherche d'une population de *Misgurnus anguillicaudatus* sur la partie aval de la rivière Orge (91). ONEMA report, 18 pp [in French]
- Chu Z, Lu G, Hu T, Wang H, Dai L, Huang W (2012) Comparative analysis on fecundity of *Misgurnus anguillicaudatus* and *Paramisgurnus dabryanus*. *Hubei Agricultural Sciences* 13: 2794–2796
- Clavero M, Suh J, Franch N, Aparicio E, Buchaca T, Caner J, Garcia-Rodriguez S, Llimona F, Pou-Rovira Q, Rocaspana R, Ventura M (2023) Invaders they are a-changing: A recent, unexpected surge of invasive loaches in Catalonia. *Freshwater Biology* 68: 621–631, <http://doi.org/10.1111/fwb.14051>
- Council of the European Union (1992) Directive 92/42/EEC of the European Parliament and the Council of 21st May 1992 on the conservation of natural habitats and of wild fauna and flora. Official Journal of the European Communities L206, 750 pp
- Cucherousset J, Olden JD (2011) Ecological impacts of non-native freshwater fishes. *Fisheries* 36: 215–230, <https://doi.org/10.1080/03632415.2011.574578>
- Denys GPJ, Manné S (2024) First record of an established population of *Misgurnus anguillicaudatus* (Cantor, 1842) (Teleostei, Cobitidae) in France. *BioInvasions Records* 13 (in press)
- Dyldin YV, Orlov AM, Hanel L, Romanov VI, Fricke R, Vasil'eva ED (2023) Ichthyofauna of the Fresh and Brackish Waters of Russia and Adjacent Areas: Annotated List with Taxonomic Comments. 2. Order Cypriniformes, Suborders Catostomoidei, Cobitoidei and Cyprinoidei. *Journal of Ichthyology*, <http://doi.org/10.1134/S0032945223040045>
- Feng X, Bing L, Chen Y, Zhu R, Jia Y, Sui X (2023) Species-level monitoring of rare and invasive fishes using eDNA metabarcoding in the middle and upper Yarlung Zangbo River, Tibet. *Water Biology and Security* 2: 100089, <https://doi.org/10.1016/j.watbs.2022.100089>
- Franch N, Clavero M, Garrido M, Gaya N, López V, Pou-Rovira Q, Queral JM (2008) On the establishment and range expansion of oriental weatherfish (*Misgurnus anguillicaudatus*) in NE Iberian Peninsula. *Biological Invasions* 10: 1327–1331, <http://doi.org/10.1007/s10530-007-9207-9>
- Freyhof J, Korte E (2005) The first record of *Misgurnus anguillicaudatus* in Germany. *Journal of Fish Biology* 66: 568–571, <http://doi.org/10.1111/j.0022-1112.2005.00606.x>
- Fuad MH, Vetešník L, Šimková A (2021) Is gynogenetic reproduction in gibel carp (*Carassius gibelio*) a major trait responsible for invasiveness? *Journal of Vertebrate Biology* 70: 21049.1–13, <https://doi.org/10.25225/jvb.21049>
- Gao X, Zeng Y, Wang J, Liu H (2010) Immediate impacts of the second impoundment on fish communities in the Three Gorges Reservoir. *Environmental Biology of Fishes* 87: 163–173, <http://doi.org/10.1007/s10641-009-9577-1>
- Gao Y, He J, He Z, Li Z, Zhao B, Mu Y, Lee JY, Chu Z (2017) Effects of fulvic acid on growth performance and intestinal health of juvenile loach *Paramisgurnus dabryanus* (Sauvage). *Fish and Shellfish Immunology* 62: 47–56, <http://doi.org/10.1016/j.fsi.2017.01.008>
- Geiger MF, Herder F, Monaghan MT, Almada V, Barbieri R, Bariche M, Berrebi P, Bohlen J, Casal-Lopez M, Delmastro GB, Denys GPJ, Dettai A, Doadrio I, Kalogianni E, Kärst H, Kottelat M, Kovačić M, Laporte M, Lorenzoni M, Marčić Z, Özuluğ M, Perdices A, Perea S, Persat H, Porcelotti S, Puzzi C, Robalo J, Šanda R, Schneider M, Šlechtová V, Stoumboudi M, Walter S, Freyhof J (2014) Spatial heterogeneity in the Mediterranean

- biodiversity hotspot affects barcoding accuracy of its freshwater fishes. *Molecular Ecology Resources* 14: 1210–1221, <http://doi.org/10.1111/1755-0998.12257>
- Gozlan RE, Britton JR, Cowx I, Copp GH (2010) Current knowledge on non-native freshwater fish introduction. *Journal of Fish Biology* 76: 751–786, <http://doi.org/10.1111/j.1095-8649.2010.02566.x>
- Huang LL, Wu ZQ, Li JH (2013) Fish fauna, biogeography and conservation of freshwater fish in Poyang Lake basin, China. *Environmental Biology of Fishes* 96: 1229–1243, <http://doi.org/10.1007/s10641-011-9806-2>
- Jerde CL (2020) Can we manage fisheries with the inherent uncertainty from eDNA? *Journal of Fish Biology* 98: 341–353, <http://doi.org/10.1111/jfb.14218>
- Kanou K, Saito S, Fuchigami S, Imamura A, Imai H, Taki Y (2007) Occurrence patterns and food habits of introduced alien loach *Paramisgurnus dabryanus* and native loach *Misgurnus anguillicaudatus* at irrigation drainages around rice fields in the Watarase river system, central Honshu, Japan. *Aquaculture Science* 55: 109–114, <https://doi.org/10.11233/aquaculturesci1953.55.109>
- Keith P, Poulet N, Denys G, Changeux T, Persat H (2020) Les poissons d'eau douce de France, second edition. Biotope, Publications scientifiques du Muséum, Mèze and Paris, 704 pp [in French]
- Keller R, Geist J, Jeschke J, Kühn I (2011) Invasive species in Europe: ecology, status, and policy. *Environmental Sciences Europe* 23: 23, <http://doi.org/10.1186/2190-4715-23-23>
- Kim I-S (2009) A Review of the Spined Loaches, Family Cobitidae (Cypriniformes) in Korea. *Korean Journal of Ichthyology* 21: 7–28
- Kirsch JE, Feeney RF, Goodbla A, Hart C, Jackson ZJ, Schreier A, Smith R (2018) First record of the large-scale loach *Paramisgurnus dabryanus* (Cobitidae) in the United States. *Journal of Fish and Wildlife Management* 9: 1–9, <http://doi.org/10.3996/012017-JFWM-008>
- Kottelat M (2001) Freshwater fishes of Northern Vietnam: a preliminary check-list of the fishes known or expected to occur in Northern Vietnam with comments on systematics and nomenclature. World Bank, Washington, 123 pp
- Kottelat M, Freyhof J (2007) Handbook of European Freshwater Fishes. Publication Kottelat, Cornell, 646 pp
- Leprieur F, Beauchard O, Blanchet S, Oberdorff T, Brosse S (2008) Fish invasions in the world's river systems: when natural processes are blurred by human activities. *PLoS Biology* 6: e28, <http://doi.org/10.1371/journal.pbio.0060028>
- Liu Y, Wang ZJ (2021) Effect of hypoxia and air-breathing restricted on respiratory physiology of air-breathing loach (*Paramisgurnus dabryanus*). *Fish Physiology and Biochemistry* 47: 251–263, <https://doi.org/10.1007/s10695-020-00908-3>
- Liu Y, Wang ZJ, Liu X (2021) Effect of intestinal air-breathing restriction on respiratory metabolism and antioxidant capability of loach (*Paramisgurnus dabryanus*). *Journal of Zoology* 52: 857–864, <http://doi.org/10.13859/j.cjz.201705015>
- Maceda-Veiga A, Escribano-Alacid J, de Sostoa A, García-Berthou E (2013) The aquarium trade as a potential source of fish introductions in southwestern Europe. *Biological Invasions* 15: 2707–2716, <https://doi.org/10.1007/s10530-013-0485-0>
- Marchesi P (2010) Histoires de loches et notes sur d'autres poissons non indigènes introduits en Valais. *Bulletin de la Murithienne* 128: 73–77 [in French]
- Mills MD, Rader RB, Belk MC (2004) Complex interactions between native and invasive fish: the simultaneous effects of multiple interactions. *Oecologia* 141: 713–721, <http://doi.org/10.1007/s00442-004-1695-z>
- Padilla DK, Williams SL (2004) Beyond ballast water: aquarium and ornamental trades as sources of invasive species in aquatic ecosystems. *Frontiers in Ecology and the Environment* 2: 131–138, [https://doi.org/10.1890/1540-9295\(2004\)002\[0131:BBWAAO\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2004)002[0131:BBWAAO]2.0.CO;2)
- Razzetti E, Nardi PA, Strosselli S, Bernini F (2001) Prima segnalazione di *Misgurnus anguillicaudatus* (Cantor, 1842) in acque interne italiane (Osteichthyes: Cobitidae). *Annali del Museo Civico di Storia Naturale di Genova* 93: 559–563
- Riffel M, Schenk M, Schreiber A (1994) Electrophoretic differentiation between European loach (*Misgurnus fossilis* L.) and oriental weatherfish (*Misgurnus mizolepis* Günther), an autochthonous and a feral species of central European freshwater fish. *Zeitschrift für angewandte Zoologie* 80: 473–483
- Sato T, Nakajima J, Huang L, Shimatani Y, Hirota SK, Wood C, Kano Y (2011) Distribution pattern of loaches (Teleostei: Cobitoidea) in the River East Tiaoxi, China. *Folia Zoologica* 60: 328–334, <http://doi.org/10.25225/fozo.v60.i4.a10.2011>
- Shedko SV, Vasil'eva ED (2022) A new species of the pond loaches *Misgurnus* (Cobitidae) from the south of Sakhalin Island. *Journal of Ichthyology* 62: 356–372, <https://doi.org/10.1134/S0032945222030158>
- Sluys R (2021) Attaching names to biological Species: the use and value of type specimens in systematic zoology and natural history collections. *Biological Theory* 16: 49–61, <https://doi.org/10.1007/s13752-020-00366-3>
- Stoeckle BC, Belle CC, Geist J, Oehm J, Effenberger M, Heiss M, Seifert K, Kuehn R (2019) Molecular confirmation of the large-scale loach *Paramisgurnus dabryanus* Dabry de Thiersant,

- 1872 (Cypriniformes, Cobitidae) in Europe. *Bioinvasions Records* 8: 419–426, <http://doi.org/10.3391/bir.2019.8.2.24>
- Strecker AL, Campbell PM, Olsen JD (2011) The aquarium trade as an invasion pathway in the Pacific Northwest. *Fisheries* 36: 74–85, <https://doi.org/10.1577/03632415.2011.10389070>
- Thomaz SM, Kovalenko KE, Havel JE, Kats LB (2015) Aquatic invasive species: general trends in the literature and introduction to the special issue. *Hydrobiologia* 756: 1–12, <http://doi.org/10.1007/s10750-014-2150-8>
- Turbelin AJ, Malamud BD, Francis RA (2017) Mapping the global state of invasive alien species: patterns of invasion and policy responses. *Global Ecology and Biogeography* 26: 78–92, <https://doi.org/10.1111/geb.12517>
- Van Kessel N, Dorenbosch M, Crombaghs B, Niemeijer B, Binnendijk E (2013) First record of Asian weather loach *Misgurnus anguillicaudatus* (Cantor, 1842) in the River Meuse basin. *BioInvasions Records* 2: 167–171, <http://doi.org/10.3391/bir.2013.2.2.14>
- Wang M, Wang WM, Yan JL (2001) Comparative studies on the age and growth of *Misgurnus anguillicaudatus* and *Paramisgurnus dabryanus*. *Reservoir Fisheries* 21: 7–9 [in Chinese]
- Wang W, Lin L, Qin Z, Sarah BV, Zhou M, Liang R, Mo J, Wu Z, Zhang X, Zhao L (2020) Histopathology of the main respiratory organs of *Paramisgurnus dabryanus* inhibiting intestinal air respiration. *Journal of Fisheries of China* 44: 845–857
- Wang YJ, Li D (2005) Research on biological characters and exploitations of *Paramisgurnus dabryanus* Sauvage. *Special Wild Economic Animal and Plant Research* 1: 60–62
- Wanzenböck J, Hopfinger M, Wanzenböck S, Fuxjäger L, Rund H, Lamatsch DK (2021) First successful hybridization experiment between native European weatherfish (*Misgurnus fossilis*) and non-native Oriental weatherfish (*M. anguillicaudatus*) reveals no evidence for postzygotic barriers. *Neobiota* 69: 29–50, <http://doi.org/10.3897/neobiota.69.67708>
- You C, Yu X, Tan D, Tong J (2008) Gynogenesis and sex determination in large-scale loach *Paramisgurnus dabryanus* (Sauvage). *Aquaculture International* 16: 203–214, <https://doi.org/10.1007/s10499-007-9138-6>
- Zangl L, Jung M, Gessl W, Koblmüller S, Ratschan C (2020) Oriental or not: First record of an alien weatherfish (*Misgurnus*) species in Austria verified by molecular data. *BioInvasions Records* 9: 375–383, <http://doi.org/10.3391/bir.2020.9.2.23>
- Zhang C, Wang Z, Zhang B, Yang H, Gu Z (2021) An investigation of zoonotic nematodes of loach *Paramisgurnus dabryanus* Dabry de Thiersant, under high-density pond farming pattern in China. *Aquaculture* 340: 736666, <https://doi.org/10.1016/j.aquaculture.2021.736666>
- Zhang H, Wang Y-X, Yang H-L, Tan H-M, Chen Y-X (2021) Taxonomic revision of Chinese species of the genera *Misgurnus* and *Paramisgurnus* (Cypriniformes: Cobitidae). *Acta Hydrobiologica Sinica* 45: 414–427, <https://doi.org/10.7541/2021.2019.166> [in Chinese]
- Zhang Y, Wang H, Qin F, Liu S, Wu T, Li M, Xu P, Zhang X, Wang X, Hu G, Yuan C, Wang Z (2012) Molecular characterization of estrogen receptor genes in loach *Paramisgurnus dabryanus* and their expression upon 17a-ethynodiol exposure in juveniles. *General and Comparative Endocrinology* 178: 194–205, <http://doi.org/10.1016/j.ygcen.2012.06.004>
- Zięba G, Copp GH, Davies GD, Stebbing P, Wesley KJ, Britton JR (2010) Recent releases and dispersal of non-native fishes in England and Wales, with emphasis on sunbleak *Leucaspis delineatus* (Heckel, 1843). *Aquatic Invasions* 5: 155–161, <http://doi.org/10.3391/ai.2010.5.2.04>

Web sites and online databases

Gargominy O, Tercerie S, Régnier C, Dupont P, Daszkiewicz P, Antonetti P, Léotard G, Ramage T, Idczak L, Vandel E, Petitteville M, Leblond S, Boullet V, Denys G, De Massary JC, Dusoulier F, Lévéque A, Jourdan H, Touroult J, Rome Q, Le Divelec R, Simian G, Savouré-Soubelet A, Page N, Barbut J, Canard A, Haffner P, Meyer C, Van Es J, Poncet R, Demerges D, Mehran B, Horellou A, Ah-Peng C, Bernard J-F, Bounias-Delacour A, Caesar M, Comolet-Tirman J, Courtecuisse R, Delfosse E, Dewynter M, Hugonnot V, Lavocat Bernard E, Lebouvier M, Lebreton E, Malécot V, Moreau PA, Moulin N, Muller S, Noblecourt T, Noël P, Pellens R, Thouvenot L, Tison JM, Robbert Gradstein S, Rodrigues C, Rouhan G, Véron S (2022) TAXREF v16.0, référentiel taxonomique pour la France. PatriNat (OFB-CNRS-MNHN), Muséum national d'Histoire naturelle, Paris. <https://inpn.mnhn.fr/telechargement/referentielEspece/taxref/16.0/menu> (accessed 11 September 2023)