



HAL
open science

Sugar modified oligonucleotides : II. Solid phase synthesis of nuclease resistant α -anomeric uridylates as potential antisense agents

Francoise Debart, Bernard Rayner, Jeon-Louis Imbach

► To cite this version:

Francoise Debart, Bernard Rayner, Jeon-Louis Imbach. Sugar modified oligonucleotides : II. Solid phase synthesis of nuclease resistant α -anomeric uridylates as potential antisense agents. *Tetrahedron Letters*, 1990, 31 (25), pp.3537-3540. 10.1016/S0040-4039(00)94436-8 . hal-04600829

HAL Id: hal-04600829

<https://hal.science/hal-04600829v1>

Submitted on 4 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**SUGAR MODIFIED OLIGONUCLEOTIDES : II. SOLID PHASE SYNTHESIS OF NUCLEASE
RESISTANT α -ANOMERIC URIDYLATES AS POTENTIAL ANTISENSE AGENTS**

Françoise DEBART, Bernard RAYNER and Jean-Louis IMBACH*

Université de MONTPELLIER II Sciences et Techniques du Languedoc, Laboratoire
de Chimie bio-organique associé au CNRS, Place Eugène Bataillon, 34095
MONTPELLIER Cedex 5, FRANCE.

Abstract : α -anomeric uridylates, α -rU₆ and α -rU₁₂, have been synthesized for the first time. These non natural α -oligoribonucleotides strongly resist to enzymatic degradation and bind to complementary RNA strands.

During the last few years, exogenous oligonucleotides binding specifically to complementary sequences of nucleic acids (RNA or DNA) through base pairing have been widely used *in vitro* as artificial regulators for gene expression¹⁻³. Conceivable chemotherapeutic applications predicated on sequence specific hybridization require antisense oligonucleotides that are resistant to *in vivo* degradation by nucleases and strongly bind to their target mRNA. Thus a new class of nuclease resistant oligodeoxynucleotides consisting exclusively of α -anomeric nucleotide units has been recently developed in our laboratory⁴. Furthermore it is generally believed that the order of increasing stability of oligonucleotide.polynucleotide complexes is DNA.DNA < DNA.RNA < RNA. RNA³. So, we anticipated that hitherto unknown α -anomeric oligoribonucleotides would combine nuclease resistance and improved binding capacity to their complementary strand.

In this communication, we describe for the first time the synthesis on a solid support of α -hexauridylate (α -rU₆) and α -dodecauridylate (α -rU₁₂) and report our results related to their stability towards exo- and endonucleases and their binding properties.

α -uridine 1 was prepared according to an already published procedure⁵. The 5'- and 2'-hydroxy functions on the ribose moiety were respectively protected by the 4,4'-dimethoxytrityl group (DMTr)⁶ and the tert-butyl-dimethylsilyl group (TBDMS)⁷ (Figure 1). For the selective monosilylation of ribonucleosides, we have applied the procedure described by Ogilvie involving nitrate ion catalysed action of 1.1 molar equivalent of tert-butyldimethylsilyl chloride (TBDMS-Cl) in THF⁸. In contrast to the results observed in the β series, the TBDMS group was shown to be selectively introduced at the 3' position of the 5'-dimethoxytrityl- α -uridine 2. When the reaction was carried out in pyridine in the presence of imidazole⁹, a mixture of 2'-TBDMS

and 3'-TBDMS derivatives **3** and **4** was obtained in nearly equal amount. These two isomers were isolated by silica gel column chromatography and characterized by high field $^1\text{H-NMR}$ spectroscopic decoupling experiments¹⁰.




Figure 1

- 1 $R_1 = R_2 = R_3 = \text{H}$
- 2 $R_1 = R_2 = \text{H}, R_3 = \text{DMTr}$
- 3 $R_1 = \text{TBDMS}, R_2 = \text{H}, R_3 = \text{DMTr}$
- 4 $R_1 = \text{H}, R_2 = \text{TBDMS}, R_3 = \text{DMTr}$
- 5 $R_1 = \text{TBDMS}, R_2 = \text{P}(\text{OCH}_3)\text{N}(\text{iPr})_2, R_3 = \text{DMTr}$
- 6 $R_1 = \text{P}(\text{OCH}_3)\text{N}(\text{iPr})_2, R_2 = \text{TBDMS}, R_3 = \text{DMTr}$
- 7 $R_1 = \text{C}(\text{O})(\text{CH}_2)_2\text{C}(\text{O})\text{O}(\text{C}_6\text{Cl}_5), R_2 = \text{TBDMS}, R_3 = \text{DMTr}$
- 8 $R_1 = \text{LCA-CPG succinyl}, R_2 = \text{TBDMS}, R_3 = \text{DMTr}$

Synthons **3** and **4** were then separately phosphitylated with methyl *N,N*-diisopropylchlorophosphoramidite¹¹ and the reaction products purified by chromatography on silica gel to give the corresponding phosphoramidites **5** and **6** respectively. The $^{31}\text{P-NMR}$ spectrum of **5** clearly showed only two signals at 152.49ppm and 151.31ppm different from those exhibited by **6** and corresponding to the expected diastereoisomeric mixture of ribonucleoside 3'-phosphoramidites. These data established that no detectable migration of the TBDMS group occurred during the phosphitylation reaction (Figure 2).




Figure 2. $^{31}\text{P-NMR}$ of α -uridine 2'- and 3'-phosphoramidites **6** and **5** obtained from separate preparations.




Figure 3. Reversed-phase HPLC analysis of the crude mixture containing U2'p5'U (a) and U3'p5'U (b) after deprotection using a C18 column.

The preparation of α -uridine derivatized support **8** was achieved via formation of a succinyl linkage between the 2'-hydroxy group of 3'-silylated α -uridine **4** and the amino group of the long chain alkylamine controlled pore glass (LCA-CPG). For this purpose the pentachlorophenyl succinate of the 3'-5'-protected α -uridine **7** was reacted with LCA-CPG according to the usual method¹².

The assembly of α -uridyates was achieved on an Applied Biosystems 381A synthesizer using 1 μ mole protected nucleoside loaded support. As in the β ribo series¹³, the condensation time required to obtain high coupling yields was 15 minutes using 0.5M tetrazole as activator and 0.15M phosphoramidite 5 in acetonitrile. Thus, the average coupling yield determined from the dimethoxytrityl cation release was 97% for the synthesis of α -rU₆ and α -rU₁₂. These two oligomers were deprotected¹⁴ and purified by HPLC (spectrophotometric purity at 260nm was better than 97%).

In order to evaluate the possible occurrence of phosphoryl migration under the deprotection conditions¹⁴, we have synthesized the dimer α -(UpU). After cleavage of the protected dimer from the support and its deprotection, the products were analysed by reversed phase HPLC. Inspection of the chromatogram (Figure 3) showed one major peak corresponding to α -uridylyl-(3'→5')- α -uridine (R_T 15.11) and a minor peak (R_T 8.67) which was coeluted with α -uridylyl-(2'→5')- α -uridine (obtained from 6) and accounted for only 0.3% of the major peak. This very low extent of isomerisation compared well with previous results obtained in the β -series¹⁵.

The substrate activity for four nucleases of rU₆ either in the all α - or all β -configuration was studied. The enzymatic hydrolysis were monitored by HPLC and the results are reported in Table 1. Under conditions where the β -hexamer was almost or completely degraded by either calf spleen phosphodiesterase, nuclease S1 or ribonuclease A, the corresponding α -hexamer remained intact. These results are consistent with the high nuclease resistance already observed with α - oligodeoxyribonucleotides¹⁶.

Substrate Nucleases	Time	α -rU ₆ *	β -rU ₆ *
Calf spleen phosphodiesterase	2 min	100%	89%
	60 min	100%	14%
Snake venom phosphodiesterase	2 min	94%	6.5%
	120 min	42%	0%
Nuclease S1	60 min	100%	68%
	300 min	100%	14%
Ribonuclease A	5 min	100%	0.3%
	35 min	100%	0%

Table 1. Time course of the cleavage of α - or β -rU₆ by four nucleases at 37°C. *Fraction of remaining hexamer after indicated time of incubation.

Base-pairing between α -rU₁₂ and poly β -rA was followed by UV absorption spectroscopy¹⁷. Upon increasing the temperature of a 1:1 (A to U) mixture formed at 2°C a hyperchromicity was observed from which a melting temperature value (T_m) of 16°C was determined. This value is lower than the calculated melting temperature¹⁸ of the duplex β -rU₁₂. β -rA₁₂ (25°C). As this first data are only related to homopolymers, we are developing the synthesis of α -oligo-

ribonucleotides presenting mixed purine-pyrimidine sequences in order to fully evaluate their binding capacities.

Acknowledgments : This work was supported by a grant from CNRS (ATIFE : "ARN non naturels") and from the Association pour la Recherche contre le Cancer (ARC) ("Oligonucleotides modifiés : nouveaux régulateurs potentiels de l'expression génétique" and for the DNA synthesizer purchase).

References and Notes.

1. "Oligonucleotides - Antisense Inhibitors of Gene Expression" (J.S.Cohen, ed.) *Mc Millan Topics in Structural and Molecular Biology Series* (S.Neidle & W.Fuller Eds.) Vol.12.
2. C.A Stein, J.Cohen, *Cancer Res.* 1988, **48**, 2659.
3. G.Zon, *Pharm. Res.* 1988, **5**, 539.
4. B.Rayner, C.Malvy, J.Paoletti, B.Lebled, C.Paoletti, J.-L.Imbach, 1989, see ref.1, pp116-136.
5. D.H.Shannahoff, R.A. Sanchez, *J. Org. Chem.* 1973, **38**, 593.
6. H.Schaller, G.Weimann, B.Lerch, H.G.Khorana, *J. Amer. Chem. Soc.* 1963, **85**, 3821.
7. N.Usman, R.T.Pon, K.K.Ogilvie, *Tetrahedron Lett.* 1985, **26**, 4567.
8. G.H.Hakimelahi, Z.A.Proba, K.K.Ogilvie, *Can. J. Chem.* 1982, **60**, 1106.
9. K.K.Ogilvie, A.L.Schifman, C.L.Penney, *Can. J. Chem.* 1979, **57**, 2230.
10. For **3** : $^1\text{H-NMR}$ (CDCl_3) δ 8.57 (s, 1H, NH), 7.59 (d, 1H, $J_{5,6} = 8.2$ Hz, H-6), 7.41-6.83 (m, 13H, ArH), 6.49 (d, 1H, $J_{1',2'} = 5.9$ Hz, H-1'), 5.69 (d, 1H, H-5), 4.85 (m, 1H, H-2'), 4.35 (m, 1H, H-4'), 4.07 (m, 1H, H-3'), 3.79 (s, 6H, O-CH₃), 3.52-3.06 (m, 2H, H-5' and H-5"), 2.54 (d, 1H, OH-3'), 0.85 (s, 9H, (CH₃)₃C), 0.06 and -0.13 (2s, 6H, (CH₃)₂Si).
For **4** : $^1\text{H-NMR}$ (CDCl_3) δ 8.73 (s, 1H, NH), 7.55 (d, 1H, $J_{5,6} = 8.2$ Hz, H-6), 7.45-6.82 (m, 13H, ArH), 6.34 (d, 1H, $J_{1',2'} = 4.4$ Hz, H-1'), 5.72 (d, 1H, H-5), 4.44 (m, 1H, H-2'), 4.35 (pseudo t, 1H, H-3'), 4.23 (m, 1H, H-4'), 3.80 (s, 6H, O-CH₃), 3.46-3.03 (m, 2H, H-5' and H-5"), 2.74 (d, 1H, OH-2'), 0.82 (s, 9H, (CH₃)₃C), 0.025 and -0.07 (2s, 6H, (CH₃)₂Si).
11. L.J.McBride, M.H.Caruthers, *Tetrahedron Lett.* 1983, **24**, 245.
12. G.R.Gough, M.J.Brunden, P.T. Gilham, *Tetrahedron Lett.* 1981, **22**, 4177.
13. N.Usman, K.K.Ogilvie, M.Y.Jiang, R.J.Cedergren, *J. Amer. Chem. Soc.* 1987, **109**, 7845.
14. Deprotection of α -oligoribonucleotides: after the removal of methyl phosphate protecting groups by a solution of thiophenol/NET₃/dioxane (1/2/2), the oligomer was cleaved from the support in ammonia 32% / ethanol (3/1 v/v) and incubated for 5 hours at 55°C (deprotection conditions for the α -oligoribonucleotides with protected bases). Then the 2'-TBDMS groups were removed by reaction with 1M tetrabutylammonium fluoride in THF. The oligonucleotides were desalted on a small ion exchange column (DEAE-Sephadex A25) and were eluted with triethylammonium bicarbonate buffer (TEAB pH=7.5) from 10⁻³M to 1M.
15. T.Wu, K.K.Ogilvie, R.T.Pon, *Nucl. Acids Res.* 1989, **17**, 3501.
16. F.Morvan, B.Rayner, J.-L.Imbach, S.Thenet, J.-R.Bertrand, J.Paoletti, C.Malvy, C.Paoletti, *Nucl. Acids Res.* 1987, **15**, 3421.
17. Concentration of each oligomer was 10 μM . Measurements were carried out in a pH 7 buffer containing 10 mM sodium cacodylate and 1 M NaCl.
18. P.N.Borer, B.Dangler, I.Tinoco Jr, O.C.Uhlenbeck, *J. Mol. Biol.* 1974, **86**, 843.