

## Comparative anatomy vs mechanistic understanding: how to interpret the diameter-vulnerability link

Frederic Lens, Sean M Gleason, Giovanni Bortolami, Craig Brodersen, Sylvain

Delzon, Steven Jansen

#### ▶ To cite this version:

Frederic Lens, Sean M Gleason, Giovanni Bortolami, Craig Brodersen, Sylvain Delzon, et al.. Comparative anatomy vs mechanistic understanding: how to interpret the diameter-vulnerability link. IAWA Journal, 2023, 44 (3-4), pp.368-380. 10.1163/22941932-bja10137 . hal-04601373

### HAL Id: hal-04601373 https://hal.science/hal-04601373v1

Submitted on 10 Jun2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

#### Comparative anatomy vs mechanistic understanding: how to interpret the diametervulnerability link

Frederic Lens<sup>1,2</sup>, Sean M. Gleason<sup>3</sup>, Giovanni Bortolami<sup>1</sup>, Craig Brodersen<sup>4</sup>, Sylvain Delzon<sup>5</sup>, Steven Jansen<sup>6</sup>

<sup>1</sup>Naturalis Biodiversity Center, P.O. Box 9517, 2300 RA Leiden, The Netherlands;
 <sup>2</sup>Leiden University, Institute of Biology Leiden, Plant Sciences, Sylviusweg 72, 2333 BE Leiden, The Netherlands;

<sup>3</sup>Water Management and Systems Research Unit, United States Department of Agriculture, Agricultural Research Service, Fort Collins, CO 80526, USA;

<sup>4</sup>School of the Environment, Yale University, New Haven, Connecticut 06511, USA;

<sup>5</sup>University of Bordeaux, INRAE, BIOGECO, 33615, Pessac, France;

<sup>6</sup>Institute of Systematic Botany and Ecology, Albert-Einstein-Allee 11, Ulm University, D-89081, Ulm, Germany.

ORCID Frederic Lens: 0000-0002-5001-0149 ORCID Sean M. Gleason: 0000-0002-5607-4741 ORCID Giovanni Bortolami: 0000-0001-7528-9644 ORCID Craig Brodersen: 0000-0002-0924-2570 ORCID Sylvain Delzon: 0000-0003-3442-1711 ORCID Steven Jansen: 0000-0002-4476-5334

#### Author for correspondence

Frederic Lens, email: frederic.lens@naturalis.nl

Short title: Role of vessel diameter on drought-induced embolism

#### Word count

abstract: 217, Main body total: 4140 Figures: 0; Tables: 0; Supporting information: 0

#### Abstract

Results from comparative and ecological wood anatomy in combination with a number of experimental studies on plant hydraulics have led to a pervasive and longstanding assumption that wider diameter vessels are more vulnerable to drought-induced embolism than narrower vessels. Although we agree that wider vessels tend to be more vulnerable than narrower vessels within stems and within roots across most species, our current understanding of the diameter-vulnerability link does not offer a mechanistic explanation for why increased vessel diameter should consistently lead to greater vulnerability or vice versa. Causes of drought-induced embolism formation and spread likely operate at the nano-level, especially at gas-liquid-surfactant interfaces inside intervessel pit membranes. We evaluate here new perspectives on drought-induced embolism and its key anatomical and physico-chemical drivers, of which vessel diameter is one of the parameters involved, although its linkage to embolism vulnerability is likely indirect. As such, the diametervulnerability link does not imply that species with on average wider vessels are consistently more susceptible to drought-induced embolism compared to species with narrower vessels. Scientific priorities for future progress should focus on more accurate predictions of how water transport in plants is affected by drought, which require a better mechanistic understanding of xylem network topology and biophysical processes at the nano-scale level in individual vessels that determine embolism formation and spread.

#### Key words

drought, embolism, pit membrane thickness, plant hydraulics, vessel diameter, wood anatomy, xylem networks, xylem sap

#### Introduction

One of the most frequently measured xylem traits in angiosperms is vessel diameter, which ranges from 10 - 700 µm both across and within species (Hacke et al. 2017; Olson 2020). This trait is commonly measured for multiple reasons, most notably because vessel diameter is easily measured from cross sections of xylem, and has proven to act as an important functional trait regulating hydraulic conductivity (Sperry et al. 2006; Lachenbruch & McCulloh 2014) and freeze-thaw-induced embolism (Sperry & Sullivan 1992; Davis et al. 1999; Sevanto et al. 2012; Lintunen et al. 2013, 2020; Charrier et al. 2014), among other plant functional traits. Broad accessibility to this trait with relatively simple methods has also provided numerous opportunities for comparative anatomy studies that allowed for insights into plant adaptation across various terrestrial habitats, many of which were initiated by Sherwin Carlquist and followed by others (e.g., Carlquist 1966, 2001; Baas et al. 1983; Lens et al. 2004; Olson et al. 2023). These comparative studies show that species from dry environments tend to have a narrower mean vessel diameter than closely related species from wetter habitats, which provides evidence for the idea that species with wider vessels are more susceptible to drought-induced embolism than species with narrower vessels (e.g., Olson et al. 2018; Stovall et al. 2019). To further highlight the role of vessel diameter in a plant's drought response, Carlquist (1975) calculated indices for Vulnerability (V; vessel diameter divided by vessel density) and Mesomorphy (V multiplied by vessel

element length) for species native to habitats with contrasting levels of aridity, and found that drought-adapted species showed *V*- and *M*-indices markedly below 1.0 and below 30, respectively (summarised in this special issue by Ewers *et al.* 2023). Yet, despite several decades of work on establishing a link between conduit diameter and vulnerability to drought-induced embolism, the current state of the underlying mechanistic understanding remains largely uncertain (Lens *et al.* 2022; Hacke *et al.* 2023).

In addition to the comparative studies of Carlquist (1966, 2001) and pioneers before him (see Olson 2020 for an historical overview), part of the theoretical background regarding the functional role of conduit diameter finds its root in the seminal work by Zimmermann (1983), which illustrates the effects of conduit diameter on idealised Hagen-Poiseuille flow in cylindrical pipes. This theoretical framework was highly influential, and set the stage for many subsequent investigations to explicitly test the hypotheses presented in Zimmermann's book. Emerging over the following decades was the idea that some inherent tradeoff must exist between optimising the xylem for hydraulic efficiency, structural support, and hydraulic safety. That hypothetical tradeoff 'triangle' (see Baas et al. 2004; Brodersen 2015) was then critically evaluated based on a large-scale data analyses, confirming (often weak) correlations between efficiency and safety (measured as P<sub>50</sub>: xylem sap water potential leading to a 50% loss of conductivity; Gleason et al. 2015), between support and safety (Hacke et al. 2001), and between conduit diameter and safety (Fig. 1 in Lens et al. 2022; Hacke et al. 2023). This latter diameter-vulnerability link was also confirmed in a number of smaller-scale experimental studies (e.g., Hargrave et al. 1994; Cai & Tyree 2010; Nardini et al. 2017; Jacobsen et al. 2019; Isasa et al. 2023). However, many other smaller-scale experimental publications did not find a diameter- $P_{50}$  link in either stems or petioles (e.g., Lens *et al.* 2011, 2016; Brodersen *et al.* 2013; Choat *et al.* 2016; Dória *et al.* 2018, 2019; Emilio *et al.* 2019; Thonglim *et al.* 2020, 2022); this link was also not retreived in some studies on conifers (Bouche *et al.* 2014; Johnson & Brodribb 2023). Two conclusions that can be drawn from these inconsistent experimental results: (1) even if a correlation between vessel diameter and  $P_{50}$  exists, this correlation is of little use without critically interpreting what it actually means in relation to total-plant drought resilience, and (2) reporting correlation between these traits does little to inform our understanding of the mechanisms underlying drought-induced embolism formation and spread, which is currently a priority for the field (Lens *et al.* 2022).

In this opinion paper, based on the ideas already published in detail in Lens *et al.* (2022) and inspired by earlier comparative wood anatomy work by Carlquist and others, we further explore the idea of a potential mechanistic relationship between vessel diameter and vulnerability to drought-induced embolism, and discuss how this impacts the interpretation of the diameter-vulnerability link. A critical assessment of this link involves two research questions: (1) are wider vessels always more prone to drought-induced embolism than narrower vessels (and vice versa), and (2) are species with a higher mean vessel diameter consistently more vulnerable to embolism - and by extension more vulnerable to drought - than species with narrower vessels? These two questions are often not separated, and we believe disentangling them will lead to a more balanced and useful understanding of the role of vessel diameter in drought-induced embolism.

Even when assuming the diameter-vulnerability link to be valid across a broad range of vascular plant species, we echo the conclusion in our previous paper stating that (1) wider

vessels are often (but not always) more vulnerable to drought-induced embolism compared to narrower vessels when the same organs are taken into account (see also Fig. 2 in Hacke *et al.* 2023), and (2) that species with on average wider vessels are not necessarily more vulnerable to drought-induced embolism than species with narrower vessels. The main reasons for these two statements are that the mechanisms leading to drought-induced embolism involve a complex interaction among multiple anatomical (incl. vessel diameter) and physico-chemical drivers operating at different scales (Lens *et al.* 2022). Unfortunately, this complexity and the fact that we do not understand exactly how plants are able to transport water under negative pressure without constant or even frequent embolism formation, makes it difficult to generalise findings related to drought-induced embolism.

To reconcile this discrepancy, there is an urgent need to keep on critically assessing the methods used to generate anatomical and experimental data, and to dive deeper into the mechanisms behind drought-induced embolism. In addition, the vulnerability discussion concerning wide vessels contributes to debates about the frequency of drought-induced embolism events *in planta*: (1) does embolism formation occur in the vulnerable wide vessels on a daily basis in the field also under normal conditions of precipitation, which indirectly suggests frequent embolism repair events (Salleo *et al.* 2004; Jacobsen *et al.* 2007; Hacke *et al.* 2009; Nardini *et al.* 2011; Jacobsen & Pratt 2012), or (2) is widespread embolism uncommon over relatively long periods in the field and limited to periods of severe drought in narrow- as well as wide-vesseled species (Cochard & Delzon 2013; Lens *et al.* 2016; Wason *et al.* 2018; Guan *et al.* 2022)? Answering these questions is crucial because embolism events could result in the loss of hydraulic conductance, tissue desiccation and/or dieback (Davis *et al.* 2002; Venturas *et al.* 2016; Pratt *et al.* 2020; Brodribb *et al.* 2021;

Mantova *et al.* 2022), and can therefore affect plant function and survival (McDowel *et al.* 2008; Allen *et al.* 2015; Anderegg *et al.* 2016; Adams *et al.* 2017; Choat *et al.* 2018; Brodribb *et al.* 2020).

# Why are wider vessels not necessarily more vulnerable to drought-induced embolism than narrower vessels?

A frequently cited argument from ecological studies on wood anatomy supporting the diameter-vulnerability link is that earlywood vessels, which are often formed during the spring in temperate regions or during the wet season in seasonally dry regions, are always wider than latewood vessels formed later in the season, often under drier conditions (Carlquist 2001; Olson et al. 2023). We agree that earlywood will be on average more vulnerable than latewood across most of these species. However, this does not necessarily indicate that all earlywood and latewood vessels differ in vulnerability to drought-induced embolism, and it is even plausible that some of the latewood vessels are considerably more vulnerable than most earlywood vessels. With respect to the tracheid-bearing conifer species Pseudotsuga menziesii, Dalla-Salda et al. (2014) indeed found that the latewood is more vulnerable to drought-induced embolism compared to earlywood, probably due to inflexible interconduit pit membranes in the latewood tracheids that facilitate embolism spread. Therefore, linking conduit diameter with vulnerability to embolism can be misleading when not considering other traits that may play an equal or more important role in drought-induced embolism formation than conduit diameter. In other words, all else being equal, wider vessels are likely more vulnerable than narrower vessels, agreeing with the large body of literature (e.g., Lens et al. 2022; Hacke et al. 2023; Olson et al. 2023).

However, all else is typically not equal in nature, implying that wider vessels can also be considerably more resistant to drought-induced embolism than narrower vessels.

Also from the point of view of xylem physiology, there are three main explanations for why the impact of vessel diameter on drought-induced embolism vulnerability should be treated in a broader perspective. First, the mechanisms behind drought-induced embolisms are complex, poorly understood, and occur at much smaller scales than whole-conduit dimensions. For example, gas bubble formation (and stabilisation), and bubble expansion may arise from nano-scale interactions occurring among the multiple interfaces between liquids, gases, and solids inside conduits. These interactions are further complicated by the presence of solutes and insoluble xylem sap compounds that change the surface tension, aka surface-active agents or surfactants, which are hypothesised to stabilise very small gas bubbles ("nanobubbles") and may result in xylem sap that is far more resilient to droughtinduced embolism than traditionally thought (Schenk et al. 2017, 2020; Yang et al. 2020; Jansen et al. 2022; Ingram et al. 2021). However, the precise combination of surfaces (cell wall, pit border, pit membrane), xylem sap, and gas properties/conditions that result in embolism formation and spread remain poorly understood, but are likely proximally related to embolism vulnerability. For example, it is possible that embolisms result from pre-existing gas bubbles, including tiny nanobubbles or surface bubbles associated with hydrophobic vessel surfaces, which expand in the xylem sap when these bubbles exceed a critical size threshold for a given pressure (Tyree et al. 1994; Hölttä et al. 2002; Lohse & Zhang 2015; Schenk et al. 2017), although experimental evidence for this idea is poor. How nanobubbles and the likelihood of embolism formation relate to vessel diameter (and length) remains to be studied: either vessels with a larger volume contain a higher total number of lipid-coated

nanobubbles that may increase embolism risk, or these wider and longer vessels could be more resistant if large neighbouring vessels during their initial stage of embolisation would extract gas from their functional neighbours in order to reach atmospheric pressure at the final stages of embolisation (Wang *et al.* 2015).

Second, correlation does not invoke causation, which obviously also applies to the diametervulnerability link. There is no known direct, mechanistic link between conduit diameter and drought-induced embolism resistance, not at the whole-plant level nor at the ultrastructural pit level. If such a mechanism were to exist, we should expect it to affect the relative vulnerabilities of xylem in leaves (narrower conduits) and stems (wider conduits) similarly within and across individuals and species, but it clearly has not. For example, although several authors have reported leaf xylem to be more resistant to embolism than branch or stem xylem (Zhu et al. 2016; Klepsch et al. 2018; Levionnois et al. 2020b; Guan et al. 2022), this result is far from consistent across studies, despite leaves having consistently narrower conduits than branches or stems. Likewise, several authors have reported that leaves are more vulnerable than perennial shoots (Pivovaroff et al. 2014; Charrier et al. 2016; Johnson et al. 2016; Creek et al. 2018; Skelton et al. 2019), or else have found no meaningful difference in xylem vulnerability between leaves, stems and/or roots of the same individuals (Skelton et al. 2017; Creek et al. 2018; Wason et al. 2018; Losso et al. 2019; Levionnois et al. 2020b; Smith-Martin et al. 2020; Wu et al. 2020, Lübbe et al. 2022). Thus, if a causal mechanism between conduit diameter and vulnerability was identified, it would need to account for the inconsistent vulnerability-diameter relationships within and across individuals. Bearing this in mind, the recent study of Isasa et al. (2023) showed that a negative vessel diameter- $P_{50}$  relationship exists at the interspecific scale, but not at the

intraspecific scale. In conclusion, a view that takes into account the entire body of research on xylem vulnerability, rather than studies that only support a vulnerability-diameter link within a single organ, does not support a mechanism operating at the scale of the whole plant.

At the pit level, researchers often cite the pit area hypothesis (Wheeler et al. 2005; Hacke et al. 2006, 2009; Christman et al. 2009, 2012) to mechanistically explain the diametervulnerability link. This hypothesis assumes that permeability to gas and embolism propagation among vessels increases with increasing intervessel pit area of a given vessel with average dimensions (Ap). While the pit area hypothesis correctly emphasises the functional importance of intervessel connectivity (Loepfe et al. 2007; Wason et al. 2021; Bouda et al. 2022), large pit membrane pores are thought to be exceptionally rare and, if observed, are most likely the result of artefacts (Jansen et al. 2008, 2009; Choat et al. 2008). Furthermore, the idea that a rare, single-layered pit membrane would be highly leaky due to a single, unusually large pore, leading to embolism propagation, is not compatible with recent theoretical insights on the three-dimensional, multi-layered nature of pit membranes with multiple pore constrictions that represent the bottlenecks for water and gas transport (Kaack et al. 2019, 2021; Yang et al. 2020; Zhang et al. 2020). These insights, based on 3Dmodelling and hydraulics measurements, suggest that P<sub>50</sub> has a stronger relationship with intervessel pit membrane thickness than Ap (Kaack et al. 2021), although Isasa et al. (2023) showed that intervessel membrane thickness alone cannot explain the  $P_{50}$  interspecific variability based on 12 temperate tree species. What emerges from these studies is that (1) the pit area hypothesis should be further tested with respect to both length-diameter scaling of conduits and connectivity (among both vessels and adjacent tracheids; Carlquist

1984; Sperry *et al.* 2006) before it can be used as a solid mechanistic explanation for a possible correlation between  $P_{50}$  and vessel diameter, (2) embolism spread is the result of a combination of different traits that may be related (e.g., vessel connectivity or Ap) or may not be related (e.g., sap composition, vessel wall chemistry, intervessel pit membrane thickness) to vessel diameter (Lens *et al.* 2022), and (3) there is an urgent need to validate the theoretical pit membrane models with ultrastructural observations of intervessel pit membranes.

Third, several different methods have been used to quantify both mean vessel diameter and P<sub>50</sub>, thereby adding noise to the diameter-vulnerability link (e.g., Brodersen et al. 2013; Cochard & Delzon 2013; Choat et al. 2016; Martin-StPaul et al. 2017; Creek et al. 2018; Emilio et al. 2019; Smith-Martin et al. 2020; Barigah et al. 2021). With respect to vessel diameter, there are multiple ways for a plant to arrive at the same mean vessel diameter (i.e., different vessel diameter frequency distributions), and "mean" vessel diameter may not be the most meaningful quantification of vessel width, as it relates to embolism vulnerability. In addition, mean vessel diameter is calculated in different ways (e.g., tangential diameter, equivalent circle diameter), and is often hydraulically weighted based on either all vessels in a cross-section or from a subset of these vessels (usually > 100) (Scholz et al. 2013). Also, the pressure gradient driving water transport through the xylem may not be uniform across all vessels in a cross section (Bouda et al. 2019), which would likely result in errors in estimated vulnerability across vessels of different diameter within the same cross section. To further complicate matters, branch age, growth rate, number of growth rings, and seasonality may affect the vulnerability curve (Charrier et al. 2018; Olson et al. 2018; Pratt et al. 2020; Sorek et al. 2020; Weithmann et al. 2022). In summary, the

interpretation of the diameter-vulnerability link would become clearer if we know the vulnerability of individual vessels, and the determinants of this vulnerability, as well as the aggregate vulnerability across many vessels.

The unexplained variation in the vulnerability-diameter correlation could also be explained by differences in  $P_{50}$  methods that may impact vulnerability curves (Cochard *et al.* 2013; Venturas et al. 2019; Gauthey et al. 2020; Hacke et al. 2023). The open vessel artefact is noteworthy in this regard, especially in studies focusing on stems of long-vesseled genera such as Quercus (Sperry et al. 2012; Martin-StPaul et al. 2014; Skelton et al. 2018, 2021; Percolla et al. 2021) and Vitis (Choat et al. 2010; Jacobsen & Pratt 2012; McElrone et al. 2012; Charrier et al. 2018). Many studies showed that this artefact occurs when embolism is artificially induced with the centrifuge method in stem (or root) segments that are considerably shorter than the maximum vessel length, thus resulting in vessels being cut at both ends ("open vessels"). These longer (and often wider) cut-open vessels embolize quickly when they are subjected to mild negative pressure, thus resulting in a rapid (exponential or "r"-shaped) decline in hydraulic conductivity with increasing xylem tension, and hence an overestimation of vulnerability (Alder et al. 1996; Cochard 2002; Choat et al. 2010; Cochard et al. 2010, 2013; McElrone et al. 2012; Martin-StPaul et al. 2014; Torres-Ruiz et al. 2014; although see Tobin et al. 2012, Jacobsen & Pratt 2012, Sperry et al. 2012, Hacke et al. 2014, and Jacobsen & Pratt 2023 for a different point of view). Other factors that are known to affect the shape of a vulnerability curve are temporal variation in "native" embolism (daily or seasonal) and the flushing of stems with water prior to taking an initial hydraulic measurement (Domec et al. 2006; Jacobsen et al. 2007; Hacke et al. 2014, 2023). Considering all these potential sources of error in embolism vulnerability measurements, a

critical examination of methods and models used to estimate xylem vulnerability remains a research priority (Cochard *et al.* 2013).

## Why are species with wider vessels not necessarily more vulnerable to drought-induced embolism than those with narrower vessels?

The mechanisms leading to drought-induced embolism involve many potential anatomical and physico-chemical drivers of drought-induced embolism. This implies that natural selection may act on combinations of traits that affect the susceptibility of xylem to drought-induced embolism at different scales, with vessel diameter only being one of these traits that may or may not have a significant impact on the drought resistance of the entire individual. The potential xylem drivers of drought-induced embolism are discussed more indepth in Lens *et al.* (2022), and are listed here in alphabetical order (see also Box 1 in Lens *et al.* 2022).

(1) macro-level (vessel) traits: 3D-conduit network (Loepfe *et al.* 2007; Brodersen 2013;
Mrad *et al.* 2018, 2021; Wason *et al.* 2021; Bouda *et al.* 2022), Ap (Wheeler *et al.* 2005;
Kaack *et al.* 2021), and conduit diameter (this paper; Isasa *et al.* 2023);

(2) nano-level xylem traits: intervessel pit membrane thickness (Jansen *et al.* 2009; Lens *et al.* 2011, 2013; Li *et al.* 2016; Dória *et al.* 2018; Levionnois *et al.* 2020a; Thonglim *et al.* 2020, 2022; Kaack *et al.* 2021), cell wall chemistry and thickness (Lens *et al.* 2016; Greenwood *et al.* 2017; Pereira *et al.* 2018; Dória *et al.* 2018, 2019; Thonglim *et al.* 2020, 2022; Frankiewicz *et al.* 2021; Liang *et al.* 2021; Zizka *et al.* 2022), wettability of the inner conduit wall (Lohse & Zhang 2015; Zwieniecki & Holbrook 2000; McCully *et al.* 2014; Schenk *et al.* 2017, 2018, 2020; Brodersen *et al.* 2018), dynamic surface tension of xylem sap and surfactant-coated nanobubbles (Schenk *et al.* 2015, 2017, 2020; Yang *et al.* 2020; Ingram *et al.* 2021; Jansen *et al.* 2015; Jansen *et al.* 2015; Jansen *et al.* 2015; Jansen *et al.* 2017; Pereira *et al.* 2015; Jansen *et al.* 2017; Pereira *et al.* 2015; Jansen *et al.* 2017; Pereira *et al.* 2015; Jansen *et* 

*al.* 2022), and gas movement and xylem sap oversaturation (Schenk *et al.* 2016; Pereira *et al.* 2020; Guan *et al.* 2021).

#### **Concluding remarks**

From the above, it is clear from both ecological wood anatomy and experimental studies that a weak but significant diameter-vulnerability link does exist when comparing all available data (mainly from stems; Fig. 1 in Lens et al. 2022). This means that there is a large body of evidence that wider vessels tend to be more vulnerable than narrower vessels within the same organ, confirming Carlquist's ideas (1966, 2001). Consequently, we agree that vessel diameter likely plays some role in drought-induced embolism formation in many clades (Lens et al. 2022; Olson et al. 2023), but most likely not within all clades (e.g., Lens et al. 2011, 2016; Brodersen et al. 2013; Choat et al. 2016; Dória et al. 2018, 2019; Emilio et al. 2019; Thonglim et al. 2020, 2022), and definitely not at the whole-plant level (e.g., Skelton et al. 2017; Creek et al. 2018; Wason et al. 2018; Losso et al. 2019; Levionnois et al. 2020b; Smith-Martin et al. 2020; Wu et al. 2020, Lübbe et al. 2022). Even if we generally accept that there is a diameter-vulnerability link, not knowing the mechanisms involved in droughtinduced embolism can lead to subtle but important differences in the interpretation of what the link actually means, i.e., how and why it has arisen and its consequences on plant function. Although we agree with Olson et al. (2023) that the salient vessel diameter patterns in nature are valuable in directing our research questions to solve how plants respond to environmental stress, we feel that this approach overemphasises the role of vessel diameter in the drought stress debate, and simplifies the relationship between vessel diameter and embolism vulnerability to the point of being misleading. The diametervulnerability link does not imply that (1) wider vessels are necessarily more susceptible to

drought-induced embolism (and vice versa) or (2) that vessels with the same width (and length) are equally vulnerable (see also Fig. 2 in Hacke *et al.* 2023), and (3) it definitely does not mean that species with on average wider vessels are more vulnerable to droughtinduced embolism or by extension are more drought-sensitive. As we all know, plants are famous for developing multiple solutions when it comes to drought response strategies (e.g., Pivovaroff *et al.* 2015; McDowell *et al.* 2022). Placing a scientific premium on the study of vessel diameter as an important and meaningful correlate of embolism vulnerability may not be the best way to prioritise future research, nor the best way to inspire research that could potentially lead to a better understanding of drought-induced embolism (Lens *et al.* 2022). Although pointing out that correlation does not imply causation may appear as a platitude, it is true that scientists still commonly cite vessel diameter in discussions about drought responses without addressing why this correlation represents a convincing argument, or not.

We suggest that a better understanding of the diameter-vulnerability link could be obtained by focusing our research efforts on three broad objectives: (1) improve our understanding of the vulnerability of individual vessels, rather than an aggregate measure of vulnerability across many vessels (although this is technically challenging, e.g., via microCT observations that span an axial segment length beyond the length of the longest vessels), and connect this information to other drought-associated traits in the xylem and beyond, (2) develop new and more powerful non-invasive techniques to identify macro and nano-scale mechanistic driver(s) of embolism initiation and spread across different environmental conditions and different plant organs, and (3) evaluate the relative importance and timing of xylem embolism among the sequence of physiological failures that occur during drought, i.e. loss of turgor, cessation of cell division and growth, plasma membrane instability, plant death (Mantova et al. 2022). Once a large body of drought-associated xylem traits are investigated in different organs of carefully selected model species that grow in contrasting environments, we can further test and fine-tune the mechanisms involved in embolism formation and spread through the development of a new family of integrative, processbased, xylem models linking nano- (Li et al. 2020; Zhang et al. 2020; Kaack et al. 2021; Weishaupt et al. 2022) to macro-scale processes (Loepfe et al. 2007; Mrad et al. 2018, 2021; Wason et al. 2021). Integrating nano- and macro-scale xylem processes would allow the evaluation of pit membrane traits, vessel network traits (e.g., connectivity among vessels and adjacent tracheids, vessel widening, segmentation or sectoriality; Carlquist 1984; Olson et al. 2020; McElrone et al. 2021) and their interactions on whole-plant transpiration, construction costs, and carbon income. This knowledge will be important to reconcile our current state-of-the-art knowledge of embolism formation and spread with observations made from comparative anatomical studies, such as the empirical correlation between embolism vulnerability and conduit diameter. Moreover, it will help us to understand how plants will respond to a warming climate, and how to manage natural and agricultural systems under these novel environmental conditions (Choat et al. 2018; Brodribb et al. 2020).

#### Acknowledgments

We warmly thank editors-in-chief Marcelo Pace and Lloyd Donaldson for their guidance through the review process of this manuscript, and their aim to provide a balanced view on the role of vessel diameter in this special issue dedicated to Sherwin Carlquist. FL and GB were supported by the Dutch Research Council NWO (grant ALWOP.488), SD by the REGION project VITIPIN and Grand Programme de Recherche Bordeaux Plant Science (BPS, WP Waterloo), and SJ acknowledges funding from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation, project number 457287575).

#### References

- Adams HD, Zeppel MJB, Anderegg WRL, Hartmann H, Landhäusser SM, *et al.* 2017. A multispecies synthesis of physiological mechanisms in drought-induced tree mortality. Nat. Ecol. Evol. 1: 1285–1291. DOI: 10.1038/s41559-017-0248-x.
- Alder NN, Sperry JS, Pockman WT. 1996. Root and stem xylem embolism, stomatal conductance, and leaf turgor in *Acer grandidentatum* populations along a soil moisture gradient. Oecologia 105: 293–301. DOI: 10.1007/BF00328731.
- Allen CD, Breshears DD, McDowell NG. 2015. On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene. Ecosphere 6: art129. DOI: 10.1890/ES15-00203.1.
- Anderegg WRL, Klein T, Bartlett M, Sack L, Pellegrini AFA, *et al.* 2016. Meta-analysis reveals that hydraulic traits explain cross-species patterns of drought-induced tree mortality across the globe. PNAS 113: 5024–5029. DOI: 10.1073/pnas.1525678113.
- Baas P, Ewers FW, Davis SD, Wheeler EA. 2004. Evolution of xylem physiology. In: AR Hemsley, I Poole (eds.), The evolution of plant physiology: 273–295. Elsevier, Oxford, UK.
- Baas P, Werker E, Fahn A. 1983. Some ecological trends in vessel characters. IAWA J. 4: 141– 159. DOI: 10.1163/22941932-90000407.

Barigah TS, Gyenge JE, Barreto F, Rozenberg P, Fernandez ME. 2021. Narrow vessels cavitate first during a simulated drought in *Eucalyptus camaldulensis*. Physiol. Plant. 173: 2081–2090. DOI: 10.1111/ppl.13556.

Bouche PS, Larter M, Domec J-C, Burlett R, Gasson P, Jansen S, Delzon S. 2014. A broad survey of hydraulic and mechanical safety in the xylem of conifers. J. Exp. Bot. 65: 4419–4431. DOI: 10.1093/jxb/eru218.

- Bouda M, Huggett BA, Prats KA, Wason JW, Wilson JP, *et al*. 2022. Hydraulic failure as a primary driver of xylem network evolution in early vascular plants. Science 378: 642–646. DOI: 10.1126/science.add2910.
- Bouda M, Windt CW, McElrone AJ, Brodersen CR. 2019. *In vivo* pressure gradient heterogeneity increases flow contribution of small diameter vessels in grapevine. Nat. Commun. 10: 5645. DOI: 10.1038/s41467-019-13673-6.

Brodersen CR. 2013. Visualizing wood anatomy in three dimensions with high-resolution Xray micro-tomography (MCT) - a review. IAWA J. 34: 408–424. DOI:

10.1163/22941932-00000033.

Brodersen CR. 2015. Finding support for theoretical tradeoffs in xylem structure and function. New Phytol. 209: 8–10. DOI: 10.1111/nph.13763.

Brodersen CR, Knipfer T, McElrone AJ. 2018. *In vivo* visualization of the final stages of xylem vessel refilling in grapevine (*Vitis vinifera*) stems. New Phytol. 217: 117–126. DOI: 10.1111/nph.14811.

Brodersen CR, McElrone AJ, Choat B, Lee EF, Shackel KA, et al. 2013. In vivo visualizations of drought-induced embolism spread in Vitis vinifera. Plant Physiol. 161: 1820–1829.
DOI: 10.1104/pp.112.212712.

# Brodribb T, Brodersen CR, Carriqui M, Tonet V, Rodriguez Dominguez C, *et al.* 2021. Linking xylem network failure with leaf tissue death. New Phytol. 232: 68–79. DOI: 10.1111/nph.17577.

- Brodribb TJ, Powers J, Cochard H, Choat B. 2020. Hanging by a thread? Forests and drought. Science 368: 261–266. DOI: 10.1126/science.aat7631.
- Cai J, Tyree MT. 2010. The impact of vessel size on vulnerability curves: data and models for within-species variability in saplings of aspen, *Populus tremuloides* Michx. Plant Cell Environ. 33: 1059–1069. DOI: 10.1111/j.1365-3040.2010.02127.x.
- Carlquist S. 1966. Wood anatomy of Compositae: a summary, with comments on factors controlling wood evolution. Aliso 6: 25–44. DOI:10.5642/ALISO.19660602.03.
- Carlquist S. 1977. Ecological factors in wood evolution: a floristic approach. Am. J. Bot. 64: 887–896. DOI:10.2307/2442382.

Carlquist S. 1984. Vessel grouping in dicotyledon wood: significance and relationship to imperforate tracheary elements. Aliso 10: 505-525. DOI: 10.5642/aliso.19841004.03.

Carlquist S. 2001. Comparative wood anatomy. Springer, Berlin.

Charrier G, Charra-Vaskou K, Kasuga J, Cochard H, Mayr S, *et al.* 2014. Freeze-thaw stress: effects of temperature on hydraulic conductivity and ultrasonic activity in ten woody angiosperms. Plant Physiol. 164: 992–998. DOI: 10.1104/pp.113.228403.

Charrier G, Delzon S, Domec J-C, Zhang L, Delmas CEL, *et al.* 2018. Drought will not leave your glass empty: Low risk of hydraulic failure revealed by long-term drought observations in world's top wine regions. Sci. Adv. 4: eaao6969. DOI: 10.1126/sciadv.aao6969. Charrier G, Torres-Ruiz JM, Badel E, Burlett R, Choat B, *et al.* 2016. Evidence for hydraulic vulnerability segmentation and lack of xylem refilling under tension. Plant Physiol. 172: 1657–1668. DOI: 10.1104/pp.16.01079.

- Choat B, Badel E, Burlett R, Delzon S, Cochard H, *et al.* 2016. Noninvasive measurement of vulnerability to drought-induced embolism by X-ray microtomography. Plant Physiol. 170: 273–282. DOI: 10.1104/pp.15.00732.
- Choat B, Brodribb TJ, Brodersen CR, Duursma RA, Lopez R, *et al.* 2018. Triggers of tree mortality under drought. Nature 558: 531–539. DOI: 10.1038/s41586-018-0240-x.
- Choat B, Cobb AR, Jansen S. 2008. Structure and function of bordered pits: new discoveries and impacts on whole-plant hydraulic function. New Phytol. 177: 608–626. DOI: 10.1111/j.1469-8137.2007.02317.x.
- Choat B, Drayton WM, Brodersen C, Matthews MA, Shackel KA, *et al.* 2010. Measurement of vulnerability to water stress-induced cavitation in grapevine: a comparison of four techniques applied to long-vesseled species. Plant Cell Environ. 33: 1502–1512. DOI: 10.1111/j.1365-3040.2010.02160.x.

Christman MA, Sperry JS, Adler FR. 2009. Testing the 'rare pit' hypothesis for xylem

10.1111/j.1469-8137.2009.02776.x.

Christman MA, Sperry JS, Smith DD. 2012. Rare pits, large vessels and extreme vulnerability to cavitation in a ring-porous tree species. New Phytol. 193: 713–720. DOI:

cavitation resistance in three species of Acer. New Phytol. 182: 664–674. DOI:

10.1111/j.1469-8137.2011.03984.x.

Cochard H. 2002. A technique for measuring xylem hydraulic conductance under high negative pressures. Plant Cell Environ. 25: 815–819. DOI: 10.1046/j.1365-3040.2002.00863.x. Cochard H, Badel E, Herbette S, Delzon S, Choat B, et al. 2013. Methods for measuring plant vulnerability to cavitation: a critical review. J. Exp. Bot. 64: 4779–4791. DOI: 10.1093/jxb/ert193.

Cochard H, Delzon S. 2013. Hydraulic failure and repair are not routine in trees. Ann. For. Sci. 70: 659-661. DOI: 10.1007/s13595-013-0317-5.

Cochard H, Herbette S, Barigah T, Vilagrosa A. 2010. Does sample length influence the shape of vulnerability to cavitation curves? A test with the Cavitron spinning technique. Plant Cell Environ. 33: 1543-1552. DOI: 10.1111/j.1365-3040.2010.02163.x.

Creek D, Blackman CJ, Brodribb TJ, Choat B, Tissue DT. 2018. Coordination between leaf, stem, and root hydraulics and gas exchange in three arid-zone angiosperms during severe drought and recovery. Plant Cell Environ. 41: 2869–2881. DOI:

10.1111/pce.13418.

Dalla-Salda G, Fernández MA, Sergeant A-S, Rozenberg P, Badel E, Martinez-Meier A. 2014. Dynamics of cavitation in a Douglas-fir tree-ring: transition-wood, the lord of the ring? J. Plant Hydr. 1: e-0005. DOI: 10.20870/jph.2014.e005.

Davis SD, Sperry JS, Hacke UG. 1999. The relationship between xylem conduit diameter and cavitation caused by freezing. Am. J. Bot. 86: 1367–1372. DOI: 10.2307/2656919.

Davis SD, Ewers FW, Sperry JS, Portwood KA, Crocker MC, et al. 2002. Shoot dieback during prolonged drought in Ceanothus (Rhamnaceae) chaparral of California: a possible case of hydraulic failure. Am. J. Bot. 89: 820–828. DOI: 10.3732/ajb.89.5.820.

Domec J-C, Scholz FG, Bucci SJ, Meinzer FC, Goldstein G, et al. 2006. Diurnal and seasonal variation in root xylem embolism in neotropical savanna woody species: impact on stomatal control of plant water status. Plant Cell Environ. 29: 26–35. DOI:

10.1111/j.1365-3040.2005.01397.x.

Dória LC, Meijs C, Podadera DS, del Arco M, Smets E, *et al.* 2019. Embolism resistance in stems of herbaceous Brassicaceae and Asteraceae is linked with differences in woodiness and precipitation. Ann. Bot. 124: 1–13. DOI: 10.1093/aob/mcy233.
Dória LC, Podadera DS, Arco M, Chauvin T, Smets E, *et al.* 2018. Insular woody daisies (*Argyranthemum*, Asteraceae) are more resistant to drought- induced hydraulic failure than their herbaceous relatives. Funct. Ecol. 32: 1467–1478. DOI:

10.1111/1365-2435.13085.

- Emilio T, Lamarque LJ, Torres-Ruiz JM, King A, Charrier G, *et al.* 2019. Embolism resistance in petioles and leaflets of palms. Ann. Bot. 124: 1173–1183. DOI: 10.1093/aob/mcz104.
- Ewers FW, Jacobsen AL, López-Portillo J. 2023. Carlquist's indices for vulnerability and mesomorphy of wood: are they relevant today? IAWA J. DOI 10.1163/22941932-bja10113.
- Frankiewicz KE, Babasiak L, Oskolski AA, Magee AR, Alsarraf M, Trzeciak P, Spalik K. 2021. Derived woodiness and annual habit evolved in African umbellifers as alternative solutions for coping with drought. BMC Plant Biol. 21: 383. DOI: 10.1186/s12870-021-03151-x.
- Gauthey A, Peters JMR, Carins-Murphy MR, Rodriguez-Dominguez CM, Li X, *et al.* 2020. Visual and hydraulic techniques produce similar estimates of cavitation resistance in woody species. New Phytol. *228*: 884–897. DOI: 10.1111/nph.16746.

Gleason SM, Westoby M, Jansen S, Choat B, Hacke UG, *et al.* 2016. Weak tradeoff between xylem safety and xylem-specific hydraulic efficiency across the world's woody plant species. New Phytol. 209: 123–136. DOI: 10.1111/nph.13646.

Greenwood S, Ruiz-Benito P, Martinez-Vilalta J, Lloret F, Kitzberger T, *et al.* 2017. Tree mortality across biomes is promoted by drought intensity, lower wood density and higher specific leaf area. Ecol. Lett. 20: 539–553. DOI: 10.1111/ele.12748.

Guan X, Pereira L, McAdam SAM, Cao K-F, Jansen. 2021. No gas source, no problem: Proximity to pre-existing embolism and segmentation affect embolism spreading in angiosperm xylem by gas diffusion. Plant Cell Environ. 44: 1329–1345. DOI: 10.1111/pce.14016.

Guan X, Werner J, Cao K-F, Pereira L, Kaack L, *et al.* 2022. Stem and leaf xylem of angiosperm trees experiences minimal embolism in temperate forests during two consecutive summers with moderate drought. Plant Biol. 24: 1208–1223.

DOI:10.1111/plb.13384.

- Hacke UG, Jacobsen AL, Pratt RB. 2009. Xylem function of arid-land shrubs from California, USA: an ecological and evolutionary analysis. Plant Cell Environ. 32: 1324–1333. DOI: 10.1111/j.1365-3040.2009.02000.x.
- Hacke UG, Jacobsen AL, Pratt RB. 2022. Vessel diameter and vulnerability to droughtinduced embolism: within-tissue and across-species patterns and the issue of survivorship bias. IAWA J. DOI: 10.1163/22941932-bja10107.
- Hacke UG, Sperry JS, Wheeler JK, Castro L. 2006. Scaling of angiosperm xylem structure with safety and efficiency. Tree Physiol. 26: 689–701. DOI: 10.1093/treephys/26.6.689.
  Hacke UG, Sperry JS, Pockman WT, Davis SD, McCulloh KA. 2001. Trends in wood density and structure are linked to prevention of xylem implosion by negative pressure. Oecologia 126: 457–461. DOI: 10.1007/s004420100628.

Hacke UG, Spicer R, Schreiber SG, Plavcová L. 2017. An ecophysiological and developmental perspective on variation in vessel diameter. Plant Cell Environ. 40: 831–845. DOI: 10.1111/pce.12777.

- Hacke UG, Venturas MD, MacKinnon ED, Jacobsen AL, Sperry JS, *et al.* 2014. The standard centrifuge method accurately measures vulnerability curves of long-vesselled olive stems. New Phytol. 205: 116–127. DOI: 10.1111/nph.13017.
- Hargrave KR, Kolb KJ, Ewers FW, Davis SD. 1994. Conduit diameter and drought-induced embolism in *Salvia mellifera* Greene (Labiatae). New Phytol. 126: 695–705. DOI: 10.1111/j.1469-8137.1994.tb02964.x.
- Hölttä T, Vesala T, Perämäki M, Nikinmaa E. 2002. Relationships between embolism, stem water tension, and diameter changes. J. Theor. Biol. 215: 23–38. DOI: 10.1006/jtbi.2001.2485.
- Ingram S, Salmon Y, Lintunen, Hölttä T, Vesala T, *et al.* 2021. Dynamic surface tension enhances the stability of nanobubbles in xylem sap. Front. Plant Sci. 12: 732701. DOI: 10.3389/fpls.2021.732701.
- Isasa E, Link RM, Jansen S, Tezeh FR, Kaack L, *et al.* 2023. Addressing controversies in the xylem embolism resistance vessel diameter relationship. New Phytol. 238: 283–296. DOI: 10.1111/nph.18731.
- Jacobsen AL, Pratt RB. 2012. No evidence for an open vessel effect in centrifuge-based vulnerability curves of a long-vesselled liana (*Vitis vinifera*). New Phytol. 194: 982–990. DOI: 10.1111/j.1469-8137.2012.04118.x.
- Jacobsen AL, Pratt RB. 2023. Vessel diameter polymorphism determines vulnerability-toembolism curve shape. IAWA J. DOI: 10.1163/22941932-bja10115

Jacobsen AL, Pratt, RB, Davis SD, Ewers FW. 2007. Cavitation resistance and seasonal hydraulics differ among three arid Californian plant communities. Plant Cell Environ. 30: 1599–1609. DOI: 10.1111/j.1365-3040.2007.01729.x.

Jacobsen AL, Pratt RB, Venturas MD, Hacke UG. 2019. Large volume vessels are vulnerable to water-stress-induced embolism in stems of poplar. IAWA J. 40: 4–22. DOI: 10.1163/22941932-40190233.

- Jansen S, Bittencourt P, Pereira L, Schenk HJ, Kunert N. 2022. A crucial phase in plants it's a gas, gas, gas! New Phytol. 233: 1556–1559. DOI: 10.1111/nph.17875.
- Jansen S, Choat B, Pletsers A. 2009. Morphological variation of intervessel pit membranes and implications to xylem function in angiosperms. Am. J. Bot. 96: 409–419. DOI: 10.3732/ajb.0800248.
- Jansen S, Pletsers A, Sano Y. 2008. The effect of preparation techniques on SEM-imaging of pit membranes. IAWA J. 29: 161–178. DOI: 10.1163/22941932-90000177.
- Johnson DM, Wortemann R, McCulloh KA, Jordan-Meille L, Ward E, *et al.* 2016. A test of the hydraulic vulnerability segmentation hypothesis in angiosperm and conifer tree species. Tree Physiol. 36: 983– 993. DOI: 10.1093/treephys/tpw031.
- Johnson KM, Brodribb TJ. 2023. Evidence for a trade-off between growth rate and xylem cavitation resistance in *Callitris rhomboidea*. Tree Physiol. 43: 1055–1065. DOI: 10.1093/treephys/tpad037.

Kaack L, Altaner CM, Carmesin C, Diaz A, Holler M, *et al*. 2019. Function and threedimensional structure of intervessel pit membranes in angiosperms: a review. IAWA
J. 40: 673–702. DOI: 10.1163/22941932-40190259.

- Kaack L, Weber M, Isasa E, Karimi Z, Li S, *et al.* 2021. Pore constrictions in intervessel pit membranes provide a mechanistic explanation for xylem embolism resistance in angiosperms. New Phytol. 230: 1829–1843. DOI: 10.1111/nph.17282.
- Klepsch M, Zhang Y, Kotowska MM, Lamarque LJ, Nolf M, *et al.* 2018. Is xylem of angiosperm leaves less resistant to embolism than branches? Insights from microCT, hydraulics, and anatomy. J. Exp. Bot. 69: 5611–5623. DOI: 10.1093/jxb/ery321.
- Lachenbruch B, McColluh KA. 2014. Traits, properties, and performance: how woody plants combine hydraulic and mechanical functions in a cell, tissue, or whole plant. New Phytol. 204: 747–764. DOI: 10.1111/nph.13035.
- Lens F, Gleason SM, Bortolami G, Brodersen C, Delzon S, *et al.* 2022. Functional xylem characteristics associated with drought-induced embolism in angiosperms. New Phytol. 236: 2019–2036. DOI: 10.1111/nph.18447.
- Lens F, Luteyn JL, Smets E, Jansen S. 2004. Ecological wood anatomy of Vaccinioideae (Ericaceae *s.l.*). Flora 199: 309–319. DOI: 10.1078/0367-2530-0058.
- Lens F, Picon-Cochard C, Delmas C, Signarbieux C, Buttler A, *et al.* 2016. Herbaceous angiosperms are not more vulnerable to drought-induced embolism than angiosperm trees. Plant Physiol. 172: 661–667. DOI: 10.1104/pp.16.00829.
- Lens F, Sperry JS, Christman MA, Choat B, Rabaey D, *et al.* 2011. Testing hypotheses that link wood anatomy to cavitation resistance and hydraulic conductivity in the genus *Acer*. New Phytol. 190: 709–723. DOI: 10.1111/j.1469-8137.2010.03518.x.
- Lens F, Tixier A, Cochard H, Sperry JS, Jansen S, *et al.* 2013. Embolism resistance as a key mechanism to understand adaptive plant strategies. Curr. Opin. Plant Biol. 16: 287–292. DOI: 10.1016/j.pbi.2013.02.005.

Levionnois S, Jansen S, Wandji RT, Beauchêne J, Ziegler C, *et al.* 2020a. Linking droughtinduced xylem embolism resistance to wood anatomical traits in Neotropical trees. New Phytol. 229: 1453–1466. DOI: 10.1111/nph.16942.

- Levionnois S, Ziegler C, Jansen S, Calvet E, Coste S, *et al.* 2020b. Vulnerability and hydraulic segmentations at the stem–leaf transition: Coordination across Neotropical trees. New Phytol. 228: 512–524. DOI: 10.1111/nph.16723.
- Li S, Klepsch M, Jansen S, Schmitt M, Lens F, *et al.* 2016. Intervessel pit membrane thickness as a key determinant of embolism resistance in angiosperm xylem. IAWA J. 37: 152– 171. DOI: 10.1163/22941932-20160128.
- Liang X, Ye Q, Lio H, Brodribb TJ. 2021. Wood density predicts mortality threshold for diverse trees. New Phytol. 229: 3053–3057. DOI: 10.1111/nph.17117.
- Lintunen A, Hölttä T, Kulmala M. 2013. Anatomical regulation of ice nucleation and cavitation helps trees to survive freezing and drought stress. Sci. Rep. 3: 2031. DOI: 10.1038/srep02031.
- Lintunen A, Losso A, Aalto J, Chan T, Hölttä T, *et al.* 2020. Propagating ice front induces gas bursts and ultrasonic acoustic emissions from freezing xylem. Tree Physiol. 40: 170– 182. DOI: 10.1093/treephys/tpz123.
- Loepfe L, Martinez-Vilalta J, Pinol J, Mencuccini M. 2007. The relevance of xylem network structure for plant hydraulic efficiency and safety. J. Theor. Biol. 247: 788–803. DOI: 10.1016/j.jtbi.2007.03.036.
- Lohse D, Zhang X. 2015. Surface nanobubbles and nanodroplets. Rev. Mod. Phys. 87: 981-1035. DOI: 10.1103/RevModPhys.87.981.
- Lübbe T, Lamarque LJ, Delzon S, Torres Ruiz JM, Burlett R, *et al.* 2022. High variation in hydraulic efficiency but not xylem safety between roots and branches in four

temperate broad-leaved tree species. Funct. Ecol. 36: 699–712. DOI: 10.1111/1365-2435.13975.

- Mantova M, Herbette S, Cochard H, Torres-Ruiz JM. 2022. Hydraulic failure and tree mortality: from correlation to causation. Trends Plant Sci. 27: 335–345. DOI: 10.1016/j.tplants.2021.10.003.
- Martin-StPaul N, Delzon S, Cochard H. 2017. Plant resistance to drought depends on timely stomatal closure. Ecol. Lett. 20: 1437–1447. DOI: 10.1111/ele.12851.
- Martin-StPaul N, Longepierre D, Huc R, Delzon S, Burtlett R, *et al.* 2014. How reliable are methods to assess xylem vulnerability to cavitation? The issue of 'open vessel' artifact in oaks. Tree Physiol. 34: 894–905. DOI: 10.1093/treephys/tpu059.
- McCully M, Canny M, Baker A, Miller C. 2014. Some properties of the walls of metaxylem vessels of maize roots, including tests of the wettability of their lumenal wall surfaces. Ann. Bot. 113: 977–989. DOI: 10.1093/aob/mcu020.
- McDowell N, Pockman WT, Allen CD, Breshears DD, Cobb N, *et al.* 2008. Mechanisms of plant survival and mortality during drought: why do some plants survive while others succumb to drought? New Phytol. 178: 719–739. DOI: 10.1111/j.1469-

8137.2008.02436.x.

McDowell N, Sapes G, Pivovaroff A, Adams HD, Allen GD, *et al.* 2022. Mechanisms of woody-plant mortality under rising drought, CO2 and vapour pressure deficit. Nature Rev. Earth & Environm. 3: 294–308. DOI: 10.1038/ s43017-022-00272-1.

McElrone AJ, Brodersen CR, Alsina MM, Drayton WM, Matthews MA, et al. 2012. Centrifuge technique consistently overestimates vulnerability to water stress-induced cavitation in grapevines as confirmed with high-resolution computed tomography. New Phytol. 196: 661–665. DOI: 10.1111/j.1469-8137.2012.04244.x.

Mrad A, Domec J-C, Huang C-W, Lens F, Katul G. 2018. A network model links wood anatomy to xylem hydraulic behaviour. Plant Cell Environ. 41: 2718–2730. DOI: 10.1111/pce.13415.

Mrad A, Johnson DM, Love DM, Domec J-C. 2021. The roles of conduit redundancy and connectivity in xylem hydraulic functions. New Phytol. 231: 996–1007. DOI: 10.1111/nph.17429.

Nardini A, Lo Gullo MA, Salleo S. 2011. Refilling embolized xylem conduits: Is it a matter of phloem unloading? Plant Sci. 180: 604–611. DOI: 10.1016/j.plantsci.2010.12.011.

Nardini A, Savi T, Losso A, Petit G, Pacile S, et al. 2017. X-ray microtomography observations of xylem embolism in stems of Laurus nobilis are consistent with hydraulic measurements of percentage loss of conductance. New Phytol. 213: 1068–1075. DOI: 10.1111/nph.14245.

- Olson ME. 2020. From Carlquist's ecological wood anatomy to Carlquist's Law: why comparative anatomy is crucial for functional xylem biology. Am. J. Bot. 107: 1328-1341. DOI: 10.1002/ajb2.1552.
- Olson ME, Anfodillo T, Gleason SM, McCulloh KA. 2020. Tip-to-base xylem conduit widening as an adaptation: causes, consequences, and empirical priorities. New Phytol. 229: 1877-1893. DOI: 10.1111/nph.16961.

Olson ME, Pace MR, Anfodillo T. 2023. The vulnerability to drought-induced embolismconduit diameter link: breaching the anatomy-physiology divide. IAWA J. DOI: 10.1163/22941932-bja10123

Olson ME, Soriano D, Rosell JA, Anfodillo T, Donoghue MJ, et al. 2018. Plant height and hydraulic vulnerability to drought and cold. PNAS 115: 7551–7556. DOI:

10.1073/pnas.1721728115.

Percolla MI, Fickle JC, Rodríguez-Zaccaro FD, Pratt RB, Jacobsen AL. 2021. Hydraulic function and conduit structure in the xylem of five oak species. IAWA J. 42: 279–298. DOI: 10.1163/22941932-bja10059.

Pereira L, Domingues-Junior AP, Jansen S, Choat B, Marrafera P. 2018. Is embolism resistance in plant xylem associated with quantity and characteristics of lignin? Trees 32: 349–358. DOI: 10.1007/s00468-017-1574-y.

Pereira L, Miranda MT, Pires GS, Pacheco VS, Guan X, *et al.* 2020. A semi-automated method for measuring xylem vessel length distribution. Theor. Exp. Plant Phys. 32: 331–340. DOI: 10.1111/pce.13647.

Pivovaroff AL, Pasquini SC, De Guzman ME, Alstad KP, Stemke JS, Santiago LS. 2015. Multiple strategies for drought survival among woody plant species. Funct. Ecol. 30: 517–526. DOI: 10.1111/1365-2435.12518.

- Pivovaroff AL, Sack L, Santiago LS. 2014. Coordination of stem and leaf hydraulic conductance in southern California shrubs: a test of the hydraulic segmentation hypothesis. New Phytol. 203: 842–850. DOI: 10.1111/nph.12850.
- Pratt RB, Castro V, Fickle JC, Madsen A, Jacobsen AL. 2020. Factors controlling drought resistance in grapevine (*Vitis vinifera*, chardonnay): application of a new micro CT method to assess functional embolism resistance. Am. J. Bot. 107: 618–627. DOI: 10.1002/ajb2.1450.

Salleo S, Lo Gullo MA, Trifilo P, Nardini A. 2004. New evidence for a role of vessel-associated cells and phloem in the rapid xylem refilling of cavitated stems of *Laurus nobilis* L. Plant Cell Environ. 27: 1065–1076. DOI: 10.1111/j.1365-3040.2004.01211.x.

Schenk HJ, Espino S, Rich-Cavazos SM, Jansen S. 2018. From the sap's perspective: The nature of vessel surfaces in angiosperm xylem. Am. J. Bot. 105: 174–187. DOI: 10.1002/ajb2.1034.

- Schenk HJ, Espino S, Romo DM, Nima N, Do AYT, *et al.* 2017. Xylem surfactants introduce a new element to the cohesion-tension theory. Plant Physiol. 173: 1177–1196. DOI: 10.1104/pp.16.01039.
- Schenk HJ, Espino S, Visser A, Esser BK. 2016. Dissolved atmospheric gas in xylem sap measured with membrane inlet mass spectrometry. Plant Cell Environ. 39: 944–950. DOI: 10.1111/pce.12678.
- Schenk HJ, Michaud JM, Espino S, Melendres T, Roth MR, et al. 2020. Lipids in xylem sap of woody plants across the angiosperm phylogeny. Plant J. 105: 1477–1494. DOI: 10.1111/tpj.15125.
- Schenk HJ, Steppe K, Jansen S. 2015. Nanobubbles: a new paradigm for air-seeding in xylem. *Trends Plant Sci.* 20: 199–205. DOI: 10.1016/j.tplants.2015.01.008.
- Scholz S, Klepsch M, Karimi Z, Jansen S. 2013. How to quantify conduits in wood? Front. Plant Sci. 4: article 56. DOI: 10.3389/fpls.2013.00056.
- Sevanto S, Holbrook NM, Ball MC. 2012. Freeze/thaw-induced embolism: probability of critical bubble formation depends on speed of ice formation. Front. Plant Sci. 3: 107. DOI: 10.3389/fpls.2012.00107.
- Skelton RP, Anderegg LDL, Diaz J, Kling M, Papper P, *et al.* 2021. Evolutionary relationships between drought-related traits and climate shape large hydraulic safety margins in western North American oaks. PNAS 118: e2008987118. DOI:
  - 10.1073/pnas.2008987118.

Skelton RP, Anderegg LDL, Papper P, Reich E, Dawson TE, et al. 2019. No local adaptation in leaf or stem xylem vulnerability to embolism, but consistent vulnerability segmentation in a North American oak. New Phytol. 223: 1296–1306. DOI: 10.1111/nph.15886.

- Skelton RP, Dawson TE, Thompson SE, Shen Y, Weitz AP, *et al.* 2018. Low vulnerability to xylem embolism in leaves and stems of North American oaks. Plant Physiol. 177: 1066–1077. DOI: 10.1104/pp.18.00103.
- Smith-Martin CM, Skelton RP, Johnson KM, Lucanu C, Brodribb TJ. 2020. Lack of vulnerability segmentation among woody species in a diverse dry sclerophyll woodland community. Funct. Ecol. 34: 777–787. DOI: 10.1111/1365-2435.13519.
- Sorek Y, Greenstein S, Netzer Y, Shtein I, Jansen S, *et al.* 2020. An increase in xylem embolism resistance of grapevine leaves during the growing season is coordinated with stomatal regulation, turgor loss point and intervessel pit membranes. New Phytol. 229: 1955–1969. DOI: 10.1111/nph.17025.
- Sperry JS, Hacke UG, Pittermann J. 2006. Size and function in conifer tracheids and angiosperm vessels. Am. J. Bot. 93: 1490–1500. DOI: 10.3732/ajb.93.10.1490.
- Sperry JS, Christman MA, Torres-Ruiz JM, Taneda H, Smith DD. 2012. Vulnerability curves by centrifugation: is there an open vessel artefact, and are 'r' shaped curves necessarily invalid? Plant Cell Environ. 35: 601–610. DOI: 10.1111/j.1365-3040.2011.02439.x.

Sperry JS, Sullivan JE. 1992. Xylem embolism in response to freeze-thaw cycles and water stress in ring-porous, diffuse-porous, and conifer species. Plant Physiol. 100: 605– 613.

Stovall AEL, Shugart H, Yang X. 2019. Tree height explains mortality risk during an intense drought. Nat. Commun. 10: 43854. DOI: 10.1038/s41467-019-12380-6.

Thonglim A, Bortolami G, Delzon S, Larter M, Offringa R, *et al.* 2022. Drought response in *Arabidopsis* displays synergistic coordination between stems and leaves. J. Exp. Bot. 74: 1004–1021. DOI 10.1093/jxb/erac446.

Thonglim A, Delzon S, Larter M, Karami O, Rahimi A, *et al.* 2020. Intervessel pit membrane thickness best explains variation in embolism resistance amongst stems of *Arabidopsis thaliana* accessions. Ann. Bot. 128: 171–182. DOI:

10.1093/aob/mcaa196.

Tobin MF, Pratt RB, Jacobsen AL, De Guzman ME. 2012. Xylem vulnerability to cavitation can be accurately characterised in species with long vessels using a centrifuge method. Plant Biol. 15: 496–504. DOI: 10.1111/j.1438-8677.2012.00678.x.

Torres-Ruiz JM, Cochard H, Mayr S, Beikircher B, Diaz-Espejo A, *et al.* 2014. Vulnerability to cavitation in *Olea europaea* current-year shoots: further evidence of an open-vessel artefact associated with centrifuge and air-injection techniques. Physiol. Plant. 152: 465–474. DOI: 10.1111/ppl.12185.

Tyree MT, Davis SD, Cochard H. 1994. Biophysical perspectives of xylem evolution: is there a tradeoff of hydraulic efficiency for vulnerability to dysfunction? IAWA J. 15: 335–360. DOI: 10.1163/22941932-90001369.

Venturas MD, MacKinnon ED, Dario HL, Jacobsen AL, Pratt RB, *et al.* 2016. Chaparral shrub hydraulic traits, size, and life history types relate to species mortality during California's historic drought of 2014. PloS ONE 11: e0159145. DOI:

10.1371/journal.pone.0159145.

Venturas MD, Pratt RB, Jacobsen AL, Castro V, Fickle JC, *et al.* 2019. Direct comparison of four methods to construct xylem vulnerability curves: Differences among techniques

are linked to vessel network characteristics. Plant Cell Environ. 42: 2422–2436. DOI: 10.1111/pce.13565.

Wang Y, Pan R, Tyree MT. 2015. Studies on the tempo of bubble formation in recently cavitated vessels: A model to predict the pressure of air bubbles. Plant Physiol. 168: 521–531. DOI: 10.1104/pp.114.256602.

- Wason JW, Anstreicher KS, Stephansky N, Huggett BA, Brodersen CR. 2018. Hydraulic safety margins and air-seeding thresholds in roots, trunks, branches and petioles of four northern hardwood trees. New Phytol. 219: 77–88. DOI: 10.1111/nph.15135.
- Wason JW, Bouda M, Lee EF, McElrone AJ, Phillips RJ, *et al.* 2021. Xylem network connectivity and embolism spread in grapevine (*Vitis vinifera* L.). Plant Physiol. 186: 373–387. DOI: 10/1093/plphys/kiab045.
- Weishaupt K, Koch T, Helmig R. 2022. A fully implicit coupled pore-network/free-flow model for the pore-scale simulation of drying processes. Drying Techn. 40: 697–718. DOI: 10.1080/07373937.2021.1955706.
- Weithmann G, Link RM, Banzragch BE, Würzberg L, Leuschner C, *et al.* 2022. Soil water availability and branch age explain variability in xylem safety of European beech in Central Europe. Oecologia 198: 629–644. DOI: 10.1007/s00442-022-05124-9.
- Wheeler JK, Sperry JS, Hacke UG, Hoang N. 2005. Inter-vessel pitting and cavitation in woody Rosaceae and other vesselled plants: a basis for a safety versus efficiency trade-off in xylem transport. Plant Cell Environ. 28: 800–812. DOI: 10.1111/j.1365-3040.2005.01330.x.
- Wu M, Zhang Y, Oya T, Marcati CR, Pereira L, *et al.* 2020. Root xylem in three woody angiosperm species is not more vulnerable to embolism than stem xylem. Plant Soil 450: 479–495. DOI: 10.1007/s11104-020-04525-0.

Yang J, Michaud JM, Jansen S, Schenk HJ, Zuo YY. 2020. Dynamic surface tension of xylem sap lipids. Tree Physiol. 40: 433–444. DOI: 10.1093/treephys/tpaa006.

- Zhang Y, Carmesin C, Kaack L, Klepsch MM, Kotowska M, et al. 2020. High porosity with tiny pore constrictions and unbending pathways characterize the 3D structure of intervessel pit membranes in angiosperm xylem. Plant Cell Environ. 43: 116–130. DOI: 10.1111/pce.13654.
- Zhu S-D, Liu H, Xu Q-Y, Cao K-F, Ye Q. 2016. Are leaves more vulnerable to cavitation than branches? Funct. Ecol. 30: 1740–1744. DOI: 10.1111/1365-2435.12656.

Zimmermann MH. 1983. Xylem structure and the ascent of sap. Springer-Verlag, Berlin-Heidelberg.

Zizka A, Onstein RE, Rozzi R, Weigelt P, Kreft H, *et al.* 2022. The evolution of insular woodiness. PNAS 119: e2208629119. DOI: 10.1073/pnas.2208629119.

Zwieniecki MA, Holbrook NM. 2000. Bordered pit structure and vessel wall surface properties: implications for embolism repair. Plant Physiol. 123: 1015–1020. DOI: 10.1104/pp.123.3.1015.