



HAL
open science

Training Family Medicine Residents in Dermoscopy Using an e-Learning Course: Pilot Interventional Study

Pauline Friche, Lionel Moulis, Aurélie Du Thanh, Olivier Dereure, Claire
Duflos, Francois Carbonnel

► **To cite this version:**

Pauline Friche, Lionel Moulis, Aurélie Du Thanh, Olivier Dereure, Claire Duflos, et al.. Training Family Medicine Residents in Dermoscopy Using an e-Learning Course: Pilot Interventional Study. JMIR Formative Research, 2024, 8, pp.e56005. 10.2196/56005 . hal-04602241

HAL Id: hal-04602241

<https://hal.science/hal-04602241>

Submitted on 5 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Original Paper

Training Family Medicine Residents in Dermoscopy Using an e-Learning Course: Pilot Interventional Study

Pauline Friche¹, MD; Lionel Moulis^{2,3}, MD, PhD; Aurélie Du Thanh^{3,4,5}, MD, PhD; Olivier Dereure^{4,5}, MD, PhD; Claire Duflos^{2,6}, MD, PhD; Francois Carbonnel^{1,7,8}, MD, PhD

¹University Department of Family Medicine, University of Montpellier, Montpellier, France

²Clinical Research and Epidemiology Unit, Department of Public Health, Montpellier University Hospital, Montpellier, France

³Pathogenesis and Control of Chronic and Emerging Infections, University of Montpellier, Institut national de la santé et de la recherche médicale, Etablissement français du sang, University of Antilles, Montpellier, France

⁴Department of Dermatology, Montpellier University Hospital, Montpellier, France

⁵Department of Dermatology, University of Montpellier, Montpellier, France

⁶Department of Public Health, University of Montpellier, Montpellier, France

⁷Desbrest Institute of Epidemiology and Public Health, Unité Mixte de Recherche, Unité d'accueil 11, University of Montpellier, Institut national de la santé et de la recherche médicale, Montpellier, France

⁸University Multiprofessional Health Center Avicenne, Montpellier, France

Corresponding Author:

Francois Carbonnel, MD, PhD

Desbrest Institute of Epidemiology and Public Health, Unité Mixte de Recherche, Unité d'accueil 11

University of Montpellier, Institut national de la santé et de la recherche médicale

Camps ADV, IURC

641 Avenue du Doyen Gaston Giraud

Montpellier, 34093

France

Phone: 33 684014834

Email: Francois.carbonnel@umontpellier.fr

Abstract

Background: Skin cancers are the most common group of cancers diagnosed worldwide. Aging and sun exposure increase their risk. The decline in the number of dermatologists is pushing the issue of dermatological screening back onto family doctors. Dermoscopy is an easy-to-use tool that increases the sensitivity of melanoma diagnosis by 60% to 90%, but its use is limited due to lack of training. The characteristics of “ideal” dermoscopy training have yet to be established. We created a Moodle (Moodle HQ)-based e-learning course to train family medicine residents in dermoscopy.

Objective: This study aimed to evaluate the evolution of dermoscopy knowledge among family doctors immediately and 1 and 3 months after e-learning training.

Methods: We conducted a prospective interventional study between April and November 2020 to evaluate an educational program intended for family medicine residents at the University of Montpellier-Nîmes, France. They were asked to complete an e-learning course consisting of 2 modules, with an assessment quiz repeated at 1 (M1) and 3 months (M3). The course was based on a 2-step algorithm, a method of dermoscopic analysis of pigmented skin lesions that is internationally accepted. The objectives of modules 1 and 2 were to differentiate melanocytic lesions from nonmelanocytic lesions and to precisely identify skin lesions by looking for dermoscopic morphological criteria specific to each lesion. Each module consisted of 15 questions with immediate feedback after each question.

Results: In total, 134 residents were included, and 66.4% (n=89) and 47% (n=63) of trainees fully participated in the evaluation of module 1 and module 2, respectively. This study showed a significant score improvement 3 months after the training course in 92.1% (n=82) of participants for module 1 and 87.3% (n=55) of participants for module 2 ($P<.001$). The majority of the participants expressed satisfaction (n=48, 90.6%) with the training course, and 96.3% (n=51) planned to use a dermatoscope in their future practice. Regarding final scores, the only variable that was statistically significant was the resident's initial scores ($P=.003$) for module 1. No measured variable was found to be associated with retention (midtraining or final evaluation) for module 2. Residents who had completed at least 1 dermatology rotation during medical school had significantly higher initial

scores in module 1 at M0 ($P=.03$). Residents who reported having completed at least 1 dermatology rotation during their family medicine training had a statistically significant higher score at M1 for module 1 and M3 for module 2 ($P=.01$ and $P=.001$).

Conclusions: The integration of an e-learning training course in dermoscopy into the curriculum of FM residents results in a significant improvement in their diagnosis skills and meets their expectations. Developing a program combining an e-learning course and face-to-face training for residents is likely to result in more frequent and effective dermoscopy use by family doctors.

(*JMIR Form Res* 2024;8:e56005) doi: [10.2196/56005](https://doi.org/10.2196/56005)

KEYWORDS

dermoscopy; dermatoscope; dermatoscopes; dermatological; skin; training; GP; family practitioner; family practitioners; family physician; family physicians; general practice; family medicine; primary health care; internship and residency; education; e-learning; eLearning; dermatology; resident; residency; intern; interns; internship; internships

Introduction

Skin cancers are the most common group of cancers diagnosed worldwide [1]. The number of new cases of cutaneous melanoma per year is expected to increase by more than 50% from 2020 to 2040 [1]. In the European Union, skin melanoma accounted for 4% of all new cancers diagnosed in 2020 and 1.3% of all deaths due to cancer [2]. Risk factors are individual (light phototype, age, male sex, personal history of skin cancer or predisposing skin lesions, and family history of melanoma) and environmental (prolonged sun exposure, artificial tanning, immunosuppression, and lower socioeconomic class) [3]. If detected early, these skin cancers have a good prognosis. Skin inspection is based on the ABCDE rule. It was the first dermoscopy algorithm used to differentiate benign from malignant melanocytic lesions [4]. Specifically, A stands for asymmetry—one half of the spot is unlike the other half. B stands for border—the spot has an irregular, scalloped, or poorly defined border. C stands for color—varying colors from one area to the next, such as shades of tan, brown or black, or areas of white, red, or blue. D stands for diameter, melanomas are usually greater than 6 mm but can be smaller. E is for evolving (the spot looks different from the rest or is changing in size, shape, or color). The ABCDE rule has a sensitivity ranging from 84% to 93% [3]. Dermoscopy relies on a handheld microscope and incident light to reveal images of the lower skin layers at magnifications of 10 to 100 times. It is said to increase the sensitivity of melanoma diagnosis by 60% to 90% [3].

In France, melanomas are traditionally diagnosed in 2 steps: a skin inspection by the family doctor (FD), during which the family doctor decides whether to refer the patient to a dermatologist, and a second stage carried out by a dermatologist, during which the decision is made whether to biopsy the lesion, based on skin inspection and dermoscopy. In total, 2981 dermatologists were active in 2023 in France (ie, 3.4 dermatologists per 100,000 inhabitants, with significant regional disparities) [5]. This is also the case for family doctors, but by 2025, their numbers could rise again. The number of dermatologists will continue to fall and is expected to reach 2.4 dermatologists per 100,000 inhabitants in 2030. Only 8% of French family doctors use a dermatoscope as part of their routine practice, compared with 40% in Australia, a country hit by a rise in melanoma [6,7]. The main obstacles limiting the use of dermatoscopes are the difficulty of accessing them in consultation and lack of training [8]. Proper training is even

more important as this technique offers nothing more than skin inspection alone if the user is untrained or inexperienced [9,10]. It also improves FDs' diagnostic accuracy and long-term performance, increases their confidence, reduces the propensity to refer benign lesions to a dermatologist, and improves the benign-to-malignant ratio of excised lesions [11].

The characteristics of “ideal” dermoscopy training have yet to be established [3]. They range from a 1-hour lecture to an intensive week-long course, often combined with digital tools [3]. It is acknowledged that e-learning is as effective as traditional teaching methods. However, there is still relatively little literature to attest to its general superiority [12]. The use of e-learning is becoming increasingly ubiquitous, especially among doctors who have recently graduated and since the COVID-19 pandemic [13,14]. Another advantage of e-learning is that it can be offered in asynchronous form, enabling participants to follow the course at their own pace. Incorporating a core training course in the family medicine (FM) specialty training program could enable these future family doctors not only to acquire dermoscopy knowledge and skills but also to become aware of the value of its use in their future practice. A survey of FM residency program directors in the United States revealed that only 6.8% of residents had 4 or more hours of dermoscopy training and asked for research to better understand how to facilitate dermoscopy training in family medicine residencies [15].

The Department of Faculty of Medicine at University of Montpellier-Nîmes designed an optional e-learning module to train second- and third-year family medicine residents in dermoscopy in 2020.

The primary objective of this study was to evaluate the evolution of dermoscopy knowledge among participants immediately after e-learning training and 1 and 3 months afterward. The secondary objectives were to determine whether sociodemographic characteristics or student satisfaction influenced e-learning results.

Methods

Study Design

This was a monocentric, nonrandomized, prospective interventional study carried out between April and November 2020. It was based on an assessment quiz hosted on Moodle (Moodle HQ), administered before (M0) and immediately after

asynchronous distance training, with the same quiz administered at 1 month (M1) and 3 months (M3) after training. This format is in line with the learning theory known as the forgetting and recovering hypothesis, according to which forgetting makes for better learning [16].

The percentages of progression and regression between M0 and M1, M1 and M3, and M0 and M3 were calculated. The primary end point was the score at different time points (M0, M1, and M3). Score progression or regression was calculated by comparing scores over time. Students progressed if the difference in scores was strictly greater than 0 and regressed if the difference in scores was strictly less than 0.

There were several secondary end points: factors associated with information retention at M1 and M3 for both modules, factors associated with scores, and factors associated with satisfaction.

Study Population

All family medicine residents who chose this optional course before April 2020 were included. The exclusion criteria were being a first-year resident (optional courses are offered only to the second or third year of the family medicine residency program). As there is an absence of an equivalent training course published in the literature, there was no basis for comparison for calculating the number of participants required. We chose to include the entire cohort of participants in this training.

Intervention Description

The e-learning course was created between December 2019 and March 2020 by PF, FC, ADT, and OD. The course was divided into 2 modules with different levels of difficulty, in accordance with the 2-step algorithm [17]. This is a method of dermoscopic analysis of pigmented skin lesions that has been internationally accepted since 2001 (Figure 1). In module 1, “Dermoscopy level 1,” participants learned to differentiate melanocytic lesions

from nonmelanocytic lesions. In module 2, “Dermoscopy level 2,” participants learned to precisely identify skin lesions by looking for dermoscopic morphological criteria specific to each lesion and to consider their specific management according to the nature of these lesions.

Each module consisted of 15 questions in the form of multiple choice, drag and drop, or short open-response questions. Each module was scored out of 15, with 1 point per question. Each question was set as a single attempt, with immediate feedback after each question. The feedback was accompanied by course notes (Figure 2). These modules served as assessment quizzes at all stages of the study. Each session lasted 15 minutes. Participants were given 3 weeks to complete each questionnaire. The images used in the clinical cases were anonymized and were freely provided by fellow dermatologists; the name of the doctor holding the image was cited.

Participants were asked to complete a prestudy survey: sex, year of family residency program, dermatology clinical rotation experience, locums in family medicine, dermatology training during medical school or family medicine residency, quality of dermatology training during family medicine residency, interest in dermatology, frequency of dermatological consultations, self-evaluation of dermatological skills and ability to use a dermatoscope. After completing the final post-test at M3, participants were invited to complete a satisfaction questionnaire. A “free comments” section was provided at the end of the questionnaire.

The course was originally designed to include a face-to-face workshop on using dermatoscopes and performing biopsies, following the e-learning modules, as part of a blended-learning approach [18]. However, due to the COVID-19 pandemic and the university shutdown, these workshops were canceled, and training videos were made available to participants upon completion of the satisfaction questionnaire.

Figure 1. The 2-step algorithm.





Figure 2. Sample multiple choice questions.

Question 7
Incorrect
Note de 0,00 sur 1,00
Marquer la question
Modifier la question

Voici les images dermatoscopiques des 3 lésions mentionnées dans la question précédente.
Selon vous, la lésion B est...



© A. Serres-Cousine

© A. Serres-Cousine

© A. Serres-Cousine

Veillez choisir une réponse :

- A. Un naevus atypique
- B. Un angiome
- C. Un mélanome
- D. Une kératose séborrhéique
- E. Un naevus dermique

✘

Votre réponse est incorrecte.

Cette lésion est parfaitement bénigne. Il s'agit d'un type de naevus très répandu et dont le risque de dégénérescence est exceptionnel, on parle de naevus « dermique » (ou parfois « tubéreux »).
Beige, rose ou brun, il présente parfois quelques poils. Il est toujours en relief voire carrément mûriforme (comme ci-dessus) ou verruqueux.
On dit qu'il est dermique car ses mélanocytes sont enfouis plus profondément, dans le derme. Son aspect peut augmenter en cas d'irritation locale (ex frottements sur un vêtement) avec l'âge ou les modifications hormonales (il se remplit de graisse ou change de pigmentation par exemple).

La lésion A correspondait à une kératose séborrhéique (KS) dont vous avez repéré les pseudo-comédons (ci-dessous)



© A. Serres-Cousine

Enfin la lésion C, avec son aspect polychromique et son réseau pigmentaire atypique correspond à un mélanome.

La réponse correcte est : Un naevus dermique

Statistical Analysis

Residents' sociodemographic characteristics were described using mean and SD for continuous variables and frequency and percentages for categorical ones. Residents' mean scores were compared regarding their characteristics using 1-tailed Student *t* tests. To evaluate the differences in scores between the different time points, Friedman tests were performed. To analyze the factors associated with knowledge retention, we used a

multivariate linear regression after checking the normality of the variable (score variations). Independent variables were sociodemographic and pathway data selected by stepwise procedures with an entry *P* of .20 and an exit *P* of .15.

The satisfaction score was the sum of two 4-item Likert scales assessing training evaluation and practical application. The satisfaction score theoretically ranged from 2 to 8. No linear regression was applicable because of its nonlinearity. Instead, we performed a median split and evaluated the factors associated

with a satisfaction score greater than or equal to 7. A logistic regression model was used in which the independent variables were the sociodemographic and academic variables and the scores at the different temporalities after the selection of the variables by a stepwise procedure with an entry P of .20 and an exit P of .15. For the free comments section, we analyzed the verbatim of the satisfaction response in free text manually, comment by comment. Postcodification was carried out using a lexical approach, and percentages of response modalities were calculated. All analyses were performed with SAS (version 9.4; SAS Institute) using 2-tailed statistical tests with α set at 5%.

Ethical Considerations

The need for consent was deemed unnecessary according to national regulations. Research into changes in practices brought about by training medical or paramedical staff for research purposes is considered noninterventional in France and therefore does not require an ethical opinion [19]. Still, the information provided at the start of the e-learning course indicated that, by taking part in the course, the participants were consenting to participate in this study. The images used in the course were anonymized. Participant data were also anonymized. They were not compensated and took part voluntarily in this program, which was valued as part of their general medicine curriculum.

Results

Participants

A flowchart details study participation (Figure 3). The study included a total of 134 residents. Eighty-nine (66.4%) and 63 (47%) trainees participated in the full evaluation (M0, M1, and M3) of module 1 and module 2, respectively. Their sociodemographic characteristics are detailed in Table 1.

Residents who reported having completed at least 1 dermatology rotation during medical school had significantly higher initial scores in module 1 at M0 than those who had not ($P=.03$). This was not the case for the other assessment periods or module 2. Residents who reported having completed at least 1 dermatology rotation during their FM training had a statistically significant higher score at M1 for module 1 and M3 for module 2 ($P=.01$ and $P=.001$). Residents who were interested in dermatology had a higher score on M1 for module 1 ($P=.01$). Residents who reported frequently seeing patients for dermatologic reasons had higher scores on M3 for module 1 ($P=.03$).

Regarding the final scores, the only variable that was statistically significant was the resident's initial scores ($P=.003$) for module 1. No measured variable was found to be associated with retention, either at midtraining or at the final evaluation for module 2.

Figure 3. Flowchart of study participation.




Table 1. The sociodemographic characteristics of the participants (n=134) and respective statistical comparisons for each module and assessment time point.

Sociodemographic characteristics	Participants, n (%)	Level 1 module score at each month, mean (SD)			Level 2 module score at each month, mean (SD)		
		0	1	2	0	1	2
Sex							
Male	43 (32.8)	10.85 (1.55)	12.63 (1.24)	13.31 (1.27)	8.23 (1.84)	10.12 (1.59)	11.18 (1.89)
Female	88 (67.2)	10.81 (1.82)	12.56 (1.43)	13.27 (1.35)	9.09 (2.13)	10.19 (2.03)	11.08 (1.97)
<i>P</i> value ^a		.89	.76	.88	.04	.85	.84
Year of FM^b residency program							
Second year	66 (50.4)	10.74 (1.85)	12.62 (1.48)	13.28 (1.25)	8.59 (2.29)	9.85 (1.91)	10.73 (2.15)
Third year	65 (49.7)	10.94 (1.41)	12.55 (1.26)	13.28 (1.4)	9.02 (1.84)	10.51 (1.85)	11.56 (1.57)
<i>P</i> value ^a		.49	.78	.98	.3	.12	.053
Hospital dermatology rotation during medical school							
Yes	50 (38.2)	11.21 (1.41)	12.85 (1.28)	13.41 (1.28)	9.22 (1.98)	10.40 (1.95)	11.29 (1.65)
No	81 (61.8)	10.61 (1.74)	12.40 (1.4)	13.19 (1.35)	9.56 (2.1)	10.03 (1.88)	10.99 (2.11)
<i>P</i> value ^a		.03	.07	.42	.11	.41	.48
6-month hospital dermatology rotation during FM residency							
Yes	18 (13.7)	11.44 (1.49)	13.18 (0.94)	13.68 (0.95)	9.6 (1.6)	11.03 (1.76)	12.52 (1.24)
No	113 (86.3)	10.74 (1.65)	12.48 (1.4)	13.21 (1.37)	8.67 (2.12)	10 (1.89)	10.88 (1.94)
<i>P</i> value ^a		.08	.01	.12	.054	.07	.001
Locums in FM during FM residency							
Yes	35 (26.9)	10.91 (1.58)	12.35 (1.22)	13.00 (1.52)	8.95 (2.05)	10.43 (1.66)	11.33 (1.14)
No	95 (72.5)	10.81 (1.68)	12.72 (1.35)	12.38 (1.23)	8.75 (2.1)	10.03 (1.98)	11.00 (2.1)
<i>P</i> value ^a		.77	.16	.26	.66	.38	.39
Use of the dermatoscope							
Acquired	26 (19.8)	10.83 (1.47)	12.71 (1.11)	13.54 (1.14)	8.81 (2.05)	10.2 (2.38)	11.43 (1.91)
Not acquired	105 (80.2)	10.84 (1.69)	12.55 (1.43)	13.21 (1.36)	8.82 (2.09)	10.16 (1.79)	11.03 (0.95)
<i>P</i> value ^a		.98	.53	.29	.98	.95	.48
Interest in dermatology							
Very interested	36 (27.5)	10.58 (1.74)	12.43 (1.34)	13.06 (1.58)	9.25 (2.72)	9.78 (1.56)	11.38 (1.77)
A little interested	80 (61.1)	10.91 (1.53)	12.82 (1.31)	13.38 (1.22)	8.74 (1.72)	10.29 (2.01)	11.22 (2.09)
Not really interested	15 (11.4)	11.98 (2.01)	11.63 (1.4)	13.27 (1.26)	8.29 (1.93)	10.26 (1.97)	10.38 (1.39)
Not interested at all	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
<i>P</i> value ^a		.50	.01	.63	.33	.62	.33
Frequency of dermatological consultations							
Often (at least 1 a day)	35 (26.7)	10.34 (1.75)	12.51 (1.64)	12.69 (1.59)	8.12 (2.46)	10.04 (1.48)	11.00 (1.25)
Regularly (at least 1 a week)	93 (71)	11.02 (1.58)	12.66 (1.24)	13.5 (1.13)	9.04 (1.86)	10.22 (2.07)	11.17 (2.19)
Seldom (less than 1 a week)	3 (2.3)	11.2 (1.14)	11.12 (1.04)	14.09 (0.83)	9.75 (3.42)	10.33 (2.12)	10.61 (1.25)
<i>P</i> value ^a		.11	.15	.02	.12	.92	.88
Dermatology skills							
Fairly comfortable	34 (26)	10.72 (1.43)	12.6 (1.28)	13.51 (0.99)	8.74 (1.87)	9.86 (1.81)	10.95 (2.15)

Sociodemographic characteristics	Participants, n (%)	Level 1 module score at each month, mean (SD)			Level 2 module score at each month, mean (SD)		
		0	1	2	0	1	2
In difficulties	86 (65.6)	10.82 (1.79)	12.64 (1.42)	13.23 (1.43)	8.94 (2.22)	10.47 (1.92)	11.28 (1.94)
Not comfortable at all	11 (8.4)	11.38 (0.85)	12.1 (1.25)	13.1 (1.29)	8.16 (1.28)	8.81 (1.44)	10.5 (1.51)
<i>P</i> value ^a		.5	.48	.65	.53	.06	.48
Evaluation of dermatology training in medical school							
Practice oriented	56 (42.8)	10.6 (1.67)	12.72 (1.36)	13.2 (1.1)	9.04 (2.11)	10.06 (1.84)	10.98 (2.22)
Not adapted to practice	75 (57.2)	11.02 (1.61)	12.47 (1.37)	13.34 (1.46)	8.66 (2.05)	10.26 (1.96)	11.20 (1.74)
<i>P</i> value ^a		.16	.33	.61	.37	.64	.65
Evaluation of dermatology training during residency							
Practice-oriented	18 (13.7)	10.46 (1.52)	12.22 (1.62)	13.47 (1.07)	9.09 (1.78)	10.22 (2.14)	10.79 (2.34)
Not adapted to practice	113 (86.3)	10.9 (1.66)	12.64 (1.32)	13.25 (1.36)	8.78 (2.12)	10.16 (1.88)	11.16 (1.88)
<i>P</i> value ^a		.27	.30	.5	.57	.4	.2
Training satisfaction							
Very satisfactory	34 (64.2)	11.1 (1.82)	12.66 (1.34)	13.42 (1.37)	9.25 (1.99)	10.44 (1.49)	11.93 (1.59)
Fairly satisfactory	14 (26.4)	11.27 (1.06)	11.67 (1.57)	13.39 (1.29)	8.44 (2.24)	11.59 (1.34)	10.56 (2.2)
Rather unsatisfactory	3 (5.7)	10.53 (0.63)	13.01 (0.34)	12.59 (2.48)	7.92 (1.86)	7.35 (0.49)	8.92 (1.09)
Very unsatisfactory	2 (3.8)	10.92 (2.71)	11.30 (0.53)	13.34 (0.23)	6.44 (1.18)	9.17 (3.54)	10.8 (0.99)
<i>P</i> value ^a		.92	.54	.87	.19	.01	.01

^aItalic formatting indicates significant differences among groups.

^bFM: family medicine.


Score Evolution

Overall, while we observed some significant differences according to the participants' profiles at some time points, there was no clear and consistent pattern on the effect of the collected characteristics on their performance. Nevertheless, the residents'

scores improved significantly with increasing assessment time (from 0 to 1 month, 1 month to 3 months, and 0 to 3 months), which is noteworthy (Table 2; Figure 4). In addition, scores increased for up to 92% (n=82) of students for the level 1 module ($P<.001$) and up to 87% (n=55) for the level 2 module ($P<.001$; Table 2; Figure 4).

Table 2. Changes in score among residents over time.

Characteristics	Residents with changed scores, n (%)		
	Between months 0 and 1	Between months 1 and 3	Between months 0 and 3
Module 1			
Increased	73 (82)	64 (71.9)	82 (92.1)
Decreased	15 (16.9)	22 (25.7)	7 (7.9)
<i>P</i> value	<.001	<.001	<.001
Module 2			
Increased	49 (77.8)	43 (68.2)	55 (87.3)
Decreased	14 (22.2)	16 (25.4)	8 (12.7)
<i>P</i> value	<.001	<.001	<.001

Figure 4. Mean (SD) scores for each e-learning training module over time (assessments at months 0, 1, and 3).

Satisfaction

Among all participants, 53 (39.6%) trainees filled out the satisfaction questionnaire (Table 1). Using the Likert scale, 90.6% (n=48) found the training very or fairly satisfactory in terms of meeting their expectations. The participants most satisfied with the training had significantly better results at M1 and M3 for module 2.

Twenty-eight (20.9%) trainees answered in the free comments section, including 2 noninformative remarks (“Nothing to report” and “Thank you”). In the end, we analyzed 26 (19.4%) free comments. The top 3 requests made by students in this section were to (1) undertake a face-to-face course to consolidate the e-learning skills (n=14, 26.4%), (2) provide access to ready-made summary sheets (organized by lesion type) and tutorials to learn how to use the equipment (n=6, 11.3%), and (3) provide access to an image bank for training purposes (n=2, 3.8%).

Discussion

Principal Findings

The e-learning training course was effective in improving the dermoscopy skills of family medicine residents. Three-month scores greatly improved in 92.1% (n=82) and 87.3% (n=55) of participants for module 1 and module 2, respectively. Dermoscopy training studies often focus on melanoma detection and are expected to improve sensitivity between 13% and 15%, with no improvement in specificity [2]. In the case of multitopic courses such as this study, a published 2-day multitopic dermoscopy training course (melanoma, angioma, seborrheic keratosis, and cutaneous carcinoma) for primary care physicians improved sensitivity by 23.6% and specificity by 21% [20]. Existing short workshops with repeated training sessions (such as this study) showed an improvement in posttraining scores (+21.3%), which correlated with the number of repeated web-based training sessions [21]. They involved a small number of participants (n=27) compared with our 134 participants.

This study also provided some interesting insights about the dermoscopy training of family medicine residents. The ones who reported knowing how to use a dermatoscope before

training did not perform any better than others. This is consistent with the fact that providing a dermatoscope without training to all family doctors would not improve skin cancer screening [2,8,10]. Students interested in dermatology and those who had completed a dermatology rotation during their medical school (which probably overlapped) performed better on the pretest of the first module. This difference disappeared after training, suggesting that their dermoscopy input exceeded their previous dermoscopy knowledge. Residents who had completed (or were in the process of completing) a dermatology rotation during their residency also performed better on the assessment at M1 of module 1 and M3 of module 2. In this study, we did not differentiate between residents who were currently completing a dermatology rotation and those who had completed one, nor did we differentiate between the type of rotation (inpatient or outpatient). Subject to further study, the lack of superiority of these students in the pretest may be interpreted as a need for time and practice beyond their rotation experience. As for their superiority at M1, but only for module 1, and at M3 for module 2 (the more complex of the 2 modules), this could show that their rotation potentiates their long-term results and confirms the interest of mixed training. We hypothesized that a replacement activity in family medicine would have a positive effect on their results as consultations related to the cutaneous system are common, estimated at 5.5% of family medicine consultations [22]. Indeed, those who reported seeing patients frequently for dermatologic causes had better results in module 1. This may reflect a higher level of investment and interest of these residents, who may be more aware of local needs.

Most of the participants (n=48, 90.6%) were satisfied with the course and 98.1% (n=52) would recommend it to their peers. Their confidence also increased over time. At the end of the course, 88.7% (n=47) declared that their dermatology skills had improved, which is consistent with the literature [20]. These excellent results are even more important since the level of satisfaction had an impact on our participants' scores. This process is thus part of a virtuous cycle: the more satisfied students are with a course, the more time and attention they will devote to it, resulting in more effective learning and skill acquisition. One of our most interesting results was that 96.2% (n=51) of participants said they would consider using a dermatoscope in their future practice, closely reflecting the

literature [3]. Practicing general practitioners have shared that they are prepared to commit up to 7 days of their personal time to training in dermatoscopy [11].

To our knowledge, this is the first study to focus on dermatoscopy training for FM residents. It also benefited from an excellent satisfaction rate, although it had been conducted in the context of the COVID-19 pandemic thanks to the use of e-learning.

Limitations

We had to make some decisions that introduced potential bias. Recruitment was voluntary, a source of selection bias. Participants were generally the most interested in the subject and the most committed, which led to better results [23]. We also excluded first-year residents, as this training is optional and therefore only accessible to second- and third-year family medicine residents at Montpellier-Nîmes Faculty of Medicine. Finally, the study was monocentric.

Although the e-learning format was well accepted by the students, there was a significant dropout rate of 33.6% (45/134) at 3 months after the start of the level 1 module and 52.9% (71/134) at 3 months after the start of the level 2 module. This phenomenon could be explained by the fact that the 3-month assessment took place during the summer (and therefore the vacations of some) and by the choice to repeat the e-learning 3 times (which could be a source of fatigue, as expressed by some comments to the study and a source of memorization bias). However, it should be noted that loss of focus rates in dermatoscopy training is in the order of 40% or more [19,24].

Perspectives

The decline over time in the diagnostic performance in dermatoscopy among family doctors is well known and could be also noted for family medicine residents [11]. Several digital tools could be deployed for our training purposes. The use of digital portfolios in medical education could guide the development of educational content adapted to each student's initial level of knowledge, needs, and possible preferences, with a view to personalized e-learning. Adaptive e-learning could lead to an increase in student satisfaction, a progression adapted to their level and preferences, and an overall improvement in their skills by encouraging them to work on topics that they find difficult [25]. This should be considered cautiously as the residents' self-assessed level of competence in dermatology in this study had no effect on their results. It will be necessary to go beyond simple self-assessment in this area to guide their education. Smartphone apps based on training modules, such as Youdermoscopy, could serve as diagnostic aid tools and help increase the user's knowledge of dermatology [26]. This course could also be a step toward the establishment of a local network of physicians practicing dermatoscopy. It is 1 of the 25 practical recommendations in primary care dermatoscopy [27]. Family doctors need to be supported in their daily practice, especially through training and the establishment of local networks (eg, in the context of coordinated practice based on teledermoscopy) [28-30]. In the short term, this could help our patients and our overburdened health care system in dermatology. In the medium term, digital tools supporting dermatologic diagnosis are expected to play a major role in benefiting patients, once the challenges for identifying dermoscopic features have been overcome [31,32].

Acknowledgments

The authors thank the teachers and residents who took part in this training course. We would also like to thank Dr Anne Serres-Cousine, MD, who helped building our clinical pictures database, and Prof Ralph Braun, MD, for permitting us to use the resources of the website Dermoscopyedia [33]. The authors especially thank Maha Baddredine, head of the Pedagogical Engineering and Audiovisual Production Department, for her help in developing the e-learning.

Data Availability

The data sets generated and analyzed during this study are available from the corresponding author on reasonable request.

Conflicts of Interest

None declared.

References

1. Arnold M, Singh D, Laversanne M, Vignat J, Vaccarella S, Meheus F, et al. Global burden of cutaneous melanoma in 2020 and projections to 2040. *JAMA Dermatol.* 2022;158(5):495-503. [FREE Full text] [doi: [10.1001/jamadermatol.2022.0160](https://doi.org/10.1001/jamadermatol.2022.0160)] [Medline: [35353115](https://pubmed.ncbi.nlm.nih.gov/35353115/)]
2. Koczkodaj P, Sulkowska U, Didkowska J, Rutkowski P, Mańczuk M. Melanoma mortality trends in 28 European countries: a retrospective analysis for the years 1960-2020. *Cancers (Basel).* 2023;15(5):1514. [FREE Full text] [doi: [10.3390/cancers15051514](https://doi.org/10.3390/cancers15051514)] [Medline: [36900305](https://pubmed.ncbi.nlm.nih.gov/36900305/)]
3. Dinnes J, Deeks JJ, Chuchu N, di Ruffano LF, Matin RN, Thomson DR, et al. Dermoscopy, with and without visual inspection, for diagnosing melanoma in adults. *Cochrane Database Syst Rev.* 2018;12(12):CD011902. [FREE Full text] [doi: [10.1002/14651858.CD011902.pub2](https://doi.org/10.1002/14651858.CD011902.pub2)] [Medline: [30521682](https://pubmed.ncbi.nlm.nih.gov/30521682/)]
4. Argenziano G, Fabbrocini G, Carli P, De Giorgi V, Sammarco E, Delfino M. Epiluminescence microscopy for the diagnosis of doubtful melanocytic skin lesions. Comparison of the ABCD rule of dermatoscopy and a new 7-point checklist based

- on pattern analysis. *Arch Dermatol*. 1998;134(12):1563-1570. [FREE Full text] [doi: [10.1001/archderm.134.12.1563](https://doi.org/10.1001/archderm.134.12.1563)] [Medline: [9875194](https://pubmed.ncbi.nlm.nih.gov/9875194/)]
5. Approche territoriale des spécialités médicales et chirurgicales. Situation au premier janvier 2023 [Territorial approach to medical and surgical specialties. Situation on 1 January 2023]. Conseil national de l'Ordre des médecins [French National Medical Council]. URL: https://www.conseil-national.medecin.fr/sites/default/files/external-package/analyse_etude/egnnt2/cnom_atlas_demographie_2023_approche_territoriale_des_specialites.pdf [accessed 2024-02-05]
 6. Chappuis P, Duru G, Marchal O, Girier P, Dalle S, Thomas L. Dermoscopy, a useful tool for general practitioners in melanoma screening: a nationwide survey. *Br J Dermatol*. 2016;175(4):744-750. [doi: [10.1111/bjd.14495](https://doi.org/10.1111/bjd.14495)] [Medline: [26914613](https://pubmed.ncbi.nlm.nih.gov/26914613/)]
 7. Youl PH, Raasch BA, Janda M, Aitken JF. The effect of an educational programme to improve the skills of general practitioners in diagnosing melanocytic/pigmented lesions. *Clin Exp Dermatol*. 2007;32(4):365-370. [doi: [10.1111/j.1365-2230.2007.02414.x](https://doi.org/10.1111/j.1365-2230.2007.02414.x)] [Medline: [17433042](https://pubmed.ncbi.nlm.nih.gov/17433042/)]
 8. Williams NM, Marghoob AA, Seiverling E, Usatine R, Tsang D, Jaimes N. Perspectives on dermoscopy in the primary care setting. *J Am Board Fam Med*. 2020;33(6):1022-1024. [FREE Full text] [doi: [10.3122/jabfm.2020.06.200238](https://doi.org/10.3122/jabfm.2020.06.200238)] [Medline: [33219084](https://pubmed.ncbi.nlm.nih.gov/33219084/)]
 9. Binder M, Puespoeck-Schwarz M, Steiner A, Kittler H, Muellner M, Wolff K, et al. Epiluminescence microscopy of small pigmented skin lesions: short-term formal training improves the diagnostic performance of dermatologists. *J Am Acad Dermatol*. 1997;36(2 Pt 1):197-202. [doi: [10.1016/s0190-9622\(97\)70280-9](https://doi.org/10.1016/s0190-9622(97)70280-9)] [Medline: [9039168](https://pubmed.ncbi.nlm.nih.gov/9039168/)]
 10. Kittler H, Pehamberger H, Wolff K, Binder M. Diagnostic accuracy of dermoscopy. *Lancet Oncol*. 2002;3(3):159-165. [doi: [10.1016/s1470-2045\(02\)00679-4](https://doi.org/10.1016/s1470-2045(02)00679-4)] [Medline: [11902502](https://pubmed.ncbi.nlm.nih.gov/11902502/)]
 11. Harkemanne E, Baeck M, Tromme I. Training general practitioners in melanoma diagnosis: a scoping review of the literature. *BMJ Open*. 2021;11(3):e043926. [FREE Full text] [doi: [10.1136/bmjopen-2020-043926](https://doi.org/10.1136/bmjopen-2020-043926)] [Medline: [33757946](https://pubmed.ncbi.nlm.nih.gov/33757946/)]
 12. Vaona A, Banzi R, Kwag KH, Rigon G, Cereda D, Pecoraro V, et al. E-learning for health professionals. *Cochrane Database Syst Rev*. 2018;1(1):CD011736. [FREE Full text] [doi: [10.1002/14651858.CD011736.pub2](https://doi.org/10.1002/14651858.CD011736.pub2)] [Medline: [29355907](https://pubmed.ncbi.nlm.nih.gov/29355907/)]
 13. Gormley GJ, Collins K, Boohan M, Bickle IC, Stevenson M. Is there a place for e-learning in clinical skills? A survey of undergraduate medical students' experiences and attitudes. *Med Teach*. 2009;31(1):e6-e12. [FREE Full text] [doi: [10.1080/01421590802334317](https://doi.org/10.1080/01421590802334317)] [Medline: [19253150](https://pubmed.ncbi.nlm.nih.gov/19253150/)]
 14. Reinhart A, Malzkorn B, Döing C, Beyer I, Jünger J, Bosse HM. Undergraduate medical education amid COVID-19: a qualitative analysis of enablers and barriers to acquiring competencies in distant learning using focus groups. *Med Educ Online*. 2021;26(1):1940765. [FREE Full text] [doi: [10.1080/10872981.2021.1940765](https://doi.org/10.1080/10872981.2021.1940765)] [Medline: [34128776](https://pubmed.ncbi.nlm.nih.gov/34128776/)]
 15. Lu MD, Usatine R, Holt J, Harper DM, Verdieck A. Dermoscopy training in family medicine residencies: a CERA survey of program directors. *Fam Med*. 2023;55(4):259-262. [FREE Full text] [doi: [10.22454/FamMed.2023.368813](https://doi.org/10.22454/FamMed.2023.368813)] [Medline: [37043187](https://pubmed.ncbi.nlm.nih.gov/37043187/)]
 16. Bjork RA, Bjork EL. Forgetting as the friend of learning: implications for teaching and self-regulated learning. *Adv Physiol Educ*. 2019;43(2):164-167. [FREE Full text] [doi: [10.1152/advan.00001.2019](https://doi.org/10.1152/advan.00001.2019)] [Medline: [30998108](https://pubmed.ncbi.nlm.nih.gov/30998108/)]
 17. Marghoob AA, Braun R. Proposal for a revised 2-step algorithm for the classification of lesions of the skin using dermoscopy. *Arch Dermatol*. 2010;146(4):426-428. [doi: [10.1001/archdermatol.2010.41](https://doi.org/10.1001/archdermatol.2010.41)] [Medline: [20404234](https://pubmed.ncbi.nlm.nih.gov/20404234/)]
 18. Vallée A, Blacher J, Cariou A, Sorbets E. Blended learning compared to traditional learning in medical education: systematic review and meta-analysis. *J Med Internet Res*. 2020;22(8):e16504. [FREE Full text] [doi: [10.2196/16504](https://doi.org/10.2196/16504)] [Medline: [32773378](https://pubmed.ncbi.nlm.nih.gov/32773378/)]
 19. Protection des personnes en matière de santé. Recherche impliquant la personne humaine [Protection of individuals in health matters. Research involving the human person]. Code de la santé publique de la république française [Public health code of the French republic]. URL: https://www.legifrance.gouv.fr/codes/section_lc/LEGITEXT000006072665/LEGISCTA00006154978/ [accessed 2024-03-15]
 20. De Bedout V, Williams NM, Muñoz AM, Londoño AM, Munera M, Naranjo N, et al. Skin cancer and dermoscopy training for primary care physicians: a pilot study. *Dermatol Pract Concept*. 2021;11(1):e2021145. [FREE Full text] [doi: [10.5826/dpc.1101a145](https://doi.org/10.5826/dpc.1101a145)] [Medline: [33614219](https://pubmed.ncbi.nlm.nih.gov/33614219/)]
 21. Seiverling EV, Li D, Stevens K, Cyr P, Dorr G, Ahrns H. Distance learning and spaced review to complement dermoscopy training for primary care. *Dermatol Pract Concept*. 2021;11(2):e2021030. [FREE Full text] [doi: [10.5826/dpc.1102a30](https://doi.org/10.5826/dpc.1102a30)] [Medline: [33954013](https://pubmed.ncbi.nlm.nih.gov/33954013/)]
 22. Letrilliart L, Supper I, Schuers M, Darmon D, Boulet P, Favre M, et al. ECOGEN: étude des Éléments de la CONsultation en médecine GENérale [ECOGEN: study of the elements of consultation in family medicine]. *Exercer*. 2014;114:148-157. [FREE Full text]
 23. Carvalho H, West CA. Voluntary participation in an active learning exercise leads to a better understanding of physiology. *Adv Physiol Educ*. 2011;35(1):53-58. [FREE Full text] [doi: [10.1152/advan.00011.2010](https://doi.org/10.1152/advan.00011.2010)] [Medline: [21386002](https://pubmed.ncbi.nlm.nih.gov/21386002/)]
 24. Menzies SW, Emery J, Staples M, Davies S, McAvoy B, Fletcher J, et al. Impact of dermoscopy and short-term sequential digital dermoscopy imaging for the management of pigmented lesions in primary care: a sequential intervention trial. *Br J Dermatol*. 2009;161(6):1270-1277. [doi: [10.1111/j.1365-2133.2009.09374.x](https://doi.org/10.1111/j.1365-2133.2009.09374.x)] [Medline: [19747359](https://pubmed.ncbi.nlm.nih.gov/19747359/)]

25. Fontaine G, Cossette S, Maheu-Cadotte MA, Mailhot T, Deschênes MF, Mathieu-Dupuis G, et al. Efficacy of adaptive e-learning for health professionals and students: a systematic review and meta-analysis. *BMJ Open*. 2019;9(8):e025252. [FREE Full text] [doi: [10.1136/bmjopen-2018-025252](https://doi.org/10.1136/bmjopen-2018-025252)] [Medline: [31467045](https://pubmed.ncbi.nlm.nih.gov/31467045/)]
26. Glines KR, Haidari W, Ramani L, Akkurt ZM, Feldman SR. Digital future of dermatology. *Dermatol Online J*. 2020;26(10):13030/qt75p7q57j. [Medline: [33147661](https://pubmed.ncbi.nlm.nih.gov/33147661/)]
27. Chuh A, Zawar V, Fölster-Holst R, Sciallis G, Rosemann T. Twenty-five practical recommendations in primary care dermoscopy. *J Prim Health Care*. 2020;12(1):10-20. [doi: [10.1071/HC19057](https://doi.org/10.1071/HC19057)] [Medline: [32223845](https://pubmed.ncbi.nlm.nih.gov/32223845/)]
28. Jones OT, Jurascheck LC, Utukuri M, Pannebakker MM, Emery J, Walter FM. Dermoscopy use in UK primary care: a survey of GPs with a special interest in dermatology. *J Eur Acad Dermatol Venereol*. 2019;33(9):1706-1712. [FREE Full text] [doi: [10.1111/jdv.15614](https://doi.org/10.1111/jdv.15614)] [Medline: [30977937](https://pubmed.ncbi.nlm.nih.gov/30977937/)]
29. Bruce AF, Mallow JA, Theeke LA. The use of teledermoscopy in the accurate identification of cancerous skin lesions in the adult population: a systematic review. *J Telemed Telecare*. 2018;24(2):75-83. [FREE Full text] [doi: [10.1177/1357633X16686770](https://doi.org/10.1177/1357633X16686770)] [Medline: [28056600](https://pubmed.ncbi.nlm.nih.gov/28056600/)]
30. Chuchu N, Dinnes J, Takwoingi Y, Matin RN, Bayliss SE, Davenport C, et al. Teledermatology for diagnosing skin cancer in adults. *Cochrane Database Syst Rev*. 2018;12(12):CD013193. [FREE Full text] [doi: [10.1002/14651858.CD013193](https://doi.org/10.1002/14651858.CD013193)] [Medline: [30521686](https://pubmed.ncbi.nlm.nih.gov/30521686/)]
31. di Ruffano LF, Takwoingi Y, Dinnes J, Chuchu N, Bayliss SE, Davenport C, et al. Computer-assisted diagnosis techniques (dermoscopy and spectroscopy-based) for diagnosing skin cancer in adults. *Cochrane Database Syst Rev*. 2018;12(12):CD013186. [FREE Full text] [doi: [10.1002/14651858.CD013186](https://doi.org/10.1002/14651858.CD013186)] [Medline: [30521691](https://pubmed.ncbi.nlm.nih.gov/30521691/)]
32. Kentley J, Weber J, Liopyris K, Braun RP, Marghoob AA, Quigley EA, et al. Agreement between experts and an untrained crowd for identifying dermoscopic features using a Gamified app: reader feasibility study. *JMIR Med Inform*. 2023;11:e38412. [FREE Full text] [doi: [10.2196/38412](https://doi.org/10.2196/38412)] [Medline: [36652282](https://pubmed.ncbi.nlm.nih.gov/36652282/)]
33. Dermoscopedia. URL: https://dermoscopedia.org/Main_Page [accessed 2024-04-22]

Abbreviations

FD: family doctor

FM: family medicine

Edited by A Mavragani; submitted 02.01.24; peer-reviewed by N Williams, J Kentley; comments to author 19.01.24; revised version received 05.02.24; accepted 20.03.24; published 13.05.24

Please cite as:

Friche P, Moulis L, Du Thanh A, Dereure O, Duflos C, Carbonnel F

Training Family Medicine Residents in Dermoscopy Using an e-Learning Course: Pilot Interventional Study

JMIR Form Res 2024;8:e56005

URL: <https://formative.jmir.org/2024/1/e56005>

doi: [10.2196/56005](https://doi.org/10.2196/56005)

PMID:

©Pauline Friche, Lionel Moulis, Aurélie Du Thanh, Olivier Dereure, Claire Duflos, Francois Carbonnel. Originally published in JMIR Formative Research (<https://formative.jmir.org>), 13.05.2024. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Formative Research, is properly cited. The complete bibliographic information, a link to the original publication on <https://formative.jmir.org>, as well as this copyright and license information must be included.