



**HAL**  
open science

## Mechanistic study of the differences in lactic acid bacteria resistance to freeze- or spray-drying and storage

Maite Gagnetten, Stéphanie Passot, Stéphanie Cenard, Sarrah Ghorbal, Carolina Schebor, Fernanda Fonseca

### ► To cite this version:

Maite Gagnetten, Stéphanie Passot, Stéphanie Cenard, Sarrah Ghorbal, Carolina Schebor, et al.. Mechanistic study of the differences in lactic acid bacteria resistance to freeze- or spray-drying and storage. *Applied Microbiology and Biotechnology*, 2024, 108 (1), pp.361. 10.1007/s00253-024-13186-3. hal-04607503

**HAL Id: hal-04607503**

**<https://hal.science/hal-04607503v1>**

Submitted on 10 Jun 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# Mechanistic study of the differences in lactic acid bacteria resistance to freeze- or spray-drying and storage

Maite Gagneten<sup>1</sup> · Stéphanie Passot<sup>2</sup> · Stéphanie Cenard<sup>2</sup> · Sarah Ghorbal<sup>2</sup> · Carolina Schebor<sup>1</sup> · Fernanda Fonseca<sup>2</sup>

Received: 16 January 2024 / Revised: 2 April 2024 / Accepted: 15 May 2024  
© The Author(s) 2024

## Abstract

*Lactobacillus delbrueckii* subsp. *bulgaricus* and *Lactiplantibacillus plantarum* are two lactic acid bacteria (LAB) widely used in the food industry. The objective of this work was to assess the resistance of these bacteria to freeze- and spray-drying and study the mechanisms involved in their loss of activity. The culturability and acidifying activity were measured to determine the specific acidifying activity, while membrane integrity was studied by flow cytometry. The glass transitions temperature and the water activity of the dried bacterial suspensions were also determined. Fourier transform infrared (FTIR) micro-spectroscopy was used to study the biochemical composition of cells in an aqueous environment. All experiments were performed after freezing, drying and storage at 4, 23 and 37 °C. The results showed that *Lb. bulgaricus* CFL1 was sensitive to osmotic, mechanical, and thermal stresses, while *Lpb. plantarum* WCFS1 tolerated better the first two types of stress but was more sensitive to thermal stress. Moreover, FTIR results suggested that the sensitivity of *Lb. bulgaricus* CFL1 to freeze-drying could be attributed to membrane and cell wall degradation, whereas changes in nucleic acids and proteins would be responsible of heat inactivation of both strains associated with spray-drying. According to the activation energy values (47–85 kJ/mol), the functionality loss during storage is a chemically limited reaction. Still, the physical properties of the glassy matrix played a fundamental role in the rates of loss of activity and showed that a glass transition temperature 40 °C above the storage temperature is needed to reach good preservation during storage.

## Key points

- Specific FTIR bands are proposed as markers of osmotic, mechanic and thermal stress
- *Lb. bulgaricus* CFL1 was sensitive to all three stresses, *Lpb. plantarum* WCFS1 to thermal stress only
- Activation energy revealed chemically limited reactions ruled the activity loss in storage

**Keywords** *Lb. bulgaricus* · *Lpb. plantarum* · Drying · FTIR spectroscopy · Glass transition temperature · Activation energy

## Introduction

Lactic acid bacteria (LAB) concentrates are widely used in the food and health industries to produce a variety of fermented products, and probiotics. Freezing is one of the

most employed LAB preservation techniques. However, it entails very low storage temperatures (commonly –80 °C), which require the right infrastructure and comprises high energy consumption (Broeckx et al. 2016; Pénicaud et al. 2018). Drying the LAB starter cultures is an interesting stabilization strategy to obtain products with longer shelf-life at higher, even non-sub-zero temperatures. Yet, the drying operation itself might cause damage and loss of viability in bacterial populations (Ermis 2021; Santivarangkna et al. 2008; Sehwat et al. 2022; Yoha et al. 2020a). Freeze-drying is the most commonly used drying method for LAB preservation since it has been shown to lead to higher survival rates than most other drying techniques (Ermis 2021; Kieps and Dembczyński 2022; Montel Mendoza et al. 2014; Sehwat et al. 2022; Yoha et al. 2020b). However, it is an

✉ Fernanda Fonseca  
fernanda.fonseca@inrae.fr

<sup>1</sup> Departamento de Industrias, Facultad de Ciencias Exactas y Naturales, ITAPROQ (UBA- CONICET), Universidad de Buenos Aires, Ciudad Autónoma de Buenos Aires, Argentina

<sup>2</sup> Université Paris-Saclay, INRAE, AgroParisTech, UMR SayFood, Palaiseau F-91120, France

energy-intensive, time-consuming, and expensive process (Champagne et al. 2012). Spray-drying is preferred in the food industry for its lower costs and higher productivity when compared to freeze-drying. However, the application of spray-drying to preserve LAB is still limited since some challenges to maintaining cell viability remain unsolved (Huang et al. 2017; Sharma et al. 2022).

During freeze-drying, the temperature decrease exposes cells to cold stress, while ice formation induces osmotic stress because of solute concentration. Mechanical stress is also present in the desorption stage when water is removed by breaking hydrogen bonds. On the other hand, during spray-drying, cells are also exposed to the osmotic and mechanical stresses related to water removal, and additional thermal stress as a consequence of the high operating temperatures employed (Liu et al. 2019; Santivarangkna et al. 2008). The osmotic stress, due to high solute concentration and loss of cell turgor, is known as a major source of cellular damage (Broeckx et al. 2016; Sehwat et al. 2022). The mechanical stress, associated with water removal, can cause loss of the lipid bilayer integrity and alteration of cell membrane functions (Perdana et al. 2013). The thermal stress due to exposure to high temperatures may cause protein denaturation and aggregation, and damage to essential structures such as ribosomes, nucleic acids, enzymes, and cell membranes (Arena et al. 2019; Capozzi et al. 2012; Perdana et al. 2013; Sehwat et al. 2022). Also, lipidic components of the cell's membrane are susceptible to oxidation, which can increase during spray-drying as a consequence of the large surface in contact with air at high temperatures (Broeckx et al. 2016). Therefore, since cell damage caused by stabilization processes can be considered a combination of these different stresses depending on the drying method (Kieps and Dembczyński 2022; Sehwat et al. 2022), identifying which kind of stress the different LAB are more sensitive to is essential and could contribute to design the most suitable dehydrating process accordingly.

To improve the cell's preservation during stabilization and storage, the addition of protective molecules to cell concentrates is applied. These molecules often include mixtures of sugars and polymers that contribute to maintaining cell integrity during freezing, drying and subsequent storage by two proposed mechanisms (Boafo et al. 2022; Crowe et al. 1988; Grasmeyer et al. 2013): (i) the water replacement theory and ii) the vitrification theory. The first one proposes that disaccharides (such as sucrose and trehalose) protect cells during drying by forming hydrogen bonds with membrane polar groups as water is removed, allowing the maintenance of a pseudo-hydrate structure. According to the second theory, the immobilization of the bacteria in a glassy matrix reduces molecular mobility and,

consequently, slows down degradation reactions during storage (Liu et al., 2019).

This work aimed to evaluate and compare different dehydration processes and storage conditions to preserve LAB and understand the mechanisms involved in their loss of activity. Using complementary analytical methods and, in particular, an original FTIR micro-spectroscopy method which enables the study of bacterial cells in aqueous conditions while avoiding dehydration that may affect cellular structures, this work seeks to contribute to identify the effect of freeze-drying and spray-drying over the specific cellular components. This FTIR method, previously used to investigate the impact of freezing and freeze-drying processes on LAB (Girardeau et al. 2022; Guerrero et al. 2022, 2023), was here applied for the first time to study the spray-drying process and compare it to freeze-drying. For this purpose, two LAB strains exhibiting different sensitivity to environmental conditions were stabilized with a protective solution containing maltodextrin, fructo-oligosaccharides (FOS) and an antioxidant, and dehydrated by freeze- or spray-drying. *Lactobacillus delbrueckii* subsp. *bulgaricus* CFL1 is known to be very sensitive to drying (Passot et al. 2012), while *Lactiplantibacillus plantarum* WCFS1 has shown more resistance to dehydration (Bensch et al. 2014; G-Alegría et al. 2004; Golowczyk et al. 2010). The FOS present in the protective mixture, not only provides prebiotic functions (Gibson et al. 2004; Rajam and Anandharamakrishnan 2015) but also contains small sugars that can replace water during drying (Romano et al. 2016), while maltodextrin contributes to increase the glass transition temperature of the matrix. To study the LAB biological activity, biochemical and physical characteristics a variety of techniques were applied. Their functional quality was assessed by culturability and acidifying activity measurements, and their membrane integrity by flow cytometry. FTIR micro-spectroscopy was applied to observe differences in the biochemical composition between both LAB and its changes during freeze- and spray-drying and storage. Differential scanning calorimetry and water activity measurements were used to determine the glass transition temperature and the thermodynamic activity of water, crucial parameters affecting the stability of dried products (Guerrero Sanchez et al. 2022, 2023; Roos 2010).

## Materials and methods

The experimental approach followed in this study and the main parameters analyzed are summarized in Fig. 1.

## Production of bacteria


### Bacterial strains and culture conditions

*Lactobacillus delbrueckii* subsp. *bulgaricus* CFL1 (CIRM-BIA; Rennes, France) and *Lactiplantibacillus plantarum* WCFS1 (NIZO Food Research B.V., The Netherlands) were chosen for this work due to their industrial relevance in the food industry.

Inocula were precultured twice before inoculating the bioreactor. The detailed information of preculture and fermentation conditions for each microorganism are provided elsewhere (<https://doi.org/10.57745/AAARUZ>).

The second preculture of *Lb. bulgaricus* CFL1 and *Lpb. plantarum* WCFS1 was used to inoculate a 20 L bioreactor containing 15 L of sterilized (121 °C for 20 min) MRS broth complemented with additional 20–40 g.L<sup>-1</sup> of glucose, respectively. The temperature was maintained at 42 °C for *Lb. bulgaricus* CFL1 and 37 °C *Lpb. plantarum* WCFS1. The pH was maintained at 5.8 with a NaOH solution (17 and 20% v/v, respectively). The consumption of NaOH was monitored and used to

calculate the acidifying rate. Cells were harvested by centrifugation (15,900 g, 10 min, 4 °C) one hour after the maximum acidifying rate was reached. Concentrated cells were re-suspended at 4 °C in a 1:2 cells: protective medium ratio (g/g). The protective medium consisted of 120 g.L<sup>-1</sup> of a commercial maltodextrin (Glucidex 6®, dextrose equivalent of 6, Roquette; Lestrem, France), 120 g.L<sup>-1</sup> of a commercial mixture of fructo-oligosaccharides (Orafti® P95, Degree of polymerization 2–8, Beneo Orafti; Tienen, Belgium), 20 g.L<sup>-1</sup> of sodium ascorbate (Royal DSM N.V.; Limburg, Netherland) and 9 g.L<sup>-1</sup> of NaCl. Two independent fermentation procedures were performed for each LAB to have two biological replicates for analysis. After resuspension in the protective medium, samples were divided into three parts: one part was directly frozen at -80 °C in Eppendorf sterile tubes to obtain the frozen samples, another one was placed in Petri dishes for freeze-drying (30 g per Petri dish, around 3–4 Petri dishes per formulation depending on the quantity of biomass obtained) and stored at -80 °C until drying and the third one was immediately spray-dried.



**Fig. 1** Diagram of the experimental approach and the main parameters investigated in this study. MD: maltodextrin, FOS: fructo-oligosaccharides

## Dehydration and storage

The spray-drying experiments were performed in a mini spray-dryer Büchi B-290 (Flawil, Switzerland). The inlet air temperature was 105 °C and the outlet air temperature 65 °C. Atomization was created by compressed air at a pressure of 6 bar and an air flux of 800 L.h<sup>-1</sup>. The nozzle diameter was 0.7 mm and the feed rate 0.156 L.h<sup>-1</sup>.

For the freeze-drying experiments, the protected bacterial concentrates frozen at -80 °C in sterile Petri dishes were transferred to pre-cooled shelves at -50 °C in a pilot-scale freeze-dryer (VirTis Genesis 35 L SQ EL-85, SP Scientific; Warminster, PA, USA). After a holding step of 1 h at -50 °C, the chamber pressure was decreased to 10 Pa and the shelf temperature was increased to -20 °C at 0.25 °C.min<sup>-1</sup> to initiate the sublimation phase. After 40 h of sublimation, the shelf temperature was increased to 25 °C at 0.25 °C.min<sup>-1</sup> to initiate the desorption phase that lasted 10 h.

The dried samples were packed in aluminium bags and stored at -80 °C. For the analysis, freeze-dried samples were milled to a powder in a chamber of very low relative humidity (around 5%). All samples were conditioned in 1.5 mL sterile Eppendorf tubes (about 0.2 g) or plastic cups (about 0.5 g), for the measurement of biological and physico-chemical properties, respectively. Storage of dried samples was performed inside vacuum sealed aluminium bags for 7, 14 and 28 days at 4, 23 and 37 °C.

## Biological studies

### Culturability

Frozen samples were thawed in a water bath at 37 °C for *Lpb. plantarum* WCFS1 and 42 °C for *Lb. bulgaricus* CFL1, and the dried samples were rehydrated in saline water (0.9% NaCl) to reach the same dry matter relation of the protected bacterial suspension before freeze-drying, and stirred gently for 5 min at ambient temperature. After serial dilutions, in-depth spread was performed in MRS agar (Biokar Diagnostics, France) and incubated for 24 h at 37 °C for *Lpb. plantarum* WCFS1 and 48 h at 42 °C under anaerobic conditions (GENbox96124, BioMérieux, Marcy l'Etoile, France) for *Lb. bulgaricus* CFL1 before counting. Plates containing between 30 and 300 colonies were kept for cell concentration evaluation. Results were expressed as colony forming units (CFU) in log(CFU.mL<sup>-1</sup>).

## Acidifying activity

Acidifying activity of the samples was measured using the Cinac System described by Meneghel et al. (2017) with some modifications. A volume of 100 µL of *Lb. bulgaricus* CFL1 or 10 µL of *Lpb. plantarum* WCFS1 cell suspensions or rehydrated samples were incubated in skim milk at 42 °C or in yeast extract (10 g.L<sup>-1</sup>) medium supplemented with glucose (10 g.L<sup>-1</sup>) at 37 °C for each bacteria strain, respectively. The time necessary to reach a pH drop of 1.5 points ( $t_{dpH=1.5}$ , in min) was used to characterize the acidifying activity of the samples. The longer the time, the lower the acidifying activity.

The acidifying activity results were then weighted by the culturability data to obtain the specific acidifying activity ( $t_{spe}$ , in min.(log(CFU.mL<sup>-1</sup>))<sup>-1</sup>) as shown in Eq. (1) according to Gautier et al. (2013).

$$t_{spe} = \frac{t_{dpH=1.5}}{\log CFU.mL^{-1}} \quad (1)$$

Culturability and acidifying activity were measured in the protected cells before and after freezing at -80 °C, after drying, and during storage of the dried samples after 7, 14 and 28 days. Since the statistical analysis demonstrated that no loss of culturability nor acidifying activity occurred because of the freezing operation (Table SM1), the frozen samples were used for further analyses and considered as the reference condition to evaluate the effect of the different stabilization and storage scenarios.

Then, the  $t_{spe}$  was plotted as a function of the storage time ( $t$ , in days) and the rate of loss of acidifying activity ( $k_{spe}$ , in min.(log(CFU.mL<sup>-1</sup>))<sup>-1</sup>.day<sup>-1</sup>) was determined from the slope of the curve for each storage temperature according to Eq. 2 .

$$t_{spe} = k_{spe} \times t + A \quad (2)$$

Subsequently, the loss rate was correlated with the inverse of the storage temperature (in K<sup>-1</sup>) and fitted to the Arrhenius equation following Eq. 3 :

$$k_{spe} = A \times e^{-E_a/RT} \quad (3)$$

Where A is a frequency factor (in days<sup>-1</sup>),  $E_a$  is the apparent energy of activation (in kJ.mol<sup>-1</sup>), R is the gas constant (8.314 J.K<sup>-1</sup>.mol<sup>-1</sup>), and T is the storage temperature (in K). From the slope of the linearization of Eq. 3 (Eq. 4) the  $E_a$  can be computed (Sosa et al. 2016).

$$\ln k_{spe} = -E_a/RT + \ln A \quad (4)$$



## Membrane integrity

Flow cytometry analyses with propidium iodide (PI) and carboxyfluorescein diacetate (cFDA) dual staining were carried out to quantify viable, injured, and dead cells. PI can only enter the cell and bind DNA if membrane damage has occurred, giving information on cell membrane integrity. cFDA only dyes cells that are still viable and gives information on intracellular esterase activity (Rault et al. 2007). A volume of 100  $\mu\text{L}$  of bacterial suspension (thawed or rehydrated cells) containing approximately  $10^7$  cells. $\text{mL}^{-1}$  was added to 900  $\mu\text{L}$  of McIlvaine's buffer (pH 7.4). The markers consisted of 10  $\mu\text{L}$  of Chemchrome V8 (Bio-Mérieux, Craonne, France) diluted to 1/10 (v/v) in acetone (Sigma Aldrich, Saint Quentin Fallavier, France) and 5  $\mu\text{L}$  of PI (10 mg PI, Sigma, diluted in 10 mL  $\text{H}_2\text{O}$  MilliQ), added to one millilitre of the diluted bacterial suspension containing approximately  $10^6$  cells. $\text{mL}^{-1}$ . The suspension was then incubated for 10 min at 40 °C before flow cytometry analysis.

The experiments were performed with a CyFlow Space cytometer (Sysmex-Partec, Villepinte, France) equipped with a solid blue laser, emitting at 488 nm and four optic filters, following the set-up described by Bouix et al. (2022). Data were collected with FloMax software (SYSMEX-PARTEC, Villepinte, France) and the numbers and percentages of stained cells determined by each detector were analyzed.

## Physical properties

### Water activity determination

The water activity ( $a_w$ ) of the dried samples was measured at 25 °C using a  $a_w$  meter labMasteraw (Novasina, Precisa, Poissy, France).

### Glass transition temperature

Glass transition temperatures were determined on the frozen cells ( $T_g'$ , in °C) as well as on the freeze- and spray-dried powders ( $T_g$ , in °C) before storage and after 28 days of storage by differential scanning calorimetry (DSC) following the calibration and measurement procedures described by Guerrero Sanchez et al. (2022) with slight modifications. A Diamond equipment with a liquid nitrogen cooling accessory (CryoFill, Perkin Elmer) was used to analyze the frozen samples that exhibited thermal events at negative temperatures; while a Pyris 1 equipment with a mechanical cooling system (Intracooler 1P, Perkin Elmer) was used for the dehydrated samples (both Perkin Elmer LLC, Norwalk,

CT, USA). Approximately 10 to 20 mg of each sample were placed in 50  $\mu\text{L}$  hermetically sealed aluminium pans. An empty pan was used as a reference. Linear heating and cooling rates of 10 °C. $\text{min}^{-1}$  were used. Freeze-thawed samples were cooled to -100 °C and then scanned up to 20 °C. For the dehydrated samples a first heating step from 5 to 110 °C was performed to erase their thermal history, followed by cooling to 5 °C and a second heating up to 130 °C. The glass transition temperature was determined from the maximum of the first derivative of the second heating curve.

## Biochemical properties by FTIR micro-spectroscopy

Frozen samples were thawed, and dried samples were rehydrated as described in previous sections. After thawing or rehydration, the suspensions were washed three times with saline water (0.9%) and centrifuged at 16,000 g for 10 min at 4 °C. Thermal source transmission FTIR micro-spectroscopy measurements were performed following the procedure described by Guerrero Sanchez et al. (2022) and Meneghel et al. (2020). A total of between 60 and 80 spectra were recorded for each sample.

### Pre-processing of raw spectra

Spectra showing Mie scattering or absorption features were sorted using the Omnic software. Then an automatic atmospheric subtraction was applied to all spectra to remove the residual contribution from water vapour and carbon dioxide.

### Procedure for water subtraction

The subtraction of bulk water contributions from spectra of bacteria was performed by using an in-house MATLAB script (version: 8.3.0.532, MathWorks, Natick, MA, USA) according to the procedure described by Meneghel et al. (2020). To calibrate the water removal program, the value of the amide I to amide II area ratio was used. The amide I area was calculated between 1727 and 1585  $\text{cm}^{-1}$  and the amide II between 1585 and 1481  $\text{cm}^{-1}$  with a baseline extending from 1727 to 1481  $\text{cm}^{-1}$ .

### Post-processing of spectra

Post-processing of the water-subtracted spectra was performed using the Unscrambler®X software package (Version 10.2, CAMO Software AS, Oslo, Norway). The analysis focused on three spectral regions: (1) from 3060 to 2800  $\text{cm}^{-1}$ , containing information on lipids, mainly fatty acyl chains of the bacterial membrane; (2) from 1800

to  $1370\text{ cm}^{-1}$  to recover information about proteins, particularly amide I and amide II bands; and (3) from  $1369$  to  $975\text{ cm}^{-1}$ , a complex region containing information on proteins (amide III band), nucleic acids, and cell wall components such as phosphorylated molecules and polysaccharides. Each spectra region was normalized and baseline was corrected using the extended multiplicative scatter correction (EMSC). The post-processed data was statistically analyzed by principal component analysis (PCA) to study the variation pattern by an unsupervised method using the Unscrambler®X software package.

To deepen the analysis, and according to Guerrero Sanchez et al. (2022, 2023), the second-order derivatives of spectra were calculated to resolve protein secondary structures, identify precise peak locations, and determine the relative variation ( $r_v$ ) of the peak's height in the dried sample compared to the frozen sample. The assignment of the principal absorption bands was performed using data from the literature.

## Statistical analysis

Nonparametric Kruskal-Wallis tests were performed over biological activity and the glass transition temperature measurements using XLSTAT 19.6 (Addinsoft, Paris, France). ANOVA analyses were carried out on biochemical composition determined by FTIR spectroscopy data. A significance level of 95% ( $p < 0.05$ ) was considered in all cases. Three technical replicates were performed for each biological replicate except for water activity.

## Results

### Biological activity

#### Culturability and acidifying activity

To study the impact of the dehydration operation on the biological activity of the LAB, the specific acidifying activity ( $t_{spe}$ ) was studied before and after drying (Table 1). An increase in  $t_{spe}$  was observed after drying for all the studied samples, as a result of the loss of culturability and decrease of the acidifying activity after this operation. Spray-drying caused a bigger increase in  $t_{spe}$  for both LAB strains than freeze-drying. Also, *Lb. bulgaricus* CFL1 presented bigger  $t_{spe}$  increases for each drying method than *Lpb. plantarum* WCFS1.

The progress of  $t_{spe}$  with the storage time for three storage temperatures is shown in Fig. 2. No significant losses were observed after 28 days of storage at  $4\text{ }^\circ\text{C}$ . At 23 and

**Table 1** Specific acidifying activity ( $t_{spe}$ ) (in  $\text{min} \cdot (\log(\text{CFU} \cdot \text{mL}^{-1}))^{-1}$ ) of the frozen, freeze-dried (FD) and spray-dried (SD) samples of *Lb. bulgaricus* CFL1 and *Lpb. plantarum* WCFS1 and the increase of  $\Delta t_{spe}$  after drying


	$t_{spe}$ frozen	$t_{spe}$ SD	$t_{spe}$ FD	$\Delta t_{spe}$ SD	$\Delta t_{spe}$ FD
CFL1	40.4 <sup>B</sup>	83 <sup>D</sup>	66 <sup>C</sup>	42 <sup>d</sup>	25 <sup>b</sup>
IQR	0.7	4	1	5	1
WCFS1	30 <sup>A</sup>	64 <sup>C</sup>	36.9 <sup>B</sup>	34 <sup>c</sup>	7 <sup>a</sup>
IQR	2	2	0.6	3	2

Data presented as median and interquartile ranges (IQR). Different superscript letters represent statistical differences between samples at a 95% confidence level: uppercase letter for absolute  $t_{spe}$  values comparison; lowercase letters for  $\Delta t_{spe}$  values comparison

$37\text{ }^\circ\text{C}$ , however, whatever the LAB and the drying method, a linear increase of  $t_{spe}$  with storage time was observed, following the relationship shown in Eq. 2. Linear relationships between the specific acidifying activity with storage time have also been described by other authors when analyzing different storage conditions (Passot et al. 2012; Streit et al. 2008). Table 2 shows the slopes of the linear regressions corresponding to specific acidifying activity loss rates ( $k_{spe}$ ) after 28 days of storage at each tested temperature for the four samples. The higher the  $k_{spe}$ , the faster the bacterial inactivation. Both spray-dried and freeze-dried samples of *Lpb. plantarum* WCFS1 showed smaller inactivation rates than *Lb. bulgaricus* CFL1 ones after 28 days of storage at the three temperatures. Also, the  $k_{spe}$  values of spray-dried samples were bigger than those for freeze-dried ones (except for samples stored at  $4\text{ }^\circ\text{C}$ , which did not exhibit significant losses). Additionally,  $k_{spe}$  values were correlated to the absolute storage temperature and fitted to the linearized Arrhenius equation (Eq. 4) to determine the activation energy  $E_{a_{spe}}$  of each sample (Fig. 3; Table 2). Although the activation energy is more commonly computed with the culturability loss rate ( $k$ ) rather than the specific acidifying activity loss rate ( $k_{spe}$ ), the activation energy values determined by this procedure were consistent with those found with the culturability results (Table SM2). Spray-dried cells of both LAB and freeze-dried ones of *Lb. bulgaricus* CFL1 showed similar activation energy values, while freeze-dried *Lpb. plantarum* WCFS1 samples showed the lowest activation energy.

### Membrane integrity and cell viability

Figure 4 shows the percentage of viable, viable and injured, and dead cells for the two LAB and the two drying methods



**Fig. 2** Evolution of the specific acidifying activity ( $t_{spe}$ ) of each biological replicate during 28 days storage at 4 °C (yellow diamonds), 23 °C (orange squares), and 37 °C (red circles). (a) *Lb. bulgaricus* CFL1 spray-dried sample, (b) *Lb. bulgaricus* CFL1 freeze-dried, (c)

*Lpb. plantarum* WCFS1 spray-dried samples and (d) *Lpb. plantarum* WCFS1 and freeze-dried samples. The regression coefficients  $R^2$  of the linear fit for the curves at 23 and 37 °C were higher than 0.75 in all cases

**Table 2** Specific acidifying activity loss rate ( $k_{spe}$ , in (min.(log (CFU.mL<sup>-1</sup>))<sup>-1</sup>.day<sup>-1</sup>) of freeze-dried (FD) and spray-dried (SD) samples of *Lb. bulgaricus* CFL1 and *Lpb. plantarum* WCFS1 during storage at 4, 23 and 37 °C and their corresponding activation energy ( $E_{a_{spe}}$ , in kJ/mol)

	$k_{spe}$ (min.(log (CFU.mL <sup>-1</sup> )) <sup>-1</sup> .day <sup>-1</sup> )			$E_{a_{spe}}$ (kJ/mol)	$R^2$
	4 °C	23 °C	37 °C		
CFL1 SD	0.04	0.9	6	105	0.999
IQR	0.00	0.1	1	6	
CFL1 FD	0.04	0.50	3.5	97	0.998
IQR	0.01	0.03	0.6	10	
WCFS1 SD	0.07	0.7	2.7	85	0.999
IQR	0.03	0.1	0.3	11	
WCFS1 FD	0.05	0.27	0.638	41	0.998
IQR	0.05	0.02	0.007	13	

Data presented as median and interquartile ranges (IQR). The correlation coefficient  $R^2$  informs on the quality of the linear regression fitting used to calculate  $E_{a_{spe}}$

before and after dehydration, and after 28 days of storage at the different temperatures. The results showed important losses of viability and membrane integrity for *Lb. bulgaricus* CFL1 after both drying methods (Fig. 4a), while *Lpb. plantarum* WCFS1 exhibited a higher tolerance to



the dehydration step (Fig. 4b). The spray-drying operation caused the death of more than 60% and 19% of *Lb. bulgaricus* CFL1 and *Lpb. plantarum* WCFS1 populations, respectively; and it increased the percentage of injured cells to 32% and 17% for each bacteria strain, respectively. In contrast, freeze-drying maintained a much higher proportion of viable and not injured *Lpb. plantarum* WCFS1 cells (93%). Storage at 23 and 37 °C amplified the damages caused after drying, increasing the percentage of dead and injured cells, except for the *Lpb. plantarum* WCFS1 freeze-dried samples, that maintained a low proportion of dead cells after 28 days of storage at 23 °C (5%) and even at 37 °C (17%).

**Physical properties**

Table 3 shows the glass transition temperature of the maximally concentrated phase ( $T_g'$ ) of the frozen samples and the glass transition temperature ( $T_g$ ) and water activity ( $a_w$ ) of the dehydrated samples before storage. As expected, no significant differences were observed between the  $T_g'$  of the two frozen bacterial suspensions since a unique protective solution, and in the same proportion was used for both samples. Freeze-dried samples showed half lower  $a_w$  values than spray-dried ones, which resulted in higher  $T_g$  values. Also, *Lpb. plantarum* WCFS1 dried samples exhibited a lower



**Fig. 3** Linearization of the Arrhenius equation (Eq. 4) of the rate of specific acidifying activity loss versus the inverse of the storage temperature. The activation energy ( $E_{a_{spe}}$ ) for each sample was determined from the slope of the linear fit and is shown in Table 2



**Fig. 4** Viable (gray bars), injured (blue bars), and dead cells (light blue bars) for the frozen samples, freeze-dried (FD) and spray-dried (SD) samples before storage, and after 28 days storage at 4, 23 and 37 °C. (a) *Lb. bulgaricus* CFL1; (b) *Lpb. plantarum* WCFS1

water activity than *Lb. bulgaricus* CFL1, and a higher  $T_g$  of the freeze-dried sample. Different drying batches and cell dehydration-rehydration properties could explain the differences observed among the strains. To evaluate the impact of the restricted mobility of the matrix in the glassy state on the specific acidifying activity loss rate,  $k_{spe}$  was plotted as a function of the difference between the storage temperature and the glass transition temperature ( $T-T_g$ ) (Fig. 5). For ( $T-T_g$ ) values under  $-40$  °C  $k_{spe}$  values were close to zero. As the ( $T-T_g$ ) parameter increased from  $-40$

**Table 3** Glass transition temperature of the maximally concentrated phase ( $T_g'$ ) of the frozen samples and the glass transition temperature ( $T_g$ ) and water activity ( $a_w$ ) of the dehydrated samples before storage

	$T_g'$ *		$T_g$		
	$a_w$		Spray-dried	Freeze-dried	
CFL1	-29.1	0.174 <sup>d</sup>	0.088 <sup>b</sup>	52.3 <sup>a</sup>	57.5 <sup>b</sup>
IQR	1.7	0.003	0.002	0.5	0.9
WCFS1	-29.5	0.141 <sup>c</sup>	0.033 <sup>a</sup>	52.6 <sup>a</sup>	67.5 <sup>c</sup>
IQR	0.1	0.003	0.004	1.9	1.5

Data presented as median and interquartile ranges (IQR). Different superscript letters within each method represent statistical differences between samples at a 95% confidence level

\* Three analytical replicates of only one biological replicate were measured for this determination

to  $-20$  °C the  $k_{spe}$  slightly increased. When the ( $T-T_g$ ) difference raised above  $-20$  °C a more abrupt increase of the  $k_{spe}$  was observed.

### Biochemical properties of cells by FTIR micro-spectroscopy

Principal component analyses (PCA) were performed in the three regions of interest for both biological replicates of the frozen, freeze-dried and spray-dried bacterial concentrates, before storage and after storage for 28 days at 4, 23 and 37 °C. In Fig. 6 and SM1 the PC1 vs. PC2 plots are displayed with their corresponding loadings. The analysis clearly discriminated between *Lb. bulgaricus* CFL1 and *Lpb. plantarum* WCFS1 bacterial populations in the three spectral regions. As displayed in Fig. 6, for *Lb. bulgaricus* CFL1 (circles) two separated clusters were observed that grouped

the frozen and the dehydrated samples. Whereas for *Lpb. plantarum* WCFS1 (crosses), it was also possible to discriminate between the freeze- and the spray-dried samples.


The spectral region between 1369 and 975  $\text{cm}^{-1}$  gives interesting information on the amide III band of proteins, nucleic acids, and phosphorylated molecules and polysaccharides from the cell wall. In this region, PC1 and PC2 explained 93% of the total variance (Fig. 6a). PC1 (71% of variance) discriminated both LAB. *Lpb. plantarum* WCFS1 cells were described by positive scores associated with bands at 1328–1317  $\text{cm}^{-1}$  (amide III bands of proteins), 1162–1147  $\text{cm}^{-1}$  (proteins from the cytoplasm and membranes, nucleic acids and sugar rings from the cell wall) and 1062–1010  $\text{cm}^{-1}$  (mostly peptidoglycan from the cell wall and nucleic acids). On the contrary, *Lb. bulgaricus* CFL1 cells were characterized by negative scores related to the band at 1110–1081  $\text{cm}^{-1}$  (carbohydrates of the membrane and cytoplasm, nucleic acids and phosphorylated proteins) (Fig. 6b). PC2 (22% of variance) showed differences between samples subjected to different process severity. Positive contributions of the bands at 1357–1334  $\text{cm}^{-1}$  (amide III band of proteins), 1145–1124  $\text{cm}^{-1}$  (carbohydrates of the membrane and cytoplasm, sugar rings of the cell wall and nucleic acids) and 1079–1072  $\text{cm}^{-1}$  (phosphodiesteres, phospholipids, and nucleic acids) were mainly associated with *Lpb. plantarum* WCFS1 frozen and freeze-dried samples and *Lb. bulgaricus* CFL1 frozen samples. Negative values at 1278–1193 (phosphodiesteres, phospholipids, and nucleic acids) and 1006–1004  $\text{cm}^{-1}$  (nucleic acids) were related to *Lpb. plantarum* WCFS1 spray-dried samples and freeze-dried samples stored at 37 °C as well as a good proportion of *Lb. bulgaricus* CFL1 dehydrated samples (Fig. 6c) (Guerrero Sanchez et al. 2022, 2023; Meneghel et al. 2020; Naumann et al. 2006).


The region between 1800 and 1370  $\text{cm}^{-1}$  is mostly related to protein components of the cell. In this spectral

region the first two PCs explained 92% of the total variance (Fig. 6d). The two LAB were discriminated by PC2 (17% of variance), while PC1 (75% of variance) discriminated the samples according to different processing and storage conditions. Frozen and freeze-dried *Lpb. plantarum* WCFS1 samples before and after storage at 4 and 23 °C, as well as *Lb. bulgaricus* CFL1 frozen samples presented positive PC1 scores, due to the contribution of bands at 1614–1562  $\text{cm}^{-1}$  (amide I and amide II bands of proteins) and 1427–1386  $\text{cm}^{-1}$  (carbohydrates of the cytoplasm and membranes, amino acids from ribosomes and other proteins and nucleic acids) (Fig. 6e). All the spray-dried samples and samples stored at higher temperatures showed negative values that were related to negative bands at 1724 to 1685  $\text{cm}^{-1}$  (amide I overlapped with esters, carboxylic acids and nucleic acids) and 1542–1430  $\text{cm}^{-1}$  (amide II bands of proteins and other vibrations of proteins from the cytoplasm and membranes) in the loading plot. On the other hand, for PC2, bands at 1646–1633  $\text{cm}^{-1}$  ( $\beta$ -pleated sheet structures of amide I) contributed to the positive scores of *Lb. bulgaricus* CFL1 samples, while bands at 1754–1729 (esters of the membrane lipids), 1558–1542 (amide II band of proteins) and 1382–1375  $\text{cm}^{-1}$  (proteins of the membranes and peptidoglycan from the cell wall) contributed to the negative ones for *Lpb. plantarum* WCFS1 (Fig. 6f) (Guerrero Sanchez et al. 2022, 2023; Meneghel et al. 2020; Naumann et al., 2006).

Finally, the region between 3060 and 2800  $\text{cm}^{-1}$  is associated with the fatty acyl chain of membrane lipids. In this region, the variance was much smaller compared to the other two regions and the first two PCs only explained 57% of the total variance (Fig. SM1.A). In this region signals related to the fatty acids chain of membrane lipids (2925–2937 and 2858–2846  $\text{cm}^{-1}$ ) contributed to PC1's negative scores associated with most of *Lb. bulgaricus* CFL1 samples (except for one biological replicate of the

**Fig. 5** Rate of loss of specific acidifying activity ( $k_{\text{spe}}$ ) as a function of ( $T-T_g$ ) for spray-dried *Lb. bulgaricus* CFL1 (orange circles), freeze-dried *Lb. bulgaricus* CFL1 (light blue squares), spray-dried *Lpb. plantarum* WCFS1 (red diamonds) and freeze-dried *Lpb. plantarum* WCFS1 (blue triangles)





**Fig. 6** Principal component analysis (PCA) of the FTIR normalized and corrected spectra of the of *Lb. bulgaricus* CFL1 (circles) and *Lpb. plantarum* WCFS1 (crosses) frozen, freeze-dried and spray-dried samples, before storage and after storage for 28 days at 4, 23 and 37 °C; in the spectral regions 1369–975  $\text{cm}^{-1}$  (a, b and c) and


1800–1370  $\text{cm}^{-1}$  (d, e and f). Both biological replicates are plotted independently in the score plots (a and d) and the discriminating bands are indicated in the loading plots (b and e for PC1; c and f for PC2)

frozen and the spray-dried sample stored at 23 °C); while signals possibly related to cyclopropane ring of cyclic FA (2995–2975  $\text{cm}^{-1}$ ) and other FA chain of membrane lipids (2941–2937  $\text{cm}^{-1}$ ) contributed to the positive scores linked to *Lpb. plantarum* WCFS1 (Fig.SM3.B) (Guerrero Sanchez et al. 2023; Kochan et al. 2018).

The second-order derivative of the spectra (Fig. SM2) made possible the identification and assignment of the peaks to specific components of the cells according to the bibliography (Table SM3). The relative variation of the peaks' height after drying compared to the frozen samples ( $r_v$ ) was calculated and the bands corresponding to the wavenumbers that suffered a variation larger than 15% for at least one of the samples are displayed in Fig. 7. A positive value means an increase of the peak's height and *vice-versa*. In the 1800–1370  $\text{cm}^{-1}$  region, the drying step caused important increases of the bands at 1751  $\text{cm}^{-1}$  ( $\nu\text{C}=\text{O}$  of esters) for all the samples (smaller change for the freeze-dried sample of *Lpb. plantarum* WCFS1), the bands at 1714

( $\nu\text{C}=\text{O}$ ,  $\nu\text{C}=\text{N}$ ,  $\nu\text{C}=\text{C}$  and  $\delta\text{NH}$  of esters, carboxylic acids and nucleic acids of DNA/RNA and ribosomes), 1548 ( $(\delta\text{N}-\text{H}) + (\nu\text{C}-\text{N})$  of amide II) for the spray-dried samples of *Lpb. plantarum* WCFS1 and *Lb. bulgaricus* CFL1, and 1515  $\text{cm}^{-1}$  ( $\nu\text{CC}$  and  $\delta\text{CH}$  of amino acids, particularly tyrosine, of the membrane and cytoplasm), especially for the spray-dried samples of *Lpb. plantarum* WCFS1. On the other hand, a decrease of the bands at 1419 ( $\delta\text{C}-\text{O}-\text{H}$  of carbohydrates, proteins and nucleic acids of membrane, cytoplasm, nucleoid and ribosomes, or  $\nu\text{C}-\text{O}$  sym of  $\text{COO}^-$  of phospholipids from the membrane) and 1400  $\text{cm}^{-1}$  ( $\nu\text{C}=\text{O}$  sym of  $\text{COO}^-$  and  $\nu\text{COO}^-$  sym of amino acids, fatty acid and peptidoglycan of the membrane, cytoplasm and cell wall) was observed for these samples. In the 1370–975  $\text{cm}^{-1}$  region, the dehydration step caused an increase of the band at 1058  $\text{cm}^{-1}$  ( $\nu\text{CO}$ ,  $\nu\text{CC}$ ,  $\delta\text{OCH}$ ,  $\nu\text{PO}_2^-$ ,  $\nu\text{C}-\text{OH}$ ,  $\nu\text{C}-\text{O}-\text{C}$  sym and  $\nu\text{P}-\text{O}-\text{C}$  sym of phospholipid phosphate, oligosaccharides, polysaccharides, sugar rings, pectin and peptidoglycan and deoxyribose of the membrane, cytoplasm, nucleoid and cell wall)

**Fig. 7** Changes in the peaks' height ( $r_v$ ) of the mean second derivative of the FTIR bands after freeze- or spray-drying compared to their frozen sample. Only peaks that showed changes larger than 15% are displayed



and a decrease of the band at  $1156\text{ cm}^{-1}$  ( $\nu\text{C-O}$  of proteins, carbohydrates and nucleic acids of the membrane and cytoplasm) more pronounced for *Lb. bulgaricus* CFL1 and the spray-dried samples. In general, spray-drying caused bigger changes in the peaks' height than freeze-drying except for the bands at  $1751$ ,  $1515$  and  $1400\text{ cm}^{-1}$  for *Lb. bulgaricus* CFL1 cells.

## Discussion

Most research on the effect of drying on LAB cannot be compared due to varying fermentation conditions, protection and/or stabilization strategies. The storage conditions often also differ or are not investigated. Moreover, different bacteria may exhibit different resistance to a drying process. The present article thus aimed to investigate the mechanisms underlying the preservation/degradation of LAB functionalities (culturability and acidifying activity) during drying and storage. For this purpose, two strains exhibiting different sensitivity were protected with the same protective solution and exposed to freeze- or spray-drying and to relevant storage temperatures for dried bacteria. Diverse techniques, from flow cytometry to FTIR spectroscopy and DSC, were used to analyze their biochemical and physical properties. Our results suggest the osmotic and mechanical stresses, occurring when freeze- or spray-drying, as the principal causes of damage to *Lb. bulgaricus* CFL1, and the thermal stress associated with spray-drying as the main stress affecting *Lpb. plantarum* WCFS1. This study also showed that, while the sensitivity of *Lb. bulgaricus* CFL1 to freeze-drying can be ascribed to loss of membrane integrity and cell wall degradation, heat inactivation of both strains associated with spray-drying can be explained by the modification of nucleic acids and proteins. The present results also substantiated the importance of the physical properties of the glassy matrix in the loss of activity rates, confirming that a glass transition temperature  $40\text{ }^{\circ}\text{C}$  lower than the storage temperature should be reached to preserve cells during storage.

The higher resistance of *Lpb. plantarum* WCFS1 compared to *Lb. bulgaricus* CFL1 was expected from previously reported work on the environmental stress responses of such subspecies (Arena et al. 2019; Capozzi et al. 2012; Mille et al. 2005; Passot et al., 2012, others). Moreover, while *Lb. bulgaricus* CFL1 was severely damaged by both drying methods, *Lpb. plantarum* WCFS1 exhibited less than three times lower loss of functionality after freeze-drying than *Lb. bulgaricus* CFL1 ( $\Delta_{\text{tspe}}$  FD 7 and 25 min. ( $\log(\text{CFU}\cdot\text{mL}^{-1})$ , respectively). This could be a consequence of the different biochemical composition between both strains induced by fermentation, as seen in the PCA results of the FTIR measurements, where both LAB exhibited important differences in the PC1 of the regions associated with cell wall

components and PC2 of the region related to proteins. By considering each drying method as a combination of three main stresses: osmotic mechanical and heat, it was possible to propose that *Lpb. plantarum* WCFS1 can tolerate better the osmotic and mechanic stress involved in both dehydration processes but is sensitive to the heat stress caused by spray-drying only. *Lb. bulgaricus* CFL1 is, in turn, sensitive to all three stresses, but certainly less sensitive to heat than *Lpb. plantarum* WCFS1. Our hypothesis agrees with previous work by Perdana et al. (2013), where from experimental and modelling studies at different inlet air temperatures, the authors concluded that the viability loss of *Lpb. plantarum* WCFS1 at high temperatures could be explained entirely by thermal inactivation. The differentiation between thermal and dehydration inactivation proposed in the present study also agrees with previous work by Lievens et al. (1992), which evidenced that thermal inactivation of *Lpb. plantarum* becomes relevant when temperatures are higher than  $50/55^{\circ}\text{C}$  on fluidised bed drying. According to Peigham-bardoust et al. (2011), the outlet air temperature is a crucial parameter, which depends on many operating variables and has a major influence on the viability of spray-dried starter cultures.

The membrane integrity measurements confirmed the cytoplasmic membrane as a site of damage during drying, as recognized by other authors for cells facing freeze-drying (Velly et al. 2015) and spray-drying (Huang et al. 2017). The high loss of membrane integrity observed for *Lb. bulgaricus* CFL1 following freeze-drying, and the comparatively lower effect of heat suggests a predominance of osmotic and mechanical effects of dehydration, inducing membrane injury. On the other hand, the percentage of injured and dead *Lpb. plantarum* WCFS1 cells significantly increased only with the temperature increase of the drying process and storage, thus revealing that heat stress also caused some membrane damage. The substantial contrast in the membrane resistance between the two strains against freeze-drying implies differences in the membrane organization that may provide *Lpb. plantarum* WCFS1 with a robust membrane little affected by osmotic and mechanical stresses.

The cell damage during drying and storage can be related to cell biochemical modifications evaluated by FTIR microspectroscopy. The differences observed for the two microorganisms were not only restricted to components of the cell membrane, but were also related to their nucleic acids, protein, and cell wall constituents. Recent works have also highlighted the modifications of many of these other cell constituents during drying processes (Romano et al. 2021; Guerrero Sanchez et al. 2022, 2023). In the present work, while the spectral region related to lipidic components showed only moderate variations, the largest variance was observed in the  $1800$  to  $1370\text{ cm}^{-1}$  and  $1369$  and  $975\text{ cm}^{-1}$  regions. Therefore, the loss of functionality during drying



and storage was mainly associated with damage to nucleic acids and proteins (Figs. 6 and 7). In particular, the bands related to nucleic acids ( $\nu\text{C}=\text{O}$  1714 and 1221  $\text{cm}^{-1}$  associated with asymmetric stretching  $\text{P}=\text{O}$ ), proteins (amide II at 1548  $\text{cm}^{-1}$  and tyrosine at 1515  $\text{cm}^{-1}$ ), and complex vibrations of lipids and proteins ( $\text{C}-\text{O}-\text{H}$  and stretching  $\text{COO}^-$  at 1419 and 1400  $\text{cm}^{-1}$ ) showed much bigger changes after spray-drying of *Lpb. plantarum* WCFS1 than after freeze-drying and may hold relevant information on the sensitivity to heat stress. Our results agree with a differential scanning calorimetry study of the damage induced by heat over *Lb. bulgaricus* cells that showed damage to the cytoplasmic membrane at temperatures below 64 °C, while at temperatures of 65 °C and above, the cell wall and proteins (specifically ribosomes) were critical sites of injury (Teixeira et al. 1997). Even though *Lb. bulgaricus* CFL1 also exhibited modifications in nucleic acid bands and proteins, the main modifications were observed in cell wall carbohydrates vibration bands (1156, 1058  $\text{cm}^{-1}$ ) for both drying techniques. These bands could thus be proposed as markers of the osmotic and mechanical stress associated with dehydration and used for characterizing the most sensitive strain. In agreement with the present work, the cell envelope of *L. helveticus* was identified as a site of injury (both by FTIR and AFM), together with the nucleic acids (DNA, by FTIR) following a low-temperature vacuum drying at 43 °C (Santivarangkana et al., 2007). Our findings also support recent work highlighting the potential role of unchanged bands at 1714 and 1058  $\text{cm}^{-1}$  (associated with nucleic acids and carbohydrates of the cell wall) as markers of *L. salivarius* (Guerrero Sanchez et al. 2022, 2023) preservation by freeze-drying (Table SM3).

The storage stability was related to the physical properties of the matrix. Since the glass transition temperature was higher than the storage temperature ( $T < T_g$ ), the protective solution ensured the immobilisation of cells in a glassy matrix. However, even in the glassy state, which is characterized by high viscosity and reduced molecular mobility, the storage stability decreased when approaching the supercooled state ( $(T-T_g)$  approaching zero). Similar results were obtained for the viability loss of freeze-dried *L. salivarius* CECT5713 (Guerrero Sanchez et al. 2022, 2023) and *Lb. bulgaricus* (Tymczyszyn et al. 2012) as the storage temperature approached the  $T_g$ . The increased molecular mobility, measured by  $^1\text{H-NMR}$  for *Lb. bulgaricus* in GOS matrices when approaching  $T_g$ , indicated that rotational, vibrational and short movement of molecules occurs even in the glassy state (Tymczyszyn et al. 2012). Moreover, the decrease of the storage stability (increase of the rate of loss of acidifying activity) with  $(T-T_g)$  showed similar non-linear trends for the two drying processes and was slightly more pronounced for *Lb. bulgaricus* CFL1 than *Lpb. plantarum* WCFS1 after spray-drying (Fig. 5). Wang et al. (2004) studied the viability

of spray- and freeze-dried LAB during storage, and also observed higher survival rates for the freeze-dried samples than for the spray-dried ones regardless of the storage conditions. Good storage stability was achieved when cells were maintained at storage temperatures 40 °C lower than the glass transition temperatures ( $(T-T_g) < -40$ ;  $k_{spe} < 0.3$ ), which corresponded to samples stored at 4 °C and freeze-dried *Lpb. plantarum* cells stored at 23 °C.

The activation energy ( $E_a$ ) values, calculated from the rate of loss of culturability during storage at different temperatures were consistent with those reported by Ziadi et al. (2005) for freeze-dried cells of *Lactococcus lactis* subsp. *lactis* protected with sucrose, and by Aschenbrenner et al. (2012) for freeze-dried *Lactobacillus paracasei* subsp. *paracasei* protected with trehalose, but about two times higher than the ones informed by Sosa et al. (2016) for spray-dried *Lpb. plantarum* CIDCA83114 protected with galacto-oligosaccharides and maltodextrin mixtures (Table SM2). The  $E_a$  and  $E_{a_{spe}}$  values found in this work (between 47 and 85 kJ/mol, and 41 to 105 kJ/mol respectively) are, according to literature, in the range of chemically limited reactions (50 to 150 kJ/mol), while for systems controlled by diffusion processes  $E_a$  values are usually below 20 kJ/mol (Aschenbrenner et al. 2012). Aschenbrenner et al. (2012) studied the inactivation of *Lb. paracasei* and also described non-linear relationships between the viability loss rate and  $(T-T_g)$ . The authors studied the effect of different protective systems and found that, while for samples protected with dextran, the inactivation constant was solely dependent on the  $(T-T_g)$  difference, for cells protected with di- or oligosaccharides this constant depended on both, absolute storage temperature and  $(T-T_g)$  parameter. The authors suggest that in the first system, the key protective mechanism is the diffusion restriction due to the high  $T_g$  provided by the protectant, whereas in systems containing oligosaccharides water replacement may have a significant contribution towards protection. In our work, the protective agent contained maltodextrin and FOS, thus, both protective mechanisms were probably involved in the preservation of the cells during drying. This hypothesis supports the activation energy findings and explains the high rates of loss of activity even in the glassy state. Also, some reactions, such as oxidation of cell components by reactive oxygen species generated during drying do not depend strongly on diffusion and early stages of non-enzymatic browning could have occurred between the small reducing sugars of FOS and glucose and proteins remaining from the culture medium, even with diffusion limitations since in a concentrated matrix the reagents may be close enough. These reactions could also explain the biological activity loss observed even if the samples were stored in a glassy state. Moreover, *Lb. bulgaricus* CFL1 exhibited similar  $E_a$  and  $E_{a_{spe}}$  values in the freeze- and spray-dried samples, while spray-dried *Lpb. plantarum* WCFS1 showed a more than 50% increase

of Ea compared to the freeze-dried form, thus confirming the sensitivity of this strain to increments in temperature.

In conclusion, this work evidenced the different resistance of two lactic acid bacteria to different types of stresses associated with freeze-drying and spray-drying. *Lb. bulgaricus* CFL1 appeared sensitive to osmotic, mechanical, and thermal stresses, while *Lpb. plantarum* WCFS1 showed higher resistance to the first two but appeared to be more sensitive to thermal stress associated with spray-drying. The results reached in this work enabled the identification of the cellular damages caused by spray- and freeze-drying and storage and a deeper understanding of the contribution of individual stress to the activity loss of two strains. The FTIR micro-spectroscopy method applied here in combination with flow cytometry to evidence the implication of the membrane in cell resistance to preservation and storage processes, showed that also other cell constituents such as nucleic acids, proteins, and cell wall components such as phosphorylated molecules and polysaccharides components have a fundamental influence. Nucleic acids and proteins were majorly affected by spray-drying, while the cell wall was the weakness of the more sensitive strain *Lb. bulgaricus* CFL1. Besides, the activation energy results suggested that regardless of the strain and drying process, chemically limited reactions ruled the functionality loss during storage. A more detailed study on this may be worth further analysis.

These results deliver practical guidance for improving stabilization and storage protocols. Overall, they suggested that freeze-drying of *Lpb. plantarum* WCFS1 could enable the storage of this lactic acid bacterium at mild ambient temperatures without major functionality loss. Besides, since freeze-drying and spray-drying caused a comparable degree of damage on *Lb. bulgaricus* CFL1 cells, spray-drying could be considered as a lower-cost and possibly more eco-friendly alternative, leading to practical, economic, and environmental advancements. The ongoing evaluation of the environmental performance of these two stabilization scenarios will shed light on (confirm or not) this hypothesis. The importance of the glass transition temperature and the difference between this parameter and the storage temperature ( $T-T_g$ ) on storage stability was also highlighted. ( $T-T_g$ ) values below  $-40\text{ }^\circ\text{C}$  guaranteed good survival while when they surpassed  $-20\text{ }^\circ\text{C}$ , an abrupt increase of the inactivation rate occurred. FTIR micro-spectroscopy in aqueous solutions represents a complementary and faster alternative to biological assays and could be applied for the screening of strains, fermentation, and stabilization conditions.

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s00253-024-13186-3>.

**Acknowledgements** We thank Maria Guerrero Sanchez and Sonia Campoy from Biosearch S.A.U (a Kerry® company) for their help in

the production of bacterial concentrates. Finally, we thank Maria Guerrero Sanchez from Biosearch for training MG in FTIR spectroscopy.

**Author contributions** Conceptualization and methodology MG, SP, FF; pilot production, freeze-drying, spray-drying of bacterial concentrates SC and SP, biological, physical and biochemical FTIR characterization after FD, SD and storage MG, flow cytometry measurements SG; formal analysis MG, SP, FF; data curation MG, SP, FF; writing original draft of the paper and writing review MG, SP, FF; final review and editing MG, SP, SC, CS, FF; supervision and project administration SP, FF; resources and funding acquisition SP, FF.

**Funding** This work has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement N° 777657.

**Data availability** The datasets generated and/or analysed during the current study are available in the Data INRAE repository (<https://doi.org/10.57745/AAARUZ>).

## Declarations

**Conflict of interest** The authors declare that they have no conflict of interest.

**Ethical approval** This article does not contain any studies with human participants or animals performed by any of the authors.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Arena MP, Capozzi V, Longo A, Russo P, Weidmann S, Rieu A, Guzzo J, Spano G, Fiocco D (2019) The phenotypic analysis of *Lactobacillus plantarum* shsp mutants reveals a potential role for hsp1 in cryotolerance. *Front Microbiol* 10:838. <https://doi.org/10.3389/fmicb.2019.00838>
- Aschenbrenner M, Kulozik U, Foerst P (2012) Evaluation of the relevance of the glassy state as stability criterion for freeze-dried bacteria by application of the Arrhenius and WLF model. *Cryobiol* 65(3):308–318. <https://doi.org/10.1016/j.cryobiol.2012.08.005>
- Bensch G, Ruger M, Wassermann M, Weinholz S, Reichl U, Cordes C (2014) Flow cytometric viability assessment of lactic acid bacteria starter cultures produced by fluidized bed drying. *Appl Microbiol Biotechnol* 98:4897–4909. <https://doi.org/10.1007/s00253-014-5592-z>
- Boafo GF, Magar KT, Ekpo MD, Qian W, Tan S, Chen C (2022) The role of cryoprotective agents in liposome stabilization and preservation. *Int J Mol Sci* 23(20):12487. <https://doi.org/10.3390/ijms232012487>

- Bouix M, Ghorbal S, Picque D, Perret B, Saulou-Bérion C (2022) A rapid method for the assessment of the vitality of microorganisms using flow cytometry. *Cytometry Part A* 101(7):577–587. <https://doi.org/10.1002/cyto.a.24553>
- Broeckx G, Vandenheuvel D, Claes IJ, Lebeer S, Kiekens F (2016) Drying techniques of probiotic bacteria as an important step towards the development of novel pharmabiotics. *Int J Pharm* 505(1–2):303–318. <https://doi.org/10.1016/j.ijpharm.2016.04.002>
- Capozzi V, Fiocco D, Weidmann S, Guzzo J, Spano G (2012) Increasing membrane protection in *Lactobacillus plantarum* cells overproducing small heat shock proteins. *Annals Microbiol* 62(2):517–522. <https://doi.org/10.1007/s13213-011-0285-7>
- Champagne CP, Raymond Y, Simon JP (2012) Effect of water activity and protective solutes on growth and subsequent survival to air-drying of *Lactobacillus* and *Bifidobacterium* cultures. *Appl Microbiol Biotechnol* 95:745–756. <https://doi.org/10.1007/s00253-012-3912-8>
- Crowe JH, Crowe LM, Carpenter JF, Rudolph AS, Wistrom CA, Spargo BJ, Anchordoguy TJ (1988) Interactions of sugars with membranes. *Biochim Biophys Acta (BBA) - Reviews on Biomembranes*, 947(2), 367–384. [https://doi.org/10.1016/0304-4157\(88\)90015-9](https://doi.org/10.1016/0304-4157(88)90015-9)
- Ermis E (2021) A review of drying methods for improving the quality of probiotic powders and characterization. *Dry Technol* 1–18. <https://doi.org/10.1080/07373937.2021.195>
- Fonseca F, Pénicaud C, Tymczyszyn EE, Gómez-Zavaglia A, Passot S (2019) Factors influencing the membrane fluidity and the impact on production of lactic acid bacteria starters. *Appl Microbiol Biotechnol* 103:6867–6883. <https://doi.org/10.1007/s00253-019-10002-1>
- G-Alegría E, López I, Ruiz JI, Sáenz J, Fernández E, Zarazaga M, Dizy M, Torres C, Ruiz-Larrea F (2004) High tolerance of wild *Lactobacillus plantarum* and *Oenococcus oeni* strains to lyophilisation and stress environmental conditions of acid pH and ethanol. *FEMS Microbiol Lett* 230(1):53–61. [https://doi.org/10.1016/S0378-1097\(03\)00854-1](https://doi.org/10.1016/S0378-1097(03)00854-1)
- Gautier J, Passot S, Pénicaud C, Guillemin H, Cenard S, Lieben P, Fonseca F (2013) A low membrane lipid phase transition temperature is associated with a high cryotolerance of *Lactobacillus delbrueckii* subspecies *bulgaricus* CFL1. *J Dairy Sci*, 96(9), 5591–5602. <https://doi.org/10.3168/jds.2013-6802>
- Gibson GR, Probert HM, Van Loo J, Rastall RA, Roberfroid MB (2004) Dietary modulation of the human colonic microbiota: updating the concept of prebiotics. *Nutr Res Rev* 17(2):259–275. <https://doi.org/10.1079/NRR200479>
- Girardeau A, Passot S, Meneghel J, Cenard S, Lieben P, Trelea IC, Fonseca F (2022) Insights into lactic acid bacteria cryoresistance using FTIR microspectroscopy. *Anal Bioanal Chem* 414(3):1425–1443. <https://doi.org/10.1007/s00216-021-03774-x>
- Golowczyc MA, Silva J, Abraham AG, De Antoni GL, Teixeira P (2010) Preservation of probiotic strains isolated from kefir by spray drying. *Lett Appl Microbiol* 50(1):7–12. <https://doi.org/10.1111/j.1472-765X.2009.02759.x>
- Grasmeijer N, Stankovic M, de Waard H, Frijlink HW, Hinrichs WL (2013) Unraveling protein stabilization mechanisms: vitrification and water replacement in a glass transition temperature controlled system. *Biochim Biophys Acta (BBA) - Proteins Proteom* 1834(4):763–769. <https://doi.org/10.1016/j.bbapap.2013.01.020>
- Guerrero Sanchez M, Passot S, Campoy S, Olivares M, Fonseca F (2022) Effect of protective agents on the storage stability of freeze-dried *Ligilactobacillus salivarius* CECT5713. *Appl Microbiol Biotechnol* 106(21):7235–7249. <https://doi.org/10.1007/s00253-022-12201-9>
- Guerrero Sanchez M, Passot S, Ghorbal S, Campoy S, Olivares M, Fonseca F (2023) Insights into the mechanisms of *L. salivarius* CECT5713 resistance to freeze-dried storage. *Cryobiol* 112:104556. <https://doi.org/10.1016/j.cryobiol.2023.104556>
- Huang S, Vignolles ML, Chen XD, Le Loir Y, Jan G, Schuck P, Jeantet R (2017) Spray drying of probiotics and other food-grade bacteria: a review. *Trends Food Sci Technol* 63:1–17. <https://doi.org/10.1016/j.tifs.2017.02.007>
- Kieps J, Dembczyński R (2022) Current trends in the production of probiotic formulations. *Foods* 11(15):2330. <https://doi.org/10.3390/foods11152330>
- Kochan K, Peng H, Gwee ES, Izgorodina E, Haritos V, Wood BR (2018) Raman spectroscopy as a tool for tracking cyclopropane fatty acids in genetically engineered *Saccharomyces cerevisiae*. *Analyst* 144(3):901–912. <https://doi.org/10.1039/C8AN01477A>
- Lievens LC, Verbeek MA, Taekema T, Meerdink G, Van't Riet K (1992) Modelling the inactivation of *Lactobacillus plantarum* during a drying process. *Chem Eng Sci* 47(1):87–97. [https://doi.org/10.1016/0009-2509\(92\)80203-O](https://doi.org/10.1016/0009-2509(92)80203-O)
- Liu H, Cui SW, Chen M, Li Y, Liang R, Xu F, Zhong F (2019) Protective approaches and mechanisms of microencapsulation to the survival of probiotic bacteria during processing, storage and gastrointestinal digestion: a review. *Crit Rev Food Sci Nutr* 59(17):2863–2878. <https://doi.org/10.1080/10408398.2017.1377684>
- Loffhagen N, Härtig C, Geyer W, Voyevoda M, Harms H (2007) Competition between cis, trans and cyclopropane fatty acid formation and its impact on membrane fluidity. *Eng Life Sci* 7(1):67–74. <https://doi.org/10.1002/elsc.200620168>
- Meneghel J, Passot S, Dupont S, Fonseca F (2017) Biophysical characterization of the *Lactobacillus delbrueckii* subsp. *bulgaricus* membrane during cold and osmotic stress and its relevance for cryopreservation. *Appl Microbiol Biotechnol* 101(4):1427–1441. <https://doi.org/10.1007/s00253-016-7935-4>
- Meneghel J, Passot S, Jamme F, Lefrançois S, Lieben P, Dumas P, Fonseca F (2020) FTIR micro-spectroscopy using synchrotron-based and thermal source-based radiation for probing live bacteria. *Anal Bioanal Chem* 412(26):7049–7061. <https://doi.org/10.1007/s00216-020-02835-x>
- Mille Y, Beney L, Gervais P (2005) Compared tolerance to osmotic stress in various microorganisms: towards a survival prediction test. *Biotechnol Bioeng* 92(4):479–484. <https://doi.org/10.1002/bit.20631>
- Montel Mendoza G, Pasteris SE, Otero MC, Fatima Nader-Macías ME (2014) Survival and beneficial properties of lactic acid bacteria from rancicure subjected to freeze-drying and storage. *J Appl Microbiol* 116(1):157–166. <https://doi.org/10.1111/jam.12359>
- Naumann D (2006) Infrared spectroscopy in Microbiology. In: Meyers RA (ed) *Encyclopedia of Analytical Chemistry*. John Wiley & Sons, Ltd, Chichester, UK, p a0117
- Passot S, Cenard S, Douania I, Trélea IC, Fonseca F (2012) Critical water activity and amorphous state for optimal preservation of lyophilised lactic acid bacteria. *Food Chem* 132(4):1699–1705. <https://doi.org/10.1016/j.foodchem.2011.06.012>
- Peighambaroust SH, Tafti AG, Hesari J (2011) Application of spray drying for preservation of lactic acid starter cultures: a review. *Trends Food Sci Technol* 22(5):215–224
- Pénicaud C, Monclus V, Perret B, Passot S, Fonseca F (2018) Life cycle assessment of the production of stabilized lactic acid bacteria for the environmentally-friendly preservation of living cells. *J Clean Prod* 184:847–858. <https://doi.org/10.1016/j.jclepro.2018.02.191>
- Perdana J, Bereschenko L, Fox MB, Kuperus JH, Kleerebezem M, Boom RM, Schutyser MAI (2013) Dehydration and thermal inactivation of *Lactobacillus plantarum* WCFS1: comparing single droplet drying to spray and freeze drying. *Food Res Int* 54(2):1351–1359. <https://doi.org/10.1016/j.foodres.2013.09.043>
- Rajam R, Anandharamkrishnan C (2015) Microencapsulation of *Lactobacillus plantarum* (MTCC 5422) with fructooligosaccharide

- as wall material by spray drying. *LWT - Food Sci. Technol.* Vol 60, Issue 2, Part 1, (773–780). <https://doi.org/10.1016/j.lwt.2014.09.062>
- Rault A, Béal C, Ghorbal S, Ogier J-C, Bouix M (2007) Multiparametric flow cytometry allows rapid assessment and comparison of lactic acid bacteria viability after freezing and during frozen storage. *Cryobiol* 55(1):35–43. <https://doi.org/10.1016/j.cryobiol.2007.04.005>
- Romano N, Schebor C, Mobili P, Gomez-Zavaglia A (2016) Role of mono- and oligosaccharides from FOS as stabilizing agents during freeze-drying and storage of *Lactobacillus delbrueckii* subsp. *bulgaricus*. *Int Food Res J* 90:251–258. <https://doi.org/10.1016/j.foodres.2016.11.003>
- Romano N, Marro M, Marsal M, Loza-Alvarez P, Gomez-Zavaglia A (2021) Fructose derived oligosaccharides prevent lipid membrane destabilization and DNA conformational alterations during vacuum-drying of *Lactobacillus delbrueckii* subsp. *bulgaricus*. *Food Res Int*, 143, 110235. <https://doi.org/10.1016/j.foodres.2021.110235>
- Roos YH (2010) Glass transition temperature and its relevance in food processing. *Annual Rev Food Sci Technol* 1:469–496. <https://doi.org/10.1146/annurev.food.102308.124139>
- Santivarangkna C, Kulozik U, Foerst P (2008) Inactivation mechanisms of lactic acid starter cultures preserved by drying processes. *J Appl Microbiol* 105(1):1–13. <https://doi.org/10.1111/j.1365-2672.2008.03744.x>
- Sehrawat R, Abdullah S, Khatri P, Kumar L, Kumar A, Mujumdar AS (2022) Role of drying technology in probiotic encapsulation and impact on food safety. *Dry Technol* 40(8):1562–1581. <https://doi.org/10.1080/07373937.2022.2044844>
- Sharma R, Rashidnejad A, Jafari SM (2022) Application of spray dried encapsulated probiotics in functional food formulations. *Food Bioprocess Technol* 15:2135–2154. <https://doi.org/10.1007/s11947-022-02803-6>
- Sosa N, Gerbino E, Golowczyc MA, Schebor C, Gómez-Zavaglia A, Tymczyszyn EE (2016) Effect of galacto-oligosaccharides: maltodextrin matrices on the recovery of *Lactobacillus plantarum* after spray-drying. *Front Microbiol* 7. <https://doi.org/10.3389/fmicb.2016.00584>
- Streit F, Delettre J, Corrieu G, Béal C (2008) Acid adaptation of *Lactobacillus delbrueckii* subsp. *bulgaricus* induces physiological responses at membrane and cytosolic levels that improves cryotolerance. *J Appl Microbiol* 105(4):1071–1080. <https://doi.org/10.1111/j.1365-2672.2008.03848.x>
- Teixeira P, Castro H, Mohácsi-Farkas C, Kirby R (1997) Identification of sites of injury in *Lactobacillus bulgaricus* during heat stress. *J Appl Microbiol* 83(2):219–226. <https://doi.org/10.1046/j.1365-2672.1997.00221.x>
- Tymczyszyn EE, Sosa N, Gerbino E, Hugo A, Gómez-Zavaglia A, Schebor C (2012) Effect of physical properties on the stability of *Lactobacillus bulgaricus* in a freeze-dried galacto-oligosaccharides matrix. *Int J Food Microbiol* 155(3):217–221. <https://doi.org/10.1016/j.ijfoodmicro.2012.02.008>
- Velly H, Bouix M, Passot S, Pénicaud C, Beinstener H, Ghorbal S, Lieben P, Fonseca F (2015) Cyclopropanation of unsaturated fatty acids and membrane rigidification improve the freeze-drying resistance of *Lactococcus lactis* subsp. *lactis* TOMSC161. *Appl Microbiol Biotechnol*, 99, 907–918. <https://doi.org/10.1007/s00253-014-6152-2>
- Wang YC, Yu RC, Chou CC (2004) Viability of lactic acid bacteria and bifidobacteria in fermented soymilk after drying, subsequent rehydration and storage. *Int J Food Microbiol* 93(2):209–217. <https://doi.org/10.1016/j.ijfoodmicro.2003.12.001>
- Yoha KS, Moses JA, Anandharamakrishnan C (2020a) Conductive hydro drying through refractance window drying—An alternative technique for drying of *Lactobacillus plantarum* (NCIM 2083). *Dry Technol* 38(5–6):610–620
- Yoha KS, Moses JA, Anandharamakrishnan C (2020b) Effect of encapsulation methods on the physicochemical properties and the stability of *Lactobacillus plantarum* (NCIM, 2083) in synbiotic powders and in-vitro digestion conditions. *J Food Eng* 283:110033. <https://doi.org/10.1016/j.jfoodeng.2020.110033>
- Ziadi M, Touhami Y, Achour M, Thonart P, Hamdi M (2005) The effect of heat stress on freeze-drying and conservation of *Lactococcus*. *Biochem Eng J* 24(2):141–145. <https://doi.org/10.1016/j.bej.2005.02.001>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.