



# Texture modeling by gaussian fields with prescribed local orientation

Kévin Polisano

## ► To cite this version:

Kévin Polisano. Texture modeling by gaussian fields with prescribed local orientation. ICIP 2014 - 21st IEEE International Conference on Image Processing, Oct 2014, Paris ( France), France. hal-04610144

HAL Id: hal-04610144

<https://hal.science/hal-04610144v1>

Submitted on 13 Jun 2024


**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



UNIVERSITÉ DE  
GRENOBLE


# TEXTURE MODELING BY GAUSSIAN FIELDS WITH PRESCRIBED LOCAL ORIENTATION



Kévin Polisano / [kevin.polisano@imag.fr](mailto:kevin.polisano@imag.fr)

joint work with

Marianne Clausel  
Valérie Perrier  
Laurent Condat



# Outline

---

## ■ Introduction

- Motivation
- General probabilistic framework

## ■ Our new stochastic model

- Definition: Locally Anisotropic Fractional Brownian Field
- Notion of tangent field


## ■ Synthesis methods

- Tangent field simulation by a turning bands method
- LAFBF simulation via tangent field formulation

## ■ Numerical experiments

## ■ Conclusion and future work

# How to synthesize natural random textures ?



Mathematical  
model ?

# How to synthesize natural random textures ?



Mathematical  
model ?

# How to synthesize natural random textures ?


Randomness  
Self-similarity



Mathematical  
model ?

# How to synthesize natural random textures ?


Randomness  
Self-similarity




Mathematical  
model ?

# How to synthesize natural random textures ?

Randomness  
Self-similarity



Mathematical  
model ?




# How to synthesize natural random textures ?

Randomness  
Self-similarity



Mathematical  
model ?





Roughness  
and regularity

# How to synthesize natural random textures ?

Randomness  
Self-similarity




Mathematical  
model ?





Roughness  
and regularity

# How to synthesize natural random textures ?

Randomness  
Self-similarity



Mathematical  
model ?




Roughness  
and regularity

# How to synthesize natural random textures ?


Randomness  
Self-similarity




Mathematical  
model ?



Orientation  
and anisotropy



Roughness  
and regularity



# The basic component :

## Fractional Brownian Field (FBF)

- $B^H$  FBF with Hurst index  $0 < H < 1$  [Mandelbrot, Van Ness, 1968]
- stationary increments :  $B^H(\cdot + \mathbf{z}) - B^H(\mathbf{z}) \stackrel{\mathcal{L}}{=} B^H(\cdot) - B^H(0)$
- self-similar :  $B^H(\lambda \cdot) \stackrel{\mathcal{L}}{=} \lambda^H B^H(\cdot)$
- isotropic :  $B^H \circ R_\theta \stackrel{\mathcal{L}}{=} B^H$
- The covariance is given by

$$\text{Cov}(B^H(\mathbf{x}), B^H(\mathbf{y})) = c_H(\|\mathbf{x}\|^{2H} + \|\mathbf{y}\|^{2H} - \|\mathbf{x} - \mathbf{y}\|^{2H})$$

# The basic component :

## Fractional Brownian Field (FBF)

■ Harmonizable representation

[Samorodnitsky,Taqqu,1997]

$$B^H(\mathbf{x}) = \int_{\mathbb{R}^2} \frac{e^{i\mathbf{x}\cdot\xi} - 1}{\|\xi\|^{H+1}} d\widehat{W}(\xi)$$

# The basic component :

## Fractional Brownian Field (FBF)

■ Harmonizable representation

[Samorodnitsky,Taqqu,1997]

$$B^H(\mathbf{x}) = \int_{\mathbb{R}^2} \frac{e^{i\mathbf{x}\cdot\xi} - 1}{\|\xi\|^{H+1}} d\widehat{W}(\xi)$$

complex Brownian measure

# The basic component :

## Fractional Brownian Field (FBF)

■ Harmonizable representation

[Samorodnitsky,Taqqu,1997]

$$B^H(x) = \int_{\mathbb{R}^2} \frac{e^{ix \cdot \xi} - 1}{\|\xi\|^{H+1}} d\widehat{W}(\xi)$$

roughness indicator

complex Brownian measure

# The basic component :

## Fractional Brownian Field (FBF)


■ Harmonizable representation

[Samorodnitsky,Taqqu,1997]

$$B^H(x) = \int_{\mathbb{R}^2} \frac{e^{ix \cdot \xi} - 1}{\|\xi\|^{H+1}} d\widehat{W}(\xi)$$

roughness indicator

complex Brownian measure



H=0.2

# The basic component :

## Fractional Brownian Field (FBF)


### ■ Harmonizable representation

[Samorodnitsky,Taqqu,1997]

$$B^H(x) = \int_{\mathbb{R}^2} \frac{e^{ix \cdot \xi} - 1}{\|\xi\|^{H+1}} d\widehat{W}(\xi)$$

roughness indicator

complex Brownian measure



## General model :

**anisotropic self-similar Gaussian fields**

$$X(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x}\cdot\xi} - 1) f^{1/2}(\mathbf{x}, \xi) d\widehat{W}(\xi)$$

## General model :

### anisotropic self-similar Gaussian fields

$$X(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x}\cdot\xi} - 1) f^{1/2}(\mathbf{x}, \xi) d\widehat{W}(\xi)$$

$$f^{1/2}(\mathbf{x}, \xi) = c(\mathbf{x}, \xi) \|\xi\|^{-h(\mathbf{x}, \xi) - 1}$$

spectral density

## General model :

### anisotropic self-similar Gaussian fields


$$X(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x}\cdot\boldsymbol{\xi}} - 1) f^{1/2}(\mathbf{x}, \boldsymbol{\xi}) d\widehat{W}(\boldsymbol{\xi})$$

$$f^{1/2}(\mathbf{x}, \boldsymbol{\xi}) = c(\mathbf{x}, \boldsymbol{\xi}) \|\boldsymbol{\xi}\|^{-h(\mathbf{x}, \boldsymbol{\xi})-1}$$

spectral density

- $c(\mathbf{x}, \boldsymbol{\xi}) \equiv 1$  and  $h(\mathbf{x}, \boldsymbol{\xi}) \equiv H \Rightarrow X = B^H$

[Mandelbrot, Van Ness, 1968]



## General model :


### anisotropic self-similar Gaussian fields

$$X(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x}\cdot\boldsymbol{\xi}} - 1) f^{1/2}(\mathbf{x}, \boldsymbol{\xi}) d\widehat{W}(\boldsymbol{\xi})$$

$$f^{1/2}(\mathbf{x}, \boldsymbol{\xi}) = c(\mathbf{x}, \boldsymbol{\xi}) \|\boldsymbol{\xi}\|^{-h(\mathbf{x}, \boldsymbol{\xi})-1}$$

spectral density

- $c(\mathbf{x}, \boldsymbol{\xi}) \equiv 1$  and  $h(\mathbf{x}, \boldsymbol{\xi}) \equiv H \Rightarrow X = B^H$  [Mandelbrot, Van Ness, 1968]
- $c(\mathbf{x}, \boldsymbol{\xi}) \equiv c(\arg \boldsymbol{\xi})$  and  $h(\mathbf{x}, \boldsymbol{\xi}) \equiv h(\arg \boldsymbol{\xi}) \Rightarrow X = AFBF$  [Bonami, Estrade, 2003]



## General model :


### anisotropic self-similar Gaussian fields

$$X(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x}\cdot\xi} - 1) f^{1/2}(\mathbf{x}, \xi) d\widehat{W}(\xi)$$

$$f^{1/2}(\mathbf{x}, \xi) = c(\mathbf{x}, \xi) \|\xi\|^{-h(\mathbf{x}, \xi) - 1}$$

spectral density

- $c(\mathbf{x}, \xi) \equiv 1$  and  $h(\mathbf{x}, \xi) \equiv H \Rightarrow X = B^H$  [Mandelbrot, Van Ness, 1968]
- $c(\mathbf{x}, \xi) \equiv c(\arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\arg \xi) \Rightarrow X = AFBF$  [Bonami, Estrade, 2003]
- Example : *elementary fields*  $c(\arg \xi) = 1_{[-\alpha, \alpha]}(\arg \xi - \alpha_0)$   
[Bierme, Richard, Moisan, 2012]



# General model :


## anisotropic self-similar Gaussian fields

$$X(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x}\cdot\xi} - 1) f^{1/2}(\mathbf{x}, \xi) d\widehat{W}(\xi)$$

spectral density

$$f^{1/2}(\mathbf{x}, \xi) = c(\mathbf{x}, \xi) \|\xi\|^{-h(\mathbf{x}, \xi) - 1}$$

- $c(\mathbf{x}, \xi) \equiv 1$  and  $h(\mathbf{x}, \xi) \equiv H \Rightarrow X = B^H$  [Mandelbrot, Van Ness, 1968]
- $c(\mathbf{x}, \xi) \equiv c(\arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\arg \xi) \Rightarrow X = AFBF$  [Bonami, Estrade, 2003]
- Example : *elementary fields*  $c(\arg \xi) = 1_{[-\alpha, \alpha]}(\arg \xi - \alpha_0)$   
[Bierme, Richard, Moisan, 2012]



# General model :


## anisotropic self-similar Gaussian fields

$$X(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x}\cdot\xi} - 1) f^{1/2}(\mathbf{x}, \xi) d\widehat{W}(\xi)$$

$$f^{1/2}(\mathbf{x}, \xi) = c(\mathbf{x}, \xi) \|\xi\|^{-h(\mathbf{x}, \xi) - 1}$$

spectral density

- $c(\mathbf{x}, \xi) \equiv 1$  and  $h(\mathbf{x}, \xi) \equiv H \Rightarrow X = B^H$  [Mandelbrot, Van Ness, 1968]
- $c(\mathbf{x}, \xi) \equiv c(\arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\arg \xi) \Rightarrow X = AFBF$  [Bonami, Estrade, 2003]
- Example : *elementary fields*  $c(\arg \xi) = 1_{[-\alpha, \alpha]}(\arg \xi - \alpha_0)$   
[Bierme, Richard, Moisan, 2012]



# General model :


## anisotropic self-similar Gaussian fields

$$X(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x}\cdot\xi} - 1) f^{1/2}(\mathbf{x}, \xi) d\widehat{W}(\xi)$$

$$f^{1/2}(\mathbf{x}, \xi) = c(\mathbf{x}, \xi) \|\xi\|^{-h(\mathbf{x}, \xi) - 1}$$

spectral density

- $c(\mathbf{x}, \xi) \equiv 1$  and  $h(\mathbf{x}, \xi) \equiv H \Rightarrow X = B^H$  [Mandelbrot, Van Ness, 1968]
- $c(\mathbf{x}, \xi) \equiv c(\arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\arg \xi) \Rightarrow X = AFBF$  [Bonami, Estrade, 2003]
- Example : elementary fields  $c(\arg \xi) = 1_{[-\alpha, \alpha]}(\arg \xi - \alpha_0)$
- $c(\mathbf{x}, \xi) \equiv c(\mathbf{x}, \arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\mathbf{x})$  [Bierme, Richard, Moisan, 2012]  
[Polisano, Clausel, Perrier, Condat, 2014]



# General model :

## anisotropic self-similar Gaussian fields

$$X(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x}\cdot\xi} - 1) f^{1/2}(\mathbf{x}, \xi) d\widehat{W}(\xi)$$

$$f^{1/2}(\mathbf{x}, \xi) = c(\mathbf{x}, \xi) \|\xi\|^{-h(\mathbf{x}, \xi) - 1}$$

spectral density

- $c(\mathbf{x}, \xi) \equiv 1$  and  $h(\mathbf{x}, \xi) \equiv H \Rightarrow X = B^H$  [Mandelbrot, Van Ness, 1968]
- $c(\mathbf{x}, \xi) \equiv c(\arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\arg \xi) \Rightarrow X = AFBF$  [Bonami, Estrade, 2003]
- Example : elementary fields  $c(\arg \xi) = 1_{[-\alpha, \alpha]}(\arg \xi - \alpha_0)$
- $c(\mathbf{x}, \xi) \equiv c(\mathbf{x}, \arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\mathbf{x})$  [Bierme, Richard, Moisan, 2012]  
[Polisano, Clausel, Perrier, Condat, 2014]



# General model :

## anisotropic self-similar Gaussian fields

$$X(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x}\cdot\xi} - 1) f^{1/2}(\mathbf{x}, \xi) d\widehat{W}(\xi)$$

$$f^{1/2}(\mathbf{x}, \xi) = c(\mathbf{x}, \xi) \|\xi\|^{-h(\mathbf{x}, \xi) - 1}$$

spectral density

- $c(\mathbf{x}, \xi) \equiv 1$  and  $h(\mathbf{x}, \xi) \equiv H \Rightarrow X = B^H$  [Mandelbrot, Van Ness, 1968]
- $c(\mathbf{x}, \xi) \equiv c(\arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\arg \xi) \Rightarrow X = AFBF$  [Bonami, Estrade, 2003]
- Example : elementary fields  $c(\arg \xi) = 1_{[-\alpha, \alpha]}(\arg \xi - \alpha_0)$
- $c(\mathbf{x}, \xi) \equiv c(\mathbf{x}, \arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\mathbf{x})$  [Bierme, Richard, Moisan, 2012]  
[Polisano, Clausel, Perrier, Condat, 2014]



# General model :


## anisotropic self-similar Gaussian fields

$$X(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x}\cdot\xi} - 1) f^{1/2}(\mathbf{x}, \xi) d\widehat{W}(\xi)$$

$$f^{1/2}(\mathbf{x}, \xi) = c(\mathbf{x}, \xi) \|\xi\|^{-h(\mathbf{x}, \xi) - 1}$$

spectral density

- $c(\mathbf{x}, \xi) \equiv 1$  and  $h(\mathbf{x}, \xi) \equiv H \Rightarrow X = B^H$  [Mandelbrot, Van Ness, 1968]
- $c(\mathbf{x}, \xi) \equiv c(\arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\arg \xi) \Rightarrow X = AFBF$  [Bonami, Estrade, 2003]
- Example : elementary fields  $c(\arg \xi) = 1_{[-\alpha, \alpha]}(\arg \xi - \alpha_0)$
- $c(\mathbf{x}, \xi) \equiv c(\mathbf{x}, \arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\mathbf{x})$  [Bierme, Richard, Moisan, 2012]  
[Polisano, Clausel, Perrier, Condat, 2014]



# General model :


## anisotropic self-similar Gaussian fields

$$X(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x}\cdot\xi} - 1) f^{1/2}(\mathbf{x}, \xi) d\widehat{W}(\xi)$$

$$f^{1/2}(\mathbf{x}, \xi) = c(\mathbf{x}, \xi) \|\xi\|^{-h(\mathbf{x}, \xi) - 1}$$

spectral density

- $c(\mathbf{x}, \xi) \equiv 1$  and  $h(\mathbf{x}, \xi) \equiv H \Rightarrow X = B^H$  [Mandelbrot, Van Ness, 1968]
- $c(\mathbf{x}, \xi) \equiv c(\arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\arg \xi) \Rightarrow X = AFBF$  [Bonami, Estrade, 2003]
- Example : elementary fields  $c(\arg \xi) = 1_{[-\alpha, \alpha]}(\arg \xi - \alpha_0)$
- $c(\mathbf{x}, \xi) \equiv c(\mathbf{x}, \arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\mathbf{x})$  [Bierme, Richard, Moisan, 2012]  
[Polisano, Clausel, Perrier, Condat, 2014]



# General model :


## anisotropic self-similar Gaussian fields

$$X(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x}\cdot\xi} - 1) f^{1/2}(\mathbf{x}, \xi) d\widehat{W}(\xi)$$

$$f^{1/2}(\mathbf{x}, \xi) = c(\mathbf{x}, \xi) \|\xi\|^{-h(\mathbf{x}, \xi) - 1}$$

spectral density

- $c(\mathbf{x}, \xi) \equiv 1$  and  $h(\mathbf{x}, \xi) \equiv H \Rightarrow X = B^H$  [Mandelbrot, Van Ness, 1968]
- $c(\mathbf{x}, \xi) \equiv c(\arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\arg \xi) \Rightarrow X = AFBF$  [Bonami, Estrade, 2003]
- Example : elementary fields  $c(\arg \xi) = 1_{[-\alpha, \alpha]}(\arg \xi - \alpha_0)$
- $c(\mathbf{x}, \xi) \equiv c(\mathbf{x}, \arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\mathbf{x})$  [Bierme, Richard, Moisan, 2012]  
[Polisano, Clausel, Perrier, Condat, 2014]



# General model :


## anisotropic self-similar Gaussian fields

$$X(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x}\cdot\xi} - 1) f^{1/2}(\mathbf{x}, \xi) d\widehat{W}(\xi)$$

$$f^{1/2}(\mathbf{x}, \xi) = c(\mathbf{x}, \xi) \|\xi\|^{-h(\mathbf{x}, \xi) - 1}$$

spectral density

- $c(\mathbf{x}, \xi) \equiv 1$  and  $h(\mathbf{x}, \xi) \equiv H \Rightarrow X = B^H$  [Mandelbrot, Van Ness, 1968]
- $c(\mathbf{x}, \xi) \equiv c(\arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\arg \xi) \Rightarrow X = AFBF$  [Bonami, Estrade, 2003]
- Example : elementary fields  $c(\arg \xi) = 1_{[-\alpha, \alpha]}(\arg \xi - \alpha_0)$
- $c(\mathbf{x}, \xi) \equiv c(\mathbf{x}, \arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\mathbf{x})$  [Bierme, Richard, Moisan, 2012]  
[Polisano, Clausel, Perrier, Condat, 2014]



# General model :


## anisotropic self-similar Gaussian fields

$$X(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x}\cdot\xi} - 1) f^{1/2}(\mathbf{x}, \xi) d\widehat{W}(\xi)$$

$$f^{1/2}(\mathbf{x}, \xi) = c(\mathbf{x}, \xi) \|\xi\|^{-h(\mathbf{x}, \xi) - 1}$$

spectral density

- $c(\mathbf{x}, \xi) \equiv 1$  and  $h(\mathbf{x}, \xi) \equiv H \Rightarrow X = B^H$  [Mandelbrot, Van Ness, 1968]
- $c(\mathbf{x}, \xi) \equiv c(\arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\arg \xi) \Rightarrow X = AFBF$  [Bonami, Estrade, 2003]
- Example : elementary fields  $c(\arg \xi) = 1_{[-\alpha, \alpha]}(\arg \xi - \alpha_0)$
- $c(\mathbf{x}, \xi) \equiv c(\mathbf{x}, \arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\mathbf{x})$  [Bierme, Richard, Moisan, 2012]  
[Polisano, Clausel, Perrier, Condat, 2014]



# General model :


## anisotropic self-similar Gaussian fields

$$X(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x}\cdot\xi} - 1) f^{1/2}(\mathbf{x}, \xi) d\widehat{W}(\xi)$$

$$f^{1/2}(\mathbf{x}, \xi) = c(\mathbf{x}, \xi) \|\xi\|^{-h(\mathbf{x}, \xi) - 1}$$

spectral density

- $c(\mathbf{x}, \xi) \equiv 1$  and  $h(\mathbf{x}, \xi) \equiv H \Rightarrow X = B^H$  [Mandelbrot, Van Ness, 1968]
- $c(\mathbf{x}, \xi) \equiv c(\arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\arg \xi) \Rightarrow X = AFBF$  [Bonami, Estrade, 2003]
- Example : elementary fields  $c(\arg \xi) = 1_{[-\alpha, \alpha]}(\arg \xi - \alpha_0)$
- $c(\mathbf{x}, \xi) \equiv c(\mathbf{x}, \arg \xi)$  and  $h(\mathbf{x}, \xi) \equiv h(\mathbf{x})$  [Bierme, Richard, Moisan, 2012]  
[Polisano, Clausel, Perrier, Condat, 2014]



# Locally Anisotropic Fractional Brownian Field (LAFBF)

■ **Definition:** Our new Gaussian model LAFBF is a local version of the elementary field

$$B_{\alpha_0, \alpha}^H(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x}\cdot\xi} - 1) \frac{\mathbb{1}_{[-\alpha, \alpha]}(\arg \xi - \alpha_0(\mathbf{x}))}{\|\xi\|^{H+1}} d\widehat{W}(\xi)$$

[Polisano et al., 2014]

# Locally Anisotropic Fractional Brownian Field (LAFBF)

- **Definition:** Our new Gaussian model LAFBF is a local version of the elementary field

$$B_{\alpha_0, \alpha}^H(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x} \cdot \xi} - 1) \frac{\mathbb{1}_{[-\alpha, \alpha]}(\arg \xi - \alpha_0(\mathbf{x}))}{\|\xi\|^{H+1}} d\widehat{W}(\xi)$$


[Polisano et al., 2014]

The orientation may varies spatially.  $\alpha_0$  is now a differentiable function on  $\mathbb{R}^2$

# Locally Anisotropic Fractional Brownian Field (LAFBF)

■ **Definition:** Our new Gaussian model LAFBF is a local version of the elementary field

$$B_{\alpha_0, \alpha}^H(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x} \cdot \xi} - 1) \frac{\mathbb{1}_{[-\alpha, \alpha]}(\arg \xi - \alpha_0(\mathbf{x}))}{\|\xi\|^{H+1}} d\widehat{W}(\xi)$$



$$\frac{\mathbb{1}_{[-\alpha, \alpha]}(\arg \xi - \alpha_0(\mathbf{x}))}{\|\xi\|^{H+1}} d\widehat{W}(\xi)$$



[Polisano et al., 2014]

$$f^{1/2}(\mathbf{x}_0, \xi) = \frac{c_{\alpha_0, \alpha}(\mathbf{x}_0, \arg \xi)}{\|\xi\|^{H+1}}$$

The orientation may varies spatially.  $\alpha_0$  is now a differentiable function on  $\mathbb{R}^2$

# Elementary field



$$\alpha_0 = 0$$



$$\alpha = -\frac{\pi}{2}$$

# Elementary field



$$\alpha_0 = 0$$



$$\alpha = 0.7$$

# Elementary field



$$\alpha_0 = 0$$



$$\alpha = 0.6$$

# Elementary field



$$\alpha_0 = 0$$



$$\alpha = 0.5$$

# Elementary field



$$\alpha_0 = 0$$



$$\alpha = 0.4$$

# Elementary field



$$\alpha_0 = 0$$



$$\alpha = 0.3$$

# Elementary field



$$\alpha_0 = 0$$



$$\alpha = 0.2$$

# Elementary field



$$\alpha_0 = 0$$



$$\alpha = 0.1$$

# Elementary field



$$\alpha_0 = 0$$



$$\alpha = 0.05$$

# Elementary field



$$\alpha_0 = -\frac{\pi}{3}$$



$$\alpha = 0.7$$

# Elementary field



$$\alpha_0 = -\frac{\pi}{3}$$



$$\alpha = 0.6$$

# Elementary field



$$\alpha_0 = -\frac{\pi}{3}$$



$$\alpha = 0.5$$

# Elementary field



$$\alpha_0 = -\frac{\pi}{3}$$



$$\alpha = 0.4$$

# Elementary field


$$\alpha_0 = -\frac{\pi}{3}$$



$$\alpha = 0.3$$

# Elementary field



$$\alpha_0 = -\frac{\pi}{3}$$



$$\alpha = 0.2$$

# Elementary field



$$\alpha_0 = -\frac{\pi}{3}$$



$$\alpha = 0.1$$

# Elementary field

$$\alpha_0 = -\frac{\pi}{3}$$



$$\alpha = 0.05$$

# Tangent field

$$B_{\alpha_0, \alpha}^H(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x} \cdot \xi} - 1) \frac{\mathbb{1}_{[-\alpha, \alpha]}(\arg \xi - \alpha_0(\mathbf{x}))}{\|\xi\|^{H+1}} d\widehat{W}(\xi)$$

## ■ Tangent field.

For a random field  $X$  locally asymptotically self-similar of order  $H$ ,

$$\frac{X(\mathbf{x}_0 + \rho \mathbf{h}) - X(\mathbf{x}_0)}{\rho^H} \xrightarrow[\rho \rightarrow 0]{\mathcal{L}} Y_{\mathbf{x}_0}$$

→  $Y_{\mathbf{x}_0}$  : tangent field of  $X$  at point  $\mathbf{x}_0 \in \mathbb{R}^2$

[Benassi, 1997]  
[Falconer, 2002]

Taylor's expansion



Tangent field

*Deterministic case*

*Stochastic case*

# Tangent field

$$B_{\alpha_0, \alpha}^H(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x} \cdot \xi} - 1) \frac{\mathbb{1}_{[-\alpha, \alpha]}(\arg \xi - \alpha_0(\mathbf{x}))}{\|\xi\|^{H+1}} d\widehat{W}(\xi)$$

## ■ Tangent field.

For a random field  $X$  locally asymptotically self-similar of order  $H$ ,

$$\frac{X(\mathbf{x}_0 + \rho \mathbf{h}) - X(\mathbf{x}_0)}{\rho^H} \xrightarrow[\rho \rightarrow 0]{\mathcal{L}} Y_{\mathbf{x}_0}$$

→  $Y_{\mathbf{x}_0}$  : tangent field of  $X$  at point  $\mathbf{x}_0 \in \mathbb{R}^2$

[Benassi, 1997]  
[Falconer, 2002]

Taylor's expansion



Tangent field

*Deterministic case*

*Stochastic case*

# Tangent field

$$B_{\alpha_0, \alpha}^H(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x} \cdot \xi} - 1) \frac{\mathbb{1}_{[-\alpha, \alpha]}(\arg \xi - \alpha_0(\mathbf{x}))}{\|\xi\|^{H+1}} d\widehat{W}(\xi)$$

■ **Theorem.** The LAFBF  $B_{\alpha_0, \alpha}^H$  admits for tangent field  $Y_{\mathbf{x}_0}$ :

$$Y_{\mathbf{x}_0}(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x} \cdot \xi} - 1) \frac{\mathbb{1}_{[-\alpha, \alpha]}(\arg \xi - \alpha_0(\mathbf{x}_0))}{\|\xi\|^{H+1}} d\widehat{W}(\xi)$$


→  $Y_{\mathbf{x}_0}$  elementary field with global orientation  $\alpha_0(\mathbf{x}_0)$

# Tangent field

$$B_{\alpha_0, \alpha}^H(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x} \cdot \xi} - 1) \frac{\mathbb{1}_{[-\alpha, \alpha]}(\arg \xi - \alpha_0(\mathbf{x}))}{\|\xi\|^{H+1}} d\widehat{W}(\xi)$$

■ **Theorem.** The LAFBF  $B_{\alpha_0, \alpha}^H$  admits for tangent field  $Y_{\mathbf{x}_0}$ :

$$Y_{\mathbf{x}_0}(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x} \cdot \xi} - 1) \frac{\mathbb{1}_{[-\alpha, \alpha]}(\arg \xi - \alpha_0(\mathbf{x}_0))}{\|\xi\|^{H+1}} d\widehat{W}(\xi)$$

constant


→  $Y_{\mathbf{x}_0}$  elementary field with global orientation  $\alpha_0(\mathbf{x}_0)$

# Tangent field


$$B_{\alpha_0, \alpha}^H(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x} \cdot \xi} - 1) \frac{\mathbb{1}_{[-\alpha, \alpha]}(\arg \xi - \alpha_0(\mathbf{x}))}{\|\xi\|^{H+1}} d\widehat{W}(\xi)$$

■ **Theorem.** The LAFBF  $B_{\alpha_0, \alpha}^H$  admits for tangent field  $Y_{\mathbf{x}_0}$ :

$$Y_{\mathbf{x}_0}(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{i\mathbf{x} \cdot \xi} - 1) \frac{\mathbb{1}_{[-\alpha, \alpha]}(\arg \xi - \alpha_0(\mathbf{x}_0))}{\|\xi\|^{H+1}} d\widehat{W}(\xi)$$

constant

→  $Y_{\mathbf{x}_0}$  elementary field with global orientation  $\alpha_0(\mathbf{x}_0)$



$$B_{\alpha_0, \alpha}^H(\mathbf{x}_0) \approx Y_{\mathbf{x}_0}(\mathbf{x} = \mathbf{x}_0)$$

# Simulation of tangent fields

■ **Continuous formulation.** Variogram of  $Y_{\mathbf{x}_0}$  : [Bierme,Richard,Moisan,2012]

$$\begin{aligned} v_{Y_{\mathbf{x}_0}}(\mathbf{x}) &= \frac{1}{2} \int_{\mathbb{R}^2} |e^{i\mathbf{x}\cdot\boldsymbol{\xi}} - 1|^2 f(\mathbf{x}_0, \boldsymbol{\xi}) d\boldsymbol{\xi} \\ &= \frac{1}{2} \gamma(H) \int_{-\pi/2}^{\pi/2} c_{\alpha_0, \alpha}(\mathbf{x}_0, \theta) |\mathbf{x} \cdot \mathbf{u}(\theta)|^{2H} d\theta \\ &= \int_{-\pi/2}^{\pi/2} \tilde{v}_\theta(\mathbf{x} \cdot \mathbf{u}(\theta)) d\theta \end{aligned}$$

# Simulation of tangent fields

■ **Continuous formulation.** Variogram of  $Y_{x_0}$  : [Bierme,Richard,Moisan,2012]

$$\begin{aligned} v_{Y_{x_0}}(\mathbf{x}) &= \frac{1}{2} \int_{\mathbb{R}^2} |e^{i\mathbf{x}\cdot\boldsymbol{\xi}} - 1|^2 f(\mathbf{x}_0, \boldsymbol{\xi}) d\boldsymbol{\xi} \\ &= \frac{1}{2} \gamma(H) \int_{-\pi/2}^{\pi/2} c_{\alpha_0, \alpha}(\mathbf{x}_0, \theta) |\mathbf{x} \cdot \mathbf{u}(\theta)|^{2H} d\theta \\ &\quad \text{in polar coordinates} \\ &= \int_{-\pi/2}^{\pi/2} \tilde{v}_\theta(\mathbf{x} \cdot \mathbf{u}(\theta)) d\theta \end{aligned}$$

# Simulation of tangent fields

■ **Continuous formulation.** Variogram of  $Y_{\mathbf{x}_0}$  : [Bierme,Richard,Moisan,2012]

$$\begin{aligned} v_{Y_{\mathbf{x}_0}}(\mathbf{x}) &= \frac{1}{2} \int_{\mathbb{R}^2} |e^{i\mathbf{x}\cdot\boldsymbol{\xi}} - 1|^2 f(\mathbf{x}_0, \boldsymbol{\xi}) d\boldsymbol{\xi} \\ &= \frac{1}{2} \gamma(H) \int_{-\pi/2}^{\pi/2} c_{\alpha_0, \alpha}(\mathbf{x}_0, \theta) |\mathbf{x} \cdot \mathbf{u}(\theta)|^{2H} d\theta \\ &\quad \text{in polar coordinates} \\ &= \int_{-\pi/2}^{\pi/2} \tilde{v}_\theta(\mathbf{x} \cdot \mathbf{u}(\theta)) d\theta \end{aligned}$$

$$\begin{aligned} \tilde{v}_\theta &= \frac{1}{2} \gamma(H) c_{\alpha_0, \alpha}(\mathbf{x}_0, \theta) |\cdot|^{2H} \\ \mathbf{u}(\theta) &= (\cos \theta, \sin \theta) \\ \gamma(H) &= \frac{\pi}{H\Gamma(2H)\sin(H\pi)} \end{aligned}$$

# Simulation of tangent fields

■ **Continuous formulation.** Variogram of  $Y_{x_0}$  : [Bierme,Richard,Moisan,2012]

$$\begin{aligned} v_{Y_{x_0}}(\mathbf{x}) &= \frac{1}{2} \int_{\mathbb{R}^2} |e^{i\mathbf{x}\cdot\boldsymbol{\xi}} - 1|^2 f(\mathbf{x}_0, \boldsymbol{\xi}) d\boldsymbol{\xi} \\ &= \frac{1}{2} \gamma(H) \int_{-\pi/2}^{\pi/2} c_{\alpha_0, \alpha}(\mathbf{x}_0, \theta) |\mathbf{x} \cdot \mathbf{u}(\theta)|^{2H} d\theta \\ &= \int_{-\pi/2}^{\pi/2} \tilde{v}_\theta(\mathbf{x} \cdot \mathbf{u}(\theta)) d\theta \end{aligned}$$

in polar  
coordinates

variogram of a fractional  
brownian motion (FBM) of order H

$$\begin{aligned} \tilde{v}_\theta &= \frac{1}{2} \gamma(H) c_{\alpha_0, \alpha}(\mathbf{x}_0, \theta) |\cdot|^{2H} \\ \mathbf{u}(\theta) &= (\cos \theta, \sin \theta) \\ \gamma(H) &= \frac{\pi}{H\Gamma(2H)\sin(H\pi)} \end{aligned}$$

# Simulation of tangent fields

■ Continuous formulation. Variogram of  $Y_{\mathbf{x}_0}$  : [Bierme,Richard,Moisan,2012]

$$\begin{aligned} v_{Y_{\mathbf{x}_0}}(\mathbf{x}) &= \frac{1}{2} \int_{\mathbb{R}^2} |e^{i\mathbf{x}\cdot\boldsymbol{\xi}} - 1|^2 f(\mathbf{x}_0, \boldsymbol{\xi}) d\boldsymbol{\xi} \\ &= \frac{1}{2} \gamma(H) \int_{-\pi/2}^{\pi/2} c_{\alpha_0, \alpha}(\mathbf{x}_0, \theta) |\mathbf{x} \cdot \mathbf{u}(\theta)|^{2H} d\theta \\ &= \int_{-\pi/2}^{\pi/2} \tilde{v}_\theta(\mathbf{x} \cdot \mathbf{u}(\theta)) d\theta \end{aligned}$$

in polar  
coordinates

variogram of a fractional  
brownian motion (FBM) of order H

$Y_{\mathbf{x}_0}$   
=

Infinite sum of independant  
rotating FBM of order H

$$\begin{aligned} \tilde{v}_\theta &= \frac{1}{2} \gamma(H) c_{\alpha_0, \alpha}(\mathbf{x}_0, \theta) |\cdot|^{2H} \\ \mathbf{u}(\theta) &= (\cos \theta, \sin \theta) \\ \gamma(H) &= \frac{\pi}{H\Gamma(2H)\sin(H\pi)} \end{aligned}$$

# Simulation of tangent fields

## ■ Discrete formulation.

[Bierme, Richard, Moisan, 2012]

- $(\theta_i)_{1 \leq i \leq n}$  are  $n$  bands orientations and  $\lambda_i = \theta_{i+1} - \theta_i$

- The **turning band field** is defined as

$$Y_{\mathbf{x}_0}^{[n]}(\mathbf{x}) = \gamma(H)^{\frac{1}{2}} \sum_{i=1}^n \sqrt{\lambda_i c_{\alpha_0, \alpha}(\mathbf{x}_0, \theta_i)} B_i^H(\mathbf{x} \cdot \mathbf{u}(\theta_i))$$

- $B_i^H$  are  $n$  independent FBM of order  $H$

- Good approximation provided  $\max_i \lambda_i \leq \varepsilon$

# Simulation of tangent fields

## ■ Discrete formulation.

[Bierme, Richard, Moisan, 2012]

- $(\theta_i)_{1 \leq i \leq n}$  are  $n$  bands orientations and  $\lambda_i = \theta_{i+1} - \theta_i$

- The **turning band field** is defined as

$$Y_{\mathbf{x}_0}^{[n]}(\mathbf{x}) = \gamma(H)^{\frac{1}{2}} \sum_{i=1}^n \sqrt{\lambda_i c_{\alpha_0, \alpha}(\mathbf{x}_0, \theta_i)} B_i^H(\mathbf{x} \cdot \mathbf{u}(\theta_i))$$

- $B_i^H$  are  $n$  independent FBM of order  $H$

- Good approximation provided  $\max_i \lambda_i \leq \varepsilon$

not equispaced 

# Simulation of tangent fields

## ■ Simulation along particular bands.

[Bierme, Richard, Moisan, 2012]

- Discrete grid  $r^{-1}\mathbb{Z}^2 \cap [0, 1]^2$  with  $r = 2^k - 1, k \in \mathbb{N}^*$
- Choose  $(\theta_i)$  such that  $\tan \theta_i = \frac{p_i}{q_i}$  and  $\max_i \lambda_i \leq \epsilon$
- Then  $B_i^H(\mathbf{x} \cdot \mathbf{u}(\theta_i))$  becomes

$$\left\{ B_i^H \left( \frac{k_1}{r} \cos \theta_i + \frac{k_2}{r} \sin \theta_i \right); 0 \leq k_1, k_2 \leq r \right\} \stackrel{\mathcal{L}}{=} \left( \frac{\cos \theta_i}{rq_i} \right)^H \{ B_i^H(k_1 q_i + k_2 p_i); 0 \leq k_1, k_2 \leq r \}$$

# Simulation of tangent fields

## ■ Simulation along particular bands.

[Bierme, Richard, Moisan, 2012]

- Discrete grid  $r^{-1}\mathbb{Z}^2 \cap [0, 1]^2$  with  $r = 2^k - 1, k \in \mathbb{N}^*$
- Choose  $(\theta_i)$  such that  $\tan \theta_i = \frac{p_i}{q_i}$  and  $\max_i \lambda_i \leq \epsilon$
- Then  $B_i^H(\mathbf{x} \cdot \mathbf{u}(\theta_i))$  becomes

Dynamic programming

$$\left\{ B_i^H \left( \frac{k_1}{r} \cos \theta_i + \frac{k_2}{r} \sin \theta_i \right); 0 \leq k_1, k_2 \leq r \right\} \stackrel{\mathcal{L}}{=} \left( \frac{\cos \theta_i}{rq_i} \right)^H \{ B_i^H(k_1 q_i + k_2 p_i); 0 \leq k_1, k_2 \leq r \}$$

# Simulation of tangent fields

## ■ Simulation along particular bands.

[Bierme, Richard, Moisan, 2012]

- Discrete grid  $r^{-1}\mathbb{Z}^2 \cap [0, 1]^2$  with  $r = 2^k - 1, k \in \mathbb{N}^*$
- Choose  $(\theta_i)$  such that  $\tan \theta_i = \frac{p_i}{q_i}$  and  $\max_i \lambda_i \leq \epsilon$
- Then  $B_i^H(\mathbf{x} \cdot \mathbf{u}(\theta_i))$  becomes

Dynamic programming

self-similarity

$$B^H(\lambda \cdot) \stackrel{\mathcal{L}}{=} \lambda^H B^H(\cdot)$$
$$\left\{ B_i^H \left( \frac{k_1}{r} \cos \theta_i + \frac{k_2}{r} \sin \theta_i \right); 0 \leq k_1, k_2 \leq r \right\} \stackrel{\mathcal{L}}{=} \left( \frac{\cos \theta_i}{rq_i} \right)^H \{ B_i^H(k_1 q_i + k_2 p_i); 0 \leq k_1, k_2 \leq r \}$$

# Simulation of tangent fields

## ■ Simulation along particular bands.


[Bierme, Richard, Moisan, 2012]

- Discrete grid  $r^{-1}\mathbb{Z}^2 \cap [0, 1]^2$  with  $r = 2^k - 1, k \in \mathbb{N}^*$
- Choose  $(\theta_i)$  such that  $\tan \theta_i = \frac{p_i}{q_i}$  and  $\max_i \lambda_i \leq \epsilon$
- Then  $B_i^H(\mathbf{x} \cdot \mathbf{u}(\theta_i))$  becomes

Dynamic programming

self-similarity 

$$B^H(\lambda \cdot) \stackrel{\mathcal{L}}{=} \lambda^H B^H(\cdot)$$
$$\left\{ B_i^H \left( \frac{k_1}{r} \cos \theta_i + \frac{k_2}{r} \sin \theta_i \right); 0 \leq k_1, k_2 \leq r \right\} \stackrel{\mathcal{L}}{=} \left( \frac{\cos \theta_i}{rq_i} \right)^H \{ B_i^H(k_1 q_i + k_2 p_i); 0 \leq k_1, k_2 \leq r \}$$

 equispaced 

# Simulation of LAFBF

## using tangent fields

[Polisano et al., 2014]

■ **Algorithm.** For each pixel  $\mathbf{x}_0 = (k_1, k_2) \in [0, r]^2$

$$B_{\alpha_0, \alpha}^H((k_1, k_2)) = \gamma(H)^{\frac{1}{2}} \sum_{i=1}^n \sqrt{\lambda_i c_{\alpha_0, \alpha}((k_1, k_2), \theta_i)} \left( \frac{\cos \theta_i}{r q_i} \right)^H B_i^H(k_1 q_i + k_2 p_i)$$

# Simulation of LAFBF

## using tangent fields

[Polisano et al., 2014]

■ **Algorithm.** For each pixel  $\mathbf{x}_0 = (k_1, k_2) \in [0, r]^2$

$$B_{\alpha_0, \alpha}^H((k_1, k_2)) = B_{\alpha_0, \alpha}^H(\mathbf{x}_0) \approx Y_{\mathbf{x}_0}(x = \mathbf{x}_0)$$
$$\gamma(H)^{\frac{1}{2}} \sum_{i=1}^n \sqrt{\lambda_i c_{\alpha_0, \alpha}((k_1, k_2), \theta_i)} \left( \frac{\cos \theta_i}{r q_i} \right)^H B_i^H(k_1 q_i + k_2 p_i)$$

# Simulation of LAFBF

## using tangent fields

[Polisano et al., 2014]

■ **Algorithm.** For each pixel  $\mathbf{x}_0 = (k_1, k_2) \in [0, r]^2$

$$B_{\alpha_0, \alpha}^H((k_1, k_2)) = \gamma(H)^{\frac{1}{2}} \sum_{i=1}^n \sqrt{\lambda_i c_{\alpha_0, \alpha}((k_1, k_2), \theta_i)} \left( \frac{\cos \theta_i}{r q_i} \right)^H B_i^H(k_1 q_i + k_2 p_i)$$


# Simulation of LAFBF

## using tangent fields

[Polisano et al., 2014]

■ **Algorithm.** For each pixel  $\mathbf{x}_0 = (k_1, k_2) \in \llbracket 0, r \rrbracket^2$

$$B_{\alpha_0, \alpha}^H((k_1, k_2)) = \gamma(H)^{\frac{1}{2}} \sum_{i=1}^n \sqrt{\lambda_i c_{\alpha_0, \alpha}((k_1, k_2), \theta_i)} \left( \frac{\cos \theta_i}{r q_i} \right)^H B_i^H(k_1 q_i + k_2 p_i)$$



# Simulation of LAFBF

## using tangent fields


[Polisano et al., 2014]

■ **Algorithm.** For each pixel  $\mathbf{x}_0 = (k_1, k_2) \in [0, r]^2$

$$B_{\alpha_0, \alpha}^H((k_1, k_2))$$

=

$$\gamma(H)^{\frac{1}{2}} \sum_{i=1}^n \sqrt{\lambda_i c_{\alpha_0, \alpha}((k_1, k_2), \theta_i)} \left( \frac{\cos \theta_i}{r q_i} \right)^H B_i^H(k_1 q_i + k_2 p_i)$$



# Simulation of LAFBF

## using tangent fields


[Polisano et al., 2014]

■ **Algorithm.** For each pixel  $\mathbf{x}_0 = (k_1, k_2) \in \llbracket 0, r \rrbracket^2$

$$B_{\alpha_0, \alpha}^H((k_1, k_2))$$

=

$$\gamma(H)^{\frac{1}{2}} \sum_{i=1}^n \sqrt{\lambda_i c_{\alpha_0, \alpha}((k_1, k_2), \theta_i)} \left( \frac{\cos \theta_i}{r q_i} \right)^H B_i^H(k_1 q_i + k_2 p_i)$$



*n* turning bands  $\theta_i$

$$\begin{aligned} 1_{[-\alpha, \alpha]}(\theta_i - \alpha_0((k_1, k_2))) &\neq 0 \\ \Leftrightarrow |\theta_i - \alpha_0((k_1, k_2))| &\leq \alpha \end{aligned}$$

# Simulation of LAFBF

## using tangent fields


[Polisano et al., 2014]

■ **Algorithm.** For each pixel  $\mathbf{x}_0 = (k_1, k_2) \in \llbracket 0, r \rrbracket^2$

$$B_{\alpha_0, \alpha}^H((k_1, k_2))$$

=

$$\gamma(H)^{\frac{1}{2}} \sum_{i=1}^n \sqrt{\lambda_i c_{\alpha_0, \alpha}((k_1, k_2), \theta_i)} \left( \frac{\cos \theta_i}{r q_i} \right)^H B_i^H(k_1 q_i + k_2 p_i)$$



*n* turning bands  $\theta_i$

$$\begin{aligned} 1_{[-\alpha, \alpha]}(\theta_i - \alpha_0((k_1, k_2))) \neq 0 \\ \Leftrightarrow |\theta_i - \alpha_0((k_1, k_2))| \leq \alpha \end{aligned}$$

# Simulation of LAFBF

## using tangent fields


[Polisano et al., 2014]

■ **Algorithm.** For each pixel  $\mathbf{x}_0 = (k_1, k_2) \in \llbracket 0, r \rrbracket^2$

$$B_{\alpha_0, \alpha}^H((k_1, k_2))$$

=

$$\gamma(H)^{\frac{1}{2}} \sum_{i=1}^n \sqrt{\lambda_i c_{\alpha_0, \alpha}((k_1, k_2), \theta_i)} \left( \frac{\cos \theta_i}{r q_i} \right)^H B_i^H(k_1 q_i + k_2 p_i)$$



$$\begin{aligned} 1_{[-\alpha, \alpha]}(\theta_i - \alpha_0((k_1, k_2))) \neq 0 \\ \Leftrightarrow |\theta_i - \alpha_0((k_1, k_2))| \leq \alpha \end{aligned}$$

# Simulation of LAFBF

## using tangent fields


[Polisano et al., 2014]

■ **Algorithm.** For each pixel  $\mathbf{x}_0 = (k_1, k_2) \in \llbracket 0, r \rrbracket^2$

$$B_{\alpha_0, \alpha}^H((k_1, k_2))$$

=

$$\gamma(H)^{\frac{1}{2}} \sum_{i=1}^n \sqrt{\lambda_i c_{\alpha_0, \alpha}((k_1, k_2), \theta_i)} \left( \frac{\cos \theta_i}{r q_i} \right)^H B_i^H(k_1 q_i + k_2 p_i)$$



$n$  turning bands  $\theta_i$

few bands  
in the cone

$$\mathbb{1}_{[-\alpha, \alpha]}(\theta_i - \alpha_0((k_1, k_2))) \neq 0$$


$\Leftrightarrow$

$$|\theta_i - \alpha_0((k_1, k_2))| \leq \alpha$$


Complexity  $O(r^2 \log n)$

# Numerical experiments


$$\vec{V}_{(x,y)}^1 = (\cos(-\pi/2 + y), \sin(-\pi/2))$$



(a)



(b)



(c)


Parameters

$$r = 255 \quad H = 0.2$$

$$\alpha = 10^{-1} \quad \epsilon = 10^{-2}$$


# Numerical experiments

$$\vec{V}_{(x,y)}^1 = (\cos(-\pi/2 + y), \sin(-\pi/2))$$



(a)

(b)



(c)

Texture with prescribed local orientation at each point  $\mathbf{x}_0$  given by a vector field

$$\vec{V}_{\mathbf{x}_0} = \mathbf{u}(\alpha_0(\mathbf{x}_0))$$


Parameters

$$r = 255 \quad H = 0.2$$


$$\alpha = 10^{-1} \quad \epsilon = 10^{-2}$$

# Numerical experiments

$$\vec{V}_{(x,y)}^1 = (\cos(-\pi/2 + y), \sin(-\pi/2))$$



(a)



(c)

Texture with prescribed local orientation at each point  $\mathbf{x}_0$  given by a vector field

$$\vec{V}_{\mathbf{x}_0} = \mathbf{u}(\alpha_0(\mathbf{x}_0))$$

A zoom around a point  $\mathbf{x}_0$  shows that locally a LAFBF behaves as an elementary field


Parameters

$$r = 255 \quad H = 0.2$$

$$\alpha = 10^{-1} \quad \epsilon = 10^{-2}$$

# Numerical experiments

$$\vec{V}_{(x,y)}^1 = (\cos(-\pi/2 + y), \sin(-\pi/2))$$




(a)

Texture with prescribed local orientation at each point  $\mathbf{x}_0$  given by a vector field

$$\vec{V}_{\mathbf{x}_0} = \mathbf{u}(\alpha_0(\mathbf{x}_0))$$

(b)

A zoom around a point  $\mathbf{x}_0$  shows that locally a LAFBF behaves as an elementary field



(c)

Regularized version of the anisotropy function


Parameters

$$r = 255 \quad H = 0.2$$

$$\alpha = 10^{-1} \quad \epsilon = 10^{-2}$$

# Numerical experiments

$$\vec{V}_{(x,y)}^1 = (\cos(-\pi/2 + y), \sin(-\pi/2))$$




(a)

Texture with prescribed local orientation at each point  $\mathbf{x}_0$  given by a vector field

$$\vec{V}_{\mathbf{x}_0} = \mathbf{u}(\alpha_0(\mathbf{x}_0))$$


A zoom around a point  $\mathbf{x}_0$  shows that locally a LAFBF behaves as an elementary field



(b)

Regularized version of the anisotropy function

$$f^{1/2}(\mathbf{x}_0, \xi) = \frac{c_{\alpha_0, \alpha}(\mathbf{x}_0, \arg \xi)}{\|\xi\|^{H+1}}$$



(c)

Parameters

$$r = 255 \quad H = 0.2$$

$$\alpha = 10^{-1} \quad \epsilon = 10^{-2}$$

# Numerical experiments

---


$$\vec{\mathbf{V}}_{(x,y)}^2 = (\cos(\cos(36xy)), \sin(\cos(36xy))) \quad \vec{\mathbf{V}}_{(x,y)}^3 = \nabla F(x, y)$$

$$F(x, y) = (4x - 2)e^{-(4x-2)^2 - (4y-2)^2}$$

# Numerical experiments

$$\vec{\mathbf{V}}_{(x,y)}^2 = (\cos(\cos(36xy)), \sin(\cos(36xy)))$$


$$\vec{\mathbf{V}}_{(x,y)}^3 = \nabla F(x, y)$$




$$F(x, y) = (4x - 2)e^{-(4x-2)^2 - (4y-2)^2}$$

# Numerical experiments

$$\vec{\mathbf{V}}_{(x,y)}^2 = (\cos(\cos(36xy)), \sin(\cos(36xy)))$$



$$\vec{\mathbf{V}}_{(x,y)}^3 = \nabla F(x, y)$$



$$F(x, y) = (4x - 2)e^{-(4x-2)^2-(4y-2)^2}$$

# Numerical experiments


H=0.2

H=0.5

$$\vec{V}_{(x,y)}^1 = (\cos(-\pi/2 + y), \sin(-\pi/2))$$

# Numerical experiments

H=0.2




H=0.5


$$\vec{V}_{(x,y)}^1 = (\cos(-\pi/2 + y), \sin(-\pi/2))$$

# Numerical experiments

H=0.2



H=0.5



$$\vec{V}_{(x,y)}^1 = (\cos(-\pi/2 + y), \sin(-\pi/2))$$

# Conclusion and future work

---

## ■ Conclusion

# Conclusion and future work

---

## ■ Conclusion

- Introduce a **new stochastic model** defined as a local version of an AFBF.

# Conclusion and future work

---

## ■ Conclusion

- Introduce a **new stochastic model** defined as a local version of an AFBF.
- Simulations based on **tangent field formulation** and the **turning bands method** produce textures with prescribed local orientations.

# Conclusion and future work

## ■ Conclusion

- Introduce a **new stochastic model** defined as a local version of an AFBF.
- Simulations based on **tangent field formulation** and the **turning bands method** produce textures with prescribed local orientations.

# Conclusion and future work

## ■ Conclusion

- Introduce a **new stochastic model** defined as a local version of an AFBF.
- Simulations based on **tangent field formulation** and the **turning bands method** produce textures with prescribed local orientations.

## ■ Future work

- Extensions of our model include Hurst index may **vary spatially**.


# Conclusion and future work

## ■ Conclusion

- Introduce a **new stochastic model** defined as a local version of an AFBF.
- Simulations based on **tangent field formulation** and the **turning bands method** produce textures with prescribed local orientations.

## ■ Future work


- Extensions of our model include Hurst index may **vary spatially**.



# Conclusion and future work


## ■ Conclusion

- Introduce a **new stochastic model** defined as a local version of an AFBF.
- Simulations based on **tangent field formulation** and the **turning bands method** produce textures with prescribed local orientations.



## ■ Future work

- Extensions of our model include Hurst index may **vary spatially**.



# Bibliography

## ■ Selected papers

- K. Polisano, M. Clausel, V. Perrier and L. Condat, "Texture modeling by Gaussian fields with prescribed local orientation", *IEEE ICIP*, 2014.
- A. Bonami and A. Estrade, "Anisotropic analysis of some Gaussian models", *Journal of Fourier Analysis and Applications*, vol. 9, no. 3, pp. 215–236, 2003.
- H. Bierme, L. Moisan, and F. Richard, "A turning-band method for the simulation of anisotropic fractional Brownian fields," *preprint MAP5 No. 2012-312012*, 2012.
- K.J. Falconer, "Tangent fields and the local structure of random fields," *Journal of Theoretical Probability*, vol. 15, no. 3, pp. 731–750, 2002.

# Questions ?

---

# Questions ?

---

**Thank you for your attention.**

■ **Dynamic programming.** The choice of the bands orientation  $(\theta_i)_{1 \leq i \leq n}$

is governed by the computational cost of the  $B_i^H$ 's within dynamic programming.

■ Let the error  $\epsilon$  fixed. Taking  $N = \lceil \frac{1}{\tan \epsilon} \rceil$  consider the following set:

$$\mathcal{V}_N = \left\{ (p, q) \in \mathbb{Z}^2 / -N \leq p \leq N, 1 \leq q \leq N, p \wedge q = 1, -\frac{\pi}{2} < \arctan \left( \frac{p}{q} \right) < \frac{\pi}{2} \right\}$$

■ The aim is to find  $n$  pairs in the set  $\mathcal{V}_N$  which minimize the following global cost:

$$C(\Theta) = \sum_{k=1}^s C(r(|p_{i_k}| + q_{i_k}))$$

where  $C(\ell)$  is the cost of one FBM  $B_i^H$  in  $O(n \log n)$ , under the constraint  $\max_i (\theta_{i+1} - \theta_i) \leq \epsilon$