



Grassmann Extrapolation of Density Matrices for Born–Oppenheimer Molecular Dynamics

Geneviève Dusson

► To cite this version:

Geneviève Dusson. Grassmann Extrapolation of Density Matrices for Born–Oppenheimer Molecular Dynamics. 16ème Colloque National en Calcul de Structures (CSMA 2024), CNRS; CSMA; ENS Paris-Saclay; CentraleSupélec, May 2024, Hyères, France. hal-04610945

HAL Id: hal-04610945

<https://hal.science/hal-04610945v1>

Submitted on 3 Dec 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Grassmann Extrapolation of Density Matrices for Born-Oppenheimer Molecular Dynamics

G. Dusson¹

¹ Laboratoire de Mathématiques de Besançon, Université de Franche-Comté, genevieve.dusson@math.cnrs.fr

Résumé — Ab-initio molecular dynamics simulations are powerful but require expensive calculations. In general, the most costly part of the computation consists of the many resolutions of the Kohn-Sham equations arising at each time-step of the simulation. Proposing good initial guesses of the density matrix for the Self-Consistent Field procedure is therefore key to reduce the overall computational cost of the simulation.

In this talk, I will first present a method to provide such guesses based on previous calculations, exploiting the geometrical structure of the manifold to which the density matrix belongs, in order to perform linear extrapolation in a vector space, while at the same time, retaining the correct physical properties of the extrapolated density. I will also show how the time-reversibility of the simulation can be obtained. I will then illustrate the performance of the method by showing numerical results on a few real-life, multiscale, polarizable QM/MM simulations. Third, I will provide a theoretical analysis of the method for a better understanding of the observed results.

Mots clés — Ab initio ; molecular dynamics ; Grassmann manifold ; molecular descriptors