



**HAL**  
open science

# Wasserstein convergence of Čech persistence diagrams for samplings of submanifolds

Charles Arnal, David Cohen-Steiner, Vincent Divol

► **To cite this version:**

Charles Arnal, David Cohen-Steiner, Vincent Divol. Wasserstein convergence of Čech persistence diagrams for samplings of submanifolds. NeurIPS 2024, Dec 2024, Vancouver (Canada), Canada. hal-04617508

**HAL Id: hal-04617508**

**<https://hal.science/hal-04617508v1>**

Submitted on 20 Jun 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# Wasserstein convergence of Čech persistence diagrams for samplings of submanifolds

Charles Arnal<sup>\*1</sup>, David Cohen-Steiner<sup>\*2</sup>, and Vincent Divol<sup>\*3</sup>

<sup>1</sup>Université Paris-Saclay, CNRS, Inria, Laboratoire de Mathématiques d’Orsay

<sup>2</sup>Université Côte d’Azur, Inria Sophia Antipolis-Méditerranée

<sup>3</sup>CEREMADE, Université Paris-Dauphine, Université PSL

June 20, 2024

## Preamble

Čech Persistence diagrams (PDs) are topological descriptors routinely used to capture the geometry of complex datasets. They are commonly compared using the Wasserstein distances  $OT_p$ ; however, the extent to which PDs are stable with respect to these metrics remains poorly understood. We partially close this gap by focusing on the case where datasets are sampled on an  $m$ -dimensional submanifold of  $\mathbb{R}^d$ . Under this manifold hypothesis, we show that convergence with respect to the  $OT_p$  metric happens exactly when  $p > m$ . We also provide improvements upon the bottleneck stability theorem in this case and prove new laws of large numbers for the total  $\alpha$ -persistence of PDs. Finally, we show how these theoretical findings shed new light on the behavior of the feature maps on the space of PDs that are used in ML-oriented applications of Topological Data Analysis.

## 1 Introduction

Topological Data Analysis (TDA) is a set of tools that aims at extracting relevant topological information from complex datasets, e.g. regarding connected components, loops, cavities, or higher dimensional features. These different notions are made formal through the use of *homology theory*, and in particular the  $i$ -th homology group  $H_i(A)$  with coefficients in some field  $\mathcal{F}$  of a set  $A$ , which captures the  $i$ -dimensional topological features of  $A$  for  $i \geq 0$ , see e.g. [Hat00]. TDA has been successfully applied in a variety of domains, including material science [KKP<sup>+</sup>16, STR<sup>+</sup>17, DKR<sup>+</sup>18, BHO18], biology [CR20, SL22, ACC<sup>+</sup>22], and neuroscience [SPCGB19, RYB<sup>+</sup>20, CPH21], to name a few. When used in conjunction with more traditional approaches such as neural networks, TDA-based methods have outperformed state of the arts methods for tasks such as graph classifications [CCI<sup>+</sup>20, KKZ<sup>+</sup>20].

The most prominent techniques in TDA rely on multiscale approaches, in particular through the use of *persistent homology* [CDSGO16]. Given a compact set  $A$  in  $\mathbb{R}^d$ , persistent homology tracks the evolution of the homology groups  $H_i(A^t)$  of the  $t$ -offset  $A^t = \bigcup_{x \in A} \overline{B}(x, t)$  of  $A$  as  $t$  goes from 0 to  $+\infty$  (where  $\overline{B}(x, t)$  is the closed ball of radius  $t$  centered at  $x$ ). The process is summarized by the *Čech persistence diagram (PD) of degree  $i$*  of the set  $A$ : the PD  $\text{dgm}_i(A)$  is

---

<sup>\*</sup>The authors contributed equally to this work.




Figure 1: The Čech PD of a point cloud  $A$  in  $\mathbb{R}^2$  for  $i = 1$  and its  $t$ -offsets. The two points far from the diagonal  $\partial\Omega$  in  $\text{dgm}_i(A)$  correspond to the two large cycles in the set  $A$ .

a multiset<sup>1</sup> of points in the half-plane  $\Omega := \{(u_1, u_2) \in \mathbb{R}^2 : u_1 < u_2\}$ , where each point  $(u_1, u_2)$  in the PD corresponds to a  $i$ -dimensional topological feature that appeared in  $A^t$  at scale  $t = u_1$  (its birth time) and disappeared at scale  $t = u_2$  (its death time), see Figure 1.<sup>2</sup> Points close to the diagonal  $\partial\Omega = \{(u_1, u_2) \in \mathbb{R}^2 : u_1 = u_2\}$  correspond to topological features of small persistence  $\text{pers}(u) = \frac{u_2 - u_1}{2}$ , which have a short lifetime in the filtration  $(A^t)_{t \geq 0}$ .

A key property of PDs is their robustness to small perturbation in the data, making them suitable for analyzing real-world datasets. This stability property is phrased in terms of the *bottleneck distance* between PDs [ELZ02]. Let  $a$  and  $b$  be two PDs. A *partial matching* between  $a$  and  $b$  is a bijection of multisets  $\gamma : a \cup \partial\Omega \rightarrow b \cup \partial\Omega$ , where each point  $(u, u)$  of  $\partial\Omega$  has infinite multiplicity. In other words, the points of  $a$  are either paired with a single point of  $b$ , or mapped to the diagonal  $\partial\Omega$  (and similarly for the points of  $b$ ). Let  $\Gamma(a, b)$  be the set of all partial matchings between  $a$  and  $b$ . The bottleneck distance is defined as

$$\text{OT}_\infty(a, b) = \inf_{\gamma \in \Gamma(a, b)} \max_{u \in a \cup \partial\Omega} \|u - \gamma(u)\|_\infty. \quad (1)$$

The Bottleneck Stability Theorem [CSEH05, CDSGO16] states that if  $A_1$  and  $A_2$  are two compact sets, then for any integer  $i \geq 0$

$$\text{OT}_\infty(\text{dgm}_i(A_1), \text{dgm}_i(A_2)) \leq \varepsilon, \quad (2)$$

where  $\varepsilon$  is the Hausdorff distance between the sets  $A_1$  and  $A_2$ , defined by

$$d_H(A_1, A_2) = \sup_{x \in \mathbb{R}^d} |d_{A_1}(x) - d_{A_2}(x)|$$

and where  $d_A$  is the distance function to a set  $A$ . An important property of the bottleneck distance is that it is blind to small-persistence topological features: if  $\text{OT}_\infty(a, b) = \varepsilon$ , then one can arbitrarily modify the PDs  $a$  and  $b$  on a slab of width  $\varepsilon$  above the diagonal (for the  $\ell_\infty$ -metric) without changing the bottleneck distance between the two PDs.

Due to this phenomenon, the bottleneck distance turns out to be too weak in many situations of interest, where some topological features of small persistence can be as important as large-scale topological features in the PD (say, with a classification or a regression task in mind). For

<sup>1</sup>A multiset is a set where each element appears with some non-zero multiplicity.

<sup>2</sup>In general, PDs can have points with infinite coordinates. For Čech PDs, this will only be the case for a single point of the diagram for  $i = 0$ , of coordinate  $(0, +\infty)$ . We discard this point in the following.

this reason, finer transport-like distances are often preferred to compare PDs. These distances, which we denote as  $OT_p$ , are defined for  $1 \leq p < \infty$  by

$$OT_p(a, b) = \inf_{\gamma \in \Gamma(a, b)} \left( \sum_{u \in a \cup \partial\Omega} \|u - \gamma(u)\|_\infty^p \right)^{1/p}, \quad (3)$$

with  $OT_p \leq OT_{p'}$  for  $1 \leq p' \leq p < \infty$ . They can be seen as modified versions of the Wasserstein distances used in optimal transport, with the diagonal  $\partial\Omega$  playing the role of a landfill of infinite mass, see e.g. [DL21].

The increased sensitivity to small perturbations of the  $OT_p$  distances is of crucial importance in the standard TDA pipeline, which we briefly recall. Starting from a sample of sets  $A_1, \dots, A_n$ , one computes a sample of PDs  $a_j = \text{dgm}_i(A_j)$ ,  $j = 1, \dots, n$ . Statistical methods to analyze this sample of PDs are typically awkward to define, due to the nonlinear geometry of the space of PDs [BW20, Wag21]. To overcome this issue, the space of PDs is mapped to a vector space through some map  $\Phi$  called a feature map. Statistical methods are then applied in the feature space on the transformed sample  $\Phi(a_1), \dots, \Phi(a_n)$ . Various feature maps have been designed [CFL+14, RHBK15, CWRW15, KHF16, CCO17, STR+18, KFH18, HKN19], important examples including persistent images [AEK+17], PersLay [CCI+20], and PLLay [KKZ+20]; in the latter two, the feature map is parametrized by a neural network. A good feature map should preserve as much as possible the geometry of the space of PDs [MV21]; in particular, Lipschitz (or Hölder) continuity of the feature map is a basic requirement. However, due to their (often desirable) sensitivity to small scale features, most common feature maps are *not* regular with respect to the  $OT_\infty$  distance on the space of diagrams. Instead, they enjoy Lipschitz regularity with respect to either the finer  $OT_1$  distance (see e.g. [DL21, Proposition 5.2]), or to the  $OT_p$  distances (for  $p > 1$ ) when restricted to diagrams  $a$  whose  $\alpha$ -total persistence

$$\text{Pers}_\alpha(a) = \sum_{u \in a} \text{pers}(u)^\alpha \quad (4)$$

is bounded for some  $\alpha > 0$  large enough, see [DP19] and [KFH18]. Boundedness assumptions on the  $\alpha$ -total persistence also yield a version of the Bottleneck Stability Theorem with respect to the finer  $OT_p$  distance: if  $A_1$  and  $A_2$  are such that  $\text{Pers}_\alpha(\text{dgm}_i(A_k)) \leq M$  ( $k = 1, 2$ ), then, for  $p \geq \alpha$ ,

$$OT_p^p(\text{dgm}_i(A_1), \text{dgm}_i(A_2)) \leq M\varepsilon^{p-\alpha}, \quad (5)$$

where  $\varepsilon = d_H(A_1, A_2)$ , see [CSEHM10]. Hence, in addition to having intrinsic theoretical interest, controlling the total persistence and convergence with respect to the  $OT_p$  distance of PDs is crucial to ensure the soundness of most methods commonly used in TDA. This is the subject of this article.

**Contributions.** We provide a deeper understanding of the structure of Čech PDs in the specific case where the underlying set  $A$  is a compact subset of an  $m$ -dimensional manifold  $M$  in  $\mathbb{R}^d$ , focusing in particular on the total persistence of the PDs and their convergence to the PDs of  $M$  with respect to the  $OT_p$  distances. The importance of this case is supported by the *manifold hypothesis*, which often serves as a fundamental principle guiding the development of algorithms and models for data analysis [GPPVW12, WB19, BCR+22]. Specifically, our main contributions are the following:

- **Theorem 2.3:** When  $A \subset M$  is a compact set satisfying  $d_H(A, M) \leq \varepsilon$  for  $\varepsilon$  small enough, we provide a quadratic improvement upon the standard Bottleneck Stability Theorem (2). Namely, we show that there exists an optimal bottleneck matching  $\gamma$  such that the distance between a coordinate of a point  $u \in \text{dgm}_i(A)$  and the coordinate of the matched point  $\gamma(u) \in \text{dgm}_i(M) \cup \partial\Omega$  is of order  $O(\varepsilon^2)$  whenever the coordinate of  $u$  is larger than  $2\varepsilon$ .
- **Theorem 3.7:** In the case where the manifold  $M$  is generic and the set  $A$  is a  $\delta$ -sparse point cloud (i.e.  $\min_{x \neq y \in A} \|x - y\| \geq \delta$ ), we provide a finer analysis by showing that the  $p$ -total persistence  $\text{Pers}_p(\text{dgm}_i(A))$  remains bounded and the distance  $\text{OT}_p(\text{dgm}_i(A), \text{dgm}_i(M))$  converges to 0 for all  $p > m$  whenever the ratio  $\varepsilon/\delta$  is upper bounded.
- **Corollary 4.17:** We then focus on a random context, by assuming that  $A = A_n$  is obtained by sampling  $n$  i.i.d. random variables with positive bounded density  $f$  on a generic manifold  $M$ . We prove that  $\text{OT}_p(\text{dgm}_i(A_n), \text{dgm}_i(M))$  converges in expectation to 0 for  $p > m$ . Furthermore, we obtain a law of large numbers for the  $\alpha$ -total persistence of  $\text{dgm}_i(A_n)$ :

$$\text{Pers}_\alpha(\text{dgm}_i(A_n)) = \text{Pers}_\alpha(\text{dgm}_i(M)) + C_i n^{1-\alpha/m} + o_{L^1}(n^{1-\alpha/m}) + O_{L^1}\left(\left(\frac{\log n}{n}\right)^{\frac{1}{m}}\right) \quad (6)$$

for all  $\alpha > 0$ , where  $C_i$  is a constant that depends explicitly on  $M$  and  $f$ . In particular, for  $0 \leq i < m$ ,  $\text{Pers}_\alpha(\text{dgm}_i(A_n))$  stays bounded if and only if  $\alpha \geq m$ .

Our contributions are to be compared to one of the only preexisting results regarding the  $\alpha$ -total persistence of a PD: in [CSEHM10], the authors proved that for all  $\alpha$  strictly greater than the ambient dimension  $d$ , the  $\alpha$ -total persistence of the Čech PD of a compact set  $A \subset B(0, R) \subset \mathbb{R}^d$  satisfies

$$\text{Pers}_\alpha(\text{dgm}_i(A)) \leq C_{\alpha,d} R^\alpha \quad (7)$$

for some constant  $C_{\alpha,d}$  depending on  $\alpha$  and  $d$ . In the two scenarios we considered (either  $\delta$ -sparse or random samples), the ambient dimension  $d$  in the constraint  $\alpha > d$  for the control of the  $\alpha$ -total persistence in (7) has been replaced by the smaller intrinsic dimension  $m$  of the problem: the manifold hypothesis has been successfully exploited.

To summarize, our work sheds light on the behaviour of PDs, provides new guarantees for commonly used ML methods (see e.g. Corollary 4.18), and suggests new heuristics (see Section 5). We also perform various experiments to illustrate the validity of our results and their relevance to the classic TDA pipeline.

**Related work.** This work is part of a long ongoing effort to understand simplicial complexes and PDs in a random context [BA14, BKS17, BW17, BK18, KM13, DC19, Owa22, BH22, HL23]. Closest to our work, Hiraoka, Shirai and Trinh gave limit laws for Čech PDs for random points in the cube  $[0, 1]^d$  [HST18], while Goel, Trinh and Tsunoda gave similar asymptotics in the case of samples on manifolds [GTT19]. Limit laws for the total persistence have been obtained by Divol and Polonik in the case of random samples in the cube [DP19]. Among other contributions, this work generalizes this result to submanifolds: unlike the cube, a manifold has a nontrivial topology; a fact which considerably complicates the situation, for we have to take into account the presence of large topological features in the Čech PDs in order to control the total persistence.

**Notation.** To simplify notation, we write  $K = K(M)$  (or  $K = K(M, a, \dots)$ ) to implicitly state that some constant  $K$  depends only on  $M$  (or only on  $M, a$ , etc.).

## 2 Čech persistence diagrams for subsets of submanifolds

Although this work is only concerned with PDs with respect to the Čech filtration, it is more natural to define PDs for the sublevel sets of proper continuous functions that are bounded below. We refer to [CDSGO16] for a thorough introduction to persistent homology and PDs in an even more general context.

Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  be a proper continuous function that is bounded below—e.g., the distance function to a compact set. For  $t \in \mathbb{R}$ , let  $X_t = f^{-1}(-\infty, t]$  be the sublevel set of  $f$  at level  $t$ . The collection  $(X_t)_{t \in \mathbb{R}}$  is called a filtration. Let  $i \geq 0$  be an integer. We let  $H_i(X_t)$  be the homology group of degree  $i$  with coefficients in any fixed field  $\mathcal{F}$  (e.g.  $\mathcal{F} = \mathbb{Z}/2\mathbb{Z}$  is a popular choice) of  $X_t$ .

The persistence diagram  $\text{dgm}_i(f)$  of degree  $i$  of  $f$  tracks the evolution of the homology groups of degree  $(H_i(X_t))_{t \in \mathbb{R}}$ . Define the extended half-plane  $\Omega_\infty = \{u = (u_1, u_2) \in (\mathbb{R} \cup \{-\infty, +\infty\})^2 : -\infty \leq u_1 < u_2 \leq +\infty\}$  and  $\Omega = \{u = (u_1, u_2) \in \mathbb{R}^2 : u_1 < u_2\}$ . Then  $\text{dgm}_i(f)$  presents itself as a multiset (a set whose elements can have multiplicity greater than 1) in  $\Omega_\infty$ . We refer to [CDSGO16] for its precise definition, but it can be intuitively understood as follows: for  $u = (u_1, u_2) \in \Omega_\infty$ , the number of points in the persistence diagram  $\text{dgm}_i(f)$  found in the upper-left quadrant  $Q_u = \{v \in (\mathbb{R} \cup \{-\infty, +\infty\})^2 : v_1 \leq u_1 < u_2 \leq v_2\}$  is informally equal to the number of  $i$ th dimensional features present in the sublevel set  $f^{-1}(-\infty, u_1]$  (captured through its  $i$ th homology group  $H_i(f^{-1}(-\infty, u_1])$ ) that are still present in the sublevel set  $f^{-1}(-\infty, u_2]$ . Note that this number of points is always finite [CDSGO16, Corollary 3.34]. Though persistence diagrams can have points with infinite coordinates, these will be of little interest in the cases considered in this article.<sup>3</sup> To simplify our notation and definitions, we let from now on  $\text{dgm}_i(f)$  denote the finite part (i.e. the points whose coordinates are finite) of the diagram of degree  $i$  of  $f$ , and we assume that every diagram considered henceforth has no point with infinite coordinates. In particular, they are all multisets of the half-plane  $\Omega := \{u = (u_1, u_2) \in \mathbb{R}^2 : u_1 < u_2\}$ , which leads to the following definition: the space  $\mathcal{D}$  of persistence diagrams is the set of all multisets  $a$  in  $\Omega$  that contain a finite number of points<sup>4</sup> in every quadrant  $Q_u$ ,  $u \in \Omega$ .

In this paper, we are interested in the PDs of the distance function  $d_A$  to various compact sets  $A$ , called the Čech persistence diagram<sup>5</sup> of  $A$  and denoted by  $\text{dgm}_i(A)$ .

Changes in the topology of the sublevels of  $d_A$  can be partially described in terms of zeros of its generalized gradient, which is defined at  $y \in \mathbb{R}^d \setminus A$  as

$$\nabla d_A(y) = \frac{y - c(\sigma_A(y))}{d_A(y)}, \quad (8)$$

where  $\sigma_A(y) = \{x \in A : \|x - y\| = d_A(y)\}$  is the set of projections of  $y$  on  $A$  and  $c(\tau)$  represents the center of the smallest enclosing ball of a set  $\tau$ . When  $y \in \mathbb{R}^d \setminus A$  satisfies  $\nabla d_A(y) = 0$ ,  $y$  is called a *differential critical point* of  $d_A$ . We let  $\text{Crit}(A)$  denote the set of differential critical points of  $d_A$ . An adapted version of the classical Isotopy Lemma for Morse functions is true for distance functions to compact sets as well, as shown in [Gro93]:

<sup>3</sup>The persistence diagram of degree  $i$  of the distance function to a set has no point with infinite coordinates if  $i > 0$ , and a single such point if  $i = 0$  whose coordinates are  $(0, +\infty)$ .

<sup>4</sup>This corresponds to the set of diagrams of  $q$ -tame persistence modules as defined in [CDSGO16].

<sup>5</sup>The distance function  $d_A$  is proper due to the compactness of  $A$ , hence its persistence module is  $q$ -tame and the associated persistence diagram is well-defined—see [CDSGO16].

**Lemma 2.1** (Isotopy Lemma for Distance Functions). *If  $0 < a < b$  are such that  $d_{\mathbf{A}}^{-1}[a, b]$  contains no differential critical point of  $d_{\mathbf{A}}$ , then  $d_{\mathbf{A}}^{-1}(-\infty, a]$  is a deformation retract of  $d_{\mathbf{A}}^{-1}(-\infty, b]$ . Consequently, any  $(u_1, u_2) \in \text{dgm}_i(\mathbf{A})$  is such that  $u_1, u_2 \notin [a, b]$ .*

A fundamental result in TDA, the Bottleneck Stability Theorem (2), states that Čech PDs are stable with respect to Hausdorff perturbations. Consider the particular setting where one has access to a set  $\mathbf{A}$ , obtained as an approximation of an unknown shape of interest  $\mathbf{S}$  through some sampling procedure, with  $\mathbf{A} \subset \mathbf{S}$  and  $\sup_{x \in \mathbf{S}} d_{\mathbf{A}}(x) \leq \varepsilon$ . The Bottleneck Stability Theorem ensures that  $\text{OT}_{\infty}(\text{dgm}_i(\mathbf{A}), \text{dgm}_i(\mathbf{S})) \leq \varepsilon$  for any  $i \geq 0$ , a bound which cannot be improved in general. However, it turns out that a finer understanding of the proximity between  $\text{dgm}_i(\mathbf{A})$  and  $\text{dgm}_i(\mathbf{S})$  can be obtained if more regularity is assumed on the shape of interest  $\mathbf{S}$ , namely in the situation where  $\mathbf{S} = \mathbf{M}$  is a compact submanifold with positive reach.

Let us first set some notation. Let  $\mathbf{M}$  be an  $m$ -dimensional compact topological submanifold of  $\mathbb{R}^d$ ; we always assume that the boundary of  $\mathbf{M}$  is empty and that  $m \geq 1$ . The orthogonal projection  $\pi_{\mathbf{M}}$  on  $\mathbf{M}$  is defined for  $x$  close enough to  $\mathbf{M}$ , and we define the *reach*  $\tau(\mathbf{M})$  as the largest  $r > 0$  such that the orthogonal projection  $\pi_{\mathbf{M}}$  is well (i.e. uniquely) defined for all  $x \in \mathbb{R}^d$  at distance strictly less than  $r$  from  $\mathbf{M}$ . The reach is a key notion to quantify the regularity of a manifold, see e.g. [Fed59] and [CCSL06] for more information. In particular, any compact  $C^2$  submanifold has positive reach. For  $x \in \mathbf{M}$ , we let  $T_x\mathbf{M}$  be the tangent space of  $\mathbf{M}$  at  $x$ , which is identified with a linear subspace of  $\mathbb{R}^d$ . We denote by  $\pi_x : \mathbb{R}^d \rightarrow T_x\mathbf{M}$  the orthogonal projection on this subspace and let  $\pi_x^{\perp} = \text{id} - \pi_x$  be the orthogonal projection on the normal space at  $x$ . A key property, that we will repeatedly use, is that the reach of  $\mathbf{M}$  controls the deviation of the manifold  $\mathbf{M}$  from its tangent space. Namely, [Fed59, Theorem 4.18] states that for all  $y \in \mathbf{M}$ ,

$$\|\pi_x^{\perp}(x - y)\| \leq \frac{\|x - y\|^2}{2\tau(\mathbf{M})}. \quad (9)$$

We also define the *weak feature size* of  $\mathbf{M}$ , denoted by  $\text{wfs}(\mathbf{M})$ , as the minimal distance between a critical point of  $d_{\mathbf{M}}$  and  $\mathbf{M}$ . As by definition, the projection is not unique at a critical point, we must have  $\text{wfs}(\mathbf{M}) \geq \tau(\mathbf{M})$ .

Let  $\mathbf{A} \subset \mathbf{M}$  be such that  $d_H(\mathbf{A}, \mathbf{M}) \leq \varepsilon$  and let  $z \in \mathbb{R}^d$ . By definition of the Hausdorff distance, it holds that  $|d_{\mathbf{A}}(z) - d_{\mathbf{M}}(z)| \leq \varepsilon$ . However, this naive bound can be quadratically improved as long as  $z$  stays far away from  $\mathbf{M}$ .

**Lemma 2.2.** *Let  $\mathbf{M} \subset \mathbb{R}^d$  be a compact submanifold with positive reach and let  $\mathbf{A} \subset \mathbf{M}$  be a compact set with  $d_H(\mathbf{A}, \mathbf{M}) \leq \varepsilon$  for some  $\varepsilon > 0$ . Let  $z \in \mathbb{R}^d \setminus \mathbf{M}$ . Then,  $|d_{\mathbf{M}}(z) - d_{\mathbf{A}}(z)| \leq \frac{\varepsilon^2}{2d_{\mathbf{M}}(z)} \left(1 + \frac{d_{\mathbf{M}}(z)}{\tau(\mathbf{M})}\right)$ .*

*Proof.* As  $\mathbf{A} \subset \mathbf{M}$ , we have  $d_{\mathbf{M}}(z) \leq d_{\mathbf{A}}(z)$ . Let  $x$  be a projection of  $z$  onto  $\mathbf{M}$ , and let  $y \in \mathbf{A}$  be a point at distance less than  $\varepsilon$  from  $x$ . Then, using (9) and the fact that  $z - x$  is orthogonal to  $T_x\mathbf{M}$ , we obtain that

$$\begin{aligned} d_{\mathbf{A}}(z)^2 &\leq \|z - y\|^2 = \|z - x\|^2 + \|x - y\|^2 + 2\langle z - x, x - y \rangle \\ &\leq d_{\mathbf{M}}(z)^2 + \varepsilon^2 + 2\langle z - x, \pi_x^{\perp}(x - y) \rangle \\ &\leq d_{\mathbf{M}}(z)^2 + \varepsilon^2 + \frac{d_{\mathbf{M}}(z)\varepsilon^2}{\tau(\mathbf{M})}. \end{aligned}$$

Hence,

$$d_{\mathbf{A}}(z) - d_{\mathbf{M}}(z) = \frac{d_{\mathbf{A}}(z)^2 - d_{\mathbf{M}}(z)^2}{d_{\mathbf{A}}(z) + d_{\mathbf{M}}(z)} \leq \frac{\varepsilon^2}{2d_{\mathbf{M}}(z)} \left(1 + \frac{d_{\mathbf{M}}(z)}{\tau(\mathbf{M})}\right). \quad \square$$




Figure 2: PDs of  $M$  (red) and of  $A$  (black).

We can build upon this basic remark to obtain a very precise control of the behavior of the Čech PD of the set  $A$ . Namely, we identify three regions in the upper halfplane  $\Omega$  (displayed in Figure 2) which contain all points in the PD  $\text{dgm}_i(A)$  (for some integer  $i \geq 0$ ). In the first region, corresponding to microscopic topological features disappearing at scales smaller than  $\varepsilon + \varepsilon^2/\tau(M)$ , the Bottleneck Stability Theorem cannot be improved. However, there exists an optimal matching (i.e. a matching  $\gamma : \text{dgm}_i(A) \cup \partial\Omega \rightarrow \text{dgm}_i(M) \cup \partial\Omega$  that realizes the bottleneck distance (1)) such that at least one of the coordinates of any point in the other two regions is larger than  $\tau(M) - \varepsilon^2/\tau(M)$ , and the proximity between a large coordinate of a point  $u \in \text{dgm}_i(A)$  and the coordinate of the matched point  $\gamma(u) \in \text{dgm}_i(M) \cup \partial\Omega$  is of order  $O(\varepsilon^2)$ . This yields a quadratic improvement upon the Bottleneck Stability Theorem.

**Theorem 2.3** (Improved Bottleneck Stability Theorem). *Let  $M \subset \mathbb{R}^d$  be a compact submanifold with positive reach and let  $A \subset M$  be a compact set such that  $d_H(A, M) \leq \varepsilon < \tau(M)/4$ . Let  $i \geq 0$  be an integer. Then  $\text{dgm}_i(A)$  is the union of three regions  $\text{dgm}_i^{(1)}(A) := \text{dgm}_i(A) \cap \{u_1, u_2 \leq \varepsilon + \frac{\varepsilon^2}{\tau(M)}\}$ ,  $\text{dgm}_i^{(2)}(A) := \text{dgm}_i(A) \cap \{u_1 \leq \varepsilon, u_2 \geq \tau(M) - \frac{\varepsilon^2}{\tau(M)}\}$  and  $\text{dgm}_i^{(3)}(A) := \text{dgm}_i(A) \cap \{u_1, u_2 \geq \tau(M) - \frac{\varepsilon^2}{\tau(M)}\}$ .*

Furthermore, let  $C = \frac{2}{\tau(M)} \left(1 + \frac{R(M)}{\tau(M)}\right)$ , where  $R(M)$  is the radius of the smallest ball that contains  $M$ . There exists an optimal matching  $\gamma : \text{dgm}_i(A) \cup \partial\Omega \rightarrow \text{dgm}_i(M) \cup \partial\Omega$  for the bottleneck distance between  $\text{dgm}_i(A)$  and  $\text{dgm}_i(M)$  such that

- *Region (1): If  $u \in \text{dgm}_i^{(1)}(A)$ , then  $\gamma(u) \in \partial\Omega$  and  $\|u - \gamma(u)\|_\infty \leq \varepsilon$ .*
- *Region (2): If  $u \in \text{dgm}_i^{(2)}(A)$ , then  $\gamma(u)$  is of the form  $(0, v_2)$  and  $|u_2 - v_2| \leq C\varepsilon^2$ . The number of such points is finite and depends only on  $M$ .*
- *Region (3): If  $u \in \text{dgm}_i^{(3)}(A)$ , then  $\|u - \gamma(u)\|_\infty \leq C\varepsilon^2$ .*

Note that for any  $i \geq d$ , the  $i$ -th PDs of  $M$  and  $A$  are actually trivial. Furthermore, under the hypotheses of Theorem 2.3, the sub-diagrams  $\text{dgm}_i^{(2)}(A)$  and  $\text{dgm}_i^{(2)}(M)$  are empty if  $i$  is strictly greater than the dimension of  $M$ .

Note also that Regions (2) and (3) can be made smaller as follows: there exists  $\delta : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  such that  $\lim_{\varepsilon \rightarrow 0} \delta(\varepsilon) = 0$  and such that  $\text{dgm}_i^{(2)}(A) \subset \text{dgm}_i(A) \cap \{u_1 \leq \varepsilon, u_2 \geq \text{wfs}(M) - \delta(\varepsilon)\}$



and  $\text{dgm}_i^{(3)}(\mathbf{A}) \subset \text{dgm}_i(\mathbf{A}) \cap \{u_1, u_2 \geq \text{wfs}(\mathbf{M}) - \delta(\varepsilon)\}$  if  $d_H(\mathbf{A}, \mathbf{M}) \leq \varepsilon$ . These inequalities are proved using the lower semicontinuity of the generalized gradient, as well as the Isotopy Lemma for distance functions. As  $\text{wfs}(\mathbf{M}) \geq \tau(\mathbf{M})$ , this can yield tighter constraints than in the statement of Proposition 2.3, but there is sadly no simple description of the function  $\delta$  (e.g. in terms of  $\tau(\mathbf{M})$  or  $\text{wfs}(\mathbf{M})$ ).

*Proof.* Proposition 5 from [AKF+22] states that if  $\varepsilon = d_H(\mathbf{A}, \mathbf{M}) < (\sqrt{2} - 1)\tau(\mathbf{M})$ , then the offset  $A^r$  deformation-retracts onto  $\mathbf{M}$  for any

$$r \in \left[ \frac{1}{2}(\tau(\mathbf{M}) + \varepsilon - \sqrt{\Delta}), \frac{1}{2}(\tau(\mathbf{M}) + \varepsilon + \sqrt{\Delta}) \right]$$

and  $\Delta = \tau(\mathbf{M})^2 - 2\varepsilon\tau(\mathbf{M}) - \varepsilon^2$ . Under the stronger assumption that  $d_H(\mathbf{A}, \mathbf{M}) < \tau(\mathbf{M})/4$ , and using elementary calculus, we find that the offset  $A^r$  deformation-retracts onto  $\mathbf{M}$  for any

$$r \in \left[ \varepsilon + \frac{\varepsilon^2}{\tau(\mathbf{M})}, \tau(\mathbf{M}) - \frac{\varepsilon^2}{\tau(\mathbf{M})} \right].$$

This means in particular that the homology type, hence the homology, of  $A^r$  does not change in that interval; as a result, there can be no birth or death of intervals in the Čech persistence diagrams of  $\mathbf{A}$  between  $\varepsilon + \frac{\varepsilon^2}{\tau(\mathbf{M})}$  and  $\tau(\mathbf{M}) - \frac{\varepsilon^2}{\tau(\mathbf{M})}$ , and all  $(u_1, u_2) \in \text{dgm}_i(\mathbf{A})$  must either be such that  $u_1, u_2 \leq \varepsilon + \frac{\varepsilon^2}{\tau(\mathbf{M})}$ , or  $u_1 \leq \varepsilon + \frac{\varepsilon^2}{\tau(\mathbf{M})}, u_2 \geq \tau(\mathbf{M}) - \frac{\varepsilon^2}{\tau(\mathbf{M})}$ , or  $u_1, u_2 \geq \tau(\mathbf{M}) - \frac{\varepsilon^2}{\tau(\mathbf{M})}$ . This almost proves that the partition of  $\text{dgm}_i(\mathbf{A})$  into  $\text{dgm}_i^{(1)}(\mathbf{A})$ ,  $\text{dgm}_i^{(2)}(\mathbf{A})$  and  $\text{dgm}_i^{(3)}(\mathbf{A})$  as defined in the statement is correct, except that the definition of  $\text{dgm}_i^{(2)}(\mathbf{A})$  requires that  $u_1 \leq \varepsilon$ , whereas we only have obtained that  $u_1 \leq \varepsilon + \frac{\varepsilon^2}{\tau(\mathbf{M})}$ .

Let  $\gamma : \text{dgm}_i(\mathbf{A}) \cup \partial\Omega \rightarrow \text{dgm}_i(\mathbf{M}) \cup \partial\Omega$  be any optimal matching for the bottleneck distance: then the Bottleneck Stability Theorem states that any point  $u = (u_1, u_2) \in \text{dgm}_i(\mathbf{A})$  is such that  $\|u - \gamma(u)\|_\infty \leq d_H(\mathbf{A}, \mathbf{M}) \leq \varepsilon$ . Suppose that  $u$  is such that  $u_1 \leq \varepsilon + \frac{\varepsilon^2}{\tau(\mathbf{M})}$  and  $u_2 \geq \tau(\mathbf{M}) - \frac{\varepsilon^2}{\tau(\mathbf{M})}$ . Its distance in the infinity norm to the diagonal  $\partial\Omega$  is equal to  $(u_2 - u_1)/2 \geq (\tau(\mathbf{M}) - \varepsilon - 2\varepsilon^2/\tau(\mathbf{M}))/2 \geq \frac{5}{16}\tau(\mathbf{M}) > \varepsilon$ , where we use the fact that  $\varepsilon < \tau(\mathbf{M})/4$ . Hence  $\gamma(u)$  must belong to  $\text{dgm}_i(\mathbf{M})$ . Let  $(v_1, v_2)$  denote  $\gamma(u)$ . The Isotopy Lemma for distance functions shows that  $\text{dgm}_i(\mathbf{M})$  only contains two types of points: points of the shape  $(0, w_2)$ , which correspond to the homology of  $\mathbf{M}$  itself, and points of the shape  $(w_1, w_2)$ , where in both cases  $w_1, w_2$  are critical values of  $d_{\mathbf{M}}$ . In particular, both  $w_1$  and  $w_2$  must be greater than  $\text{wfs}(\mathbf{M})$ . Let  $\text{dgm}_i^{(2)}(\mathbf{M})$  denote the multiset of all points of the first type, and  $\text{dgm}_i^{(3)}(\mathbf{M})$  denote the multiset of all points of the second type. If  $v_1$  was non-zero, it would have to be greater than  $\text{wfs}(\mathbf{M}) \geq \tau(\mathbf{M})$ , and we would have  $\|u - \gamma(u)\|_\infty \geq |u_1 - v_1| \geq \tau(\mathbf{M}) - \varepsilon - \frac{\varepsilon^2}{\tau(\mathbf{M})} \geq \tau(\mathbf{M})/2 > \varepsilon$ , which would be a contradiction (we once again use that  $\varepsilon < \tau(\mathbf{M})/4$ ). Hence  $v_1$  must be 0 (i.e.  $\gamma(u) \in \text{dgm}_i^{(2)}(\mathbf{M})$ ), and  $u_1 = |u_1 - v_1| \leq \|u - \gamma(u)\|_\infty \leq \varepsilon$ . This proves the correctness of the partition into regions from the statement.

Consider now  $u = (u_1, u_2) \in \text{dgm}_i^{(1)}(\mathbf{A})$ . All  $v = (v_1, v_2) \in \text{dgm}_i(\mathbf{M})$  are such that

$$\|u - v\|_\infty \geq v_2 - u_2 \geq \text{wfs}(\mathbf{M}) - \varepsilon - \frac{\varepsilon^2}{\tau(\mathbf{M})} \geq \tau(\mathbf{M}) \left( 1 - \frac{1}{4} - \frac{1}{16} \right) > \varepsilon \geq \|u - \gamma(u)\|_\infty,$$

hence  $\gamma(u)$  must belong to  $\partial\Omega$ . This completes the proof of the first bullet point of the statement.

We have already shown that if  $\gamma$  is an optimal matching and  $u \in \text{dgm}_i^{(2)}(\mathbf{A})$ , then  $\gamma(u) \in \text{dgm}_i(\mathbf{M})$  is of the form  $(0, v_2)$ . As  $\gamma$  maps at most a single point of  $\text{dgm}_i^{(2)}(\mathbf{A})$  to each point of  $\text{dgm}_i(\mathbf{M})$ , the number of such points is upper bounded by the number of points of the form  $(0, v_2)$  with  $v_2 \geq \text{wfs}(\mathbf{M})$  in  $\text{dgm}_i(\mathbf{M})$ . Though  $\text{dgm}_i(\mathbf{M})$  need not be finite, applying Corollary 3.34 from [CDSGO16] to  $d_{\mathbf{M}}$  shows that  $\text{dgm}_i(\mathbf{M})$  is q-tame, and in particular that it contains only a finite number  $N$  of points  $(v_1, v_2)$  with  $v_1 \leq \text{wfs}(\mathbf{M})/4$  and  $v_2 \geq \text{wfs}(\mathbf{M})/2$ . Hence the cardinality of  $\text{dgm}_i^{(2)}(\mathbf{A})$  is bounded by  $N$ .

It only remains to show that  $\gamma$  can be chosen such that if  $u = (u_1, u_2) \in \text{dgm}_i^{(2)}(\mathbf{A})$ , respectively  $u' \in \text{dgm}_i^{(3)}(\mathbf{A})$ , then  $|u_2 - \gamma(u)_2| \leq C\varepsilon^2$ , respectively  $\|u' - \gamma(u')\|_\infty \leq C\varepsilon^2$ . To that end, remember first that as shown above, our starting optimal matching  $\gamma$  must be such that points in  $\text{dgm}_i^{(2)}(\mathbf{A})$  must be matched to points in  $\text{dgm}_i^{(2)}(\mathbf{M})$ . Conversely and for the same reasons, points in  $\text{dgm}_i^{(2)}(\mathbf{M})$  must be matched to points in  $\text{dgm}_i^{(2)}(\mathbf{A})$ . Similarly, points  $u \in \text{dgm}_i^{(3)}(\mathbf{A})$  can only be matched to points in  $\text{dgm}_i^{(3)}(\mathbf{M})$  or to the diagonal  $\partial\Omega$ ; otherwise,  $\|u - \gamma(u)\|_\infty \geq |u_1 - \gamma(u)_1| = u_1$  would be too large. Likewise, points in  $\text{dgm}_i^{(3)}(\mathbf{M})$  can only be matched to points in  $\text{dgm}_i^{(3)}(\mathbf{A})$  or to the diagonal. Hence  $\gamma$  defines disjoint submatchings

$$\gamma^{(2)} : \text{dgm}_i^{(2)}(\mathbf{A}) \rightarrow \text{dgm}_i^{(2)}(\mathbf{M})$$

and

$$\gamma^{(3)} : \text{dgm}_i^{(3)}(\mathbf{A}) \cup \partial\Omega \rightarrow \text{dgm}_i^{(3)}(\mathbf{M}) \cup \partial\Omega.$$

Now let  $R(\mathbf{M})$  be the radius of the smallest ball that contains  $\mathbf{M}$ , and consider the functions

$$a : \mathbb{R}^d \rightarrow \mathbb{R}, x \mapsto \min(\max(d_{\mathbf{A}}(x), \tau(\mathbf{M})/2), R(\mathbf{M}))$$

and

$$m : \mathbb{R}^d \rightarrow \mathbb{R}, x \mapsto \min(\max(d_{\mathbf{M}}(x), \tau(\mathbf{M})/2), R(\mathbf{M})).$$

Let us compare the persistence diagrams  $\text{dgm}_i(a)$  and  $\text{dgm}_i(m)$  of the sublevel sets filtration of  $a$  and  $m$  and the Čech persistence diagrams  $\text{dgm}_i(\mathbf{A})$  and  $\text{dgm}_i(\mathbf{M})$  respectively (which are by definition the persistence diagrams of the sublevel sets filtration of  $d_{\mathbf{A}}$  and  $d_{\mathbf{M}}$ ).

Note first that  $d_{\mathbf{A}}$  and  $d_{\mathbf{M}}$  can have no critical values strictly greater than  $R(\mathbf{M})$ , as a critical point must belong to the convex hull of its projections. Note also that for any  $t \in [\tau(\mathbf{M})/2, R(\mathbf{M})]$ , the sublevel set  $a^{-1}(-\infty, t]$  is exactly equal to  $d_{\mathbf{A}}^{-1}(-\infty, t] = \mathbf{A}^t$ . Consequently,  $\text{dgm}_i(a)$  contains exactly two disjoint types of points. The first type are points of the form  $(\tau(\mathbf{M})/2, u_2)$ , which are in bijection with the points  $(u_1, u_2) \in \text{dgm}_i(\mathbf{A})$  with  $u_1 \leq \tau(\mathbf{M})/2$  (the bijection maps  $(u_1, u_2) \mapsto (\tau(\mathbf{M})/2, u_2)$ ); those are exactly the points in  $\text{dgm}_i^{(2)}(\mathbf{A})$ . The second type are points of the form  $(u_1, u_2)$  with  $u_1 \geq \tau(\mathbf{M})/2$ , which are in trivial bijection (the bijection is the identity) with the points in  $\text{dgm}_i(\mathbf{A})$  that satisfy the same condition; those are exactly the points of  $\text{dgm}_i^{(3)}(\mathbf{A})$ . The points of  $\text{dgm}_i^{(1)}(\mathbf{A})$  cannot be “seen” in  $\text{dgm}_i(a)$ . We will call  $\text{dgm}_i^{(2)}(a)$  the subdiagram comprised of the points of the first type, and  $\text{dgm}_i^{(3)}(a)$  the subdiagram of  $\text{dgm}_i(a)$  comprised of the points of the second type.

Similarly,  $\text{dgm}_i(m)$  contains two types of points: the first type are points of the form  $(\tau(\mathbf{M})/2, v_2)$ , which are in bijection with the points  $(0, v_2) \in \text{dgm}_i(\mathbf{M})$  (the bijection maps  $(0, v_2) \mapsto (\tau(\mathbf{M})/2, v_2)$ ); those are exactly the points in  $\text{dgm}_i^{(2)}(\mathbf{M})$ . The second type are points of the form  $(v_1, v_2)$  with  $v_1 \geq \tau(\mathbf{M})/2$ , which are in trivial bijection (the bijection is the identity) with the points in  $\text{dgm}_i(\mathbf{M})$  that satisfy the same condition; those are exactly the points

of  $\text{dgm}_i^{(3)}(\mathbf{M})$ . We will call  $\text{dgm}_i^{(2)}(m)$  the subdiagram of  $\text{dgm}_i(m)$  comprised of the points of the first type, and  $\text{dgm}_i^{(3)}(m)$  the subdiagram comprised of the points of the second type.

Recall that Lemma 2.2 states that  $|d_{\mathbf{M}}(z) - d_{\mathbf{A}}(z)| \leq \frac{\varepsilon^2}{2d_{\mathbf{M}}(z)} \left(1 + \frac{d_{\mathbf{M}}(z)}{\tau(\mathbf{M})}\right)$  for any  $z \in \mathbb{R}^d \setminus \mathbf{M}$ . If  $z \in \mathbb{R}^d$  is such that  $d_{\mathbf{A}}(z) \leq \tau(\mathbf{M})/2$ , then  $a(z) = m(z) = \tau(\mathbf{M})/2$ ; if it is such that  $d_{\mathbf{M}}(z) \geq R(\mathbf{M})$ , then  $a(z) = m(z) = R(\mathbf{M})$ . Otherwise,  $d_{\mathbf{M}}(z) \geq d_{\mathbf{A}}(z) - d_H(\mathbf{A}, \mathbf{M}) \geq \tau(\mathbf{M})/4$  and  $d_{\mathbf{M}}(z) \leq R(\mathbf{M})$ , hence

$$|d_{\mathbf{M}}(z) - d_{\mathbf{A}}(z)| \leq \frac{\varepsilon^2}{2d_{\mathbf{M}}(z)} \left(1 + \frac{d_{\mathbf{M}}(z)}{\tau(\mathbf{M})}\right) \leq \frac{2\varepsilon^2}{\tau(\mathbf{M})} \left(1 + \frac{R(\mathbf{M})}{\tau(\mathbf{M})}\right) = C\varepsilon^2,$$

where  $C$  is as defined in the proposition. This means that

$$\|m - a\|_{\infty} \leq C\varepsilon^2.$$

Due to the Bottleneck Stability Theorem, the diagrams  $\text{dgm}_i(a)$  and  $\text{dgm}_i(m)$  must be at bottleneck distance less than  $C\varepsilon^2$ .

Furthermore, let  $\delta$  denote  $\max_{u \in \text{dgm}_i(\mathbf{A}) \cup \partial\Omega} \|u - \gamma(u)\|_{\infty}$ . The matching  $\gamma$  (and in particular the submatchings  $\gamma^{(2)}$  and  $\gamma^{(3)}$ ) also induces (through the correspondence detailed above between the points of  $\text{dgm}_i(a)$  and a subset of the points of  $\text{dgm}_i(\mathbf{A})$ ) a matching  $\gamma'$  between  $\text{dgm}_i(a)$  and  $\text{dgm}_i(m)$  such that  $\max_{u \in \text{dgm}_i(a) \cup \partial\Omega} \|u - \gamma'(u)\|_{\infty} \leq \delta$ . Hence the bottleneck distance between  $\text{dgm}_i(a)$  and  $\text{dgm}_i(m)$  is at most  $\min(\delta, C\varepsilon^2)$ .

Let  $\beta : \text{dgm}_i(a) \cup \partial\Omega \rightarrow \text{dgm}_i(m) \cup \partial\Omega$  be an optimal matching for the bottleneck distance. For similar reasons as for  $\gamma$ , the matching  $\beta$  can also be decomposed into two disjoint submatchings

$$\beta^{(2)} : \text{dgm}_i^{(2)}(a) \rightarrow \text{dgm}_i^{(2)}(m)$$

and

$$\beta^{(3)} : \text{dgm}_i^{(3)}(a) \cup \partial\Omega \rightarrow \text{dgm}_i^{(3)}(m) \cup \partial\Omega.$$

We can use the two matchings  $\beta^{(2)}$  and  $\beta^{(3)}$  to define a new optimal matching  $\tilde{\gamma} : \text{dgm}_i(\mathbf{A}) \cup \partial\Omega \rightarrow \text{dgm}_i(\mathbf{M}) \cup \partial\Omega$  as follows:

- The points in  $\text{dgm}_i^{(1)}(\mathbf{A})$  are matched by  $\tilde{\gamma}$  to  $\partial\Omega$  as with  $\gamma$ .
- Given  $u = (u_1, u_2) \in \text{dgm}_i^{(2)}(\mathbf{A})$ , let  $u' = (\tau(\mathbf{M})/2, u_2)$  be the point of  $\text{dgm}_i^{(2)}(a)$  with which  $u$  is in bijection. We let  $\tilde{\gamma}$  match  $u$  with the point  $v = (0, v_2) \in \text{dgm}_i^{(2)}(\mathbf{M})$  which is in bijection with  $\beta^{(2)}(u') = (\tau(\mathbf{M})/2, v_2) \in \text{dgm}_i^{(2)}(m)$ . Then  $|u_2 - v_2| \leq \min(\delta, C\varepsilon^2)$  due to the optimality of  $\beta$ , and this defines a bijective matching  $\tilde{\gamma}^{(2)} : \text{dgm}_i^{(2)}(\mathbf{A}) \rightarrow \text{dgm}_i^{(2)}(\mathbf{M})$ . Note also that  $\max_{u \in \text{dgm}_i^{(2)}(\mathbf{A})} |u_1 - \tilde{\gamma}^{(2)}(u)_1| = \max_{u \in \text{dgm}_i^{(2)}(\mathbf{A})} u_1 = \max_{u \in \text{dgm}_i^{(2)}(\mathbf{A})} |u_1 - \gamma^{(2)}(u)_1|$ , hence  $\max_{u \in \text{dgm}_i^{(2)}(\mathbf{A})} \|u - \tilde{\gamma}^{(2)}(u)\|_{\infty} \leq \delta$ .
- We have seen that  $\text{dgm}_i^{(3)}(a) = \text{dgm}_i^{(3)}(\mathbf{A})$  and  $\text{dgm}_i^{(3)}(m) = \text{dgm}_i^{(3)}(\mathbf{M})$ . We simply define the restriction and corestriction  $\tilde{\gamma}^{(3)} : \text{dgm}_i^{(3)}(\mathbf{A}) \cup \partial\Omega \rightarrow \text{dgm}_i^{(3)}(\mathbf{M}) \cup \partial\Omega$  of  $\tilde{\gamma}$  as being equal to  $\beta^{(3)} : \text{dgm}_i^{(3)}(a) \cup \partial\Omega \rightarrow \text{dgm}_i^{(3)}(m) \cup \partial\Omega$ . The optimality of  $\beta$  implies that  $\max_{u \in \text{dgm}_i^{(3)}(\mathbf{A})} \|u - \tilde{\gamma}^{(3)}(u)\|_{\infty} \leq \min(\delta, C\varepsilon^2)$ .

Thus the global matching  $\tilde{\gamma} : \text{dgm}_i(\mathbf{A}) \cup \partial\Omega \rightarrow \text{dgm}_i(\mathbf{M}) \cup \partial\Omega$  is well-defined, is optimal for the bottleneck distance, and satisfies the conditions stated in the proposition. This completes the proof.  $\square$

### 3 Čech persistence diagrams for subsets of generic submanifolds

The improved Bottleneck Stability Theorem (Theorem 2.3) yields information relative to the location of points in the Čech PD of  $A$ , but not about their numbers. However, both the  $\alpha$ -total persistence of  $\text{dgm}_i(A)$  and the distance  $\text{OT}_p(\text{dgm}_i(A), \text{dgm}_i(M))$  for  $p < \infty$  crucially depend on the number of points in  $\text{dgm}_i(A)$  having small persistence. Unfortunately, no control on, say, the total persistence, can exist without additional assumptions. Indeed, in general, even the  $\alpha$ -total persistence of the Čech PD of the submanifold  $M$  can be infinite.

**Example 3.1.** Let  $f : x \in \mathbb{R} \mapsto 1 + x^4 \sin(1/x)^2$ . Consider the  $C^2$  curve  $M$  in  $\mathbb{R}^2$  defined as the union of the graphs of the functions  $f$  and  $-f$  on  $[-2, 2]$ . Being  $C^2$ , the curve has a positive reach [Fed59]. For  $i = 1$ , the Čech PD of  $M$  contains a sequence of points  $(1, \ell_n)$  for  $n \geq 1$ , where  $\ell_n = 1 + \Theta(n^{-4})$ . In particular, as  $\sum_{n \geq 1} (\ell_n - 1)^\alpha = +\infty$  for  $\alpha < 1/4$ , the  $\alpha$ -total persistence of  $\text{dgm}_1(M)$  is infinite for such a value of  $\alpha$ . By considering the product  $M^m \subset \mathbb{R}^{2m}$ , one can also build an  $m$ -dimensional  $C^2$  compact submanifold without boundary such that  $\text{dgm}_1(M)$  has an infinite  $\alpha$ -total persistence for  $\alpha < m/4$ .

The existence of such counterexamples is explained by the fact that the distance function  $d_M$  to a set  $M$  is not well-behaved in general, even when the set  $M$  is smooth. In contrast to this bleak general case, Song, Yim & Monod (in the case of surfaces in  $\mathbb{R}^3$ ) and the authors (in the general case) studied the distance function  $d_M$  to  $M$  when  $M$  is a *generic* submanifold [ACSD23, SYM23]. Their findings indicate that, although counterexamples such as the one presented in Example 3.1 exist, they are extremely uncommon in a sense which can be made precise. To do so, we first need to introduce topological Morse functions.

The theory of persistent homology was historically developed for Morse functions  $f : \mathbb{R}^d \rightarrow \mathbb{R}$ . A Morse function  $f$  is a  $C^2$  function whose critical points  $x$  (points for which  $d_x f = 0$ ) are non-degenerate (meaning that the Hessian of  $f$  at  $x$  is non-degenerate). The index of the critical point  $x$  is equal to the number of negative eigenvalues of the corresponding Hessian. The changes of topology of the sublevel sets of such a function are perfectly understood. First, the isotopy lemma states that two sublevel sets  $f^{-1}(-\infty, u_1]$  and  $f^{-1}(-\infty, u_2]$  are isotopic if no critical values are found in the interval  $[u_1, u_2]$  and if  $f^{-1}[u_1, u_2]$  is compact. Second, if  $f^{-1}[u_1, u_2]$  is compact and contains the critical points  $x_1, \dots, x_K$ , then  $f^{-1}(-\infty, u_2]$  has the homotopy type of  $f^{-1}(-\infty, u_1]$  with cells  $e_k$  of dimension equal to the index of  $x_k$  attached along their boundaries (see e.g. [Mil63] for a much more in-depth treatment).

In such a situation, the PD  $\text{dgm}_i(f)$  has a clear interpretation: the coordinates  $(u_1, u_2)$  of a point  $u \in \text{dgm}_i(f)$  correspond to the critical value of a critical point of index  $i$  and  $i + 1$  respectively. Informally, the corresponding  $i$ -dimensional topological feature appears with the attachment of a  $i$ -dimensional cell at value  $u_1$ , and is “killed” by the attachment of a  $(i + 1)$ -dimensional cell at value  $u_2$ .

The notion of Morse function extends to continuous functions with the following definition.

**Definition 3.2** (Topological Morse functions [Mor59]). Let  $U \subset \mathbb{R}^d$  be an open set and let  $f : U \rightarrow \mathbb{R}$  be a continuous function.

- A point  $z \in U$  is said to be a topological regular point of  $f$  if there is a homeomorphism  $\phi : V_1 \rightarrow V_2$  between open neighborhoods  $V_1$  of 0 in  $\mathbb{R}^d$  and  $V_2$  of  $U$  in  $\mathbb{R}^d$  with  $\phi(0) = z$  and such that for all  $x = (x_1, \dots, x_d) \in V_1$ ,

$$f \circ \phi(x) = f(z) + x_d. \tag{10}$$




Figure 3: A generic torus.

- A point  $z \in U$  is said to be a topological critical point of  $f$  if it is not a topological regular point of  $f$ .
- A point  $z \in U$  is said to be a non-degenerate topological critical point of  $f$  of index  $i$  if there exist an integer  $0 \leq i \leq d$  and a homeomorphism  $\phi : V_1 \rightarrow U_2$  between open neighborhoods  $V_1$  of 0 in  $\mathbb{R}^d$  and  $V_2$  of  $U$  in  $\mathbb{R}^d$  with  $\phi(0) = z$  such that for all  $x = (x_1, \dots, x_d) \in V_1$ ,

$$f \circ \phi(x) = f(z) - \sum_{j=1}^i x_j^2 + \sum_{j=i+1}^d x_j^2. \quad (11)$$

- The function  $f$  is said to be a topological Morse function if all its topological critical points are non-degenerate.

For topological Morse functions, both the isotopy lemma and the handle attachment lemma stay valid:

**Lemma 3.3** (Isotopy Lemma). *Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  be a proper topological Morse function. Let  $a < b$  be such that  $f^{-1}[a, b]$  contains no topological critical point. Then  $f^{-1}(-\infty, a]$  is a deformation retract of  $f^{-1}(-\infty, b]$ .*

**Lemma 3.4** (Handle Attachment Lemma). *Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  be a proper topological Morse function. Let  $c \in \mathbb{R}$  and  $\varepsilon > 0$  be such that  $f^{-1}[c - \varepsilon, c + \varepsilon]$  contains no topological critical point except for  $z_1, \dots, z_k \in f^{-1}(c)$ , with  $z_j$  of index  $i_j$ . Then  $f^{-1}(-\infty, c + \varepsilon]$  is homotopically equivalent to  $f^{-1}(-\infty, c - \varepsilon]$  with cells  $B^{i_1}, \dots, B^{i_k}$  of dimension  $i_1, \dots, i_k$  attached, i.e.*

$$f^{-1}(-\infty, c + \varepsilon] \simeq f^{-1}(-\infty, c - \varepsilon] \cup B^{i_1} \cup \dots \cup B^{i_k}.$$

Their proofs are roughly the same as for smooth Morse functions –see [SYM23, Theorems 4 and 5] for details. As a consequence, the description of PDs for Morse functions also stays valid for topological Morse functions.

It is not true in general that the distance function  $d_A$  to a compact set  $A$  is a topological Morse function, as shown by Example 3.1. Though the Isotopy Lemma for distance functions (Lemma 2.1) is an analogue of the Isotopy Lemma for Morse functions, there is no equivalent to the Handle Attachment Lemma to control the changes occurring at critical values, and as

a result little can be said regarding the topology of the sublevel sets of  $d_M$  without further assumptions. However, the distance function  $d_M$  to a compact  $C^2$  submanifold  $M \subset \mathbb{R}^d$  turns out to be a topological Morse function in a “generic” sense, as was proven by the authors.

**Theorem 3.5** (Genericity Theorem [ACSD23]). *Let  $M$  be a compact  $C^2$  (abstract) manifold. Then the set of  $C^2$  embeddings  $i : M \rightarrow \mathbb{R}^d$  such that*

1. *the distance function  $d_{i(M)} : \mathbb{R}^d \setminus i(M) \rightarrow \mathbb{R}$  is a topological Morse function,*
2. *for every  $z \in \text{Crit}(M)$ , the projections  $\sigma_M(z)$  are the vertices of a non-degenerate simplex of  $\mathbb{R}^d$  and  $z$  belongs to its relative interior,*
3. *the set  $\text{Crit}(M)$  is finite,*
4. *for every  $z \in \text{Crit}(M)$  and every  $x \in \sigma_M(z)$ , the sphere  $S(z, d_M(z))$  is non-osculating  $M$  at  $x$ , in the sense that there exist  $\delta > 0$  and  $\alpha > 0$  such that for all  $y \in M \cap B(x, \delta)$ ,*

$$\|y - z\|^2 \geq \|x - z\|^2 + \alpha \|y - x\|^2, \quad (12)$$

5. *there exist constants  $C > 0$  and  $\mu_0 \in (0, 1)$  such that for every  $\mu \in [0, \mu_0)$ , any point  $x$  such that  $\|\nabla d_M(x)\| \leq \mu$  is at distance at most  $C\mu$  from  $\text{Crit}(M)$ ,*

*is open and dense in the set of  $C^2$  embeddings  $M \rightarrow \mathbb{R}^d$  for the Whitney  $C^2$ -topology.*

Put another way, given an abstract manifold  $M$ , “almost all”  $C^2$  embeddings  $M$  of  $M$  into  $\mathbb{R}^d$  are such that  $d_M$  is a topological Morse function, and satisfies other regularity conditions whose usefulness will become apparent in some of the coming proofs. In what follows, we will simply describe a  $C^2$  submanifold  $M$  that satisfies Conditions 1-5 from Theorem 3.5 as *generic*. When  $M$  is generic, the topological critical points of  $d_M$  coincide with its differential critical points (see [ACSD23, Tmh 1.8]), and the Čech PD  $\text{dgm}_i(M)$  can be related to the critical points of  $d_M$  in the same way as for smooth Morse function. We summarize this fact in the next proposition.

**Proposition 3.6.** *Let  $M$  be a generic compact submanifold of  $\mathbb{R}^d$ .*

- *The set  $\text{Crit}(M)$  is finite, and so is  $\text{dgm}_i(M)$  for any  $i \geq 0$ . In particular, for all  $\alpha > 0$ ,*

$$\text{Pers}_\alpha(\text{dgm}_i(M)) < +\infty.$$

- *If  $z \in \text{Crit}(M)$  is of index  $i$ , the set of its projections  $\sigma_M(z)$  forms a non-degenerate simplex of dimension at most  $i$ .*
- *For any  $i \geq 1$ , the multiset of critical values  $d_M(z)$  of critical points  $z$  of  $M$  of index  $i$  is equal to the multiset*

$$\{u_1 : (u_1, u_2) \in \text{dgm}_i(M), u_1 \neq 0\} \cup \{u_2 : (u_1, u_2) \in \text{dgm}_{i-1}(M)\}. \quad (13)$$

Note that as for the Čech persistence diagram of any compact set,  $\text{dgm}_0(M)$  contains only points of the form  $(0, u_2)$  for some  $u_2 > 0$ . Note also that the proof below shows that the Čech persistence modules of  $M$  are point-wise finite dimensional, and can be decomposed into intervals.

*Proof.* The proof is similar to that used in the case of smooth Morse functions, with the Isotopy Lemma and the Handle Attachment Lemma playing the same role; we briefly summarize it for completeness nonetheless. We do not distinguish between differential and topological critical values and points, as they coincide.

The set  $\text{Crit}(\mathbf{M})$  is finite: this is simply Condition 3. from the Genericity Theorem. The Isotopy Lemma shows that there is no change in homotopy type between  $\mathbf{M}^a$  and  $\mathbf{M}^b$  if  $0 < a < b$  and  $[a, b]$  contains no critical value. Similarly,  $\mathbf{M}$  has the same homotopy type as  $\mathbf{M}^a$  if  $a > 0$  is small enough. Hence changes in homology in the offsets can only occur at 0, when the entire submanifold appears in the filtration, and at critical values of  $d_{\mathbf{M}}$ , and there can be no birth or death of interval between them.

Let us now consider a critical value  $c > 0$  and  $0 < \varepsilon < c$  such that  $d_{\mathbf{M}}^{-1}[c - \varepsilon, c + \varepsilon]$  contains no critical point except for  $z_1, \dots, z_k \in d_{\mathbf{M}}^{-1}(c)$ , where  $z_j$  is of index  $i_j$ . Then the Handle Attachment Lemma states that  $\mathbf{M}^{c+\varepsilon}$  is homotopically equivalent to  $\mathbf{M}^{c-\varepsilon}$  with cells  $B^{i_1}, \dots, B^{i_k}$  of dimension  $i_1, \dots, i_k$  attached, i.e.

$$\mathbf{M}^{c+\varepsilon} \simeq \mathbf{M}^{c-\varepsilon} \cup B^{i_1} \cup \dots \cup B^{i_k}.$$

Let  $i \geq 1$ , and let  $D_{i,b}$  be the dimension of the cokernel of  $H_i(\mathbf{M}^{c-\varepsilon}) \rightarrow H_i(\mathbf{M}^{c+\varepsilon})$  (where the map is induced by the inclusion): it is precisely the number of births of intervals between  $c - \varepsilon$  and  $c + \varepsilon$  (hence precisely at  $c$ ) in the persistence module of degree  $i$  of the filtration. Similarly, the dimension  $D_{i-1,d}$  of the kernel of  $H_{i-1}(\mathbf{M}^{c-\varepsilon}) \rightarrow H_{i-1}(\mathbf{M}^{c+\varepsilon})$  is the number of deaths of intervals at  $c$  in the persistence module of degree  $i - 1$  of the filtration. A straightforward application of the Mayer-Vietoris exact sequence yields that  $D_{i,b} + D_{i-1,d}$  is exactly equal to the number of  $i$ -dimensional cells among  $B^{i_1}, \dots, B^{i_k}$ , meaning that each  $i$ -cell corresponds exactly either to the birth of an interval for the homology of degree  $i$ , or to the death of an interval for the homology of degree  $i - 1$  (in particular, an  $i - 1$ -cell and an  $i$ -cell cannot “cancel each other out”). This proves that for any  $i \geq 1$ , the multiset of critical values  $d_{\mathbf{M}}(z)$  of critical points  $z$  of  $\mathbf{M}$  of index  $i$  is equal to the multiset

$$\{u_1 : (u_1, u_2) \in \text{dgm}_i(\mathbf{M}), u_1 \neq 0\} \cup \{u_2 : (u_1, u_2) \in \text{dgm}_{i-1}(\mathbf{M})\}. \quad (14)$$

This fact proves in turn that  $\text{dgm}_i(\mathbf{M})$  is finite for any  $i \geq 0$ , which immediately implies that  $\text{Pers}_\alpha(\text{dgm}_i(\mathbf{M})) < +\infty$  for all  $\alpha > 0$ .

Finally, the bound on the dimension of a critical simplex associated to a critical point of index  $i$  is a consequence of Remark 7.3 in [ACSD23] –we give some additional details further below.  $\square$

Considering a well-chosen ellipse and a hyperbole gives a simple illustration of the fact that the index of a critical point (2 for the center of the ellipse, 1 for the “center” of the hyperbole) cannot be directly deduced from the dimension of the associated simplex (1 in both cases). Instead, and as explained in Section 7 of [ACSD23], the index of a critical point  $z \in \text{Crit}(\mathbf{M})$  can be further decomposed as follows: the restriction of the distance function  $d_{\mathbf{M}}$  to a small neighborhood of  $z$  in  $\{x \in \mathbb{R}^d \setminus \mathbf{M} : \dim(\sigma_{\mathbf{M}}(x)) = \dim(\sigma_{\mathbf{M}}(z))\}$  is a  $C^2$  Morse function, and  $z$  is a critical point of it. Let  $j$  be its index with respect to this restriction; then the index  $i$  of  $z$  with respect to  $d_{\mathbf{M}}$  satisfies

$$i = j + \dim(\sigma_{\mathbf{M}}(z)).$$

When  $\mathbf{M}$  is a generic submanifold, it becomes a reasonable task to control the number of points in  $\text{dgm}_i(\mathbf{A})$  where  $\mathbf{A} \subset \mathbf{M}$  is an approximation of  $\mathbf{M}$  with  $d_H(\mathbf{A}, \mathbf{M}) \leq \varepsilon$ . The Bottleneck Stability Theorem implies that when  $\varepsilon$  is small enough (compared to the smallest persistence of

a point in  $\text{dgm}_i(\mathbb{M})$ ), every point of  $\text{dgm}_i(\mathbb{M})$  is mapped to a point in  $\text{dgm}_i(\mathbb{A})$  by the optimal bottleneck matching, leaving the points of  $\text{dgm}_i(\mathbb{A})$  at  $\ell_\infty$ -distance to  $\partial\Omega$  less than  $\varepsilon$  unmatched; those will be mapped to the diagonal  $\partial\Omega$ . The Improved Bottleneck Stability Theorem 2.3 (see Figure 2) shows that these points are of two kinds: those in Region (1), corresponding to small topological features in the set  $\mathbb{A}$  (of size of order  $O(\varepsilon)$ ), and those in Region (3), corresponding to large topological features. There are many points in Region (1) (in fact, our proofs show that when  $\mathbb{A} = \mathbb{A}_n$  is a random sample of  $n$  points, the number of points in Region (1) is of order  $O(n)$ ). In contrast, the genericity hypothesis allows us to show that the number of points in Region (3) is small under reasonable sampling assumptions.

We say that a point cloud  $\mathbb{A} \subset \mathbb{M}$  is  $(\delta, \varepsilon)$ -dense in  $\mathbb{M}$  if  $d_H(\mathbb{A}, \mathbb{M}) \leq \varepsilon$  and  $\min_{x \neq y \in \mathbb{A}} \|x - y\| \geq \delta$ . Such point clouds naturally occur, e.g. as products of the farthest point sampling algorithm [AL18].

**Theorem 3.7.** *Let  $\mathbb{M} \subset \mathbb{R}^d$  be a generic compact submanifold and  $\mathbb{A} \subset \mathbb{M}$  be such that  $d_H(\mathbb{A}, \mathbb{M}) \leq \varepsilon$ . Let  $i \geq 0$  be an integer. There exist  $\varepsilon_0 = \varepsilon_0(\mathbb{M}) > 0$  and  $C_0 = C_0(\mathbb{M})$  such that if  $\varepsilon \leq \varepsilon_0$ , then:*

1. *For each  $u = (u_1, u_2) \in \text{dgm}_i^{(3)}(\mathbb{A})$  with  $\gamma(u) \in \partial\Omega$ , one of the finitely many critical values  $c$  of  $d_{\mathbb{M}}$  is such that  $u_1, u_2 \in [c - C_0\varepsilon^2, c + C_0\varepsilon^2]$ .*

*Furthermore, assume that  $\mathbb{A}$  is a  $(\delta, \varepsilon)$ -dense set in  $\mathbb{M}$  for some  $\delta > 0$ . Let  $a \geq \varepsilon/\delta$ . Then, there exist constants  $C_1, C_2, C_3, C_4$  depending on  $\mathbb{M}$  and  $a$  such that:*

2. *The persistence diagram  $\text{dgm}_i^{(3)}(\mathbb{A})$  has at most  $C_1$  points.*
3. *For all  $p \geq 1$ ,  $\alpha \geq 0$ ,*

$$\begin{aligned} \text{OT}_p^p(\text{dgm}_i(\mathbb{A}), \text{dgm}_i(\mathbb{M})) &\leq C_2\varepsilon^{p-m} \\ \text{Pers}_\alpha(\text{dgm}_i(\mathbb{A})) &\leq C_3(C_4^\alpha + \varepsilon^{\alpha-m}). \end{aligned} \tag{15}$$

In particular, as long as the ratio  $\varepsilon/\delta$  is larger than some constant  $a > 0$ , the  $\text{OT}_p$  distance between  $\text{dgm}_i(\mathbb{A})$  and  $\text{dgm}_i(\mathbb{M})$  converges to 0 for all  $p > m$  as  $\varepsilon \rightarrow 0$ , while the  $p$ -total persistence  $\text{Pers}_p(\text{dgm}_i(\mathbb{A}))$  stays bounded.

**Example 3.8.** *Consider two parallel line segments  $\mathbb{M}$  in  $\mathbb{R}^2$ , and a finite set  $\mathbb{A}$  consisting of two parallel grids of step  $2\varepsilon$ : the set  $\mathbb{A}$  is  $(2\varepsilon, \varepsilon)$ -dense in  $\mathbb{M}$ . Then, there are  $O(\varepsilon^{-1})$  points in  $\text{dgm}_1(\mathbb{A})$  with birth coordinates  $u_1$  equal to  $1/2$  and persistence of order  $O(\varepsilon^2)$ ; they all belong to  $\text{dgm}_1^{(3)}(\mathbb{A})$ , whose cardinality is thus not bounded by some  $C_0 = C_0(\mathbb{M})$ . This example can be easily modified to make  $\mathbb{M}$  a compact  $C^2$  1-dimensional manifold. This shows that the first conclusion of Theorem 3.7 cannot hold without a genericity assumption on  $\mathbb{M}$ .*

*Proof.* 1. Though the distance function to  $\mathbb{A}$  need not be topologically Morse, it remains true (due to the Isotopy Lemma for distance functions) that the changes in topology of the sublevel sets of  $d_{\mathbb{A}}$  occur at its critical values. Hence  $u = (u_1, u_2) \in \text{dgm}_i^{(3)}(\mathbb{A})$  is such that  $u_1, u_2$  are critical values of  $d_{\mathbb{A}}$ , with  $u_1, u_2 \geq \tau(\mathbb{M})/2$ . Theorem 1.6 from [ACSD23] states that there exists  $K_1 = K_1(\mathbb{M}) > 0$  such that if  $\varepsilon$  is smaller than some  $\varepsilon_1$ , then each point in  $\text{Crit}(\mathbb{A})$  at distance more than  $\tau(\mathbb{M})/2$  from  $\mathbb{A}$  must be at distance at most  $K_1\varepsilon$  from one of the finitely many points of  $\text{Crit}(\mathbb{M})$ . Let us consider  $z_{\mathbb{A}} \in \text{Crit}(\mathbb{A})$  and  $z_{\mathbb{M}} \in \text{Crit}(\mathbb{M})$  such that  $\|z_{\mathbb{A}} - z_{\mathbb{M}}\| \leq K_1\varepsilon$  and  $d_{\mathbb{A}}(z_{\mathbb{A}}) \geq \tau(\mathbb{M})/2$ ; we are going to show that there exists  $C_2 = C_2(\mathbb{M}) > 0$  such that



$|d_A(z_A) - d_M(z_M)| \leq C_2\varepsilon^2$ . Let us first consider  $x_M \in \sigma_M(z_M)$ ; the proof of Theorem 1.6 from [ACSD23] also shows that there exists  $x_A \in \sigma_A(z_A)$  such that  $\|x_A - x_M\| \leq K_2\varepsilon$  for some constant  $K_2 = K_2(M)$ . As  $z_M - x_M$  is orthogonal to  $T_{x_M}M$ , Inequality (9) states that  $\|\pi_{x_M}^\perp(x_A - x_M)\| \leq \frac{\|x_A - x_M\|^2}{2\tau(M)} \leq \frac{K_2^2\varepsilon^2}{2\tau(M)}$ , hence

$$\begin{aligned} \|z_M - x_A\|^2 &= \|z_M - x_M\|^2 + 2\langle z_M - x_M, x_M - x_A \rangle + \|x_M - x_A\|^2 \\ &\leq d_M(z_M)^2 + \frac{K_2^2\varepsilon^2}{\tau(M)}d_M(z_M) + K_2^2\varepsilon^2 \leq d_M(z_M)^2 + \varepsilon^2K_3, \end{aligned}$$

where  $K_3 = K_2^2\left(\frac{R(M)}{\tau(M)} + 1\right)$ ,  $R(M)$  is the radius of the smallest ball that contains  $M$ , and  $d_M(z_M) \leq R(M)$  because a critical point must belong to the convex hull of its projections. As the same bound applies to each of the projections  $x'_A \in \sigma_A(z_A)$ , we find that the closed ball  $\overline{B}\left(z_M, \sqrt{d_M(z_M)^2 + \varepsilon^2K_3}\right)$  contains  $\sigma_A(z_A)$ . Since  $z_A$  is the center of the smallest ball that contains  $\sigma_A(z_A)$ , whose radius is  $d_A(z_A)$ , we find that

$$d_A(z_A)^2 \leq d_M(z_M)^2 + \varepsilon^2K_3. \quad (16)$$

Similarly, the proof of Theorem 1.6 from [ACSD23] shows that there exists  $y \in \sigma_M(z_A)$  such that  $\|y - x_M\|, \|y - x_A\| \leq O(\varepsilon)$ , and the same reasoning as above yields that  $\|z_A - x_A\|^2 = \|z_A - y\|^2 + O(\varepsilon^2)$  and  $\|z_A - x_M\|^2 = \|z_A - y\|^2 + O(\varepsilon^2)$  (with all big  $O$  constants depending only on  $M$ ), hence that  $\|z_A - x_M\|^2 \leq \|z_A - x_A\|^2 + \varepsilon^2K_4 = d_A(z_A)^2 + \varepsilon^2K_4$  for some  $K_4 = K_4(M)$ . As before, this shows that the closed ball  $\overline{B}\left(z_A, \sqrt{d_A(z_A)^2 + \varepsilon^2K_4}\right)$  contains  $\sigma_M(z_M)$ , hence that

$$d_M(z_M)^2 \leq d_A(z_A)^2 + \varepsilon^2K_4. \quad (17)$$

This, together with (16) and the fact that  $d_A(z_A) \geq \tau(M)/2$ , shows that  $|d_A(z_A) - d_M(z_M)| \leq C_2\varepsilon^2$  for some  $C_2 = C_2(M)$ . We have shown that any critical value of  $d_A$  greater than  $\tau(M)/2$  is at distance less than  $C_2\varepsilon^2$  from a critical value of  $d_M$ , which concludes the proof of the first point (by taking  $\varepsilon_0$  small enough for  $2C_2\varepsilon_0^2$  to be smaller than the minimum distance between two distinct critical values of  $d_M$ ).

2. Let us now prove the bound on the cardinality of  $\text{dgm}_i^{(3)}(A)$ . As  $M$  is generic, Theorem 1.6 from [ACSD23] states that if  $\varepsilon$  is smaller than some  $\varepsilon_0 = \varepsilon_0(M)$ , then each point in  $\text{Crit}(A)$  at distance more than  $\tau(M)/2$  from  $A$  must be at distance at most  $K_5\varepsilon$  from one of the finitely many points of  $\text{Crit}(M)$  for some  $K_5 = K_5(M)$ . Corollary 1.7 from the same article then states that the number of points in  $\text{Crit}(A)$  at distance less than  $K_1\varepsilon$  from a given point of  $\text{Crit}(M)$  is upper bounded by some constant that depends on  $M$  and the ratio  $\varepsilon/\delta$ , and is decreasing in this ratio; hence it is upper bounded by some constant that depends on  $M$  and  $a$ . The proof of this corollary also shows that the maximum number of projections on  $A$  of each of these points of  $\text{Crit}(A)$  is also upper bounded by some constant that depends on  $M$  and  $a$ . Hence there exist constants  $K_6 = K_6(M, a)$  and  $K_7 = K_7(M, a)$  such that if  $\varepsilon \leq \varepsilon_0$ , then there are at most  $K_6$  points in  $\text{Crit}(A)$  at distance more than  $\tau(M)/2$  from  $A$ , and each has at most  $K_7$  projections on  $A$ .

Lemma A.1, applied to the interval  $[\tau(M)/2, \infty)$  and the set  $A$ , then states that the number of points in  $\text{dgm}_i(A)$  such that at least one of their coordinates is greater than  $\tau(M)/2$  is bounded by  $K_6\binom{K_7}{i+2}$ . Hence there are at most  $C_0 := K_62^{K_7} \geq K_6\binom{K_7}{i+2}$  points in  $\text{dgm}_i^{(3)}(A)$  when  $\varepsilon \leq \varepsilon_0$ .

3. Let us bound  $\text{OT}_p^p(\text{dgm}_i(\mathbf{A}), \text{dgm}_i(\mathbf{M}))$  and  $\text{Pers}_\alpha(\text{dgm}_i(\mathbf{A}))$ . As stated in Theorem 2.3, which applies as  $\varepsilon_0 < \tau(\mathbf{M})/4$ , each point  $u = (u_1, u_2) \in \text{dgm}_i^{(1)}(\mathbf{A})$  is such that its coordinates satisfy  $0 \leq u_1, u_2 \leq \varepsilon + \varepsilon^2/\tau(\mathbf{M}) \leq 2\varepsilon$ . In particular, they must correspond to the birth or the death of an interval of the Čech persistence module of  $\mathbf{A}$  that occurs before filtration time  $2\varepsilon$ . The homology of the offsets  $\mathbf{A}^t$  can be computed using the Čech simplicial complex of  $\mathbf{A}$  (see e.g. [EH10]). In particular, each change in the homology of the offsets, hence each birth or death in the Čech persistence module of  $\mathbf{A}$ , is induced by the apparition of some simplex  $\sigma$  at the corresponding filtration value in the Čech complex, and each such apparition causes at most a single death or birth. If a simplex  $\sigma$  appears before filtration time  $2\varepsilon$ , it is by definition contained in a ball of radius  $2\varepsilon$ , hence it is of diameter at most  $4\varepsilon$ . Let us assume from now on that  $\varepsilon_0 \leq \tau(\mathbf{M})/16$ . Consider  $x \in \mathbf{A}$ ; then [AL18, Proposition 8.7] states that the intersection  $\bar{B}(x, 4\varepsilon) \cap \mathbf{A}$  contains at most  $K_8(\varepsilon/\delta)^m \leq K_8 a^m$  points for some constant  $K_8 = K_8(\mathbf{M})$ . Hence  $x$  belongs to at most  $2^{K_8 a^m}$  simplices that appear before  $\varepsilon$ , and there are at most  $\#\mathbf{A} \cdot 2^{K_8 a^m}$  such simplices. As the cardinality  $\#\mathbf{A}$  can be bounded by  $K_9/\delta^m$  for some  $K_9 = K_9(\mathbf{M})$ , we find that  $\#(\text{dgm}_i^{(1)}(\mathbf{A})) \leq K_{10}/\delta^m$  for some  $K_{10} = K_{10}(\mathbf{M}, a)$ .

Furthermore, when  $\mathbf{M}$  is generic, Proposition 3.6 states that its PD  $\text{dgm}_i(\mathbf{M})$  has a finite number of points. Let  $\gamma$  be an optimal matching between  $\text{dgm}_i(\mathbf{A})$  and  $\text{dgm}_i(\mathbf{M})$  for the bottleneck distance that satisfies the conclusions of Theorem 2.3. We find that any point  $u \in \text{dgm}_i^{(1)}(\mathbf{A})$  is matched to a point of  $\partial\Omega$  at distance at most  $\varepsilon$  from  $u$ . Moreover, the number of points in  $\text{dgm}_i^{(2)}(\mathbf{A}) \cup \text{dgm}_i^{(3)}(\mathbf{A})$  is bounded by some constant  $K_{11} = K_{11}(\mathbf{M}, a)$ , and they are all matched to a point of  $\text{dgm}_i(\mathbf{M})$  or  $\partial\Omega$  at distance at most  $\varepsilon$ . In particular, these finitely many points are at distance at most  $K_{12} = K_{12}(\mathbf{M})$  from  $\partial\Omega$ . Furthermore, this matching is surjective, in the sense that  $\gamma$  matches all points of  $\text{dgm}_i(\mathbf{M})$  to a point of  $\text{dgm}_i(\mathbf{A})$ . As a result, for any  $p \geq 1$ , we find that

$$\begin{aligned} \text{OT}_p^p(\text{dgm}_i(\mathbf{A}), \text{dgm}_i(\mathbf{M})) &\leq \sum_{u \in \text{dgm}_i^{(1)}(\mathbf{A})} \|u - \gamma(u)\|_\infty^p + \sum_{u \in \text{dgm}_i^{(2)}(\mathbf{A}) \cup \text{dgm}_i^{(3)}(\mathbf{A})} \|u - \gamma(u)\|_\infty^p \\ &\leq K_{10} \delta^{-m} \varepsilon^p + K_{11} \varepsilon^p \leq K_{10} a^m \varepsilon^{p-m} + K_{11} \varepsilon^p \leq C_1 \varepsilon^{p-m} \end{aligned}$$

for some  $C_1 = C_1(\mathbf{M}, a)$ . Likewise, for any  $\alpha \geq 0$ , we have that

$$\begin{aligned} \text{Pers}_\alpha(\text{dgm}_i(\mathbf{A})) &= \sum_{u \in \text{dgm}_i^{(1)}(\mathbf{A})} \text{pers}(u)^\alpha + \sum_{u \in \text{dgm}_i^{(2)}(\mathbf{A}) \cup \text{dgm}_i^{(3)}(\mathbf{A})} \text{pers}(u)^\alpha \\ &\leq K_{10} \delta^{-m} \varepsilon^\alpha + K_{11} K_{12}^\alpha \leq K_{10} a^m \varepsilon^{\alpha-m} + K_{11} K_{12}^\alpha \leq C_2 (C_3^\alpha + \varepsilon^{\alpha-m}) \end{aligned}$$

for some  $C_2 = C_2(\mathbf{M}, a)$ ,  $C_3 = C_3(\mathbf{M})$ . This completes the proof.  $\square$

## 4 Random samplings of submanifolds

We now turn to the case of random samplings of (non-generic and generic) submanifolds. They tend to adopt configurations that are more regular than what can be expected from e.g. a general  $\varepsilon$ -dense sampling, yet their randomness gives rise to new technical difficulties. Let  $P$  be a probability measure having a density  $f$  (with respect to the volume measure) on a compact submanifold  $\mathbf{M}$  of dimension  $m \geq 1$ . Let  $\mathbf{A} = \mathbf{A}_n = \{X_1, \dots, X_n\}$ , where  $X_1, \dots, X_n$  is an i.i.d. sample from distribution  $P$ . Let  $i \geq 0$  be an integer; we consider the three regions described

in Figure 2 and in the statement of Theorem 2.3, and write again  $\text{dgm}_i^{(1)}(\mathbf{A}_n)$ ,  $\text{dgm}_i^{(2)}(\mathbf{A}_n)$  and  $\text{dgm}_i^{(3)}(\mathbf{A}_n)$  for the three corresponding PDs. This section is devoted to the study of the probabilistic asymptotic behaviour of these three random PDs, which can be decomposed into two almost independent questions:  $\text{dgm}_i^{(1)}(\mathbf{A}_n)$  only depends on small-scale phenomena and can essentially be reduced to the case of a cube, even if  $\mathbf{M}$  is non-generic, while  $\text{dgm}_i^{(2)}(\mathbf{A}_n)$  and  $\text{dgm}_i^{(3)}(\mathbf{A}_n)$  are tightly connected to the macroscopic geometry of the submanifold and can be further controlled using genericity assumptions on  $\mathbf{M}$ .

Describing the limit behavior of the random PD  $\text{dgm}_i^{(1)}(\mathbf{A}_n)$  requires extending the  $\text{OT}_p$  metrics between PDs to more general Radon measures. Indeed, a PD can equivalently be seen as an integer-valued discrete Radon measure on  $\Omega$ , by identifying a multiset  $a$  with the Radon measure  $\sum_{u \in a} \delta_u$ . Let  $\mathcal{M}$  denote the space of Radon measures on  $\Omega$ , that is the space of Borel measures on  $\Omega$  which give finite mass to every compact set  $K \subset \Omega$ .<sup>6</sup> The space of Radon measures is endowed with the *vague topology*, where a sequence  $(\mu_n)_n$  of measures in  $\mathcal{M}$  is said to converge vaguely to  $\mu \in \mathcal{M}$  if  $\int_{\Omega} \phi d\mu_n \rightarrow \int_{\Omega} \phi d\mu$  as  $n \rightarrow \infty$  for all continuous functions  $\phi : \Omega \rightarrow \mathbb{R}$  with compact support.

The  $\alpha$ -total persistence is defined for  $\mu \in \mathcal{M}$  by  $\text{Pers}_{\alpha}(\mu) = \int_{\Omega} \text{pers}(u)^{\alpha} d\mu(u)$ . For  $p \geq 1$ , we let  $\mathcal{M}^p = \{\mu \in \mathcal{M} : \text{Pers}_p(\mu) < +\infty\}$ . The distance  $\text{OT}_p$ , defined between PDs in (3), can be extended to the set  $\mathcal{M}^p$  as follows (see [DL21] for details): let us define  $\bar{\Omega} := \{u = (u_1, u_2) \in \mathbb{R}^2 : u_1 \leq u_2\}$ . We call  $\pi$  an admissible transport plan between  $\nu_1, \nu_2 \in \mathcal{M}$  if it is a Radon measure on  $\bar{\Omega} \times \bar{\Omega}$  such that for all Borel sets  $A, B \subset \Omega$ ,

$$\pi(A \times \bar{\Omega}) = \nu_1(A) \quad \text{and} \quad \pi(\bar{\Omega} \times B) = \nu_2(B). \quad (18)$$

For  $p \in [1, +\infty)$ , we define

$$\text{OT}_p^p(\nu_1, \nu_2) = \inf_{\pi \in \text{Adm}(\nu_1, \nu_2)} \iint \|u - v\|_{\infty}^p d\pi(u, v) \in \mathbb{R} \cup \{+\infty\}, \quad (19)$$

where  $\text{Adm}(\nu_1, \nu_2)$  is the set of all admissible transport plans between  $\nu_1$  and  $\nu_2$ . We also define

$$\text{OT}_{\infty}(\nu_1, \nu_2) = \inf_{\pi \in \text{Adm}(\nu_1, \nu_2)} \sup\{\|u - v\|_{\infty} : (u, v) \in \text{Support}(\pi)\} \quad (20)$$

For all  $p \in [1, \infty]$ , the infimum is in fact a minimum. We call  $\text{OT}_p(\nu_1, \nu_2)$  the  $p$ -Wasserstein distance between  $\nu_1$  and  $\nu_2$ , though it differs from the usual Wasserstein distance, which is defined between measures with equal finite mass, while  $\text{OT}_p$  is defined between measures that can have different (and even infinite) masses. Intuitively,  $\text{OT}_p$  allows for some of the mass of  $\nu_1$  and  $\nu_2$  to be transported to the diagonal  $\partial\Omega := \{(u, u) \in \mathbb{R}^2\}$ , which acts as an infinitely deep landfill. The  $p$ -Wasserstein distance is a distance on the space  $\mathcal{M}_p = \{\nu \in \mathcal{M} : \text{OT}_p(\nu, 0) < \infty\}$ , where 0 denotes the null measure, and its restriction to the space of PDs coincides with the  $\text{OT}_p$  distance defined earlier.

We require the following lemma from [DL21]:

**Lemma 4.1.** *A sequence of measures  $(\nu_n)_{n \geq 1}$  converges with respect to  $\text{OT}_p$  to some measure  $\nu$  if and only if the sequence  $(\nu_n)_{n \geq 1}$  converges vaguely towards  $\nu$  and  $\text{Pers}_p(\nu_n) \rightarrow \text{Pers}_p(\nu)$  as  $n \rightarrow \infty$ .*

<sup>6</sup>A compact set  $K \subset \Omega$  is at *positive distance* from the diagonal. Hence, a measure  $\mu \in \mathcal{M}$  can have an accumulation of mass close to  $\partial\Omega$ .

For  $q > 0$ , given a function  $f : \mathbb{N} \rightarrow \mathbb{R}$  and a sequence of (non-necessarily measurable) real maps  $(Y_n)_n$  defined on some probabilistic space, the notation  $Y_n = O_{L^q}(f(n))$  means that  $\mathbb{E}^*[|Y_n|^q] = O(f(n)^q)$ , where  $\mathbb{E}^*$  denotes the outer expectation [VDVW97, p.6] (and similarly for the little  $o$  notation).

## 4.1 Region (1)

Consider the rescaled Radon measure  $\mu_{n,i} = \frac{1}{n} \sum_{u \in \text{dgm}_i^{(1)}(\mathbf{A}_n)} \delta_{n^{1/m}u}$ —this rescaling is natural, as the number of points in  $a_{n,i}^{(1)}$  is typically of order  $n$ , while their coordinates are of order  $n^{-1/m}$ —and note that  $\mu_{n,i}$  is a random measure, owing to the randomness of the set  $\mathbf{A}_n$ . Goel, Trinh and Tsunoda studied the vague convergence of the sequence  $(\mu_{n,i})_n$  [GTT19, Remark 4.2]. Namely, assuming that the density  $f$  satisfies  $\int_{\mathbb{M}} f^j < \infty$  for all  $j \geq 0$ , they show that with probability 1 the sequence  $(\mu_{n,i})_n$  converges vaguely to some (non-random) Radon measure  $\mu_{f,i}$ . The limit measure  $\mu_{f,i}$  has support  $\{0\} \times \mathbb{R}_+$  if  $i = 0$  and  $\Omega$  if  $0 < i < m$ ; it is the zero measure if  $i \geq m$ . We can further describe it as follows: let  $\mu_{\infty,i,m}$  be the limit of the sequence  $(\mu_{n,i})$  in the case where the sample  $\mathbf{A}_n$  is uniform on the unit cube  $[0, 1]^m$  (see [DP19]). Then, for any continuous function  $\phi : \Omega \rightarrow \mathbb{R}$  with compact support,

$$\int_{\Omega} \phi(u) d\mu_{f,i}(u) = \int_{\Omega} \int_{\mathbb{M}} f(x) \phi(f(x)^{-1/m}u) dx d\mu_{\infty,i,m}(u). \quad (21)$$

Note that the vague convergence of Radon measures is only defined with respect to compactly supported functions; as such, it is blind to phenomena located increasingly close to the diagonal  $\partial\Omega$  as  $n$  goes to infinity. In particular, and except in the case of the uniform distribution on the unit cube  $[0, 1]^m$  (see [DP19]), it was not known whether  $\mu_{n,i}$  converges to  $\mu_{f,i}$  for the  $\text{OT}_p$  distance as well, nor whether the sequences of total persistence  $(\text{Pers}_{\alpha}(\mu_{n,i}))$  converge. We close this gap with the following result.

**Theorem 4.2** (Law of large numbers). *Assume that  $P$  has a density  $f$  on  $\mathbb{M}$  bounded away from 0 and  $\infty$ . Let  $i \geq 0$  be an integer and let  $1 \leq p < \infty$ . Then  $\mu_{f,i} \in \mathcal{M}_p$  and*

$$\mathbb{E}[\text{OT}_p^p(\mu_{n,i}, \mu_{f,i})] \xrightarrow{n \rightarrow \infty} 0.$$

Furthermore, for all  $\alpha > 0$ ,  $\text{Pers}_{\alpha}(\text{dgm}_i^{(1)}(\mathbf{A}_n)) n^{\frac{\alpha}{m}-1} = \text{Pers}_{\alpha}(\mu_{n,i}) = \text{Pers}_{\alpha}(\mu_{f,i}) + o_{L^1}(1)$ .

Note that the statement becomes clearly wrong when  $p = \infty$ , as the support of  $\mu_{n,i}$  is bounded for all  $n$ , while that of  $\mu_{f,i}$  is not. Note also that without the assumption from Theorem 4.2 that  $f$  be bounded away from 0, it is easy to find examples where  $\int_{\mathbb{M}} f(x)^{1-\frac{p}{m}} dx = \infty$  for  $p$  large enough, hence  $\text{Pers}_p(\mu_{f,i}) = \infty$  and  $\mu_{f,i} \notin \mathcal{M}_p$ ; consider e.g. the segment  $[-1/2, 1/2]$ ,  $f(x) = x$  and  $p \geq 2$  (boundaryless examples can be similarly constructed).

*Proof of Theorem 4.2.* Recall that Goel, Trinh and Tsunoda [GTT19] have shown that almost surely, the sequence of Radon measures  $(\mu_{n,i})_n$  vaguely converges to the Radon measure  $\mu_{f,i}$ . We start by showing that, using this vague convergence and Lemma 4.1, it is enough to prove the convergence of the  $p$ -total persistence.

**Lemma 4.3.** *Let  $(\nu_n)_{n \geq 1}$  be a sequence of random measures in  $\mathcal{M}_p$  that converges vaguely almost surely to a Radon measure  $\nu \in \mathcal{M}_p$ , and such that  $\mathbb{E}[|\text{Pers}_p(\nu_n) - \text{Pers}_p(\nu)|] \rightarrow_{n \rightarrow \infty} 0$ . Then,  $\mathbb{E}[\text{OT}_p^p(\nu_n, \nu)] \rightarrow_{n \rightarrow \infty} 0$ .*

*Proof.* Let us first show that  $(D_n)_{n \geq 1} = (\text{OT}_p^p(\nu_n, \nu))_{n \geq 1}$  converges in probability to 0. We use the following standard result: if for every subsequence  $(Z_{n_k})_{k \geq 1}$  of  $(Z_n)_{n \geq 1}$ , one can extract a subsequence  $(Z_{n_{k_l}})_{l \geq 1}$  that converges almost surely to 0, then the sequence  $(Z_n)_{n \geq 1}$  converges in probability to 0. Let  $(D_{n_k})_{k \geq 1}$  be a subsequence of  $(D_n)_{n \geq 1} = (\text{OT}_p^p(\nu_n, \nu))_{n \geq 1}$ . Then, as  $(\text{Pers}_p(\nu_n))_{n \geq 1}$  converges in  $L^1$  to  $\text{Pers}_p(\nu)$ , it also converges in probability. In particular, there exists a subsequence  $(n_{k_l})_{l \geq 1}$  such that  $(\text{Pers}_p(\nu_{n_{k_l}}))_{l \geq 1}$  converges almost surely to  $\text{Pers}_p(\nu)$ . When restricting ourselves to this subsequence, we have both vague convergence of the measures and convergence of the  $p$ -total persistence. Hence, according to Lemma 4.1, we have  $D_{n_{k_l}} = \text{OT}_p^p(\nu_{n_{k_l}}, \nu) \rightarrow_{l \rightarrow \infty} 0$  almost surely, proving that we actually have that  $(D_n)_{n \geq 1}$  converges in probability to 0. To prove that  $\mathbb{E}[D_n] \rightarrow_{n \rightarrow \infty} 0$ , it remains to show that the sequence  $(D_n)_{n \geq 1}$  is uniformly integrable. By considering the trivial transport plan that sends all probability mass to  $\partial\Omega$ , we have for all  $n \geq 1$

$$D_n \leq \text{Pers}_p(\nu_n) + \text{Pers}_p(\nu).$$

But the sequence  $(\text{Pers}_p(\nu_n))_{n \geq 1}$  is uniformly integrable, as it converges in  $L^1$ . Hence, so is the sequence  $(D_n)_{n \geq 1}$ , concluding the proof.  $\square$

Using Lemma 4.3, Theorem 4.2 would follow from the facts that  $\mu_{f,i} \in \mathcal{M}_p$  and that  $\mathbb{E}[|\text{Pers}_p(\mu_{n,i}) - \text{Pers}_p(\mu_{f,i})|]$  converges to 0.

Recall that  $C_c(\Omega)$  is the set of continuous functions  $f : \Omega \rightarrow \mathbb{R}$  with compact support (i.e. the support is bounded and at positive distance from  $\partial\Omega$ ). For  $s \geq 0$ , let  $T_s = \{(u_1, u_2) \in \Omega : u_2 \geq s\}$ .

**Lemma 4.4.** *Let  $\alpha > 0$ . Let  $(\nu_n)_{n \geq 1}$  be a sequence of random measures in  $\mathcal{M}_\alpha$  that converges vaguely almost surely to a Radon measure  $\nu \in \mathcal{M}$ . Assume that the sequence of random variables  $(\nu_n(\Omega))_{n \geq 1}$  is uniformly integrable and that*

$$\sup_n \mathbb{E}[\text{Pers}_\alpha(\nu_n)] < +\infty \text{ and } \lim_{s \rightarrow +\infty} \limsup_n \mathbb{E}\left[\int_{T_s} \text{pers}^\alpha(u) d\nu_n(u)\right] = 0.$$

*Then,  $\nu \in \mathcal{M}_\alpha$  and  $\mathbb{E}[|\text{Pers}_\alpha(\nu_n) - \text{Pers}_\alpha(\nu)|] \rightarrow_{n \rightarrow \infty} 0$ .*

*Proof.* We divide the proof into several steps.

1. Let  $\phi \in C_c(\Omega)$ . We first show that  $(\int \phi d\nu_n)_{n \geq 1}$  converges in  $L^1$  to  $\int \phi d\nu$ . By assumption, the convergence holds almost surely. Furthermore, as  $\phi$  is bounded and as the sequence  $(\nu_n(\Omega))_{n \geq 1}$  is uniformly integrable, so is the sequence  $(\int \phi d\nu_n)_{n \geq 1}$ . Hence,  $\mathbb{E}[|\int \phi d(\nu_n - \nu)|] \rightarrow_{n \rightarrow \infty} 0$ .
2. Let  $(\phi_k)_{k \geq 1}$  be an increasing sequence of functions in  $C_c(\Omega)$  that converge pointwise to the function  $\text{pers}_\alpha$ . Then, almost surely,

$$\int \phi_k d\nu \leq \liminf_{n \rightarrow \infty} \int \phi_k d\nu_n \leq \liminf_{n \rightarrow \infty} \int \text{pers}_\alpha d\nu_n.$$

By Fatou's lemma,  $\mathbb{E}[\liminf_{n \rightarrow \infty} \int \text{pers}_\alpha d\nu_n] \leq \liminf_{n \rightarrow \infty} \mathbb{E}[\text{Pers}_\alpha(\nu_n)] = C < +\infty$  by assumption. Hence, by letting  $k \rightarrow \infty$  and applying the monotone convergence theorem, we obtain  $\text{Pers}_\alpha(\nu) \leq C$ , proving that  $\nu \in \mathcal{M}_\alpha$ .

3. The same argument can be applied to the constant function equal to 1, showing that  $\nu(\Omega) < +\infty$ .

4. Let  $s \geq 1$ . The function  $\text{pers}_\alpha$  can be decomposed into a sum of three positive continuous functions  $\text{pers}_\alpha = \phi_s^{(1)} + \phi_s^{(2)} + \phi_s^{(3)}$ , where  $\phi_s^{(1)}$  has compact support, the support of  $\phi_s^{(2)}$  is included in the band  $\{u \in \Omega : \text{pers}(u) \leq 1/s\}$  and the support of  $\phi_s^{(3)}$  is included in  $T_s$ . Hence,

$$\begin{aligned} \limsup_{n \rightarrow +\infty} \mathbb{E}[|\text{Pers}_\alpha(\nu_n) - \text{Pers}_\alpha(\nu)|] &\leq \limsup_{n \rightarrow +\infty} \mathbb{E}\left[\left|\int \phi_s^{(1)} d(\nu_n - \nu)\right|\right] \\ &+ \limsup_{n \rightarrow +\infty} \mathbb{E}\left[\left|\int \phi_s^{(2)} d(\nu_n - \nu)\right|\right] + \limsup_{n \rightarrow +\infty} \mathbb{E}\left[\left|\int \phi_s^{(3)} d(\nu_n - \nu)\right|\right]. \end{aligned}$$

The first term in the above sum is equal to zero because of the first item, the second one is smaller than  $s^{-\alpha}(\sup_n \mathbb{E}[\nu_n(\Omega)] + \nu(\Omega))$ , and the third one is smaller than

$$\limsup_n \mathbb{E}\left[\int_{T_s} \text{pers}_\alpha(u) d\nu_n(u)\right] + \int_{T_s} \text{pers}_\alpha(u) d\nu(u).$$

Using the hypotheses of the lemma, the second and the third term converges to 0 as  $s$  goes to  $\infty$ . We obtain that  $\limsup_{n \rightarrow +\infty} \mathbb{E}[|\text{Pers}_\alpha(\nu_n) - \text{Pers}_\alpha(\nu)|] = 0$ .  $\square$

Our goal is to show that the conditions of Lemma 4.4 holds for the sequence  $(\mu_{n,i})_{n \geq 1}$  to conclude. Remark that for any Radon measure  $\nu \in \mathcal{M}_\alpha$  and  $s \geq 0$

$$\begin{aligned} \int_{T_s} \text{pers}^\alpha(u) d\nu(u) &= \alpha \int_0^\infty t^{\alpha-1} \nu(T_s \cap \{u : \text{pers}(u) \geq t\}) dt \\ &\leq \alpha \int_{s/2}^\infty t^{\alpha-1} \nu(T_{2t}) dt + \alpha \int_0^{s/2} t^{\alpha-1} \nu(T_s) dt \\ &\leq \alpha \int_{s/2}^\infty t^{\alpha-1} \nu(T_{2t}) dt + (s/2)^\alpha \nu(T_s), \end{aligned} \tag{22}$$

where we use Fubini's theorem for the first equality and the fact that  $\{u : \text{pers}(u) \geq t\} \subset T_{2t}$  for the first inequality. We also have  $\nu(\Omega) = \nu(T_0)$ . Hence, the different conditions of Lemma 4.4 can all be obtained by controlling the random variable  $\mu_{n,i}(T_s)$  for  $s \geq 0$ .

**Proposition 4.5.** *Let  $M$  be a compact submanifold with positive reach. Assume that  $P$  has a density  $f$  on  $M$  satisfying  $f_{\min} \leq f \leq f_{\max}$  for two positive constants  $f_{\min}, f_{\max}$ . Then there exist  $c, C > 0$  that depend on  $M, i, f_{\min}$  and  $f_{\max}$  such that for all integer  $n \geq 1$  and all  $s \geq 0$ ,*

$$\mathbb{E}[\mu_{n,i}(T_s)^2] \leq C \exp(-cs^m). \tag{23}$$

Before proving Proposition 4.5, let us show how to use it to conclude the proof of Theorem 4.2. First, it implies that the random variables  $\mu_{n,i}(\Omega) = \mu_{n,i}(T_0)$  for  $n \geq 1$  have a uniformly bounded second moment, and are therefore uniformly integrable. Second, we have  $\mathbb{E}[\mu_{n,i}(T_s)] \leq \mathbb{E}[\mu_{n,i}(T_s)^2]^{1/2}$  using Hölder's inequality. Hence, (22) implies that for any  $\alpha > 0$ , we have

$$\begin{aligned} \mathbb{E}[\text{Pers}_\alpha(\mu_{n,i})] &\leq \alpha \int_0^\infty t^{\alpha-1} \mathbb{E}[\mu_{n,i}(T_{2t})] dt \\ &\leq \alpha \sqrt{C} \int_0^\infty t^{\alpha-1} e^{-c2^{m-1}t^m} dt. \end{aligned}$$

In particular,  $\sup_n \mathbb{E}[\text{Pers}_\alpha(\mu_{n,i})] < +\infty$ . Likewise,

$$\begin{aligned} \sup_n \mathbb{E} \left[ \int_{T_s} \text{pers}_\alpha(u) d\mu_{n,i}(u) \right] &\leq \alpha \int_{s/2}^\infty t^{\alpha-1} \mathbb{E}[\mu_{n,i}(T_{2t})] dt + (s/2)^\alpha \mathbb{E}[\mu_{n,i}(T_s)] \\ &\leq \alpha \sqrt{C} \int_0^\infty t^{\alpha-1} e^{-c2^{m-1}t^m} dt + (s/2)^\alpha \sqrt{C} \exp(-c/2s^m) \end{aligned}$$

so that  $\lim_{s \rightarrow +\infty} \sup_n \mathbb{E}[\int_{T_s} \text{pers}_\alpha(u) d\mu_{n,i}(u)] = 0$ .

We are therefore in position to apply Lemma 4.4, proving the convergence of the  $\alpha$ -total persistence. Together with Lemma 4.3 with  $\alpha = p \geq 1$ , we also obtain the  $\text{OT}_p$ -convergence of  $\mu_{n,i}$ . It remains to prove Proposition 4.5.

*Proof of Proposition 4.5.* Write  $\varepsilon_n = d_H(\mathbf{A}_n, \mathbf{M})$ . Let  $s \geq 0$  and let  $\varepsilon_0 < \tau(\mathbf{M})/2$  be a small parameter, to be fixed later. Recall that we write  $\#S$  for the cardinality of a multiset  $S$ . Notice that  $\mu_{n,i}(T_s) = 0$  if  $sn^{-1/m} > \varepsilon_n + \varepsilon_n^2/\tau(\mathbf{M})$ . In particular, we may assume without loss of generality that  $s \leq n^{1/m} \text{diam}(\mathbf{M})(1 + \text{diam}(\mathbf{M})/\tau(\mathbf{M})) = n^{1/m} \varepsilon_{\max}$ , for otherwise there is nothing to prove. Consider the event  $E = \{\varepsilon_n + \varepsilon_n^2/\tau(\mathbf{M}) < \varepsilon_0\}$ . By definition of Region (1), if  $E$  is satisfied, then all the coordinates of points of the PD  $\text{dgm}_i^{(1)}(\mathbf{A}_n)$  are smaller than  $\varepsilon_0$ . Notice that the cardinality of  $\text{dgm}_i(\mathbf{A}_n)$  is smaller than the number of  $i$ -dimensional simplices in the Čech complex of  $\mathbf{A}_n$ , which is itself smaller than  $n^{i+1}$  (as each simplex corresponds uniquely to a choice of  $i+1$  vertices of  $\mathbf{A}_n$ ). Hence

$$\mathbb{E}[\mu_{n,i}(T_s)^2 \mathbf{1}\{E^c\}] \leq n^{-2} n^{2i+2} \mathbb{P}(\varepsilon_n + \varepsilon_n^2/\tau(\mathbf{M}) \geq \varepsilon_0).$$

We require the following lemma, which bounds the upper tail of the random variable  $\varepsilon_n = d_H(\mathbf{A}_n, \mathbf{M})$  and which we prove in the Appendix.

**Lemma 4.6.** *If  $r \leq \tau(\mathbf{M})/2$ , then*

$$\mathbb{P}(d_H(\mathbf{A}_n, \mathbf{M}) > r) \leq \frac{C}{f_{\min} r^m} \exp(-nc f_{\min} r^m) \quad (24)$$

for two positive constants  $c = c(m), C = C(m)$ . In particular, for any  $q \geq 1$ ,  $d_H(\mathbf{A}_n, \mathbf{M}) = O_{L^q}((\ln n/n)^{1/m})$ .

Note that  $\varepsilon_n \leq \text{diam}(\mathbf{M})$ , so  $\mathbb{P}(\varepsilon_n + \varepsilon_n^2/\tau(\mathbf{M}) \geq \varepsilon_0) \leq \mathbb{P}(\varepsilon_n \geq c_0 \varepsilon_0)$ , with  $c_0 = (1 + \text{diam}(\mathbf{M})/\tau(\mathbf{M}))^{-1}$ . Apply Lemma 4.6 with  $r = c_0 \varepsilon_0$  to obtain that

$$\mathbb{E}[\mu_{n,i}(T_s)^2 \mathbf{1}\{E^c\}] \leq n^{2i} \frac{C}{f_{\min} r^m} \exp(-nc f_{\min} r^m) \leq C_0 \exp(-c_1 n)$$

for some positive constants  $c_1, C_0$ . Furthermore, recall that  $s \leq n^{1/m} \varepsilon_{\max}$ . Hence,

$$\mathbb{E}[\mu_{n,i}(T_s)^2 \mathbf{1}\{E^c\}] \leq C_0 \exp(-c_1 \varepsilon_{\max}^{-m} s^m) = C_0 \exp(-c_2 s^m)$$

for some positive constant  $c_2$ .

It remains to bound  $\mathbb{E}[\mu_{n,i}(T_s)^2 \mathbf{1}\{E\}]$ . For each  $j = 1, \dots, n$ , consider the set  $\Xi_{n,s}^j$  of critical points  $z \in \text{Crit}(\mathbf{A}_n)$  such that  $\sigma_{\mathbf{A}_n}(z)$  contains the point  $X_i$  and  $sn^{-1/m} \leq d_{\mathbf{A}_n}(z) \leq \varepsilon_n + \varepsilon_n^2/\tau(\mathbf{M})$ .

**Lemma 4.7.** *It holds that  $\mu_{n,i}(T_s)$  is smaller than*

$$\frac{1}{n} \sum_{j=1}^n L_j^{i+2}, \quad (25)$$

where  $L_j$  is the cardinality of the set  $\bigcup_{z \in \Xi_{n,s}^j} \sigma_{A_n}(z)$ .

*Proof.* Lemma A.1 applied to a realization of  $A_n$  and the interval  $[sn^{-1/m}, \varepsilon_n + \varepsilon_n^2/\tau(\mathbf{M})]$  yields that the number of points in  $\text{dgm}_i(A_n)$  with at least one of their coordinates in  $[sn^{-1/m}, \varepsilon_n + \varepsilon_n^2/\tau(\mathbf{M})]$  is upper bounded by  $\sum_{j=1}^n L_j^{i+2}$ . By definition, this means that  $\mu_{n,i}(T_s) \leq \frac{1}{n} \sum_{j=1}^n L_j^{i+2}$ , as desired.  $\square$

One can show that the set  $\Xi_{n,s}^j$  is localized, in the sense that it is included in a ball centered at  $X_j$ , with a radius depending on the sample  $A_n$ , that is small with high probability. Hence, the number  $L_j$  is controlled by the number of points in  $A_n$  found in a small (random) neighborhood of  $X_j$ . Let us make this idea rigorous.

For  $r \geq 0$ , define the shape

$$\mathcal{C}(r) = \{y = (y_m, y_{d-m}) \in \mathbb{R}^m \times \mathbb{R}^{d-m} : \|y\| \leq r, \|y_{d-m}\| \leq \frac{\|y\|^2}{2\tau(\mathbf{M})}\}. \quad (26)$$

We build a partition of  $\mathcal{C}(r)$  in the following way. Consider a finite partition  $\mathcal{W}$  of the unit sphere in  $\mathbb{R}^m \times \{0\}^{d-m}$  into sets of diameters smaller than  $\theta = \pi/4$  (for the geodesic distance on the sphere), which we fix for a given dimension  $m$ . Let  $\mathcal{W}(r)$  be the partition of  $\mathcal{C}(r)$  consisting of the sets

$$\{y = (y_m, y_{d-m}) \in \mathcal{C}(r) : y_m/\|y_m\| \in W\},$$

where  $W$  is an element of the partition  $\mathcal{W}$ .

For  $x \in \mathbf{M}$ , consider an isometry  $\iota_x : \mathbb{R}^d \rightarrow \mathbb{R}^d$  sending  $\mathbb{R}^m \times \{0\}^{d-m}$  to  $T_x\mathbf{M}$ . Let  $\mathcal{C}(x, r) = x + \iota_x(\mathcal{C}(r))$ . Likewise, we define a partition  $\mathcal{W}(x, r)$  by applying the affine transformation  $W \mapsto x + \iota_x(W)$  to each  $W \in \mathcal{W}(r)$ .

For  $j = 1, \dots, n$ , let  $R_{jn}$  be the smallest radius  $r \leq \varepsilon_0$  such that every  $W \in \mathcal{W}(X_j, r)$  contains a point of  $A_n$  other than  $X_j$ . By convention, we let  $R_{jn} = \varepsilon_0$  if such a radius does not exist.  $R_{jn}$  can be made measurable with a good choice of  $x \mapsto \iota_x$ ; we assume it to be the case henceforth.

**Lemma 4.8.** *Let  $\varepsilon_0 \leq \tau(\mathbf{M})/\sqrt{2}$ . For all  $j = 1, \dots, n$ , if  $z \in \text{Crit}(A_n)$  is such that  $X_j \in \sigma_{A_n}(z)$  and  $d_{A_n}(z) \leq \varepsilon_0$ , then  $\|X_j - z\| \leq c_0 R_{jn}$  for some positive absolute constant  $c_0$ .*

*Proof.* Recall that for  $x \in \mathbf{M}$ ,  $\pi_x$  is the orthogonal projection on  $T_x\mathbf{M}$  while  $\pi_x^\perp$  is the orthogonal projection on the normal space at  $x$ . Let  $j = 1, \dots, n$  and let  $z \in \text{Crit}(A_n)$  be such that  $X_j \in \sigma_{A_n}(z)$ . The direction  $e = \pi_{X_j}(z - X_j)/\|\pi_{X_j}(z - X_j)\|$  belongs to the unit sphere in  $T_{X_j}\mathbf{M}$ ; note that  $\pi_{X_j}(z - X_j) \neq 0$  due to Lemma B.2 below, which applies as  $d_{A_n}(z) \leq \varepsilon_0 \leq \tau(\mathbf{M})/\sqrt{2}$ . Hence,  $\iota_x^{-1}(e)$  belongs to an element  $W_0$  of the partition  $\mathcal{W}$ . Consider the corresponding element  $W$  of the partition  $\mathcal{W}(X_j, R_{jn})$ .

If  $R_{jn} = \varepsilon_0$ , then the conclusion of the lemma holds (for  $c_0 = 1$ ): indeed we have  $\|X_j - z\| = d_{A_n}(z) \leq \varepsilon_0 \leq R_{jn}$ . Otherwise, by assumption, there exists a point  $X_k \in W$  for some  $k \neq j$ . As  $X_j \in \sigma_{A_n}(z)$ , it holds that  $\|X_j - z\| \leq \|X_k - z\|$ . Hence,

$$\|X_j - z\|^2 \leq \|X_k - z\|^2 = \|X_j - z\|^2 + \|X_j - X_k\|^2 + 2\langle X_k - X_j, X_j - z \rangle$$



and

$$\langle X_k - X_j, z - X_j \rangle \leq \frac{\|X_j - X_k\|^2}{2}. \quad (27)$$

We write

$$\langle X_k - X_j, z - X_j \rangle = \langle \pi_{X_j}(X_k - X_j), \pi_{X_j}(z - X_j) \rangle + \langle \pi_{X_j}^\perp(X_k - X_j), \pi_{X_j}^\perp(z - X_j) \rangle$$

By construction, as the diameter of  $W_0$  is less than  $\theta$ , we have  $\langle \pi_{X_j}(X_k - X_j), \pi_{X_j}(z - X_j) \rangle \geq \cos(\pi/4) \|\pi_{X_j}(X_k - X_j)\| \|\pi_{X_j}(z - X_j)\|$ . On the other hand, according to [Fed59, Theorem 4.18],

$$\|\pi_{X_j}^\perp(X_k - X_j)\| \leq \frac{\|X_k - X_j\|^2}{2\tau(\mathbf{M})}, \quad (28)$$

which also implies that  $\|\pi_{X_j}(X_k - X_j)\| \geq \|X_j - X_k\| \sqrt{1 - \frac{\varepsilon_0^2}{4\tau(\mathbf{M})^2}}$ . Similarly, Lemma B.2 from the Appendix states that  $\|\pi_{X_j}(z - X_j)\| \geq \|z - X_j\|/\sqrt{2}$  and  $\|\pi_{X_j}^\perp(z - X_j)\| \leq \|z - X_j\|^2/\tau(\mathbf{M})$  (using our assumption that  $d_{\mathbf{A}_n}(z) \leq \varepsilon_0 \leq \tau(\mathbf{M})/\sqrt{2}$ ).

Hence we obtain that

$$\begin{aligned} \cos(\pi/4) \|X_j - X_k\| \|z - X_j\| \frac{\sqrt{1 - \frac{\varepsilon_0^2}{4\tau(\mathbf{M})^2}}}{\sqrt{2}} &\leq \cos(\pi/4) \|\pi_{X_j}(X_k - X_j)\| \|\pi_{X_j}(z - X_j)\| \\ &\leq \frac{\|X_k - X_j\|^2}{2} + \frac{\|X_k - X_j\|^2 \|z - X_j\|^2}{2\tau(\mathbf{M})^2} \\ &\leq \|X_k - X_j\|^2 \left( \frac{1}{2} + \frac{\varepsilon_0^2}{2\tau(\mathbf{M})^2} \right). \end{aligned}$$

Dividing by  $\|X_j - X_k\|$  and using that  $\|X_j - X_k\| \leq R_{jn}$  and that  $\varepsilon_0 \leq \tau(\mathbf{M})/\sqrt{2}$ , we see that  $\|z - X_j\|$  is smaller than  $R_{jn}$  up to an absolute multiplicative constant.  $\square$

Let us now show that the random variable  $R_{jn}$  has controlled tails.

**Lemma 4.9.** *For all  $x \in \mathbf{M}$ ,  $r \geq 0$ ,  $B(x, r) \cap \mathbf{M} \subset \mathcal{C}(x, r)$ .*

*Proof.* Let  $y \in \mathbf{M}$  be such that  $\|\pi_x(y - x)\| \leq r$ . Then, according to [Fed59, Theorem 4.18],  $\|\pi_x^\perp(y - x)\| \leq \frac{\|y - x\|^2}{2\tau(\mathbf{M})}$ . In particular,  $B(x, r) \cap \mathbf{M} \subset \mathcal{C}(x, r)$ .  $\square$

**Lemma 4.10.** *For  $0 < t \leq \tau(\mathbf{M})/4$  and  $j = 1, \dots, n$ , we have  $\mathbb{P}(R_{jn} > t) \leq C e^{-c f_{\min(n-1)} t^m}$  for some positive constants  $c = c(m)$ ,  $C = C(m)$ .*

*Proof.* If  $R_{jn}$  is larger than  $t$ , then there exists at least one set  $W \in \mathcal{W}(X_j, t)$  such that its intersection with  $\mathbf{A}_n$  contains only  $X_j$ . Hence,

$$\mathbb{P}(R_{jn} > t | X_j) \leq \sum_{W \in \mathcal{W}(X_j, t)} \mathbb{P}(\mathbf{A}_n \cap W = X_j | X_j) = \sum_{W \in \mathcal{W}(X_j, t)} (1 - P(W))^{n-1}.$$

Let  $\pi_0$  be the orthogonal projection from  $\mathbb{R}^d$  to  $\mathbb{R}^m \times \{0\}^{d-m}$ . The image of a set  $W \in \mathcal{W}(X_j, t)$  by the projection  $y \mapsto \pi_{X_j}(y - X_j)$  is equal to  $\iota_{X_j}(\pi_0(W_0)) \subset T_{X_j} \mathbf{M}$  for some  $W_0 \in \mathcal{W}(t)$ . For  $t \leq \tau(\mathbf{M})/4$ , the orthogonal projection  $y \in B(X_j, t) \cap \mathbf{M} \mapsto \pi_{X_j}(y - X_j) \in T_{X_j} \mathbf{M}$  is a diffeomorphism on its image, with Jacobian lower bounded by a constant  $c(m)$  that depends only on  $m$ , see e.g.

[Div21, Lemma 2.2]. According to Lemma 4.9, the preimage of  $\iota_{X_j}(\pi_0(W_0)) \subset T_{X_j}\mathbf{M}$  by this diffeomorphism is equal to  $W \cap M$ . Hence, by a change of variable,

$$P(W) = P(W \cap M) \geq f_{\min} c(m) \text{Vol}_m(W_0) \geq f_{\min} c'(m) t^m$$

for some  $c'(m) > 0$ . Hence,

$$\mathbb{P}(R_{jn} > t | X_j) \leq \#\mathcal{W} e^{-f_{\min} c'(m)(n-1)t^m}.$$

We conclude by taking the expectation.  $\square$

Let us now control the number of points found in a ball  $B(X_j, \kappa R_{jn})$  for some  $\kappa \geq 0$ . Let  $\rho_0 > 0$  be small enough such that  $\int_{B(x, \rho_0) \cap \mathbf{M}} f < 1/2$  for any  $x \in \mathbf{M}$ .

**Lemma 4.11.** *Let  $l \geq 0, \kappa > 0$  be such that  $\kappa \varepsilon_0 \leq \min(\rho_0, \tau(\mathbf{M})/4)$ . For  $j = 1, \dots, n$ , let  $K_{jn}(\kappa)$  be the number of elements of  $\mathbf{A}_n$  found in  $B(X_j, \kappa R_{jn})$ . Then,  $\mathbb{E}[K_{jn}(\kappa)^l] \leq C(1 + \left(\frac{f_{\max}}{f_{\min}}\right)^l \kappa^{lm})$  for some constant  $C = C(m, l)$ .*

*Proof.* Let us write  $\{W_1, \dots, W_K\} = \mathcal{W}(X_j, \kappa R_{jn})$ . Without loss of generality, we can assume that  $n \geq K + 1$ , as otherwise the bound is trivial. As in [DP19, Lemma 5], we remark that there is at least one sample point in every  $W_i$ , and that there is (almost surely) one single element  $W_{i^*}$  of the partition with exactly one sample point on its boundary. Let  $N_i$  be the cardinality of  $(W_i \cap \mathbf{A}_n) \setminus \{X_j\}$ , and  $N_{-1}$  be the cardinality of  $\mathbf{A}_n \setminus B(X_j, \kappa R_{jn})$ . Define

$$\tilde{\alpha}_i := \int_{W_i} f$$

and  $\alpha_i := \frac{\tilde{\alpha}_i}{1 - \tilde{\alpha}_{i^*}}$  for all  $i \neq i^*$ , as well as  $\alpha_{-1} = \frac{1 - \sum_{i=1}^K \tilde{\alpha}_i}{1 - \tilde{\alpha}_{i^*}}$ . Note that as  $\kappa R_{jn} \leq \rho_0$ , we have  $\tilde{\alpha}_i < 1/2$  for all  $i = 1, \dots, K$ . As  $\kappa R_{jn} \leq \kappa \varepsilon_0 \leq \tau(\mathbf{M})/4$ , we may use once again that the orthogonal projection  $y \in B(X_j, \kappa R_{jn}) \cap \mathbf{M} \mapsto \pi_{X_j}(y - X_j) \in T_{X_j}\mathbf{M}$  is a diffeomorphism on its image, with Jacobian upper and lower bounded by constants depending only on  $m$  (see [Div21, Lemma 2.2]) to also obtain that

$$c f_{\min}(\kappa R_{jn})^m \leq \tilde{\alpha}_i \leq C f_{\max}(\kappa R_{jn})^m$$

for all  $i = 1, \dots, K$  and some constants  $c = c(m), C = C(m)$ . Hence there exists  $C' = C'(m) > 0$  such that

$$c f_{\min}(\kappa R_{jn})^m \leq \alpha_i \leq C' f_{\max}(\kappa R_{jn})^m$$

for all  $i = 1, \dots, K$ . Consider a multinomial random variable  $L = (L_1, \dots, \widehat{L_{i^*}}, \dots, L_K, L_{-1})$  of parameters  $n - 2$  and  $(\alpha_1, \dots, \widehat{\alpha_{i^*}}, \dots, \alpha_K, \alpha_{-1})$ , and let  $E$  denote the event

$$\{L_i \geq 1 \forall i \in \{1, \dots, K\} \setminus \{i^*\}\}.$$

Then conditionally on  $X_j, R_{jn}$  and  $i^*$ , the variable  $N = (N_1, \dots, \widehat{N_{i^*}}, \dots, N_K, N_{-1})$  follows the same distribution as  $L | E$ . Thus, conditionally on  $X_j, R_{jn}$  and  $W_{i^*}$ , the variable  $K_{jn}(\kappa) = 1 + \sum_{i=1}^K N_i = 2 + \sum_{i=1, i \neq i^*}^K N_i$  (where the initial 1 comes from  $X_j$ ) has the same distribution as

$$2 + \sum_{i=1, i \neq i^*}^K L_i | E.$$

Note that as  $\kappa R_{jn} \leq \rho_0$ , we have  $\alpha_{-1} \geq 1/2$ .

As a result, Lemma B.3 in the Appendix yields that

$$\begin{aligned}
\mathbb{E}[K_{jn}(\kappa)^l | X_j, R_{jn}, W_{i^*}] &= \mathbb{E}[(2 + \sum_{i=1, i \neq i^*}^K L_i)^l | E] \leq 2^l (2^l + \mathbb{E}[(\sum_{i=1, i \neq i^*}^K L_i)^l | E]) \\
&\leq 2^l (2^l + C(K, l)(1 + (n-2) \sum_{i=1, i \neq i^*}^K \alpha_i)^l) \\
&\leq C'(K, l)(1 + (n \sum_{i=1, i \neq i^*}^K \alpha_i)^l) \leq C'(K, l)(1 + (nKC' f_{\max}(\kappa R_{jn})^m)^l) \\
&\leq C(m, l)(1 + \kappa^{ml} f_{\max}^l n^l R_{jn}^{ml}).
\end{aligned}$$

We can now conclude by considering

$$\mathbb{E}[K_{jn}(\kappa)^l] = \mathbb{E}_{X_j, R_{jn}, W_{i^*}}[\mathbb{E}[K_{jn}(\kappa)^l | X_j, R_{jn}, W_{i^*}]] \leq \mathbb{E}_{R_{jn}}[C(m, l)(1 + \kappa^{ml} f_{\max}^l n^l R_{jn}^{ml})]$$

The quantity  $n^l \mathbb{E}[R_{jn}^{ml}]$  is bounded by a constant, which is proved by integrating the tail bound found in Lemma 4.10. Indeed, Lemma 4.10 implies that the random variable  $nR_{jn}^m$  is subexponential with a subexponential norm  $m$  independent of  $n$ , of order  $O(1/f_{\min})$ ; the moment of order  $l$  of such a random variable is bounded by  $C_l m^l$ , see [Ver18, Section 2.7].  $\square$

Let us wrap things up. Recall Lemma 4.7: it holds that

$$\mu_{n,i}(T_s) \leq \frac{1}{n} \sum_{j=1}^n L_j^{i+2}, \quad (29)$$

where  $L_j$  is the cardinality of the set  $\bigcup_{z \in \Xi_{n,s}^j} \sigma_{A_n}(z)$ . But according to Lemma 4.8, if  $z \in \Xi_{n,s}^j$ , then  $\|X_j - z\| \leq c_0 R_{jn}$ . In particular,  $d_{A_n}(z) \leq c_0 R_{jn}$ , and any point  $y \in \sigma_{A_n}(z)$  is at distance less than  $2c_0 R_{jn}$  from  $X_j$ . Hence,  $L_j$  is smaller than  $K_{jn}(\kappa)$  for  $\kappa = 2c_0$ . Choose  $\varepsilon_0$  so that  $\kappa \varepsilon_0 \leq \min(\rho_0, \tau(\mathbf{M})/4)$ . We are in position to apply Lemma 4.11. We further remark that Lemma 4.8 implies that  $\Xi_{n,s}^j$  is empty if  $c_0 R_{jn} < sn^{-1/m}$ .

Hence, by Jensen's inequality,

$$\begin{aligned}
\mathbb{E}[\mu_{n,i}(T_s)^2 \mathbf{1}\{E\}] &\leq \mathbb{E} \left[ \left( \frac{1}{n} \sum_{j=1}^n \mathbf{1}\{2c_0 R_{jn} \geq sn^{-1/m}\} K_{jn}(2c_0)^{i+2} \right)^2 \right] \\
&\leq \mathbb{E} \left[ \mathbf{1}\{2c_0 R_{1n} \geq sn^{-1/m}\} K_{1n}(2c_0)^{2i+4} \right] \\
&\leq \sqrt{\mathbb{P}(2c_0 R_{jn} \geq sn^{-1/m}) \mathbb{E}[K_{1n}(2c_0)^{4i+8}]} \\
&\leq C(m, i) \exp(-c(m) f_{\min} s^m),
\end{aligned}$$

where we apply Lemma 4.10 and Lemma 4.11 at the last line.  $\square$

This completes the proof of Theorem 4.2.  $\square$

## 4.2 Regions (2)-(3)

It is a well-known fact that the Hausdorff distance  $\varepsilon = d_H(\mathbf{A}_n, \mathbf{M})$  between a random sample and  $\mathbf{M}$  is of order  $(\log n/n)^{1/m}$  whenever the underlying density  $f$  is bounded away from zero and  $\infty$  on  $\mathbf{M}$ , see e.g. [FLR<sup>+</sup>14]. Hence the PDs  $\text{dgm}_i^{(2)}(\mathbf{A}_n)$  and  $\text{dgm}_i^{(3)}(\mathbf{A}_n)$  can be described using Theorem 2.3. However, in the case where  $\mathbf{M}$  is a generic submanifold, one can actually obtain tighter results. We let  $\#E$  denote the cardinality of a multiset  $E$ .

**Proposition 4.12.** *Let  $\mathbf{M}$  be a generic  $m$ -dimensional submanifold. Assume that  $P$  has a density  $f$  on  $\mathbf{M}$  bounded away from 0 and  $\infty$ . Let  $i \geq 0$  be an integer. There exists an optimal matching  $\gamma_n : \text{dgm}_i(\mathbf{A}_n) \cup \partial\Omega \rightarrow \text{dgm}_i(\mathbf{M}) \cup \partial\Omega$  for the bottleneck distance between  $\text{dgm}_i(\mathbf{A}_n)$  and  $\text{dgm}_i(\mathbf{M})$  such that for any  $q \geq 1$ :*

- *Region (2): It holds that  $\max_{u \in \text{dgm}_i^{(2)}(\mathbf{A}_n)} |u_2 - \gamma_n(u)_2| = O_{L^q}(n^{-2/m})$ .*
- *Region (3): It holds that  $\max_{u \in \text{dgm}_i^{(3)}(\mathbf{A}_n)} \|u - \gamma_n(u)\|_\infty = O_{L^q}(n^{-2/m})$  and  $\#(\text{dgm}_i^{(3)}(\mathbf{A}_n)) = O_{L^q}(1)$ .*

We remark that the same bounds can be obtained almost surely (e.g. ‘‘a.s. there exists  $C > 0$  such that  $\max_{u \in \text{dgm}_i^{(3)}(\mathbf{A}_n)} \|u - \gamma_n(u)\|_\infty \leq Cn^{-2/m}$ ’), rather than in expectation, using similar arguments.

Proposition 4.12 yields two distinct improvements upon direct applications of Theorem 2.3 and Theorem 3.7 to the random case. First, we obtain bounds of order  $n^{-2/m}$  instead of bounds of order  $(\log n/n)^{2/m}$ . Second, the random sample  $\mathbf{A}_n$  is in general only  $\delta$ -sparse for  $\delta$  of order  $n^{-2/m}$ . Hence,  $\mathbf{A}_n$  is  $(\delta, \varepsilon)$ -dense in  $\mathbf{M}$ , but with a diverging ratio  $\varepsilon/\delta$ . Therefore, Theorem 3.7 cannot be applied to control the total number of points in  $\text{dgm}_i(\mathbf{A}_n)$  in Region (3).

*Proof.* Let  $\varepsilon_n = d_H(\mathbf{A}_n, \mathbf{M})$ . Let  $\Pi_{\mathbf{M}} := \{x \in \sigma_{\mathbf{M}}(z) : z \in \text{Crit}(\mathbf{M})\}$  be the (finite) set of projections of critical points  $z \in \text{Crit}(\mathbf{M})$ . The proof of Theorem 3.7 relied on the use of Theorem 1.6 in [ACSD23]. This theorem states roughly that both critical points  $z \in \text{Crit}(\mathbf{A}_n)$  far from  $\mathbf{M}$  and their projections  $x \in \sigma_{\mathbf{A}_n}(z)$  are stable with respect to the Hausdorff distance, meaning that every such point  $z$  is at distance  $O(\varepsilon_n)$  from a critical point  $z' \in \text{Crit}(\mathbf{M})$ , with  $x$  being at distance  $O(\varepsilon_n)$  from a point  $x' \in \Pi_{\mathbf{M}}$ . Thus, the number of critical points of  $\mathbf{A}_n$  located close to a given  $z' \in \text{Crit}(\mathbf{M})$  is crudely upper bounded by the number of subsets which can be formed by selecting elements in neighborhoods of size  $O(\varepsilon_n)$  around  $x' \in \sigma_{\mathbf{M}}(z')$ . We used the same idea to bound the cardinality of  $\text{dgm}_i^{(3)}(\mathbf{A})$  in Theorem 3.7.

In a random setting, the distance  $\varepsilon_n$  is of order  $(\ln n/n)^{1/m}$ , as suggested by Lemma 4.6, while the number of points found in a ball of radius  $r$  is typically of order  $nr^m$ , yielding a logarithmic number of elements in a neighborhood of size  $O(\varepsilon_n)$  around a given point  $x \in \mathbf{M}$ . Hence, our earlier strategy is not tight enough to bound in expectation the cardinality of  $\text{dgm}_i^{(3)}(\mathbf{A})$  by a constant.

We improve upon this strategy with the following intuition: the maximal distance between a point  $x$  of  $\mathbf{M}$  and the point cloud  $\mathbf{A}_n$  does not really matter in [ACSD23, Theorem 1.6], but only the density of the point cloud  $\mathbf{A}_n$  around the (finitely many) projections  $x \in \Pi_{\mathbf{M}}$ . Although  $\varepsilon_n$  is of order  $(\ln n/n)^{1/m}$ , the distance between a fixed point  $x \in \mathbf{M}$  and  $\mathbf{A}_n$  is known to be of order  $n^{-1/m}$  (see (40)). This remark explains how we can intuitively replace neighborhoods of radii  $(\ln n/n)^{1/m}$  by radii of size  $n^{-1/m}$  in the previous arguments.

We now make these ideas rigorous. Let  $\text{Crit}_>(\mathbf{A}_n) = \{z \in \text{Crit}(\mathbf{A}_n) : d_{\mathbf{A}_n}(z) \geq \tau(\mathbf{M})/2\}$  denote the set of critical points of  $\mathbf{A}_n$  “far” from  $\mathbf{M}$  and let  $\Pi_{\mathbf{A}_n} := \{x \in \sigma_{\mathbf{A}_n}(z) : z \in \text{Crit}_>(\mathbf{A}_n)\}$  be the corresponding set of projections. We let  $\varepsilon_0$  be a small constant to be fixed later. Theorem 1.6 from [ACSD23] states that, thanks to the genericity of  $\mathbf{M}$ , there exist  $K_1, K_2 > 0$  such that for  $\varepsilon_0$  small enough, if  $\varepsilon_n < \varepsilon_0$ :

- there exists a map  $\phi : \text{Crit}_>(\mathbf{A}_n) \rightarrow \text{Crit}(\mathbf{M})$  such that  $d(z, \phi(z)) \leq K_1 \varepsilon_n$  for all  $z \in \text{Crit}_>(\mathbf{A}_n)$ , and
- there exists a map  $\psi : \Pi_{\mathbf{A}_n} \rightarrow \Pi_{\mathbf{M}}$  such that  $d(x, \psi(x)) \leq K_2 \varepsilon_n$  for all  $x \in \Pi_{\mathbf{A}_n}$ .

Furthermore, the map  $\psi$  is such that for each  $z \in \text{Crit}_>(\mathbf{A}_n)$  and each  $x' \in \sigma_{\mathbf{M}}(\phi(z))$ , there exists  $x \in \sigma_{\mathbf{A}_n}(z)$  such that  $\psi(x) = x'$  (if  $\varepsilon_0$  is chosen small enough). Indeed, due to the genericity of  $\mathbf{M}$ , each critical point  $z' \in \text{Crit}(\mathbf{M})$  belongs to the relative interior of  $\sigma_{\mathbf{M}}(z')$  (as stated in Theorem 3.5). In particular, it cannot belong to the convex hull of any strict subset of  $\sigma_{\mathbf{M}}(z')$ . By continuity of the convex hull for the Hausdorff distance, as soon as  $z \in \text{Crit}_>(\mathbf{A}_n)$  is close enough to  $\phi(z)$  and its projections  $x \in \sigma_{\mathbf{A}_n}(z)$  are close enough to  $\sigma_{\mathbf{M}}(\phi(z))$  (i.e. as soon as  $\varepsilon_n$  is small enough), the point  $z$  cannot belong to the convex hull of any subset  $S \subset \sigma_{\mathbf{A}_n}(z)$  that does not contain for each  $x' \in \sigma_{\mathbf{M}}(\phi(z))$  at least one point  $x$  such that  $\psi(x) = x'$ . As  $z$  must belong to the convex hull of  $\sigma_{\mathbf{A}_n}(z)$ , we conclude that each  $x' \in \sigma_{\mathbf{M}}(\phi(z))$  has at least one preimage by  $\psi$  in  $\sigma_{\mathbf{A}_n}(z)$ . As  $\text{Crit}(\mathbf{M})$  is finite, taking the minimum distance such that this property holds for  $z'$  over all  $z' \in \text{Crit}(\mathbf{M})$  proves the claim.

We introduce the random function defined as

$$\forall r \geq 0, E(r) := \min(\sup\{d_{\mathbf{A}_n}(y) : y \in \mathbf{M} \cap \Pi_{\mathbf{M}}^r\}, \varepsilon_0) =: \min(F(r), \varepsilon_0), \quad (30)$$

where  $\Pi_{\mathbf{M}}^r$  is as usual the  $r$ -offset of  $\Pi_{\mathbf{M}}$ . This random variable measures the density of the point cloud  $\mathbf{A}_n$  in neighborhoods of size  $r$  of points  $x' \in \Pi_{\mathbf{M}}$ .

Let

$$\rho_n := \sup\{d_{\Pi_{\mathbf{M}}}(x) : x \in \Pi_{\mathbf{A}_n}\} \quad (31)$$

give (when  $\varepsilon_n < \varepsilon_0$  for  $\varepsilon_0$  small enough) the largest distance between a projection  $x \in \Pi_{\mathbf{A}_n}$  and the corresponding projection  $\psi(x) \in \Pi_{\mathbf{M}}$ . We also define

$$\eta_n := \sup\{d_{\Pi_{\mathbf{M}}}(x) : x = \pi_{\mathbf{M}}(z), z \in \text{Crit}_>(\mathbf{A}_n)\}. \quad (32)$$

We require the following controls on the random variables  $\rho_n$  and  $\eta_n$ .

**Lemma 4.13.** *There exist positive constants  $\varepsilon_0 = \varepsilon_0(\mathbf{M})$ ,  $K_3 = K_3(\mathbf{M})$ ,  $K_4 = K_4(\mathbf{M})$  and  $K_5 = K_5(\mathbf{M})$  such that if  $\varepsilon_n < \varepsilon_0$ , it holds that for any  $z \in \text{Crit}_>(\mathbf{A}_n)$*

$$|d_{\mathbf{A}_n}(z) - d_{\mathbf{M}}(\phi(z))| \leq K_3 E(\eta_n)^2, \quad (33)$$

$$\eta_n \leq K_4 E(\eta_n), \quad (34)$$

$$\rho_n \leq K_5 E(\eta_n). \quad (35)$$

*Proof.* The proof mostly follows from a careful read of the proof of Theorem 1.6 in [ACSD23]. First, remark that in the proof of Lemma 5.1 in [ACSD23] (applied with  $r = \tau(\mathbf{M})/2$ ,  $R = \text{diam}(\mathbf{M})$ ), the Hausdorff distance  $\varepsilon = d_H(\mathbf{A}_n, \mathbf{M})$  can be replaced by  $E(\eta_n)$ . Hence, if  $z \in \text{Crit}_>(\mathbf{A}_n)$ , there exists a  $\mu$ -critical point  $z'$  of  $\mathbf{M}$  at distance less than  $E(\eta_n)$ , with  $\mu \leq L_1(\mathbf{M})E(\eta_n)$ . By genericity, for a choice of  $\varepsilon_0$  small enough, this point  $z'$  is at distance  $L_2(\mathbf{M})\mu$

from a critical point  $z_0 \in \text{Crit}(\mathbf{M})$ . For  $\varepsilon_0$  small enough, this point  $z_0$  is necessarily equal to  $\phi(z)$ , with  $\|z - \phi(z)\| \leq (1 + L_2L_1)E(\eta_n)$ . Due to the Lipschitz property of the projection onto  $\mathbf{M}$  around  $z_0$  (see the arguments found at the bottom of p. 19 in [ACSD23]), any point  $x \in \pi_{\mathbf{M}}(z)$  is such that  $\|x - x'\| \leq L_3(\mathbf{M})\|z - \phi(z)\|$  for some  $x' \in \sigma_{\mathbf{M}}(\phi(z))$ . By taking the supremum over all such points  $x$ , we obtain that

$$\eta_n \leq L_3(1 + L_2L_1)E(\eta_n),$$

proving (34).

One can also check that  $\varepsilon_n$  can be replaced by  $E(\eta_n)$  in the end of the proof of Theorem 1.6 in [ACSD23], so that (35) holds.

The proof of (33) is essentially the same as that of the first point of Theorem 3.7—we omit it.  $\square$

Consider a (random) optimal matching  $\gamma_n : \text{dgm}_i(\mathbf{A}_n) \cup \partial\Omega \rightarrow \text{dgm}_i(\mathbf{M}) \cup \partial\Omega$  such that  $\max_{u \in \text{dgm}_i^{(2)}(\mathbf{A}_n)} |u_2 - \gamma_n(u)_2| \leq C\varepsilon_n^2$  and  $\max_{u \in \text{dgm}_i^{(3)}(\mathbf{A}_n)} \|u - \gamma_n(u)\|_{\infty} \leq C\varepsilon_n^2$  whose existence is guaranteed by Theorem 2.3 (for some  $C = C(\mathbf{M}) > 0$ ). For  $r \geq 0$ , we let  $N(r)$  be the number of points of  $\mathbf{A}_n$  found in  $\Pi_{\mathbf{M}}^r$  (i.e. at distance less than  $r$  from a point in  $\Pi_{\mathbf{M}}$ ).

**Lemma 4.14.** *There exists  $\varepsilon_0 = \varepsilon_0(\mathbf{M})$  such that if  $\varepsilon_n \leq \varepsilon_0$ , then*

$$\begin{aligned} \max_{u \in \text{dgm}_i^{(2)}(\mathbf{A}_n)} |u_2 - \gamma_n(u)_2| &\leq K_3E(\eta_n)^2, \\ \max_{u \in \text{dgm}_i^{(3)}(\mathbf{A}_n)} \|u - \gamma_n(u)\|_{\infty} &\leq K_3E(\eta_n)^2. \end{aligned}$$

Furthermore,  $\#(\text{dgm}_i^{(3)}(\mathbf{A}_n))$  is smaller than  $N(\rho_n)^{i+2}$ .

*Proof.* Let us assume from now on that  $\varepsilon_0$  is small enough that  $2(K_3 + C)\varepsilon_0^2$  is smaller than the smallest difference between two distinct critical values of  $d_{\mathbf{M}}$ .

Consider a point  $(u_1, u_2) \in \text{dgm}_i^{(3)}(\mathbf{A}_n)$  that is mapped by  $\gamma_n$  to the diagonal. The coordinates  $u_1$  and  $u_2$  differ by at most  $C\varepsilon_n^2$ , hence they are critical values of  $d_{\mathbf{A}_n}$  that correspond to critical points that are mapped by  $\phi$  to two critical points of  $\mathbf{M}$  that have the same critical value (due to  $2(K_3 + C)\varepsilon^2$  being smaller than the smallest difference between two distinct critical values of  $d_{\mathbf{M}}$ ). But as stated in Equation (33) of Lemma 4.13, these critical values must then be  $K_3E(\eta_n)^2$ -close to that of the two critical points of  $\mathbf{M}$ , hence  $\|u - \gamma_n(u)\|_{\infty} = d(u, \partial\Omega) = (u_2 - u_1)/2 \leq K_3E(\eta_n)^2$ , as desired.

Likewise, consider a point  $(u_1, u_2)$  of  $\text{dgm}_i^{(3)}(\mathbf{A}_n)$  that is mapped by  $\gamma_n$  to a point  $(v_1, v_2)$  of  $\text{dgm}_i^{(3)}(\mathbf{M})$ . The birth coordinates  $u_1$  and  $v_1$  differ by at most  $C\varepsilon^2$ , hence the associated critical values must correspond to a critical point  $z$  of  $\mathbf{A}_n$  and a critical point  $z'$  of  $\mathbf{M}$  such that  $\phi(z)$  has the same filtration value as  $z'$  (again due to  $2(K_3 + C)\varepsilon^2$  being smaller than the smallest difference between two distinct critical values of  $d_{\mathbf{M}}$ ). But as above, we have  $|u_1 - v_1| = |d_{\mathbf{A}_n}(z) - d_{\mathbf{M}}(z')| = |d_{\mathbf{A}_n}(z) - d_{\mathbf{M}}(\phi(z))| \leq K_3E(\eta_n)^2$ . As the same applies to  $u_2$  and  $v_2$ , we find that  $\|u - \gamma_n(u)\|_{\infty} \leq K_3EE(\eta_n)^2$  and  $\max_{u \in \text{dgm}_i^{(3)}(\mathbf{A}_n)} \|u - \gamma_n(u)\|_{\infty} \leq K_3E(\eta_n)^2$ . The same reasoning shows that  $\max_{u \in \text{dgm}_i^{(2)}(\mathbf{A}_n)} |u_2 - \gamma_n(u)_2| \leq K_3E(\eta_n)^2$ .

Finally, and by definition,  $\Pi_{\mathbf{A}_n} = \{x \in \sigma_{\mathbf{A}_n}(z) : z \in \text{Crit}(\mathbf{A}_n), d_{\mathbf{A}_n}(z) \geq \tau(\mathbf{M})/2\}$  is included in  $\mathbf{A}_n \cap \Pi_{\mathbf{M}}^{\rho_n}$ , hence  $\#(\Pi_{\mathbf{A}_n}) \leq N(\rho_n)$ . Lemma A.1 applied to  $\mathbf{A}_n$  and the interval  $[\tau(\mathbf{M})/2, +\infty)$  then yields that the number of points in  $\text{dgm}_i^{(3)}(\mathbf{A}_n)$  is smaller than  $N(\rho_n)^{i+2}$ .  $\square$

**Lemma 4.15.** *Let  $\varepsilon_0$  be the parameter defined in Lemma 4.14. It holds that for all  $q \geq 1$ ,  $E(\eta_n)\mathbf{1}\{\varepsilon_n \leq \varepsilon_0\} = O_{L^q}(n^{-1/m})$  and  $N(\rho_n)\mathbf{1}\{\varepsilon_n \leq \varepsilon_0\} = O_{L^q}(1)$ .*

Assume for a moment that Lemma 4.15 holds. Then, we write (for a choice of  $\varepsilon_0$  small enough)

$$\begin{aligned} \max_{u \in \text{dgm}_i^{(2)}(\mathbf{A}_n)} |u_2 - \gamma_n(u)_2| &\leq K_3 E(\eta_n)^2 \mathbf{1}\{\varepsilon_n \leq \varepsilon_0\} + R(\mathbf{M}) \mathbf{1}\{\varepsilon_n > \varepsilon_0\} \\ &\leq K_3 E(\eta_n)^2 \mathbf{1}\{\varepsilon_n \leq \varepsilon_0\} + R(\mathbf{M}) \mathbf{1}\{\varepsilon_n > \varepsilon_0\}. \end{aligned} \quad (36)$$

where we recall that  $R(\mathbf{M})$  is the radius of the smallest enclosing ball of  $\mathbf{M}$ . Because of Lemma 4.6, the random variable  $\mathbf{1}\{\varepsilon_n > \varepsilon_0\}$  is a  $O_{L^q}(n^{-2/m})$  for all  $q \geq 1$ . But then, because of Lemma 4.15, we obtain that the right-hand side in (36) is a  $O_{L^q}(n^{-2/m})$  for all  $q \geq 1$ . Likewise, we obtain that  $\max_{u \in \text{dgm}_i^{(3)}(\mathbf{A}_n)} \|u - \gamma_n(u)\|_\infty = O_{L^q}(n^{-2/m})$  for all  $q \geq 1$ . At last, we have

$$\#(\text{dgm}_i^{(3)}(\mathbf{A}_n)) \leq N(\rho_n)^{i+2} \mathbf{1}\{\varepsilon_n \leq \varepsilon_0\} + n^{i+2} \mathbf{1}\{\varepsilon_n > \varepsilon_0\}.$$

The first term is a  $O_{L^q}(1)$  for all  $q \geq 1$  because of Lemma 4.15, while the second one is a  $O_{L^q}(1)$  for all  $q \geq 1$  because of Lemma 4.6. This concludes the proof of Proposition 4.12. Let us now prove Lemma 4.15.  $\square$

*Proof of Lemma 4.15.* Let  $t > 0$ . The key observation to obtain Lemma 4.15 is that (34) implies that if  $\eta_n > t$ , then there exists  $r > t$  with  $r \leq K_4 E(r)$ .

**Lemma 4.16.** *For all  $\lambda > 0$ , there exist positive constants  $C_\lambda, c_\lambda$  (depending on  $\lambda, \mathbf{M}$  and  $f_{\min}$ ) such that for all  $r > 0$ ,  $\mathbb{P}(r \leq \lambda E(r)) \leq C_\lambda \exp(-c_\lambda n r^m)$ .*

*Proof.* Remark that if  $r \leq \lambda E(r)$ , then  $r \leq \lambda F(r)$ . The set  $\mathbf{M} \cap \Pi_{\mathbf{M}}^r$  is the union of a finite number of balls of radius  $r$ . Hence, it can be covered by  $C_\lambda = C_\lambda(\mathbf{M})$  open balls of radius  $r/(2\lambda)$ , with centers  $x_1, \dots, x_{C_\lambda} \in \mathbf{M}$ . Note that if all these balls intersect  $\mathbf{A}_n$ , then for all  $y \in \mathbf{M} \cap \Pi_{\mathbf{M}}^r$ ,  $d_{\mathbf{A}_n}(y) < r/\lambda$ . Hence, if  $\lambda F(r) \geq r$ , then the intersection of one of these balls with  $\mathbf{A}_n$  is empty. Hence, according to Lemma B.1,

$$\mathbb{P}(r \leq \lambda F(r)) \leq \sum_{k=1}^{C_\lambda} \mathbb{P}(d_{\mathbf{A}_n}(x_k) \geq r/(2\lambda)) \leq C_\lambda \exp(-C(\mathbf{M}) f_{\min} n (r/\lambda)^m). \quad \square$$

Remark that the fonction  $E$  is nondecreasing and 1-Lipschitz continuous: we have for  $r < s$ ,  $E(r) \leq E(s) \leq E(r) + (s - r)$ . Fix  $t > 0$  and consider the sequence  $t_k = a^k t$  for some  $a > 1$  to fix. Assume that  $\eta_n > t$  and that  $\varepsilon_n \leq \varepsilon_0$ . Then  $\eta_n$  is between two values  $t_k < t_{k+1}$ . But then, according to Lemma 4.14,

$$\begin{aligned} \frac{E(t_k)}{t_k} &\geq \frac{E(\eta_n) - (\eta_n - t_k)}{t_k} \geq \frac{1}{a} \frac{E(\eta_n)}{\eta_n} \frac{t_{k+1}}{t_k} - (a - 1) \geq \frac{1}{K_4} - (a - 1) \\ \frac{E(t_{k+1})}{t_{k+1}} &\geq \frac{1}{a} \frac{E(\eta_n)}{t_k} \geq \frac{1}{a} \frac{E(\eta_n)}{\eta_n} \geq \frac{1}{K_4 a}. \end{aligned}$$

Choose  $a > 1$  such that  $\frac{1}{K_4} - (a - 1) > 0$ , and let

$$\lambda = \min\left(\frac{1}{K_4} - (a - 1), \frac{1}{K_4 a}\right) > 0$$

We have proven that if  $\varepsilon_n \leq \varepsilon_0$  and  $\eta_n > t$ , then there exists  $k \geq 0$  with  $E(t_k) \geq \lambda t_k$ .

Hence,

$$\mathbb{P}(\eta_n > t, \varepsilon_n \leq \varepsilon_0) \leq C_\lambda \sum_{k \geq 0} \exp(-c_\lambda a^{km} n t^m).$$

A standard comparison between this sum and an integral shows that this sum is at most of order

$$K_6 (n t^m)^{-1} \exp(-K_7 n t^m). \quad (37)$$

for two positive constants  $K_6, K_7$  depending on  $M$  and  $f_{\min}$ . But when  $n t^m \leq 1$ , we can simply use the bound  $\mathbb{P}(\eta_n > t, \varepsilon_n \leq \varepsilon_0) \leq 1$ . Hence,

$$\mathbb{P}(\eta_n > t, \varepsilon_n \leq \varepsilon_0) \leq \min(1, K_6 (n t^m)^{-1} \exp(-K_7 n t^m)) \leq K_8 \exp(-K_9 n t^m) \quad (38)$$

for two other constants  $K_8, K_9$ .

To summarize, we have shown that the random variable  $n \eta_n^m \mathbf{1}\{\varepsilon_n \leq \varepsilon_0\}$  is subexponential, with subexponential norm depending only on  $M$  and  $f_{\min}$ , see e.g. [Ver18, Section 2.7]. We will now simply say that a random variable is subexponential to indicate that it is subexponential with a norm depending only on  $M$  and  $f_{\min}$ .

Lemma B.1 shows that the random variable  $n d_{\mathbf{A}_n}(x)^m$  for a fixed  $x \in M$  is also subexponential. Thus, so is  $n E(0)^m = n \max_{x \in \Pi_M} d_{\mathbf{A}_n}(x)^m$ , as a maximum of a finite number of subexponential random variables. As  $n E(\eta_n)^m \leq n(\eta_n + E(0))^m \leq 2^{m-1}(n \eta_n^m + n E(0)^m)$ , the random variable  $n E(\eta_n)^m \mathbf{1}\{\varepsilon_n \leq \varepsilon_0\}$  is also subexponential. In particular, we have  $E(\eta_n) \mathbf{1}\{\varepsilon_n \leq \varepsilon_0\} = O_{L^q}(n^{-1/m})$  for all  $q \geq 1$ .

It remains to bound  $N(\rho_n)$ . First, because of (35), the random variable  $n \rho_n^m \mathbf{1}\{\varepsilon_n \leq \varepsilon_0\}$  is subexponential. Also, for a fixed  $t$ ,  $N(t)$  follows a binomial distribution of parameter  $n$  and  $P(\Pi_M^t)$ . As long as  $t \leq \tau(M)/4$ , a ball of radius  $t$  is of mass smaller than  $C_m f_{\max} t^m$  (see Lemma B.1). Let  $0 \leq k \leq n$  be an integer. For any  $t \leq \tau(M)/4$ , we bound

$$\begin{aligned} \mathbb{P}(N(\rho_n) \geq k, \varepsilon_n \leq \varepsilon_0) &\leq \mathbb{P}(N(t) \geq k, \rho_n \leq t) + \mathbb{P}(\rho_n > t, \varepsilon_n \leq \varepsilon_0) \\ &\leq \mathbb{P}(N(t) \geq k) + 2 \exp(-K_{10} n t^m), \end{aligned}$$

where  $K_{10}$  is proportional to the subexponential norm of  $n \rho_n^m \mathbf{1}\{\varepsilon_n \leq \varepsilon_0\}$ , see [Ver18, Section 2.7]. Let  $t = (k/n)^{1/m} \min(\tau(M)/4, 1/(\#\Pi_M \cdot 2C_m f_{\max})^{1/m})$ . This choice of  $t$  ensures that  $t \leq \tau(M)/4$  and that  $\mathbb{E}[N(t)] \leq n(\#\Pi_M) C_m f_{\max} t^m \leq k/2$ . Then, by Bernstein's inequality [Ver18, Theorem 2.8.4],

$$\begin{aligned} \mathbb{P}(N(t) \geq k) &\leq \mathbb{P}(N(t) - \mathbb{E}[N(t)] \geq k/2) \leq \exp\left(-\frac{k^2/8}{n(\#\Pi_M) C_m f_{\max} t^m + k/6}\right) \\ &\leq \exp(-K_{11} k) \end{aligned}$$

for some constant  $K_{11}$  depending on  $m, M$  and  $f_{\max}$ . We have proven that for all  $k \geq 0$ ,

$$\begin{aligned} \mathbb{P}(N(\rho_n) \geq k, \varepsilon_n \leq \varepsilon_0) &\leq 2 \exp(-K_{11} k) + 2 \exp(-K_{10} n t^m) \\ &\leq 2 \exp(-K_{11} k) + 2 \exp(-K_{12} k) \end{aligned}$$

for some constant  $K_{12}$  depending on  $m, M, f_{\min}$  and  $f_{\max}$ . Hence, the random variable  $N(\rho_n) \mathbf{1}\{\varepsilon_n \leq \varepsilon_0\}$  is subexponential, with a subexponential norm depending on  $M, f_{\min}$  and  $f_{\max}$ . This implies in particular that  $N(\rho_n) \mathbf{1}\{\varepsilon_n \leq \varepsilon_0\} = O_{L^q}(1)$  for all  $q \geq 1$ .  $\square$



### 4.3 Consequences for the Wasserstein convergence of persistence diagrams

As a simple consequence of Theorem 4.2 and Proposition 4.12, we obtain that for  $i < m$ , the  $p$ -Wasserstein convergence of  $(\text{dgm}_i(\mathbf{A}_n))$  to  $\text{dgm}_i(\mathbf{M})$  holds if and only if  $p > m$ , as well as precise asymptotics for the total persistence of  $(\text{dgm}_i(\mathbf{A}_n))$ .

**Corollary 4.17.** *Let  $p \geq 1$  and let  $0 \leq i < d$  be an integer. Under the same assumptions as in Proposition 4.12, the following holds:*

- If  $p > m$ , then  $\mathbb{E}[\text{OT}_p^p(\text{dgm}_i(\mathbf{A}_n), \text{dgm}_i(\mathbf{M}))] \rightarrow 0$  as  $n \rightarrow \infty$ .
- If  $p = m$ ,  $\mathbb{E}[\text{OT}_p^p(\text{dgm}_i(\mathbf{A}_n), \text{dgm}_i(\mathbf{M}))] \rightarrow \text{Pers}_p(\mu_{\infty, i, m}) \text{Vol}(\mathbf{M})$  as  $n \rightarrow \infty$ , where  $\text{Vol}(\mathbf{M})$  is the volume of  $\mathbf{M}$ .
- If  $p < m$  and  $i < m$ , then  $\mathbb{E}[\text{OT}_p^p(\text{dgm}_i(\mathbf{A}_n), \text{dgm}_i(\mathbf{M}))] \rightarrow +\infty$  as  $n \rightarrow \infty$ .

Furthermore, for all  $\alpha > 0$ ,  $\text{Pers}_\alpha(\text{dgm}_i(\mathbf{A}_n))$  is equal to

$$\text{Pers}_\alpha(\text{dgm}_i(\mathbf{M})) + n^{1-\frac{\alpha}{m}} \text{Pers}_\alpha(\mu_{\infty, i, m}) \int_{\mathbf{M}} f(x)^{1-\frac{\alpha}{m}} dx + o_{L^1}(n^{1-\frac{\alpha}{m}}) + O_{L^1}\left(\left(\frac{\log n}{n}\right)^{\frac{1}{m}}\right).$$

As noted earlier, both  $\text{dgm}_i(\mathbf{A}_n)$  and  $\text{dgm}_i(\mathbf{M})$  are trivial if  $i \geq d$ .

This corollary, whose proof can be found in the Appendix, gives a precise answer to the questions raised in the introduction. First, when  $\mathbf{A}_n$  is a random subset of a  $m$ -dimensional generic manifold in  $\mathbb{R}^d$ , the  $\alpha$ -total persistence of  $\text{dgm}_i(\mathbf{A}_n)$  is not only bounded for  $\alpha > d$  (as was shown by Cohen-Steiner & al. [CSEHM10]), but for all  $\alpha \geq m$ . Moreover, the sequence  $\text{dgm}_i(\mathbf{A}_n)$  converges for the  $\text{OT}_p$  distance if  $p > m$ . A curious phenomenon can be observed in the case  $p = m$ : the sequence does not converge to  $\text{dgm}_i(\mathbf{M})$  as one would expect, but its distance to the power  $p$  to  $\text{dgm}_i(\mathbf{M})$  converges to some constant—in that case, the cost to the power  $p$  of matching all the points in Region (1) to the diagonal  $\partial\Omega$  neither converges to 0 nor diverges, but is asymptotically equal to this constant.

Using these asymptotics for the total persistence, we obtain regularity guarantees for a large family of feature maps, called *linear feature maps*, which includes feature maps introduced in [CFL<sup>+</sup>14, AEK<sup>+</sup>17, RHBK15, CWRW15, KHF16, STR<sup>+</sup>18, CCI<sup>+</sup>20]. Let  $(V, \|\cdot\|)$  be a normed vector space, and let  $\phi : \Omega \rightarrow V$  be a continuous bounded map. For  $\alpha \geq 0$ , the linear feature map  $\Phi_\alpha$  associated to  $\phi$  and defined on the space  $\mathcal{D}_f$  of PDs having a finite number of points is defined for all  $a \in \mathcal{D}_f$  by  $\Phi_\alpha(a) = \sum_{u \in a} \text{pers}(u)^\alpha \phi(u) \in V$ .

**Corollary 4.18.** *Let  $\alpha \geq 1$  and let  $0 \leq i < d$  be an integer. Under the same assumptions as in Proposition 4.12, it holds that  $\Phi_\alpha(\text{dgm}_i(\mathbf{A}_n))$  converges in probability to  $\Phi_\alpha(\text{dgm}_i(\mathbf{M}))$  whenever  $\alpha > m$ .*

The proof is given in the Appendix. Remark that other weighting schemes are possible. For instance, [KFH18] argued for using linear feature maps of the form

$$\Phi_\alpha(a) = \sum_{u \in a} \arctan(\text{pers}(u)^\alpha) \phi(u).$$

Similar results would hold for such feature maps, as the map  $u \mapsto \arctan(\text{pers}(u)^\alpha) \phi(u) / \text{pers}(u)^\alpha$  is continuous and bounded whenever  $\phi$  is.




Figure 4: Left: the Čech PD  $\text{dgm}_1(A_n)$  of a sample of  $n = 10^4$  points sampled on a generic torus, with points in Regions (1), (2) and (3) highlighted in different colors. Right: the persistence images of  $\text{dgm}_1(A_n)$  with weight  $\text{pers}^p$  for different values of  $p$ .

## 5 Numerical experiments

We illustrate our results with synthetic experiments. We create a generic submanifold of dimension  $m$  by applying a random diffeomorphism  $\Psi$  to a given  $m$ -dimensional submanifold  $M_0$  (e.g. a torus). We then draw a sample of  $n$  i.i.d. observations sampled according to the pushforward  $P$  of the uniform distribution on  $M$  by  $\Psi$ .




Figure 5: Plot in log-log scale of  $\text{Pers}_p(\text{dgm}_i^{(1)}(A_n))$  as a function of  $n$  for points sampled on a circle,  $i = 0$  (left), points sampled on a torus,  $i = 0$  (center), points sampled on a torus,  $i = 1$  (right). Dashed lines have slopes equal to  $1 - p/m$ .

**Continuity of feature maps.** As a first experiment, we test the continuity of a feature map, the persistence image [AEK<sup>+</sup>17]. In Figure 4, we plot the persistence image of  $\text{dgm}_1(A_n)$  where  $A_n$  is a sample of  $n = 10^4$  points on a generic torus. We observe that the map is discontinuous for  $p < 2$ : the two points with large persistence corresponding to the PD of the underlying torus are nonapparent in the image. For  $p > 2$ , the two points are apparent, and the contribution of points with small persistence (close to the lower edge) has vanished. In the limit situation  $p = 2$ , we see the contribution of both points with large and small persistence. This phenomenon suggests the following heuristics: **when in presence of multiple datasets on  $m$ -dimensional objects whose global geometries need to be distinguished, feature maps with weights  $\text{pers}^p$  with  $p > m$  should be used; when the relevant information is the underlying density of the datasets, the choice  $p < m$  should be preferred.**

**Convergence of total persistences.** We verify the rate of convergence of the total per-

sistence predicted by Theorem 4.2. For values of  $n$  ranging from  $10^2$  to  $10^4$ , we compute  $\text{Pers}_p(\text{dgm}_i^{(1)}(\mathbf{A}_n))$  in three scenarios: points sampled on a circle for  $i = 0$ , and points sampled on a torus for  $i = 0$  and  $i = 1$ . The correct rates of convergence are observed on a log-log plot, see Figure 5. For  $i = 1$ , we remark that the asymptotic regime starts at larger values of  $n$ , above  $n = 10^3$ .

**Convergence of  $\mu_{n,i}$ .** We sample  $n$  points on a torus by uniformly sampling the two angles  $(\theta, \phi)$  parametrizing the torus. We obtain a (nonuniform) probability measure, having density  $f$ . We then compute, for various values of  $n$ , the measure  $\mu_{n,1}$ . The measure is approximated by kernel density estimation (see Figure 6). We approximate in a similar manner the measure  $\mu_{\infty,1,2}$  by sampling  $n = 10^5$  points on a square. We then apply the change of variable formula (21) to compute the theoretical limit  $\mu_{f,1}$ . The distance  $\text{OT}_2(\mu_{n,1}, \mu_{f,1})$  is then computed by approximating the measures on a grid: the distance converges to 0 as predicted by Theorem 4.2. See also Figure 6.




Figure 6: Left: Heatmap of  $\mu_{n,1}$  for  $n = 5 \cdot 10^4$  points sampled on the torus with density  $f$ . Center: Heatmap of  $\mu_{f,1}$ . Right:  $\text{OT}_2$  distance between  $\mu_{n,1}$  and  $\mu_{f,1}$  (normalized by  $\text{OT}_2(0, \mu_{f,i})$ ) for  $n$  ranging from  $10^2$  to  $10^5$ .

## 6 Conclusion

Under the manifold hypothesis, we have greatly refined earlier work regarding the persistent homology of subsamples of compact sets, with especially strong results when the sampling is either random or well-behaved. In particular, we have precisely described the PDs of such samplings, and provided new convergence guarantees w.r.t. the  $p$ -Wassertein distances, as well as detailed asymptotics for their total  $\alpha$ -persistence. The main limitations of our work were the assumptions that the data is sampled from a submanifold, and without any noise. Relaxing those assumptions, as well as establishing similar guarantees for Vietoris-Rips complexes, could be the subject of future research. We also plan on exploring the consequences of our findings regarding the persistent homology dimension [AAF<sup>+</sup>20, SSDE20, BLGS21, Sch21] of submanifolds.

## A Proofs of additional lemmas in Section 3

The goal of this section is to prove the following lemma, used in the proof of Theorem 3.7.

**Lemma A.1.** Let  $A \subset \mathbb{R}^d$  be a finite set, and let  $a < b \in \mathbb{R} \cup \{-\infty, +\infty\}$  and  $i \geq 0$ . Let

$$\bigcup_{j=1}^L C_j = \text{Crit}(A) \cap d_A^{-1}[a, b]$$

be a covering of the set of critical points of  $d_A$  whose critical value belongs to  $[a, b]$ ,<sup>7</sup> and let

$$N_j := \# \left( \bigcup_{z \in C_j} \sigma_A(z) \right)$$

be the cardinality of the union of the projections of the critical points of  $C_j$ . Then the number of points in  $\text{dgm}_i(A)$  such that at least one of their coordinates belongs to  $[a, b]$  is upper-bounded by  $\sum_{j=1}^L \binom{N_j}{i+2}$ , hence by  $\sum_{j=1}^L N_j^{i+2}$ .

We first require an intermediate result. Following [BE17, Definition 4.2], we say that a finite set  $B \subset \mathbb{R}^d$  is in *general position* if for every  $S \subset B$  of at most  $d + 1$  points,  $S$  is affinely independent in  $\mathbb{R}^d$  and no point of  $B \setminus S$  lies on the smallest circumsphere of  $S$ .

**Lemma A.2.** Let  $A \subset \mathbb{R}^d$  be a finite set. For any  $\varepsilon > 0$ , there exists a finite set  $B \subset \mathbb{R}^d$  such that:

- $B$  is in general position,
- the distance function  $d_B$  is topologically Morse,
- there exists a bijection  $\phi : B \rightarrow A$  with  $d(\phi(b), b) \leq \varepsilon$  for all  $b \in B$ , and
- there exists a map  $\psi : \text{Crit}(B) \rightarrow \text{Crit}(A)$  such that  $|d_B(z) - d_A(\psi(z))| \leq \varepsilon$  and  $\phi(\sigma_B(z)) \subset \sigma_A(\psi(z))$  for all  $z \in \text{Crit}(B)$ .

*Proof.* Let  $N$  be the cardinality of  $A = \{p_1, \dots, p_N\}$ . As being in general position is a dense property, we can find for any  $\delta > 0$  a set  $B := \{q_1, \dots, q_N\} \subset \mathbb{R}^d$  such that  $d(q_i, p_i) \leq \delta$  and such that the set  $B$  is in general position in  $\mathbb{R}^d$ . If  $\delta < \varepsilon$  is small enough, this defines a bijection  $\phi : B \rightarrow A, q_i \mapsto p_i$  that satisfies the second condition of the statement. As  $B$  is in general position, the distance function  $d_B$  is then topologically Morse (see [GR97]).

Moreover, Theorem 3.4 from [CCSL06], combined with the lower semicontinuity of the norm of the generalized gradient  $\nabla d_A$  and the compactness of  $A$ , implies that for any  $\rho > 0$ , each critical point of  $B$  must be at distance less than  $\rho$  from one of the critical points of  $A$  as long as  $\delta > 0$  is small enough. This defines a map  $\psi : \text{Crit}(B) \rightarrow \text{Crit}(A)$ . If  $\delta + \rho < \varepsilon$ , then  $|d_B(z) - d_A(\psi(z))| \leq |d_B(z) - d_A(z) + d_A(z) - d_A(\psi(z))| \leq \delta + \rho < \varepsilon$  and the first condition on  $\psi$  is satisfied.

Furthermore, for each of the finitely many critical point  $z' \in \text{Crit}(A)$ , there exists  $\lambda(z') > 0$  such that  $B(z', d_A(z') + \lambda(z')) \cap A = \sigma_A(z')$ . Assume now that  $\delta > 0$  is small enough that  $\delta < \min_{z' \in \text{Crit}(A)} \lambda(z')/4$  and that each critical point  $z$  of  $B$  is at distance less than  $\min_{z' \in \text{Crit}(A)} \lambda(z')/4$  from  $\psi(z) \in \text{Crit}(A)$ . Then such a  $z$  is necessarily such that  $\sigma_B(z) \subset \phi^{-1}(\sigma_A(\psi(z)))$ . Hence  $\psi$  is as required for  $\delta$  small enough, and the lemma is proved.  $\square$

<sup>7</sup>When  $a = -\infty$ , we commit a minor abuse of notation by writing  $[a, b]$  rather than  $(a, b]$ , and similarly when  $b = +\infty$ .

*Proof of Lemma A.1.* Lemma A.2 states that for any  $\delta > 0$ , there exists a point cloud  $\mathbf{B}$  in general position, a bijection  $\phi : \mathbf{B} \rightarrow \mathbf{A}$  with  $d(\phi(b), b) \leq \delta$  for all  $b \in \mathbf{B}$ , and a map  $\psi : \text{Crit}(\mathbf{B}) \rightarrow \text{Crit}(\mathbf{A})$  such that  $|d_{\mathbf{B}}(z) - d_{\mathbf{A}}(\psi(z))| \leq \delta$  and  $\phi(\sigma_{\mathbf{B}}(z)) \subset \sigma_{\mathbf{A}}(\psi(z))$  for all  $z \in \text{Crit}(\mathbf{B})$  and such that  $d_{\mathbf{B}}$  is topologically Morse. Let  $\rho > 0$  be such that all critical values of  $d_{\mathbf{A}}$  in  $[a - \rho, b + \rho]$  belong to  $[a, b]$ , and let  $\mathbf{B}$  be as described for some  $\delta < \rho/2$ . Since the distance function  $d_{\mathbf{B}}$  is topologically Morse, and as explained in the proof of Proposition 3.6, the critical points  $z \in \text{Crit}(\mathbf{B})$  are topological critical points and are in bijection (via their critical values) with the non-zero coordinates of the points in (the union over all degrees of) the Čech persistence diagrams of  $\mathbf{B}$ . In particular, the number of points in  $\text{dgm}_i(\mathbf{B})$  such that at least one of their coordinates belongs to  $[a - \rho/2, b + \rho/2]$  is upper bounded by the number of points  $z \in \text{Crit}(\mathbf{B})$  of topological Morse index  $i$  or  $i + 1$  such that  $d_{\mathbf{B}}(z) \in [a - \rho/2, b + \rho/2]$ . For any such  $z$ , the point  $\psi(z) \in \text{Crit}(\mathbf{A})$  is such that  $d_{\mathbf{A}_n}(\psi(z)) \in [a, b]$ , and as such belongs to  $C_j$  for some  $j \in \{1, \dots, L\}$ . Furthermore,  $\phi(\sigma_{\mathbf{B}}(z)) \subset \sigma_{\mathbf{A}}(\psi(z)) \subset \bigcup_{z' \in C_j} \sigma_{\mathbf{A}}(z')$ . As  $\mathbf{B}$  is in general position, each critical point of  $\mathbf{B}$  of index  $i$  is uniquely identified by its  $i + 1$  projections on  $\mathbf{B}$  (respectively  $i + 2$  projections for points of index  $i + 1$ ), and this bounds the total number of  $z \in \text{Crit}(\mathbf{B})$  such that  $d_{\mathbf{B}}(z) \in [a - \rho/2, b + \rho/2]$  by  $\sum_{j=1}^L \binom{N_j}{i+2}$ .

We have shown that  $\text{dgm}_i(\mathbf{B})$  does not have more than  $\sum_{j=1}^n \binom{N_j}{i+2}$  points with at least one of their coordinates greater  $\in [a - \rho/2, b + \rho/2]$ . By the Bottleneck Stability Theorem, there is a surjection from this set to the set of points in  $\text{dgm}_i(\mathbf{A})$  with at least one of their coordinates in  $[a, b]$  as soon as  $\delta > 0$  is small enough. Hence there are at most  $\sum_{j=1}^n \binom{N_j}{i+2}$  points in  $\text{dgm}_i(\mathbf{A})$  with at least one of their coordinates in  $[a, b]$ , as desired.  $\square$

## B Proofs of additional lemmas in Section 4

In this section, we give additional lemmas needed to prove Theorem 4.2. First, we state two simple lemmas on properties of probability measures and of samples on  $\mathbf{M}$ .

**Lemma B.1.** *Let  $\mathbf{M}$  be a compact submanifold with positive reach. Let  $P$  be a probability measure having a density  $f$  on  $\mathbf{M}$  satisfying  $f_{\min} \leq f \leq f_{\max}$  for two strictly positive constants  $f_{\min}, f_{\max}$ . There exist constants  $c = c(m), C = C(m)$  such that for all  $0 \leq r \leq \tau(\mathbf{M})/4$*

$$cf_{\min}r^m \leq P(\overline{B}(x, r)) \leq Cf_{\max}r^m. \quad (39)$$

*Let  $\mathbf{A}_n$  be a sample of  $n$  i.i.d. observations of law  $P$ . Then, there exists  $C_0 = C_0(\mathbf{M})$  depending on  $\mathbf{M}$  such that for all  $x \in \mathbf{M}$  and all  $r > 0$ ,*

$$\mathbb{P}(d(x, \mathbf{A}_n) \geq r) \leq \exp(-nC_0f_{\min}r^m). \quad (40)$$

*Proof.* For the first statement, see [AL18, Proposition 31]. Let us prove the second one. Remark that the probability is zero for  $r > \text{diam}(\mathbf{M})$ . Hence, we can assume that  $r \leq \text{diam}(\mathbf{M})$ . When  $r \leq \tau(\mathbf{M})/4$ , it holds that

$$\mathbb{P}(d(x, \mathbf{A}_n) \geq r) = (1 - P(\overline{B}(x, r)))^n \leq \exp(-nc_mf_{\min}r^m).$$

When  $\tau(\mathbf{M})/4 \leq r \leq \text{diam}(\mathbf{M})$ , we write

$$\begin{aligned} \mathbb{P}(d(x, \mathbf{A}_n) \geq r) &\leq \mathbb{P}(d(x, \mathbf{A}_n) \geq \tau(\mathbf{M})/4) \leq \exp(-ncf_{\min}(\tau(\mathbf{M})/4)^m) \\ &\leq \exp(-ncf_{\min} \frac{(\tau(\mathbf{M})/4)^m}{\text{diam}(\mathbf{M})^m} r^m). \end{aligned}$$

Hence, the result holds with  $C_0 = c \min\left(1, \frac{(\tau(\mathbf{M})/4)^m}{\text{diam}(\mathbf{M})^m}\right)$ .  $\square$

**Lemma 4.6.** *If  $r \leq \tau(\mathbf{M})/2$ , then*

$$\mathbb{P}(d_H(\mathbf{A}_n, \mathbf{M}) > r) \leq \frac{C}{f_{\min} r^m} \exp(-nc f_{\min} r^m) \quad (24)$$

for two positive constants  $c = c(m), C = C(m)$ . In particular, for any  $q \geq 1$ ,  $d_H(\mathbf{A}_n, \mathbf{M}) = O_{L^q}((\ln n/n)^{1/m})$ .

*Proof.* The bound  $\mathbb{P}(d_H(\mathbf{A}_n, \mathbf{M}) > r)$  is given in [Aam17, Lemma III.23]. Furthermore, [Aam17, Lemma III.23] also states that for any  $q > 0$ , there exists  $C(q)$  depending on  $f_{\min}$  and  $m$  such that, with probability at least  $1 - n^{-q/m}$ ,  $d_H(\mathbf{A}_n, \mathbf{M}) \leq C(q) \left(\frac{\ln n}{n}\right)^{1/m}$ . In particular, we obtain that

$$\mathbb{E}[d_H(\mathbf{A}_n, \mathbf{M})^q] \leq C(q)^q \left(\frac{\ln n}{n}\right)^{q/m} + \text{diam}(\mathbf{M})^q n^{-q/m},$$

proving the second claim of the lemma.  $\square$

Next, we state a simple geometric lemma.

**Lemma B.2.** *Let  $\mathbf{A} \subset \mathbf{M}$ ,  $z \in \text{Crit}(\mathbf{A})$  and  $x \in \sigma_{\mathbf{A}}(z)$ . Then  $\|\pi_x^\perp(z - x)\| \leq \|z - x\|^2/\tau(\mathbf{M})$ . Furthermore, if  $d_{\mathbf{A}}(z) \leq \tau(\mathbf{M})/\sqrt{2}$ , then  $\|\pi_x(z - x)\| \geq \|z - x\|/\sqrt{2}$ .*

*Proof.* The point  $z$  can be written as a convex combination  $z = \sum_k \lambda_k y_k$  where the points  $y_k$  are in  $\mathbf{A} \subset \mathbf{M}$ . Then, using [Fed59, Theorem 4.18],

$$\begin{aligned} \|\pi_x^\perp(z - x)\| &\leq \sum_k \lambda_k \|\pi_x^\perp(y_k - x)\| \leq \sum_k \lambda_k \frac{\|y_k - x\|^2}{2\tau(\mathbf{M})} \\ &= \sum_k \lambda_k \frac{\|y_k - z\|^2 + \|z - x\|^2 + 2\langle y_k - z, z - x \rangle}{2\tau(\mathbf{M})} = \frac{\|z - x\|^2}{\tau(\mathbf{M})}, \end{aligned}$$

as  $\|y_k - z\|^2 = \|x - z\|^2$  and  $\sum_k \lambda_k (y_k - z) = 0$ . The second inequality  $\|\pi_x(z - x)\| \geq \|z - x\|/\sqrt{2}$  from the statement follows from the first one through a direct computation.  $\square$

At last, the following property of the multinomial distribution was used in the proof of Theorem 4.2.

**Lemma B.3.** *Let  $K \geq 1$  and  $L = (L_1, \dots, L_K, L_{K+1})$  be a random multinomial variable of parameters  $n$  and  $\alpha_1, \dots, \alpha_K, \alpha_{K+1}$ . Then there exists  $C = C(K, l)$  such that*

$$\mathbb{E}\left[\left(\sum_{i=1}^K L_i\right)^l \mid L_i \geq 1 \ \forall i = 1, \dots, K\right] \leq C_l \left(1 + \left(n \sum_{i=1}^K \alpha_i\right)^l\right).$$

*Proof.* Let  $X_1, \dots, X_n$  be i.i.d. categorical variables of parameters  $\alpha_1, \dots, \alpha_K, \alpha_{K+1}$ . Define  $L_k(p) = \sum_{r \leq p} \mathbf{1}_{X_r = k}$ ; then  $(L_1(n), \dots, L_K(n), L_{K+1}(n))$  has the same distribution as  $(L_1, \dots, L_K, L_{K+1})$ , and we identify the two in our notation. For a fixed  $n$ , and for any injective function  $\iota : \{1, \dots, k\} \rightarrow \{1, \dots, n\}$ , consider the event

$$E_\iota := \{L_1(\iota(1)) = 1, \dots, L_1(\iota(K)) = 1\},$$

i.e.  $\iota(i)$  is the first appearance of  $i$  among the variables  $X_1, \dots, X_n$ . Note that  $E := \{L_1, \dots, L_K \geq 1\} = \bigsqcup_{\iota} E_{\iota}$  where the sum is taken over all such injective functions. Then

$$\mathbb{E}[(\sum_{i=1}^K L_i)^l | E] = \sum_{\iota} \mathbb{P}(E_{\iota} | E) \mathbb{E}[(\sum_{i=1}^K L_i)^l | E_{\iota}].$$

Fix a function  $\iota$ , and assume without loss of generality that  $\iota(1) < \iota(2) < \dots < \iota(K)$ . Conditioned by  $A_{\iota}$ , the variable  $Y_i := \sum_{r=\iota(i)+1}^{\iota(i+1)-1} \mathbf{1}_{X_r \neq K+1}$  is a binomial variable of parameters  $\iota(i+1) - \iota(i) - 2 \leq n$  and  $\frac{\sum_{i=1}^K \alpha_i}{\alpha_1 + \dots + \alpha_i + \alpha_{K+1}} \leq \sum_{i=1}^K \alpha_i$ . Hence  $\mathbb{E}[Y_i^l | E_{\iota}] \leq C_1(l)(n \sum_{i=1}^K \alpha_i)^l$  using classical bounds on the  $l$ -th moment of a binomial variable, and we see that

$$\begin{aligned} \mathbb{E}[(\sum_{i=1}^K L_i)^l | E_{\iota}] &= \mathbb{E}[(\sum_{r=1}^n \mathbf{1}_{X_r \neq K+1})^l | E_{\iota}] = \mathbb{E}[(K + \sum_{i=0}^K \sum_{r=\iota(i)+1}^{\iota(i+1)-1} \mathbf{1}_{X_r \neq K+1})^l | E_{\iota}] \\ &\leq C_2(K, l)(K^l + \sum_{i=0}^K \mathbb{E}[Y_i^l | E_{\iota}]) \leq C_3(K, l)(1 + (n \sum_{i=1}^K \alpha_i)^l) \end{aligned}$$

where we write  $\iota(0) = 0$  and  $\iota(K+1) = n$  to simplify notation.  $\square$

## C Proofs of Corollary 4.17 and Corollary 4.18

We restate and prove two corollaries from Section 4.

**Corollary 4.17.** *Let  $p \geq 1$  and let  $0 \leq i < d$  be an integer. Under the same assumptions as in Proposition 4.12, the following holds:*

- If  $p > m$ , then  $\mathbb{E}[\text{OT}_p^p(\text{dgm}_i(\mathbf{A}_n), \text{dgm}_i(\mathbf{M}))] \rightarrow 0$  as  $n \rightarrow \infty$ .
- If  $p = m$ ,  $\mathbb{E}[\text{OT}_p^p(\text{dgm}_i(\mathbf{A}_n), \text{dgm}_i(\mathbf{M}))] \rightarrow \text{Pers}_p(\mu_{\infty, i, m}) \text{Vol}(\mathbf{M})$  as  $n \rightarrow \infty$ , where  $\text{Vol}(\mathbf{M})$  is the volume of  $\mathbf{M}$ .
- If  $p < m$  and  $i < m$ , then  $\mathbb{E}[\text{OT}_p^p(\text{dgm}_i(\mathbf{A}_n), \text{dgm}_i(\mathbf{M}))] \rightarrow +\infty$  as  $n \rightarrow \infty$ .

Furthermore, for all  $\alpha > 0$ ,  $\text{Pers}_{\alpha}(\text{dgm}_i(\mathbf{A}_n))$  is equal to

$$\text{Pers}_{\alpha}(\text{dgm}_i(\mathbf{M})) + n^{1-\frac{\alpha}{m}} \text{Pers}_{\alpha}(\mu_{\infty, i, m}) \int_{\mathbf{M}} f(x)^{1-\frac{\alpha}{m}} dx + o_{L^1}(n^{1-\frac{\alpha}{m}}) + O_{L^1}\left(\left(\frac{\log n}{n}\right)^{\frac{1}{m}}\right).$$

*Proof.* Let  $\varepsilon_n = d_H(\mathbf{A}_n, \mathbf{M})$ . Consider the event  $E = \{\varepsilon_n < \tau(\mathbf{M})/2\}$ . According to Theorem 2.3, on  $E$ , the number of points of  $\text{dgm}_i^{(2)}(\mathbf{A}_n)$  is bounded by some constant  $N_0$  depending only on  $\mathbf{M}$ .

- Consider the optimal matching  $\gamma_n$  given by Proposition 4.12. On the event  $E$ , this optimal matching sends all the points of  $\text{dgm}_i^{(1)}(\mathbf{A}_n)$  to the diagonal  $\partial\Omega$ . Thus, we have (when  $E$  is satisfied)

$$\begin{aligned} &|\text{OT}_p^p(\text{dgm}_i(\mathbf{A}_n), \text{dgm}_i(\mathbf{M})) - \text{Pers}_p(\text{dgm}_i^{(1)}(\mathbf{A}_n))| \\ &\leq N_0 \max_{u \in \text{dgm}_i^{(2)}(\mathbf{A}_n)} \|u - \gamma_n(u)\|_{\infty}^p + \#(\text{dgm}_i^{(3)}(\mathbf{A}_n)) \max_{u \in \text{dgm}_i^{(3)}(\mathbf{A}_n)} \|u - \gamma_n(u)\|_{\infty}^p. \end{aligned} \quad (41)$$

Furthermore, note that

$$\begin{aligned} \max_{u \in \text{dgm}_i^{(2)}(\mathbf{A}_n)} \|u - \gamma_n(u)\|_\infty^p &\leq \max \left( \max_{(u_1, u_2) \in \text{dgm}_i^{(2)}(\mathbf{A}_n)} |u_2 - \gamma_n(u)_2|^p, \varepsilon_n^p \right) \\ &\leq \max_{(u_1, u_2) \in \text{dgm}_i^{(2)}(\mathbf{A}_n)} |u_2 - \gamma_n(u)_2|^p + \varepsilon_n^p. \end{aligned}$$

We take the expectation and apply the Cauchy-Schwarz inequality to obtain (together with Theorem 4.2, Proposition 4.12 and Lemma 4.6):

$$\mathbb{E}[\text{OT}_p^p(\text{dgm}_i(\mathbf{A}_n), \text{dgm}_i(\mathbf{M})) \mathbf{1}\{E\}] \leq n^{1-p/m} \text{Pers}_p(\mu_{f,i}) + o(1).$$

When  $E$  is not realized, we crudely bound  $\text{OT}_p^p(\text{dgm}_i(\mathbf{A}_n), \text{dgm}_i(\mathbf{M}))$  by considering the matching sending every point to the diagonal. The cost of this matching is bounded by  $C_M(1 + n^{i+1})$ , where  $n^{i+1}$  is an upper bound on the number of points in  $\text{dgm}_i(\mathbf{A}_n)$  and  $C_M$  is some constant depending on  $\mathbf{M}$ . Hence,

$$\mathbb{E}[\text{OT}_p^p(\text{dgm}_i(\mathbf{A}_n), \text{dgm}_i(\mathbf{M})) \mathbf{1}\{E^c\}] \leq C_M(1 + n^{i+1}) \mathbb{P}(\varepsilon_n > \tau(\mathbf{M})/2) = o(1),$$

according to Lemma 4.6. This proves the first bullet point.

- The second bullet point is proved likewise from (41). Indeed, in that case, it holds that

$$\mathbb{E}[|\text{OT}_p^p(\text{dgm}_i(\mathbf{A}_n), \text{dgm}_i(\mathbf{M})) - \text{Pers}_p(\text{dgm}_i^{(1)}(\mathbf{A}_n))| \mathbf{1}\{E\}] = o(1). \quad (42)$$

Also, using the same crude bound, we obtain that

$$\mathbb{E}[|\text{OT}_p^p(\text{dgm}_i(\mathbf{A}_n), \text{dgm}_i(\mathbf{M})) - \text{Pers}_p(\text{dgm}_i^{(1)}(\mathbf{A}_n))| \mathbf{1}\{E^c\}] = o(1). \quad (43)$$

So far, we have proved that  $\text{OT}_p^p(\text{dgm}_i(\mathbf{A}_n), \text{dgm}_i(\mathbf{M})) = \text{Pers}_p(\text{dgm}_i^{(1)}(\mathbf{A}_n)) + o_{L^1}(1)$ . According to Theorem 4.2, it holds that  $\text{Pers}_p(\text{dgm}_i^{(1)}(\mathbf{A}_n)) = \text{Pers}_p(\mu_{f,i}) + o_{L^1}(1)$  for  $p = m$ , proving the second bullet point.

- For the third bullet point, we use that on the event  $E$ ,

$$\text{OT}_p^p(\text{dgm}_i(\mathbf{A}_n), \text{dgm}_i(\mathbf{M})) \geq \text{Pers}_p(\text{dgm}_i^{(1)}(\mathbf{A}_n)). \quad (44)$$

The latter is equal to  $n^{1-p/m} \text{Pers}_p(\mu_{f,i}) + o_{L^1}(n^{1-p/m})$ , with  $\text{Pers}_p(\mu_{f,i}) > 0$ : indeed, the support of the measure  $\mu_{f,i}$  is nontrivial for  $i < m$ , see [HST18, GTT19]. Hence,

$$\begin{aligned} \mathbb{E}[\text{OT}_p^p(\text{dgm}_i(\mathbf{A}_n), \text{dgm}_i(\mathbf{M}))] &\geq \mathbb{E}[\text{OT}_p^p(\text{dgm}_i(\mathbf{A}_n), \text{dgm}_i(\mathbf{M})) \mathbf{1}\{E\}] \\ &\geq \mathbb{E}[\text{Pers}_p(\text{dgm}_i^{(1)}(\mathbf{A}_n))] - \mathbb{E}[\text{Pers}_p(\text{dgm}_i^{(1)}(\mathbf{A}_n)) \mathbf{1}\{E^c\}]. \end{aligned}$$

The second term goes to zero (use the crude bound  $\text{Pers}_p(\text{dgm}_i^{(1)}(\mathbf{A}_n)) \leq C_M(1 + n^{i+1})$ ), while the first one diverges. This proves the third bullet point.

At last, we prove the formula for the asymptotic expansion of

$$\text{Pers}_\alpha(\text{dgm}_i(\mathbf{A}_n)) = \text{Pers}_\alpha(\text{dgm}_i^{(1)}(\mathbf{A}_n)) + \text{Pers}_\alpha(\text{dgm}_i^{(2)}(\mathbf{A}_n)) + \text{Pers}_\alpha(\text{dgm}_i^{(3)}(\mathbf{A}_n)). \quad (45)$$



The first term is equal to  $n^{1-\alpha/m} \text{Pers}_\alpha(\mu_{\infty,i,m}) \int_{\mathbf{M}} f(x)^{1-\frac{\alpha}{m}} dx + o_{L^1}(n^{1-\alpha/m})$  according to Theorem 4.2. Remark that for  $u, v \in \Omega$ ,  $|\text{pers}_\alpha(u) - \text{pers}_\alpha(v)| \leq 2\alpha(\text{pers}_\alpha(u) + \text{pers}_\alpha(v))\|u - v\|_\infty$ . Hence, on the event  $E = \{\varepsilon_n < \tau(\mathbf{M})/2\}$ ,

$$\begin{aligned} & |\text{Pers}_\alpha(\text{dgm}_i^{(2)}(\mathbf{A}_n)) + \text{Pers}_\alpha(\text{dgm}_i^{(3)}(\mathbf{A}_n)) - \text{Pers}_\alpha(\text{dgm}_i(\mathbf{M}))| \\ & \leq 2\alpha\varepsilon_n(\text{Pers}_\alpha(\text{dgm}_i^{(2)}(\mathbf{A}_n)) + \text{Pers}_\alpha(\text{dgm}_i^{(3)}(\mathbf{A}_n)) + \text{Pers}_\alpha(\text{dgm}_i(\mathbf{M}))). \end{aligned}$$

We crudely bound the persistence of a point in  $\text{dgm}_i(\mathbf{A}_n)$  by  $R(\mathbf{M})$ , the radius of the smallest enclosing ball of  $\mathbf{M}$ , to obtain that

$$\begin{aligned} & |\text{Pers}_\alpha(\text{dgm}_i^{(2)}(\mathbf{A}_n)) + \text{Pers}_\alpha(\text{dgm}_i^{(3)}(\mathbf{A}_n)) - \text{Pers}_\alpha(\text{dgm}_i(\mathbf{M}))| \\ & \leq 2\alpha\varepsilon_n(R(\mathbf{M})^\alpha + 1)(\#(\text{dgm}_i^{(2)}(\mathbf{A}_n)) + \#(\text{dgm}_i^{(3)}(\mathbf{A}_n)) + \text{Pers}_\alpha(\text{dgm}_i(\mathbf{M}))). \end{aligned}$$

We have already established that  $\#(\text{dgm}_i^{(2)}(\mathbf{A}_n)) = O_{L^2}(1)$  (actually, it is larger than  $N_0$  with only exponentially small probability). Furthermore, Proposition 4.12 states that  $\#(\text{dgm}_i^{(3)}(\mathbf{A}_n)) = O_{L^2}(1)$ . Lemma 4.6 also states that  $\varepsilon_n = O_{L^2}(((\log n)/n)^{1/m})$ . Hence,

$$|\text{Pers}_\alpha(\text{dgm}_i^{(2)}(\mathbf{A}_n)) + \text{Pers}_\alpha(\text{dgm}_i^{(3)}(\mathbf{A}_n)) - \text{Pers}_\alpha(\text{dgm}_i(\mathbf{M}))| = O_{L^1}(((\log n)/n)^{1/m}).$$

This concludes the proof.  $\square$

**Corollary 4.18.** *Let  $\alpha \geq 1$  and let  $0 \leq i < d$  be an integer. Under the same assumptions as in Proposition 4.12, it holds that  $\Phi_\alpha(\text{dgm}_i(\mathbf{A}_n))$  converges in probability to  $\Phi_\alpha(\text{dgm}_i(\mathbf{M}))$  whenever  $\alpha > m$ .*

*Proof.* According to [DL21, Proposition 5.1], the feature map  $\Phi_\alpha$  is continuous with respect to the  $\text{OT}_\alpha$  distance. But we have shown in Corollary 4.17 that  $\mathbb{E}[\text{OT}_\alpha^\alpha(\text{dgm}_i(\mathbf{A}_n), \text{dgm}_i(\mathbf{M}))] \rightarrow 0$  as  $n \rightarrow \infty$ . In particular,  $\text{OT}_\alpha(\text{dgm}_i(\mathbf{A}_n), \text{dgm}_i(\mathbf{M}))$  converges in probability to 0. By continuity, so does  $\|\Phi_\alpha(\text{dgm}_i(\mathbf{A}_n)) - \Phi_\alpha(\text{dgm}_i(\mathbf{M}))\|$ .  $\square$

## References

- [AAF<sup>+</sup>20] Henry Adams, Manuchehr Aminian, Elin Farnell, Michael Kirby, Joshua Mirth, Rachel Neville, Chris Peterson, and Clayton Shonkwiler. A fractal dimension for measures via persistent homology. In Topological Data Analysis: The Abel Symposium 2018, pages 1–31. Springer, 2020.
- [Aam17] Eddie Aamari. Rates of convergence for geometric inference. PhD thesis, 2017.
- [ACC<sup>+</sup>22] Andrew Aukerman, Mathieu Carrière, Chao Chen, Kevin Gardner, Raúl Rabadán, and Rami Vanguri. Persistent homology based characterization of the breast cancer immune microenvironment: A feasibility study. Journal of Computational Geometry, 12(2):183–206, 2022.
- [ACSD23] Charles Arnal, David Cohen-Steiner, and Vincent Divol. Critical points of the distance function to a generic submanifold. arXiv preprint arXiv:2312.13147, 2023.

- [AEK<sup>+</sup>17] Henry Adams, Tegan Emerson, Michael Kirby, Rachel Neville, Chris Peterson, Patrick Shipman, Sofya Chepushtanova, Eric Hanson, Francis Motta, and Lori Ziegelmeier. Persistence images: A stable vector representation of persistent homology. Journal of Machine Learning Research, 18(8):1–35, 2017.
- [AKF<sup>+</sup>22] Dominique Attali, Hana Dal Poz Kouřimská, Christopher Fillmore, Ishika Ghosh, André Lieutier, Elizabeth Stephenson, and Mathijs Wintraecken. Tight Bounds for the Learning of Homotopy à la Niyogi, Smale, and Weinberger for Subsets of Euclidean Spaces and of Riemannian Manifolds. <https://arxiv.org/abs/2206.10485>, 2022.
- [AL18] Eddie Aamari and Clément Levrard. Stability and minimax optimality of tangential Delaunay complexes for manifold reconstruction. Discrete & Computational Geometry, 59:923–971, 2018.
- [BA14] Omer Bobrowski and Robert Adler. Distance functions, critical points, and the topology of random Čech complexes. Homology, Homotopy & Applications, 16(2), 2014.
- [BCR<sup>+</sup>22] Bradley CA Brown, Anthony L Caterini, Brendan Leigh Ross, Jesse C Cresswell, and Gabriel Loaiza-Ganem. Verifying the union of manifolds hypothesis for image data. arXiv preprint arXiv:2207.02862, 2022.
- [BE17] Ulrich Bauer and Herbert Edelsbrunner. The Morse theory of Čech and Delaunay complexes. Transactions of the American Mathematical Society, 369(5):pp. 3741–3762, 2017.
- [BH22] Magnus B Botnan and Christian Hirsch. On the consistency and asymptotic normality of multiparameter persistent Betti numbers. Journal of Applied and Computational Topology, pages 1–38, 2022.
- [BHO18] Mickaël Buchet, Yasuaki Hiraoka, and Ippei Obayashi. Persistent homology and materials informatics. Nanoinformatics, pages 75–95, 2018.
- [BK18] Omer Bobrowski and Matthew Kahle. Topology of random geometric complexes: a survey. Journal of applied and Computational Topology, 1:331–364, 2018.
- [BKS17] Omer Bobrowski, Matthew Kahle, and Primoz Skraba. Maximally persistent cycles in random geometric complexes. The Annals of Applied Probability, pages 2032–2060, 2017.
- [BLGS21] Tolga Birdal, Aaron Lou, Leonidas J Guibas, and Umut Simsekli. Intrinsic dimension, persistent homology and generalization in neural networks. Advances in Neural Information Processing Systems, 34:6776–6789, 2021.
- [BW17] Omer Bobrowski and Shmuel Weinberger. On the vanishing of homology in random Čech complexes. Random Structures & Algorithms, 51(1):14–51, 2017.
- [BW20] Peter Bubenik and Alexander Wagner. Embeddings of persistence diagrams into Hilbert spaces. Journal of Applied and Computational Topology, 4(3):339–351, 2020.

- [CCI<sup>+</sup>20] Mathieu Carrière, Frédéric Chazal, Yuichi Ike, Théo Lacombe, Martin Royer, and Yuhei Umeda. Perslay: A neural network layer for persistence diagrams and new graph topological signatures. In International Conference on Artificial Intelligence and Statistics, pages 2786–2796. PMLR, 2020.
- [CCO17] Mathieu Carriere, Marco Cuturi, and Steve Oudot. Sliced Wasserstein kernel for persistence diagrams. In International conference on machine learning, pages 664–673. PMLR, 2017.
- [CCSL06] Frédéric Chazal, David Cohen-Steiner, and André Lieutier. A sampling theory for compact sets in Euclidean space. Discrete & Computational Geometry, 41:461–479, 06 2006.
- [CDSGO16] Frédéric Chazal, Vin De Silva, Marc Glisse, and Steve Oudot. The structure and stability of persistence modules, volume 10. Springer, 2016.
- [CFL<sup>+</sup>14] Frédéric Chazal, Brittany Terese Fasy, Fabrizio Lecci, Alessandro Rinaldo, and Larry Wasserman. Stochastic convergence of persistence landscapes and silhouettes. In Proceedings of the thirtieth annual symposium on Computational geometry, pages 474–483, 2014.
- [CPH21] Luigi Caputi, Anna Pidnebesna, and Jaroslav Hlinka. Promises and pitfalls of topological data analysis for brain connectivity analysis. NeuroImage, 238:118245, 2021.
- [CR20] Mathieu Carrière and Raúl Rabadán. Topological data analysis of single-cell hi-c contact maps. In Topological Data Analysis: The Abel Symposium 2018, pages 147–162. Springer, 2020.
- [CSEH05] David Cohen-Steiner, Herbert Edelsbrunner, and John Harer. Stability of persistence diagrams. In Proceedings of the twenty-first annual symposium on Computational geometry, pages 263–271, 2005.
- [CSEHM10] David Cohen-Steiner, Herbert Edelsbrunner, John Harer, and Yuriy Mileyko. Lipschitz functions have  $L_p$ -stable persistence. Foundations of Computational Mathematics, 10(2):127–139, Apr 2010.
- [CWRW15] Yen-Chi Chen, Daren Wang, Alessandro Rinaldo, and Larry Wasserman. Statistical analysis of persistence intensity functions. arXiv preprint arXiv:1510.02502, 2015.
- [DC19] Vincent Divol and Frédéric Chazal. The density of expected persistence diagrams and its kernel based estimation. Journal of Computational Geometry, 10(2):127–153, 2019.
- [Div21] Vincent Divol. Minimax adaptive estimation in manifold inference. Electronic Journal of Statistics, 15(2):5888–5932, 2021.
- [DKR<sup>+</sup>18] Joshua A Dijkstra, Lenka Kovalcinova, Jie Ren, Robert P Behringer, Miroslav Kramar, Konstantin Mischaikow, and Lou Kondic. Characterizing granular networks using topological metrics. Physical Review E, 97(4):042903, 2018.

- [DL21] Vincent Divol and Théo Lacombe. Understanding the topology and the geometry of the space of persistence diagrams via optimal partial transport. Journal of Applied and Computational Topology, 5:1–53, 2021.
- [DP19] Vincent Divol and Wolfgang Polonik. On the choice of weight functions for linear representations of persistence diagrams. Journal of Applied and Computational Topology, 3(3):249–283, 2019.
- [EH10] Herbert Edelsbrunner and John Harer. Computational Topology: An Introduction. 01 2010.
- [ELZ02] Edelsbrunner, Letscher, and Zomorodian. Topological persistence and simplification. Discrete & computational geometry, 28:511–533, 2002.
- [Fed59] Herbert Federer. Curvature measures. Transactions of the American Mathematical Society, 93(3):418–491, 1959.
- [FLR<sup>+</sup>14] Brittany Terese Fasy, Fabrizio Lecci, Alessandro Rinaldo, Larry Wasserman, Sivaraman Balakrishnan, and Aarti Singh. Confidence sets for persistence diagrams. The Annals of Statistics, 42(6):2301, 2014.
- [GPPVW12] Christopher R Genovese, Marco Perone-Pacifico, Isabella Verdinelli, and Larry Wasserman. Manifold estimation and singular deconvolution under Hausdorff loss. The Annals of Statistics, 40(2):941–963, 2012.
- [GR97] Vladimir Gershkovich and Hyam Rubinstein. Morse theory for min-type functions. The Asian Journal of Mathematics, 1, 01 1997.
- [Gro93] Karsten Grove. Critical point theory for distance functions. Proceedings of Symposia in Pure Mathematics, pages 357–385, 1993.
- [GTT19] Akshay Goel, Khanh Duy Trinh, and Kenkichi Tsunoda. Strong law of large numbers for Betti numbers in the thermodynamic regime. Journal of Statistical Physics, 174(4):865–892, 2019.
- [Hat00] Allen Hatcher. Algebraic topology. Cambridge Univ. Press, Cambridge, 2000.
- [HKN19] Christoph D Hofer, Roland Kwitt, and Marc Niethammer. Learning representations of persistence barcodes. Journal of Machine Learning Research, 20(126):1–45, 2019.
- [HL23] Olympio Hacquard and Vadim Lebovici. Euler characteristic tools for topological data analysis. arXiv preprint arXiv:2303.14040, 2023.
- [HST18] Yasuaki Hiraoka, Tomoyuki Shirai, and Khanh Duy Trinh. Limit theorems for persistence diagrams. The Annals of Applied Probability, 28(5):2740–2780, 2018.
- [KFH18] Genki Kusano, Kenji Fukumizu, and Yasuaki Hiraoka. Kernel method for persistence diagrams via kernel embedding and weight factor. Journal of Machine Learning Research, 18(189):1–41, 2018.

- [KHF16] Genki Kusano, Yasuaki Hiraoka, and Kenji Fukumizu. Persistence weighted gaussian kernel for topological data analysis. In International conference on machine learning, pages 2004–2013. PMLR, 2016.
- [KKP<sup>+</sup>16] L Kondic, M Kramár, Luis A Pugnali, C Manuel Carlevaro, and K Mischaikow. Structure of force networks in tapped particulate systems of disks and pentagons. ii. persistence analysis. Physical Review E, 93(6):062903, 2016.
- [KKZ<sup>+</sup>20] Kwangho Kim, Jisu Kim, Manzil Zaheer, Joon Kim, Frederic Chazal, and Larry Wasserman. Pllay: Efficient topological layer based on persistent landscapes. In H. Larochelle, M. Ranzato, R. Hadsell, M.F. Balcan, and H. Lin, editors, Advances in Neural Information Processing Systems, volume 33, pages 15965–15977. Curran Associates, Inc., 2020.
- [KM13] Matthew Kahle and Elizabeth Meckes. Limit theorems for Betti numbers of random simplicial complexes. Homology, Homotopy and Applications, 15(1):343–374, 2013.
- [Mil63] J.W. Milnor. Morse Theory. Annals of mathematics studies. Princeton University Press, 1963.
- [Mor59] Marston Morse. Topologically non-degenerate functions on a compact n-manifold m. Journal d’Analyse Mathématique, 7(1):189–208, 1959.
- [MV21] Atish Mitra and Žiga Virk. The space of persistence diagrams on  $n$  points coarsely embeds into Hilbert space. Proceedings of the American Mathematical Society, 149(6):2693–2703, 2021.
- [Owa22] Takashi Owada. Convergence of persistence diagram in the sparse regime. The Annals of Applied Probability, 32(6):4706–4736, 2022.
- [RHBK15] Jan Reininghaus, Stefan Huber, Ulrich Bauer, and Roland Kwitt. A stable multi-scale kernel for topological machine learning. In Proceedings of the IEEE conference on computer vision and pattern recognition, pages 4741–4748, 2015.
- [RYB<sup>+</sup>20] Bastian Rieck, Tristan Yates, Christian Bock, Karsten Borgwardt, Guy Wolf, Nicholas Turk-Browne, and Smita Krishnaswamy. Uncovering the topology of time-varying fMRI data using cubical persistence. Advances in neural information processing systems, 33:6900–6912, 2020.
- [Sch21] Benjamin Schweinhart. Persistent homology and the upper box dimension. Discrete & Computational Geometry, 65(2):331–364, 2021.
- [SL22] Yara Skaf and Reinhard Laubenbacher. Topological data analysis in biomedicine: A review. Journal of Biomedical Informatics, 130:104082, 2022.
- [SPCGB19] Ann E Sizemore, Jennifer E Phillips-Cremins, Robert Ghrist, and Danielle S Bassett. The importance of the whole: topological data analysis for the network neuroscientist. Network Neuroscience, 3(3):656–673, 2019.

- [SSDE20] Umut Simsekli, Ozan Sener, George Deligiannidis, and Murat A Erdogdu. Hausdorff dimension, heavy tails, and generalization in neural networks. Advances in Neural Information Processing Systems, 33:5138–5151, 2020.
- [STR<sup>+</sup>17] Mohammad Saadatfar, Hiroshi Takeuchi, Vanessa Robins, Nicolas Francois, and Yasuaki Hiraoka. Pore configuration landscape of granular crystallization. Nature communications, 8(1):15082, 2017.
- [STR<sup>+</sup>18] Anirudh Som, Kowshik Thopalli, Karthikeyan Natesan Ramamurthy, Vinay Venkataraman, Ankita Shukla, and Pavan Turaga. Perturbation robust representations of topological persistence diagrams. In Proceedings of the European Conference on Computer Vision (ECCV), pages 617–635, 2018.
- [SYM23] Anna Song, Ka Man Yim, and Anthea Monod. Generalized morse theory of distance functions to surfaces for persistent homology. arXiv preprint arXiv:2306.14716, 2023.
- [VDVW97] Aad W Van Der Vaart and Jon A Wellner. Weak convergence and empirical processes: with applications to statistics. Springer New York, 1997.
- [Ver18] Roman Vershynin. High-dimensional probability: An introduction with applications in data science, volume 47. Cambridge university press, 2018.
- [Wag21] Alexander Wagner. Nonembeddability of persistence diagrams with  $p > 2$  Wasserstein metric. Proceedings of the American Mathematical Society, 149(6):2673–2677, 2021.
- [WB19] Jonathan Weed and Francis Bach. Sharp asymptotic and finite-sample rates of convergence of empirical measures in Wasserstein distance. Bernoulli, 25(4A):2620–2648, 2019.