

Polyvinyltriazole-functionalized SWCNT ink for printed Arsenic (III) chemistors for drink water monitoring applications

Leshan Usgodaarachchi, Balakumara Vignesh Muppidathi, Benoit Piro, Senthilmurugan Subbiah, Eric Cloutet, Bérengère Lebental


▶ To cite this version:

Leshan Usgodaarachchi, Balakumara Vignesh Muppidathi, Benoit Piro, Senthilmurugan Subbiah, Eric Cloutet, et al.. Polyvinyltriazole-functionalized SWCNT ink for printed Arsenic (III) chemistors for drink water monitoring applications. 5th Workshop of the IEEE Sensors France Chapter, Grenoble, Jun 2024, Grenoble, France. hal-04618688

HAL Id: hal-04618688 https://hal.science/hal-04618688

Submitted on 21 Jun2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Polyvinyltriazole (PVT)-functionalized SWCNT ink for printed Arsenic (III) chemistors for drink water monitoring applications


Leshan USGODAARACHCHI ^{1,2}, Balakumara VIGNESH M ^{1,3}, Benoit PIRO ², Senthilmurugan SUBBIAH ³, Eric CLOUTET ⁴, Bérengère LEBENTAL ^{1,*}

¹ IMSE (Laboratoire Instrumentation, Modelisation, Simulation Et Experimentation) - Université Gustave Eiffel ² ITODYS (Interfaces – Traitement – Organisation et Dynamique des Systèmes) – Université Paris Cité ³ Water and Energy Nexus lab, Department of Chemical Engineering, Indian Institute of Technology Guwahati 4 University of Bordeaux, CNRS, Bordeaux INP, LCPO, UMR 5629, F-33600 Pessac, France


Health consequences

1. Arsenic in water and Toxicology

Arsenic: top twentieth abundant element on Earth).(WHO drinking water quality guidelines: maximum 10ppb).(




2. PVT functionalized ink and chemoresitive transduction



- The Triazole group (C=N-N)).(present in the PVT molecule has an affinity for the As (III) ions

3.1. Chemoresistive device (chemistors) fabrication




Chemistors operate on the principle of chemoresistive sensing, where changes in electrical resistance are measured

PVT functionalized ink deposited on




interdigitated electrodes by ink dispersion printing (IDP) using the KELEN DMD 100


demonstration




The SEM morphology of the fabricated).(chemistors showed the well oriented SWCNT network on the surface of the interdigitated electrode



3. 2. Assembling to the prototype sensor and excitation





).(



2 SWCNT chemistors


chemistors

- 2 PVT-118 functionalized).(chemistors
- 2 Polyfluorene based).(polymer functionalized







Square wave used for the excitation with a duty cycle of 50 1.28 µA used for the excitation





3. 3. Pristine SWCNT chemistors (refernce) response


3. 5. Future perspectives













ur l'analyse d'un fluide comprenant une portion de conduite (EU Patent, 2023) Dispositif pour l'analyse d'un fluide comprenant une tête de sonde (EU Patent, 2023)

- The adsorption capacity of multilayer PVT for SWCNT was determined using adsorption isotherm analysis
-).(Experimental data are well established for the Toth isotherm model and the maximal surface coverage (Q_m) is 3684.54 mg/g and r² is **0.96**
- The LOTUS printed circuit bord).((LPCB) will be used for the future characterization experiments
-)<u>×</u>(It consists of 12 chemistors, 2 temperature sensors and 2 conductivity sensors
- Based on the Q_m calculations different stoichiometries will be further investigated to obtain the best sensitivity
- A pH and chlorine sensor will be).(integrated into the LPCB flatform



- After 20ppb SWCNT sensor response saturated at).(400kΩ
- R_m = measured resistance (Ω)
- 1ppb 20ppb From to sensor response is recognized as linear
- Up to 20ppb, **0.33% / ppb** and and r² is **0.98**).($R_B = Baseline \ resistance \ (\Omega)$

3. 4. PVT functionalized SWCNT chemistor response variation



- From 1 ppb to 65 ppb of As (III) addition, the sensor shows a linear response)....
- After 65ppb, the sensor does not shows a linear response).(
- Linear regression slope is **0.77%** /ppb and r² is **0.98**

4. Conclusion

- **PVT f-SWCNT** chemistor offers a **MLOD of 5ppb** in tap water with the presence of the active),(chlorine (ClO⁻) alone with the minor impurities of other ions (Na⁺, Ca²⁺, Mg²⁺, etc.)
- **PVT f-SWCNT** chemistor showed a dynamic range of 5 to 50 ppb, sensitivity of **0.77% /ppb**).(with reversibility
- SWCNT chemistor used as a reference sensor to compare the performance improvements due).(to functionalization, exhibited a dynamic range of 1 to 20 ppb with a sensitivity of 0.33% /ppb

5. Acknowledgement	6. Bibilography
The authors acknowledge support from the European Union for HS4U project under Grant Agreement N°101069937 and for the project LOTUS under Grant Agreement N° 820881. LOTUS is also financed by the Indian Government, Ministry of Science and Technology	J. H. T. Luong, E. Lam, and K. B. Male, 'Recent advances in electrochemical detection of arsenic in drinking and ground waters', Anal. Methods, vol. 6, no. 16, pp. 6157–6169, Jul. 2014, doi: <u>10.1039/C4AY00817K</u> . G. ZUCCHI et al., 'Chemical sensors based on carbon nanotubes functionalised by conjugated polymers for analysis in aqueous medium', US11697707B2, Jul. 11, 2023 Accessed: Sep. 06, 2023. [Online]. Available: <u>https://patents.google.com/patent/US11697707B2/en</u>