



HAL
open science

Uniform estimates for transmission problems in electromagnetism with high contrast in magnetic permeabilities

Victor Péron

► **To cite this version:**

Victor Péron. Uniform estimates for transmission problems in electromagnetism with high contrast in magnetic permeabilities. 2024. hal-04620526

HAL Id: hal-04620526

<https://hal.science/hal-04620526>

Preprint submitted on 21 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

UNIFORM ESTIMATES FOR TRANSMISSION PROBLEMS IN ELECTROMAGNETISM WITH HIGH CONTRAST IN MAGNETIC PERMEABILITIES

VICTOR PÉRON

ABSTRACT. We consider the time-harmonic Maxwell equations set on a domain made up of two subdomains that represent a magnetic conductor and a non-magnetic material, and we assume that the relative magnetic permeability μ_r between the two materials is high. We prove uniform a priori estimates for Maxwell solutions when the interface between the two subdomains is supposed to be Lipschitz. Assuming smoothness for the interface between the subdomains, we prove also that the magnetic field possesses a multiscale expansion in powers of $\varepsilon = 1/\sqrt{\mu_r}$ with *profiles* rapidly decaying inside the magnetic conductor.

Keywords: *Maxwell equations, High relative permeability, Lipschitz domains, Asymptotic expansions*

1. INTRODUCTION

The numerical computation of the electromagnetic field in strongly heterogeneous media is crucial in many subject areas in real-world applications, see, *e.g.*, [14, 26, 15, 23, 9, 25, 29, 18, 27, 3, 19, 4, 1] where additional references may also be found. The computation can be made particularly difficult due in particular to very high contrasts of magnetic permeabilities, see, *e.g.*, [14, 23, 25, 29, 1]. In several fields of applied geophysics, for instance, the computation of the controlled source electromagnetic induction (CSEM) response of conductive and permeable bodies buried in a conductive and permeable medium is widely used, but the presence of buried objects, such as steel pipes, concrete rebar, cables, drums, unexploded ordnance, or storage tanks introduce significant complexity into the CSEM response, especially since these objects often present significant relative magnetic permeability, see, *e.g.*, [14, 23, 25]. High contrasts in magnetic permeabilities can also pose a significant challenge for developing homogenization techniques to deduce effective properties of heterogeneous materials. These techniques are particularly necessary for the design of optimal electromagnetic devices based on soft magnetic composites in many electrical engineering applications, see, *e.g.*, [29].

Many works in the literature are related to issues of uniform estimates for stiff transmission problems in electromagnetism. We refer the reader, for instance, on scattering problems of an electromagnetic plane wave by a medium of high index (*i.e.*, with high complex permeability, and high complex permittivity) and of class \mathcal{C}^2 (smooth), [6, 7].

In particular, the authors provided a rigorous proof of Leontovich conditions and proved accurate error estimates in Ref. [7], and they tackled the scattering problem by a thin layer of high index in Ref. [6]. We also refer the reader to the work in Ref. [17] in which the authors provide uniform a priori estimates for static Maxwell interface problems, and to the works in Ref. [15, 10, 9] for transmission problems with high contrast of electric conductivities.

In this work we investigate a transmission problem in materials presenting high contrast of magnetic permeabilities. The domain of interest is made up of two subdomains that represent a magnetic conductor and a non-magnetic material, and the relative magnetic permeability μ_r between the materials can be very high. Our interest is the solvability of the time-harmonic Maxwell equations together with uniform energy estimates in \mathbf{L}^2 for the electromagnetic field when μ_r tends to infinity. This issue of uniform estimates is not obvious, and this is especially true since we do not assume that the interface between the subdomains is smooth.

The main result of this paper is uniform a priori estimates for Maxwell solutions when the interface between the two subdomains is supposed to be Lipschitz, cf. Theorem 2.1 in Sec. 2. This new result shows in particular that the magnetic field tends to 0 in energy norm inside the magnetic conductor as μ_r tends to infinity. Our result is valid for a right-hand side in \mathbf{L}^2 . This is notably a difference with the uniform \mathbf{L}^2 estimates at high conductivity [10, Theorem 2.1], which is valid for data in \mathbf{L}^2 with divergence in \mathbf{L}^2 .

The proof of our main result is based on an appropriate decomposition of the magnetic field into a vector field in the Sobolev space \mathbf{H}^1 and a gradient field which is piecewise $\mathbf{H}^{\frac{1}{2}}$ (does not belong to the energy space), and the gradient part is estimated thanks to uniform estimates for scalar transmission problems with constant coefficients on two subdomains. Note that the uniform estimates for these scalar transmission problems (cf. Lemma 3.4 in Sec. 3.1) are a consequence of a more general result [10, Theorem 2.1], and also that the decomposition of the magnetic field based on a vector potential technique is rather classical [5]. However, the proof of the main result on uniform estimates for the electromagnetic field (Theorem 2.1) is new. In particular, the proof of uniform \mathbf{L}^2 estimates for the electric field at high conductivity [10, Lemma 4.1] does not seem to be easily adaptable to our problem because we consider that the relative permeability can be unbounded.

As an application of our uniform estimates, we develop also an argument for the convergence of an asymptotic expansion. Assuming *smoothness for the interface* between the subdomains, we prove rigorously that the magnetic field possesses a multiscale expansion in powers of $\varepsilon = \frac{1}{\sqrt{\mu_r}}$ with *profile* terms rapidly decaying inside the magnetic conductor, cf. Sec. 4.

We also refer the reader to several works which are devoted to asymptotic expansions of the electromagnetic field at high relative magnetic permeability ([28, 2, 1]) or at high conductivity (see, e.g., [30, 21, 22, 15, 10, 13, 9]) in a domain made up of two subdomains when the interface is smooth: see, e.g., [30, 21, 22, 28, 2, 1] for plane interface

and eddy current approximations, and [15, 10, 13, 9] for a three-dimensional model of *skin effect* in non-magnetic materials.

This paper is organized as follows. In Section 2 we introduce the framework and we state the main result. In Section 3 we prove uniform estimates for Maxwell solutions when the relative magnetic permeability is high. In Section 4 we apply these uniform estimates for proving the convergence of an asymptotic expansion of the magnetic field at high relative magnetic permeability, and we provide the first terms of this expansion. We give also elements of proof for this expansion in Appendix A.

2. FRAMEWORK AND MAIN RESULT

Let Ω be a smooth bounded simply connected domain in \mathbb{R}^3 , and $\Omega_- \subset\subset \Omega$ be a Lipschitz connected subdomain of Ω . We denote by Γ the boundary of Ω , and by Σ the boundary of Ω_- . Finally, we denote by Ω_+ the complementary of $\overline{\Omega_-}$ in Ω , cf. Figure 1.

We consider the time-harmonic Maxwell equations given by Faraday's and Ampère's laws in Ω :

$$(2.1) \quad \operatorname{curl} \mathbf{E} - i\omega \underline{\mu} \mathbf{H} = 0 \quad \text{and} \quad \operatorname{curl} \mathbf{H} + (i\omega \varepsilon_0 - \underline{\sigma}) \mathbf{E} = \mathbf{j} \quad \text{in } \Omega.$$

Here, (\mathbf{E}, \mathbf{H}) represents the electromagnetic field, ε_0 is the electric permittivity, ω is the angular frequency ($\omega \neq 0$ and the time convention is $\exp(-i\omega t)$), \mathbf{j} represents a current density and is supposed to belong to $\mathbf{L}^2(\Omega)$, $\underline{\sigma}$ is the electric conductivity, and $\underline{\mu}$ is the magnetic permeability. We assume that the domain Ω is made of two connected subdomains Ω_+ and Ω_- in which the coefficients $\underline{\sigma}$ and $\underline{\mu}$ take different values ($\sigma_+ > 0, \sigma_- > 0$) and ($\mu_+ > 0, \mu_- > 0$), respectively, and we denote by μ_r the relative magnetic permeability between the subdomains Ω_- and Ω_+ , and which is defined as:

$$(2.2) \quad \mu_r = \mu_- / \mu_+.$$

In this work we are particularly interested in the case where this parameter can be high,




FIGURE 1. The domain Ω and its subdomains Ω_- and Ω_+

which is often the case when the domain Ω_- represents a linear magnetic conductive material and the domain Ω_+ represents a non-magnetic material.

To complement the Maxwell harmonic equations (2.1), we consider either the perfectly insulating electric boundary conditions

$$(2.3) \quad \mathbf{E} \cdot \mathbf{n} = 0 \quad \text{and} \quad \mathbf{H} \times \mathbf{n} = 0 \quad \text{on} \quad \Gamma,$$

or the perfectly conducting electric boundary conditions

$$(2.4) \quad \mathbf{E} \times \mathbf{n} = 0 \quad \text{and} \quad \mathbf{H} \cdot \mathbf{n} = 0 \quad \text{on} \quad \Gamma.$$

2.1. Main result: uniform estimates in Lipschitz domains. Hereafter, we denote by $\|\cdot\|_{0,\mathcal{O}}$ the norm in $\mathbf{L}^2(\mathcal{O}) = \mathbf{L}^2(\mathcal{O})^3$. In the framework above it is possible to prove uniform a priori estimates for the Maxwell transmission problem:

Theorem 2.1. *Let $\sigma_{\pm} > 0$, and $\mu_+ > 0$. If the interface Σ is Lipschitz, there are constants μ_* and $C > 0$ such that for all $\mu_r \geq \mu_*$ the Maxwell problem (2.1) with boundary conditions (2.3) and data $\mathbf{j} \in \mathbf{L}^2(\Omega)$ has a unique solution (\mathbf{E}, \mathbf{H}) in $\mathbf{L}^2(\Omega)^2$, which satisfies:*

$$(2.5) \quad \|\mathbf{H}\|_{0,\Omega} + \|\mathbf{E}\|_{0,\Omega} + \sqrt{\mu_r} \|\mathbf{H}\|_{0,\Omega_-} \leq C \|\mathbf{j}\|_{0,\Omega}.$$

A similar result holds for boundary conditions (2.4).

This result is proved in Section 3. The proof is based on an appropriate decomposition of the magnetic field into a regular field in $\mathbf{H}^1(\Omega)$ and a gradient field. The gradient part is estimated thanks to piecewise $H^{3/2}$ uniform estimates for a scalar transmission problem with constant coefficients on two subdomains (cf. Lemma 3.4 in Section 3.1).

In that follows, we introduce for convenience the following parameter in the Maxwell equations (2.1)

$$(2.6) \quad \varepsilon = \frac{1}{\sqrt{\mu_r}}.$$

Hence, ε tends to 0 as μ_r tends to infinity. Then for $\mu_r \geq \mu_*$, where the constant μ_* is given by Theorem 2.1, we denote by $(\mathbf{E}_{(\varepsilon)}, \mathbf{H}_{(\varepsilon)})$ the solution of the system (2.1) – (2.3). As an application of uniform estimates (2.5) we will develop an argument for the convergence of an asymptotic expansion of $(\mathbf{E}_{(\varepsilon)}, \mathbf{H}_{(\varepsilon)})$ as $\varepsilon \rightarrow 0$, see Section 4.2.

2.2. Magnetic formulation. We deduce from the Maxwell system (2.1)-(2.3) the following variational formulation for the magnetic field $\mathbf{H}_{(\varepsilon)}$. The variational space is the Hilbert space $\mathbf{H}_0(\text{curl}, \Omega)$:

$$(2.7) \quad \mathbf{H}_0(\text{curl}, \Omega) = \{\mathbf{u} \in \mathbf{L}^2(\Omega) \mid \text{curl} \mathbf{u} \in \mathbf{L}^2(\Omega), \mathbf{u} \times \mathbf{n} = 0 \text{ on } \Gamma\},$$

endowed with its graph norm, and the variational problem writes

Find $\mathbf{H}_{(\varepsilon)} \in \mathbf{H}_0(\text{curl}, \Omega)$ such that for all $\mathbf{K} \in \mathbf{H}_0(\text{curl}, \Omega)$

$$(2.8) \quad \int_{\Omega} \left(\left(1 + i \frac{\sigma}{\omega \varepsilon_0} \right)^{-1} \text{curl} \mathbf{H}_{(\varepsilon)} \cdot \text{curl} \bar{\mathbf{K}} - \kappa_+^2 \mu_r(\varepsilon) \mathbf{H}_{(\varepsilon)} \cdot \bar{\mathbf{K}} \right) dx = \int_{\Omega} \left(1 + i \frac{\sigma}{\omega \varepsilon_0} \right)^{-1} \mathbf{j} \cdot \text{curl} \bar{\mathbf{K}} dx .$$

Here we have set

$$(2.9) \quad \mu_r(\varepsilon) = \mathbf{1}_{\Omega_+} + \frac{1}{\varepsilon_2} \mathbf{1}_{\Omega_-} , \quad \text{and} \quad \kappa_+ := \omega \sqrt{\varepsilon_0 \mu_+} .$$

3. PROOF OF UNIFORM ESTIMATES FOR MAXWELL SOLUTIONS AT HIGH RELATIVE MAGNETIC PERMEABILITY

We consider the harmonic Maxwell system (2.1)-(2.3). We are going to prove the following statement:

Lemma 3.1. *Let $\sigma_{\pm} > 0$, and $\mu_{\pm} > 0$. There are constants μ_{\star} and $C_0 > 0$ such that if $\mu_r \geq \mu_{\star}$ any solution $(\mathbf{E}, \mathbf{H}) \in \mathbf{L}^2(\Omega)^2$ of problem (2.1) with boundary condition (2.3) and data $\mathbf{j} \in \mathbf{L}^2(\Omega)$ satisfies the estimate*

$$(3.1) \quad \|\mathbf{H}\|_{0,\Omega} \leq C_0 \|\mathbf{j}\|_{0,\Omega} .$$

A similar statement holds for boundary conditions (2.4).

This lemma is the key for the proof of Theorem 2.1 and is going to be proved in the next subsection. As a consequence of this lemma, we will obtain estimates (2.5):

Corollary 3.2. *Let $\sigma_{\pm} > 0$, and $\mu_{\pm} > 0$. Let $(\mathbf{E}, \mathbf{H}) \in \mathbf{L}^2(\Omega)^2$ be solution of problem (2.1) with boundary condition (2.3) and data $\mathbf{j} \in \mathbf{L}^2(\Omega)$. Then we have $\text{div}(\mu \mathbf{H}) = 0$ in Ω , and if \mathbf{H} satisfies estimate (3.1), then setting $m = \min(\sqrt{\omega^2 \varepsilon_0^2 + \sigma_-^2}, \sqrt{\omega^2 \varepsilon_0^2 + \sigma_+^2})$,*

$$C_1 = \frac{\max\left(\frac{\omega^2 \varepsilon_0^2 + \sigma_-^2}{\sigma_-}, \frac{\omega^2 \varepsilon_0^2 + \sigma_+^2}{\sigma_+}\right)}{m}, \quad \text{and} \quad C_2 = \sqrt{\frac{\varepsilon_0 C_1^2}{m^2} + \frac{C_1}{\omega m}},$$

there holds

$$(3.2) \quad \|\mathbf{H}\|_{0,\Omega} + \|\text{curl} \mathbf{H}\|_{0,\Omega} + \sqrt{\mu_-} \|\mathbf{H}\|_{0,\Omega_-} \leq (C_0 + C_1 + C_2) \|\mathbf{j}\|_{0,\Omega} .$$

A similar estimate holds for boundary conditions (2.4).

This result is going to be proved in Section 3.3. Finally, estimate (3.1) implies existence and uniqueness of solutions.

Corollary 3.3. *Let $\sigma_{\pm} > 0$, and $\mu_{\pm} > 0$. We assume that estimate (3.1) holds for any solution $(\mathbf{E}, \mathbf{H}) \in \mathbf{L}^2(\Omega)^2$ of problem (2.1)-(2.3) with $\mathbf{j} \in \mathbf{L}^2(\Omega)$. Then for any $\mathbf{j} \in \mathbf{L}^2(\Omega)$, there exists a unique solution $(\mathbf{E}, \mathbf{H}) \in \mathbf{L}^2(\Omega)^2$ of problem (2.1)-(2.3). A similar result holds for boundary conditions (2.4).*

This result is going to be proved in Section 3.4. Finally we deduce Theorem 2.1 from the previous three statements.

3.1. Preliminary lemma. In this section we provide a uniform piecewise estimate for the solution of a scalar transmission problem with constant coefficients on two subdomains. We will use this result in the proof of Lemma 3.1 (cf. Section 3.2) for obtaining a uniform L^2 estimate for the magnetic field.

For any given function $\underline{a} = (a_+, a_-)$ determined by the two constants a_{\pm} on Ω_{\pm} , we consider the following variational problem : Find $\varphi \in H_0^1(\Omega)$ such that

$$(3.3) \quad \forall \psi \in H_0^1(\Omega), \quad \int_{\Omega_+} a_+ \nabla \varphi^+ \cdot \nabla \bar{\psi}^+ \, d\mathbf{x} + \int_{\Omega_-} a_- \nabla \varphi^- \cdot \nabla \bar{\psi}^- \, d\mathbf{x} = (a_- - a_+) \int_{\Sigma} g \bar{\psi} \, ds,$$

where the right-hand side g satisfies the regularity assumption

$$(3.4) \quad g \in L^2(\Sigma)$$

and the extra compatibility condition

$$(3.5) \quad \int_{\Sigma} g \, ds = 0.$$

In the framework of Section 2 (we recall that the surface Σ is Lipschitz), we have the following uniform piecewise estimate for the solution of the problem (3.3).

Lemma 3.4. *Let us assume that $a_+ \neq 0$. There exist a constant $\rho_0 > 0$ independent of a_+ such that for all $a_- \in \{z \in \mathbb{C} \mid |z| \geq \rho_0 |a_+|\}$, the problem (3.3) with a right-hand side g satisfying (3.4)-(3.5) has a unique solution $\varphi \in H_0^1(\Omega)$ which moreover is piecewise $H^{3/2}$ and satisfies the uniform estimate*

$$(3.6) \quad \|\varphi^+\|_{\frac{3}{2}, \Omega_+} + \|\varphi^-\|_{\frac{3}{2}, \Omega_-} \leq C_{\rho_0} \|g\|_{0, \Sigma},$$

with a constant $C_{\rho_0} > 0$, independent of a_+ , a_- , and g .

This result is obtained as a consequence of [10, Theorem 2.1].

Remark 3.5. The assumption on the interface Σ , that is a Lipschitz surface, is necessary in Lemma 3.4. Indeed, there exists non-Lipschitz surfaces such that estimate (3.6) does not hold, for instance in the case of a checkerboard configuration in a two dimensional domain, cf. [12, Theorem 8.1] and [10, Rem 2.3].

3.2. Proof of Lemma 3.1: Uniform L^2 estimate of the magnetic field. Reductio ad absurdum: We assume that there is a sequence $(\mathbf{E}_m, \mathbf{H}_m) \in L^2(\Omega)^2$, $m \in \mathbb{N}$, of solutions of the Maxwell system (2.1)-(2.3) associated with a magnetic permeability $\underline{\mu}_m = (\mu_+, \mu_m, \mu_+)$ and a right hand side $\mathbf{j}_m \in L^2(\Omega)$:

$$(3.7a) \quad \operatorname{curl} \mathbf{E}_m - i\omega \underline{\mu}_m \mathbf{H}_m = 0 \quad \text{in } \Omega,$$

$$(3.7b) \quad \operatorname{curl} \mathbf{H}_m + (i\omega \varepsilon_0 - \underline{\sigma}) \mathbf{E}_m = \mathbf{j}_m \quad \text{in } \Omega,$$

$$(3.7c) \quad \mathbf{E}_m \cdot \mathbf{n} = 0 \quad \text{and} \quad \mathbf{H}_m \times \mathbf{n} = 0 \quad \text{on } \partial\Omega,$$

satisfying the following conditions

$$(3.8a) \quad \mu_m \rightarrow \infty \quad \text{as } m \rightarrow \infty,$$

$$(3.8b) \quad \|\mathbf{H}_m\|_{0,\Omega} = 1 \quad \forall m \in \mathbb{N},$$

$$(3.8c) \quad \|\mathbf{j}_m\|_{0,\Omega} \rightarrow 0 \quad \text{as } m \rightarrow \infty.$$

First, we particularize the magnetic variational formulation (2.8) for the sequence $\{\mathbf{H}_m\}$: For all $\mathbf{H}' \in \mathbf{H}_0(\text{curl}, \Omega)$:

$$(3.9) \quad \int_{\Omega} \left(\frac{1}{i\omega\varepsilon_0 - \underline{\sigma}} \text{curl } \mathbf{H}_m \cdot \text{curl } \overline{\mathbf{H}'} + i\omega\underline{\mu}_m \mathbf{H}_m \cdot \overline{\mathbf{H}'} \right) d\mathbf{x} = \int_{\Omega} \frac{\mathbf{j}_m}{i\omega\varepsilon_0 - \underline{\sigma}} \cdot \text{curl } \overline{\mathbf{H}'} d\mathbf{x}.$$

Choosing $\mathbf{H}' = \mathbf{H}_m$ in (3.9) and taking the real part, and then using the Cauchy-Schwarz inequality, we obtain the following uniform bound on the curl of the magnetic field in Ω :

$$(3.10) \quad \|\text{curl } \mathbf{H}_m\|_{0,\Omega} \leq \frac{\max\left(\frac{\omega^2\varepsilon_0^2 + \sigma_-^2}{\sigma_-}, \frac{\omega^2\varepsilon_0^2 + \sigma_+^2}{\sigma_+}\right)}{\min(\sqrt{\omega^2\varepsilon_0^2 + \sigma_-^2}, \sqrt{\omega^2\varepsilon_0^2 + \sigma_+^2})} \|\mathbf{j}_m\|_{0,\Omega}.$$

3.2.1. *Decomposition of the magnetic field and bound in $H^{\frac{1}{2}}$.* We recall that we have assumed that the domain Ω is simply connected and has a smooth connected boundary. Relying to Theorem 2.12 and Theorem 3.17 in [5], for all $m \in \mathbb{N}$ there exists a unique $\mathbf{v}_m \in \mathbf{H}^1(\Omega)$ such that

$$(3.11) \quad \text{curl } \mathbf{v}_m = \text{curl } \mathbf{H}_m, \quad \text{div } \mathbf{v}_m = 0 \text{ in } \Omega, \quad \text{and } \mathbf{v}_m \times \mathbf{n} = 0 \text{ on } \partial\Omega.$$

Moreover, we have the estimate

$$(3.12) \quad \|\mathbf{v}_m\|_{1,\Omega} \leq C \|\text{curl } \mathbf{H}_m\|_{0,\Omega},$$

where C is independent of m . As a consequence of the equality $\text{curl } \mathbf{v}_m = \text{curl } \mathbf{H}_m$ and the simple connectedness of Ω , we obtain that there exists $\varphi_m \in H_0^1(\Omega)$ such that

$$(3.13) \quad \mathbf{H}_m = \mathbf{v}_m + \nabla\varphi_m.$$

Then using (3.13) we write equation (3.7a) as

$$\text{curl } \mathbf{E}_m - i\omega\underline{\mu}_m (\mathbf{v}_m + \nabla\varphi_m) = 0 \quad \text{in } \Omega.$$

Let $\psi \in H_0^1(\Omega)$ be a test function. Multiplying the above equality by $\nabla\overline{\psi}$ and integrating over Ω , we obtain, using that $\text{div } \mathbf{v}_m = 0$:

$$(3.14) \quad \int_{\Omega} \underline{\mu}_m \nabla\varphi_m \cdot \nabla\overline{\psi} d\mathbf{x} = (\mu_m - 1)\mu_+ \int_{\Sigma} \mathbf{v}_m \cdot \mathbf{n} \Big|_{\Sigma} \overline{\psi} ds,$$

where \mathbf{n} denotes the unit normal vector on Σ , inwardly oriented to Ω_- . Note that the boundary value $\psi = 0$ on $\partial\Omega$ has been used in (3.14).

Thus φ_m is solution of the Dirichlet problem defined by the variational equation (3.14). Since

$$\mathbf{v}_m \cdot \mathbf{n} \in L^2(\Sigma) \quad \text{and} \quad \int_{\Sigma} \mathbf{v}_m \cdot \mathbf{n} \Big|_{\Sigma} ds = \int_{\Omega_-} \text{div } \mathbf{v}_m d\mathbf{x} = 0,$$

the Dirichlet problem defined by (3.14) satisfies the assumptions of Lemma 3.4 (cf. Section 3.1) with $a_- = \mu_m \mu_+$ and $a_+ = \mu_+$, and $g = \mathbf{v}_m \cdot \mathbf{n}|_{\Sigma}$. Therefore we obtain the following uniform estimate for μ_m large enough (i.e. for m large enough, cf. (3.8a))

$$\|\varphi_m^+\|_{\frac{3}{2}, \Omega_+} + \|\varphi_m^-\|_{\frac{3}{2}, \Omega_-} \leq C_0 \|\mathbf{v}_m \cdot \mathbf{n}\|_{0, \Sigma}.$$

Since $\|\mathbf{v}_m \cdot \mathbf{n}\|_{0, \Sigma}$ is bounded by $\|\mathbf{v}_m\|_{1, \Omega}$, the last inequality implies

$$(3.15) \quad \|\varphi_m^+\|_{\frac{3}{2}, \Omega_+} + \|\varphi_m^-\|_{\frac{3}{2}, \Omega_-} \leq C_0 \|\mathbf{v}_m\|_{1, \Omega}.$$

Finally (3.10), (3.12), and (3.15) implies that

$$(3.16) \quad \|\varphi_m^+\|_{\frac{3}{2}, \Omega_+} + \|\varphi_m^-\|_{\frac{3}{2}, \Omega_-} + \|\mathbf{v}_m\|_{1, \Omega} \leq B$$

for a constant $B > 0$ independent of m . With (3.13), (3.16) gives that the sequence $\{\mathbf{H}_m\}$ is bounded in $H^{\frac{1}{2}}$ on Ω_- and Ω_+ :

$$\|\mathbf{H}_m^+\|_{\frac{1}{2}, \Omega_+} + \|\mathbf{H}_m^-\|_{\frac{1}{2}, \Omega_-} \leq B.$$

Combining the above bound with (3.10), we obtain the uniform bound

$$(3.17) \quad \|\mathbf{H}_m^+\|_{\frac{1}{2}, \Omega_+} + \|\mathbf{H}_m^-\|_{\frac{1}{2}, \Omega_-} + \|\operatorname{curl} \mathbf{H}_m\|_{0, \Omega} \leq C.$$

3.2.2. Limit of the sequence and conclusion. The domains Ω_{\pm} being bounded, the embedding of $H^{\frac{1}{2}}(\Omega_{\pm})$ in $L^2(\Omega_{\pm})$ is compact. Hence as a consequence of (3.17), we can extract a subsequence of $\{\mathbf{H}_m\}$ (still denoted by $\{\mathbf{H}_m\}$) which is converging in $L^2(\Omega)$. By the Banach-Alaoglu theorem, we can assume that the sequence $\{\operatorname{curl} \mathbf{H}_m\}$ is weakly converging in $L^2(\Omega)$: We deduce that there is $\mathbf{H} \in L^2(\Omega)$ such that

$$(3.18) \quad \begin{cases} \operatorname{curl} \mathbf{H}_m \rightharpoonup \operatorname{curl} \mathbf{H} & \text{in } L^2(\Omega) \\ \mathbf{H}_m \rightarrow \mathbf{H} & \text{in } L^2(\Omega). \end{cases}$$

A consequence of the strong convergence of $\{\mathbf{H}_m\}$ in $L^2(\Omega)$ and (3.8b) is that

$$(3.19) \quad \|\mathbf{H}\|_{0, \Omega} = 1.$$

We are going to prove that

$$(3.20) \quad \mathbf{H} = 0 \quad \text{in } \Omega,$$

which will contradict (3.19), and finally prove estimate (3.1).

Choosing $\mathbf{H}' = \mathbf{H}_m$ in the magnetic variational formulation (3.9) for the sequence $\{\mathbf{H}_m\}$, then taking the imaginary part and using the Cauchy-Schwarz inequality, we obtain

$$(3.21) \quad \omega \mu_m \mu_+ \|\mathbf{H}_m^-\|_{0, \Omega_-}^2 + \omega \mu_+ \|\mathbf{H}_m^+\|_{0, \Omega_+}^2 \leq \frac{\omega \varepsilon_0}{\omega^2 \varepsilon_0^2 + \sigma_-^2} \|\operatorname{curl} \mathbf{H}_m^-\|_{0, \Omega_-}^2 + \frac{\omega \varepsilon_0}{\omega^2 \varepsilon_0^2 + \sigma_+^2} \|\operatorname{curl} \mathbf{H}_m^+\|_{0, \Omega_+}^2 + \left\| \frac{\mathbf{j}_m}{i\omega \varepsilon_0 - \sigma} \right\|_{0, \Omega} \|\operatorname{curl} \mathbf{H}_m\|_{0, \Omega}.$$

Hence, according to (3.10)-(3.8c) we obtain that the right-hand side in (3.21) tends to zero and we infer the convergence results

$$(3.22) \quad \sqrt{\mu_m} \|\mathbf{H}_m^-\|_{0, \Omega_-} \rightarrow 0 \quad \text{as } m \rightarrow \infty, \quad \text{and} \quad \|\mathbf{H}_m^+\|_{0, \Omega_+} \rightarrow 0 \quad \text{as } m \rightarrow \infty.$$

Finally, according to (3.8a)-(3.18), we deduce that $\mathbf{H}^- := \mathbf{H}|_{\Omega_-}$ and $\mathbf{H}^+ := \mathbf{H}|_{\Omega_+}$ satisfy

$$(3.23) \quad \mathbf{H}^- = 0 \quad \text{in } \Omega_-, \quad \text{and} \quad \mathbf{H}^+ = 0 \quad \text{in } \Omega_+,$$

which proves (3.20), contradicts (3.19) and finally ends the proof of Lemma 3.1.

3.3. Proof of Corollary 3.2. Let $(\mathbf{E}, \mathbf{H}) \in \mathbf{L}^2(\Omega)^2$ be a solution of the Maxwell problem (2.1) with boundary condition (2.3) and data $\mathbf{j} \in \mathbf{L}^2(\Omega)$, and we assume that

$$(3.24) \quad \|\mathbf{H}\|_{0,\Omega} \leq C_0 \|\mathbf{j}\|_{0,\Omega}.$$

First, taking the divergence of equation $\text{curl } \mathbf{E} - i\omega \underline{\mu} \mathbf{H} = 0$ in Ω , we immediately obtain

$$(3.25) \quad \text{div}(\underline{\mu} \mathbf{H}) = 0 \quad \text{in } \Omega.$$

Second, $\mathbf{H} \in \mathbf{H}_0(\text{curl}, \Omega)$ is solution of the variational problem (2.8). Taking as test function \mathbf{H} itself, we obtain the identity

$$(3.26) \quad \int_{\Omega} \left(\frac{1}{i\omega \varepsilon_0 - \underline{\sigma}} \text{curl } \mathbf{H} \cdot \text{curl } \overline{\mathbf{H}} + i\omega \underline{\mu} \mathbf{H} \cdot \overline{\mathbf{H}} \right) dx = \int_{\Omega} \frac{\mathbf{j}}{i\omega \varepsilon_0 - \underline{\sigma}} \cdot \text{curl } \overline{\mathbf{H}} dx.$$

Taking the real part of (3.26), we obtain

$$\frac{\sigma_-}{\omega^2 \varepsilon_0^2 + \sigma_-^2} \|\text{curl } \mathbf{H}\|_{0,\Omega_-}^2 + \frac{\sigma_+}{\omega^2 \varepsilon_0^2 + \sigma_+^2} \|\text{curl } \mathbf{H}\|_{0,\Omega_+}^2 = \text{Re} \left(\frac{\mathbf{j}}{\underline{\sigma} - i\omega \varepsilon_0}, \text{curl } \mathbf{H} \right)_{0,\Omega}$$

hence, using Cauchy-Schwarz inequality, and setting $C_1 = \frac{\max\left(\frac{\omega^2 \varepsilon_0^2 + \sigma_-^2}{\sigma_-}, \frac{\omega^2 \varepsilon_0^2 + \sigma_+^2}{\sigma_+}\right)}{\min(\sqrt{\omega^2 \varepsilon_0^2 + \sigma_-^2}, \sqrt{\omega^2 \varepsilon_0^2 + \sigma_+^2})}$, we obtain

$$(3.27) \quad \|\text{curl } \mathbf{H}\|_{0,\Omega} \leq C_1 \|\mathbf{j}\|_{0,\Omega}.$$

Then, taking the imaginary part of (3.26), since $\omega \neq 0$, we obtain

$$\begin{aligned} \mu_- \|\mathbf{H}\|_{0,\Omega_-}^2 + \mu_+ \|\mathbf{H}\|_{0,\Omega_+}^2 &= \frac{\varepsilon_0}{\omega^2 \varepsilon_0^2 + \sigma_-^2} \|\text{curl } \mathbf{H}\|_{0,\Omega_-}^2 + \frac{\varepsilon_0}{\omega^2 \varepsilon_0^2 + \sigma_+^2} \|\text{curl } \mathbf{H}\|_{0,\Omega_+}^2 \\ &\quad + \frac{1}{\omega} \text{Im} \left(\frac{\mathbf{j}}{i\omega \varepsilon_0 - \underline{\sigma}}, \text{curl } \mathbf{H} \right)_{0,\Omega}, \end{aligned}$$

hence, using Cauchy-Schwarz inequality and inequality (3.27), we infer

$$(3.28) \quad \sqrt{\mu_-} \|\mathbf{H}\|_{0,\Omega_-} \leq \sqrt{\frac{\varepsilon_0 C_1^2}{\min(\omega^2 \varepsilon_0^2 + \sigma_-^2, \omega^2 \varepsilon_0^2 + \sigma_+^2)} + \frac{C_1}{\omega \min(\sqrt{\omega^2 \varepsilon_0^2 + \sigma_-^2}, \sqrt{\omega^2 \varepsilon_0^2 + \sigma_+^2})}} \|\mathbf{j}\|_{0,\Omega}.$$

Formulae (3.24)-(3.27)-(3.28) yield Corollary 3.2.

3.4. Proof of Corollary 3.3. Let μ_{\pm} , σ_{\pm} and ω (i.e., κ_{+}) be fixed. Let us define $\boldsymbol{\mu}_r$ the piecewise constant function on Ω as (we recall $\mu_r = \mu_-/\mu_+$):

$$(3.29) \quad \boldsymbol{\mu}_r = \mathbf{1}_{\Omega_+} + \mu_r \mathbf{1}_{\Omega_-}.$$

Then, according to (2.6)-(2.9) the sesquilinear form in the left hand side of (2.8) writes

$$(3.30) \quad \int_{\Omega} \left(\left(1 + i \frac{\sigma}{\omega \varepsilon_0}\right)^{-1} \operatorname{curl} \mathbf{H} \cdot \operatorname{curl} \overline{\mathbf{H}'} - \kappa_+^2 \boldsymbol{\mu}_r \mathbf{H} \cdot \overline{\mathbf{H}'} \right) \mathrm{d}\mathbf{x}.$$

The proof of Corollary 3.3 relies on a regularization procedure: We consider the functional space

$$\mathbf{X}_{\mathbf{N}}(\Omega, \boldsymbol{\mu}_r) = \{\mathbf{H} \in \mathbf{H}_0(\operatorname{curl}, \Omega) \mid \operatorname{div}(\boldsymbol{\mu}_r \mathbf{H}) \in L^2(\Omega)\}.$$

Let $s > 0$ be a real number which will be fixed later. Let us introduce the sesquilinear forms A_s and $B: \mathbf{X}_{\mathbf{N}}(\Omega, \boldsymbol{\mu}_r) \times \mathbf{X}_{\mathbf{N}}(\Omega, \boldsymbol{\mu}_r) \rightarrow \mathbb{C}$

(3.31a)

$$A_s(\mathbf{H}, \mathbf{H}') = \int_{\Omega} \left(\left(1 + i \frac{\sigma}{\omega \varepsilon_0}\right)^{-1} \operatorname{curl} \mathbf{H} \cdot \operatorname{curl} \overline{\mathbf{H}'} + s \operatorname{div} \boldsymbol{\mu}_r \mathbf{H} \operatorname{div} \overline{\boldsymbol{\mu}_r \mathbf{H}'} \right) \mathrm{d}\mathbf{x}$$

(3.31b)

$$B(\mathbf{H}, \mathbf{H}') = \int_{\Omega} \boldsymbol{\mu}_r \mathbf{H} \cdot \overline{\mathbf{H}'} \mathrm{d}\mathbf{x}.$$

The regularized variational formulation is: Find $\mathbf{H} \in \mathbf{X}_{\mathbf{N}}(\Omega, \boldsymbol{\mu}_r)$ such that

(3.32)

$$\forall \mathbf{H}' \in \mathbf{X}_{\mathbf{N}}(\Omega, \boldsymbol{\mu}_r), \quad A_s(\mathbf{H}, \mathbf{H}') - \kappa_+^2 B(\mathbf{H}, \mathbf{H}') = \int_{\Omega} \left(1 + i \frac{\sigma}{\omega \varepsilon_0}\right)^{-1} \mathbf{j} \cdot \operatorname{curl} \overline{\mathbf{H}'} \mathrm{d}\mathbf{x}$$

As a consequence of [11, Th. 7.1], we obtain that if

$$(3.33) \quad \frac{\kappa_+^2}{s} \text{ is not an eigenvalue of the Dirichlet problem for the operator } -\operatorname{div} \boldsymbol{\mu}_r \nabla,$$

then any solution $(\mathbf{E}, \mathbf{H}) \in \mathbf{L}^2(\Omega)^2$ of problem (2.1)-(2.3) with $\mathbf{j} \in \mathbf{L}^2(\Omega)$ provides a solution of problem (3.32), and conversely, any solution \mathbf{H} of (3.32) provides a solution of (2.1)-(2.3) by setting $\mathbf{E} = (i\omega\varepsilon_0 - \sigma)^{-1}(\mathbf{j} - \operatorname{curl} \mathbf{H})$.

Thus, we fix s such that (3.33) holds.

Since the form A_s is coercive on $\mathbf{X}_{\mathbf{N}}(\Omega, \boldsymbol{\mu}_r)$ and the embedding of $\mathbf{X}_{\mathbf{N}}(\Omega, \boldsymbol{\mu}_r)$ in $\mathbf{L}^2(\Omega)$ is compact, we obtain that the Fredholm alternative is valid: If the kernel of the adjoint problem to (3.32)

$$(3.34) \quad \text{Find } \mathbf{H}' \in \mathbf{X}_{\mathbf{N}}(\Omega, \boldsymbol{\mu}_r), \quad \forall \mathbf{H} \in \mathbf{X}_{\mathbf{N}}(\Omega, \boldsymbol{\mu}_r), \quad A_s(\mathbf{H}, \mathbf{H}') - \kappa_+^2 B(\mathbf{H}, \mathbf{H}') = 0,$$

is reduced to $\{0\}$, then problem (3.32) is solvable.

We see that the assumption of Corollary 3.3 implies that (3.34) has only the zero solution, and that the same holds for the direct problem

$$(3.35) \quad \text{Find } \mathbf{H} \in \mathbf{X}_{\mathbf{N}}(\Omega, \boldsymbol{\mu}_r), \quad \forall \mathbf{H}' \in \mathbf{X}_{\mathbf{N}}(\Omega, \boldsymbol{\mu}_r), \quad A_s(\mathbf{H}, \mathbf{H}') - \kappa_+^2 B(\mathbf{H}, \mathbf{H}') = 0.$$

All this implies the unique solvability of problem (2.1)-(2.4) with $\mathbf{j} \in \mathbf{L}^2(\Omega)$.

4. MULTISCALE EXPANSION OF THE MAGNETIC FIELD IN SMOOTH DOMAINS

This section is concerned with a rigorous multiscale expansion for the magnetic field at high relative magnetic permeability. In Section 4.1 we provide the first terms of this expansion. Then in Section 4.2 we prove error estimates for this expansion.

For the sake of completeness we give elements of proof for the multiscale expansion in Appendix A. We expand the Maxwell operator in power series of ε , cf. §A.1. Then we deduce the equations satisfied by the coefficients of the magnetic field, cf. §A.2. We derive explicitly the first terms of this expansion in §A.3.

In that follows we assume that right-hand side \mathbf{j} in (2.1) is smooth and that Assumption 4.1 holds:

Assumption 4.1. *We assume that the surfaces Σ (interface) and Γ (external boundary) are smooth.*

Let $\sigma_{\pm} > 0$, $\mu_{+} > 0$, and recall $\varepsilon = \frac{1}{\sqrt{\mu_r}} > 0$ (cf. (2.6)). By Theorem 2.1 there exists $\varepsilon_{*} > 0$ such that for all $\varepsilon \in (0, \varepsilon_{*})$, the Maxwell problem (2.1) with boundary conditions (2.3) has a unique solution $(\mathbf{E}_{\varepsilon}, \mathbf{H}_{\varepsilon})$. In that follows, the magnetic field $\mathbf{H}_{(\varepsilon)}$ is denoted by $\mathbf{H}_{(\varepsilon)}^{+}$ in the non-magnetic part Ω_{+} , and by $\mathbf{H}_{(\varepsilon)}^{-}$ in the magnetic conducting part Ω_{-} . Then both parts possess series expansions in powers of ε :

$$(4.1) \quad \mathbf{H}_{(\varepsilon)}^{+}(\mathbf{x}) \approx \sum_{j \geq 0} \varepsilon^j \mathbf{H}_j^{+}(\mathbf{x}), \quad \mathbf{x} \in \Omega_{+},$$

$$(4.2) \quad \mathbf{H}_{(\varepsilon)}^{-}(\mathbf{x}) \approx \sum_{j \geq 0} \varepsilon^j \mathbf{H}_j^{-}(\mathbf{x}; \varepsilon), \quad \mathbf{x} \in \Omega_{-}, \quad \text{with} \quad \mathbf{H}_j^{-}(\mathbf{x}; \varepsilon) = \chi(y_3) \underline{\mathfrak{H}}_j(y_{\alpha}, \frac{y_3}{\varepsilon}).$$


In (4.2), (y_{α}, y_3) is a *normal coordinate system* to the surface Σ which is defined in \mathcal{U}_{-} , a tubular neighborhood of the surface Σ in the domain Ω_{-} (cf. Figure 2): y_{α} ($\alpha = 1, 2$) are tangential coordinates on Σ , and y_3 is the normal coordinate to Σ , cf. e.g., [9]. The function $\mathbf{y} \mapsto \chi(y_3)$ is a smooth cut-off with support in $\bar{\mathcal{U}}_{-}$ and equal to 1 in a smaller tubular neighborhood of Σ . The vector fields $\underline{\mathfrak{H}}_j : (y_{\alpha}, Y_3) \mapsto \underline{\mathfrak{H}}_j(y_{\alpha}, Y_3)$ are *profiles* defined on $\Sigma \times \mathbb{R}^{+}$: They are exponentially decreasing with respect to Y_3 and are smooth in all variables.

4.1. First terms of the multiscale expansion. In this section, we provide the construction of the first profiles $\underline{\mathfrak{H}}_j = (\mathfrak{H}_j, \mathfrak{h}_j)$ and of the first terms \mathbf{H}_j^{+} . The first profile $\underline{\mathfrak{H}}_0$ in the magnetic conductor is zero:

$$(4.3) \quad \underline{\mathfrak{H}}_0 = 0.$$

Then, the first term of the magnetic field in the non-magnetic region solves Maxwell equations with perfectly insulating electric boundary conditions on $\partial\Omega_{+} = \Sigma \cup \Gamma$ (we recall $\kappa_{+} = \omega \sqrt{\varepsilon_0 \mu_{+}}$, cf. (2.9)):

$$\begin{cases} \operatorname{curl} \operatorname{curl} \mathbf{H}_0^{+} - \kappa_{+}^2 (1 + \frac{i}{\delta_{+}^2}) \mathbf{H}_0^{+} = \operatorname{curl} \mathbf{j} & \text{in } \Omega_{+} \\ \mathbf{H}_0^{+} \times \mathbf{n} = 0 & \text{on } \Sigma \cup \Gamma. \end{cases}$$

FIGURE 2. A tubular neighbourhood of the surface Σ

The first profile in the magnetic region is a tangential field which is exponential with a complex rate λ :

$$(4.4) \quad \mathfrak{H}_1(y_\alpha, Y_3) = -j_0(y_\alpha) e^{-\lambda Y_3}.$$

Here $j_0(y_\alpha) = \lambda^{-1}(1 + \frac{i}{\delta_-^2})(1 + \frac{i}{\delta_+^2})^{-1} (\text{curl } \mathbf{H}_0^+ \times \mathbf{n})(y_\alpha, 0)$, and λ is given by:

$$(4.5) \quad \lambda = \kappa_+ \sqrt[4]{1 + \frac{1}{\delta_-^4}} e^{i \frac{\theta(\delta_-) - \pi}{2}} \quad \text{with} \quad \delta_- = \sqrt{\omega \varepsilon_0 / \sigma_-}, \quad \text{and} \quad \theta(\delta_-) = \arctan \frac{1}{\delta_-^2}.$$

Note that $\text{Re } \lambda > 0$, and if j_0 is not identically 0, there exists a constant $C > 0$ independent of ε such that

$$(4.6) \quad C^{-1} \sqrt{\varepsilon} \leq \| \mathbf{H}_1^-(\cdot; \varepsilon) \|_{0, \Omega_-} \leq C \sqrt{\varepsilon}.$$

Note also that the normal component of the profile \mathfrak{H}_1 is zero:

$$\mathfrak{h}_1 = 0.$$

The next term in the non-magnetic region solves:

$$\begin{cases} \text{curl curl } \mathbf{H}_1^+ - \kappa_+^2 (1 + \frac{i}{\delta_+^2}) \mathbf{H}_1^+ = 0 & \text{in } \Omega_+ \\ \mathbf{H}_1^+ \times \mathbf{n} = -j_0 \times \mathbf{n} & \text{on } \Sigma \\ \mathbf{H}_1^+ \times \mathbf{n} = 0 & \text{on } \Gamma. \end{cases}$$

Like above, define j_1 as $j_1(y_\alpha) = \lambda^{-1}(1 + \frac{i}{\delta_-^2})(1 + \frac{i}{\delta_+^2})^{-1} (\text{curl } \mathbf{H}_1^+ \times \mathbf{n})(y_\alpha, 0)$ on the interface Σ . Then, the tangential components of the profile \mathfrak{H}_2 are given by the tangential field \mathfrak{H}_2 :

$$(4.7) \quad \mathfrak{H}_2(y_\alpha, Y_3) = \left[-j_1 + (\lambda^{-1} + Y_3) (C - \mathcal{H}) j_0 \right] (y_\alpha) e^{-\lambda Y_3}.$$

Here $\mathcal{H} = \frac{1}{2} b_\alpha^\alpha$ is the *mean curvature* of the surface Σ^1 , and \mathcal{C} is the curvature tensor field on Σ defined by

$$(4.8) \quad (\mathcal{C}j)_\alpha = b_\alpha^\beta j_\beta,$$

with $b_\alpha^\beta = a^{\beta\gamma} b_{\gamma\alpha}$, and $a^{\beta\gamma}$ is the inverse of the metric tensor $a_{\beta\gamma}$ in Σ , and $b_{\gamma\alpha}$ is the curvature tensor in Σ . The next term which is determined is the normal component \mathfrak{h}_2 of the profile $\underline{\mathfrak{H}}_2$:

$$\mathfrak{h}_2(y_\alpha, Y_3) = -\lambda^{-1} \operatorname{div}_\Sigma j_0(y_\alpha) e^{-\lambda Y_3}.$$

Here $\operatorname{div}_\Sigma$ is the surface divergence operator on Σ . The next term in the non-magnetic region solves :

$$\begin{cases} \operatorname{curl} \operatorname{curl} \mathbf{H}_2^+ - \kappa_+^2 (1 + \frac{i}{\delta_+^2}) \mathbf{H}_2^+ = 0 & \text{in } \Omega_+ \\ \mathbf{n} \times \mathbf{H}_2^+ \times \mathbf{n} = -\mathbf{j}_1 + \lambda^{-1} (\mathcal{C} - \mathcal{H}) j_0 & \text{on } \Sigma \\ \mathbf{H}_2^+ \times \mathbf{n} = 0 & \text{on } \Gamma. \end{cases}$$

4.2. Validation of the multiscale expansion. The validation of the multiscale expansion (4.1)-(4.2) for the magnetic field $\mathbf{H}_{(\varepsilon)}$ consist in proving estimates for remainders $\mathbf{R}_{m;\varepsilon}$ which are defined as

$$(4.9) \quad \mathbf{R}_{m;\varepsilon} = \mathbf{H}_{(\varepsilon)} - \sum_{j=0}^m \varepsilon^j \mathbf{H}_j \quad \text{in } \Omega.$$

This is done by an evaluation of the right hand side when the Maxwell operator is applied to $\mathbf{R}_{m;\varepsilon}$. By construction (cf. Sec. A.2, Appendix A), we obtain

$$(4.10) \quad \begin{cases} \operatorname{curl} \alpha_+^{-1} \operatorname{curl} \mathbf{R}_{m;\varepsilon}^+ - \kappa_+^2 \mathbf{R}_{m;\varepsilon}^+ = 0 & \text{in } \Omega_+ \\ \operatorname{curl} \alpha_-^{-1} \operatorname{curl} \mathbf{R}_{m;\varepsilon}^- - \varepsilon^{-2} \kappa_+^2 \mathbf{R}_{m;\varepsilon}^- = \mathbf{j}_{m;\varepsilon}^- & \text{in } \Omega_- \\ [\mathbf{R}_{m;\varepsilon} \times \mathbf{n}]_\Sigma = 0 & \text{on } \Sigma \\ [\underline{\alpha}^{-1} \operatorname{curl} \mathbf{R}_{m;\varepsilon} \times \mathbf{n}]_\Sigma = \mathbf{g}_{m;\varepsilon} & \text{on } \Sigma \\ \mathbf{R}_{m;\varepsilon}^+ \times \mathbf{n} = 0 & \text{on } \partial\Omega. \end{cases}$$

Here, $\underline{\alpha} = (\alpha_+, \alpha_-)$, $\alpha_+ = 1 + i/\delta_+^2$ and $\alpha_- = 1 + i/\delta_-^2$, and $[\mathbf{H} \times \mathbf{n}]_\Sigma$ denotes the jump of $\mathbf{H} \times \mathbf{n}$ across Σ . The right hand sides (residues) $\mathbf{j}_{m;\varepsilon}^-$ and $\mathbf{g}_{m;\varepsilon}$ are, roughly, of the order ε^m : we have $\mathbf{j}_{0;\varepsilon}^- = 0$ and $\mathbf{g}_{0;\varepsilon} = \mathcal{O}(1)$; for all $m \in \mathbb{N} \setminus \{0\}$, we have $\mathbf{j}_{m;\varepsilon}^- = \mathcal{O}(\varepsilon^{m-1})$ and $\mathbf{g}_{m;\varepsilon} = \mathcal{O}(\varepsilon^m)$, and we have the following estimates

$$(4.11) \quad \|\mathbf{j}_{m;\varepsilon}^-\|_{0,\Omega_-} \leq C_m \varepsilon^{m-1} \quad \text{and} \quad \|\mathbf{g}_{m;\varepsilon}\|_{\frac{1}{2},\Sigma} \leq C_m \varepsilon^m,$$

where $C_m > 0$ is independent of ε . The main result of this section is the following.

Theorem 4.2. *In the framework above, for all $m \in \mathbb{N}$ and $\varepsilon \in (0, \varepsilon_0]$, the remainder $\mathbf{R}_{m;\varepsilon}$ (4.9) satisfies the optimal estimate*

$$(4.12) \quad \|\mathbf{R}_{m;\varepsilon}^+\|_{0,\Omega_+} + \|\operatorname{curl} \mathbf{R}_{m;\varepsilon}^+\|_{0,\Omega_+} + \varepsilon^{-\frac{1}{2}} \|\mathbf{R}_{m;\varepsilon}^-\|_{0,\Omega_-} + \varepsilon^{\frac{1}{2}} \|\operatorname{curl} \mathbf{R}_{m;\varepsilon}^-\|_{0,\Omega_-} \leq C_m \varepsilon^{m+1}.$$

¹In particular, the sign of \mathcal{H} depends on the orientation of the surface Σ . As a convention, the unit normal vector \mathbf{n} on the surface Σ is inwardly oriented to Ω_- , see Figure 2.

Proof. STEP 1. Since $\mathbf{g}_{m;\varepsilon} \neq 0$, according to (4.10) the vector field $\underline{\alpha}^{-1} \operatorname{curl} \mathbf{R}_{m;\varepsilon}$ does not define an element of the Hilbert space $\mathbf{H}(\operatorname{curl}, \Omega)$:

$$\mathbf{H}(\operatorname{curl}, \Omega) = \{ \mathbf{u} \in \mathbf{L}^2(\Omega) \mid \operatorname{curl} \mathbf{u} \in \mathbf{L}^2(\Omega) \} .$$

Hence we can not apply Theorem 2.1 directly because the vector field $\operatorname{curl} \underline{\alpha}^{-1} \operatorname{curl} \mathbf{R}_{m;\varepsilon} - \kappa_+^2 \boldsymbol{\mu}_r(\varepsilon) \mathbf{R}_{m;\varepsilon}$ does not define an element of $\mathbf{L}^2(\Omega)$. We are going to introduce a corrector $\mathbf{C}_{m;\varepsilon}$ satisfying suitable estimates and so that

$$(4.13) \quad \begin{cases} [(\mathbf{R}_{m;\varepsilon} - \mathbf{C}_{m;\varepsilon}) \times \mathbf{n}]_{\Sigma} & = 0 & \text{on } \Sigma \\ [\underline{\alpha}^{-1} \operatorname{curl}(\mathbf{R}_{m;\varepsilon} - \mathbf{C}_{m;\varepsilon}) \times \mathbf{n}]_{\Sigma} & = 0 & \text{on } \Sigma \\ (\mathbf{R}_{m;\varepsilon} - \mathbf{C}_{m;\varepsilon}) \times \mathbf{n} & = 0 & \text{on } \partial\Omega . \end{cases}$$

Construction of $\mathbf{C}_{m;\varepsilon}$: We take $\mathbf{C}_{m;\varepsilon} = 0$ in Ω_- and use a trace lifting to define $\mathbf{C}_{m;\varepsilon}$ in Ω_+ . It suffices that

$$(4.14) \quad \begin{cases} \mathbf{C}_{m;\varepsilon}^+ \times \mathbf{n} & = 0 & \text{on } \Sigma \\ \alpha_+^{-1} \operatorname{curl} \mathbf{C}_{m;\varepsilon}^+ \times \mathbf{n} & = \mathbf{g}_{m;\varepsilon} & \text{on } \Sigma \\ \mathbf{C}_{m;\varepsilon}^+ \times \mathbf{n} & = 0 & \text{on } \partial\Omega . \end{cases}$$

Denoting by C_β and C_3 the tangential and normal components of $\mathbf{C}_{m;\varepsilon}^+$ associated with a system of normal coordinates $\mathbf{y} = (y_\beta, y_3)$, and by g_β the components of $\mathbf{g}_{m;\varepsilon}$ the above system becomes (cf. [10, Sec. 5.2, Eq. (5.12)])

$$(4.15) \quad \begin{cases} C_\beta & = 0 & \text{on } \Sigma \\ \alpha_+^{-1} (\partial_3 C_\beta - \partial_\beta C_3) & = g_\beta & \text{on } \Sigma \\ C_\beta & = 0 & \text{on } \partial\Omega . \end{cases}$$

It can be solved in $\mathbf{H}^2(\Omega_+)$ choosing $C_3 = 0$ and a standard lifting of the first two traces on Σ and $\partial\Omega$ with the estimate

$$(4.16) \quad \|\mathbf{C}_{m;\varepsilon}^+\|_{2,\Omega_+} \leq C \|\mathbf{g}_{m;\varepsilon}\|_{\frac{1}{2},\Sigma} .$$

We deduce from assumption (4.11), and (4.16)

$$(4.17) \quad \|\mathbf{C}_{m;\varepsilon}^+\|_{2,\Omega_+} \leq C \varepsilon^m ,$$

where C may depend on m . We set

$$(4.18) \quad \tilde{\mathbf{R}}_{m;\varepsilon} := \mathbf{R}_{m;\varepsilon} - \mathbf{C}_{m;\varepsilon} \quad \text{and} \quad \tilde{\mathbf{J}}_{m;\varepsilon} := \operatorname{curl} \underline{\alpha}^{-1} \operatorname{curl} \tilde{\mathbf{R}}_{m;\varepsilon} - \kappa_+^2 \boldsymbol{\mu}_r(\varepsilon) \tilde{\mathbf{R}}_{m;\varepsilon} .$$

Hence by construction, we have $\tilde{\mathbf{J}}_{m;\varepsilon} \in \mathbf{L}^2(\Omega)$ with the estimates

$$(4.19) \quad \|\tilde{\mathbf{J}}_{m;\varepsilon}\|_{0,\Omega} \leq C \varepsilon^{m-1} .$$

STEP 2. We can apply Theorem 2.1 to the couple $(\mathbf{H}, \mathbf{E}) = (\tilde{\mathbf{R}}_{m;\varepsilon}, (\underline{\sigma} - i\omega\varepsilon_0)^{-1} \operatorname{curl} \tilde{\mathbf{R}}_{m;\varepsilon})$ and, thanks to (4.18), we obtain

$$\|\tilde{\mathbf{R}}_{m;\varepsilon}\|_{0,\Omega} + \|\operatorname{curl} \tilde{\mathbf{R}}_{m;\varepsilon}\|_{0,\Omega} \leq C \|\tilde{\mathbf{J}}_{m;\varepsilon}\|_{0,\Omega} .$$

Combining this estimate with (4.17) and (4.19), we deduce

$$(4.20) \quad \|\mathbf{R}_{m;\varepsilon}\|_{0,\Omega} + \|\operatorname{curl} \mathbf{R}_{m;\varepsilon}\|_{0,\Omega} \leq C\varepsilon^{m-1},$$

where C may depend on m .

STEP 3. In order to have an optimal estimate for $\mathbf{R}_{m;\varepsilon}$, we use (4.20) for $m + 2$. We obtain

$$(4.21) \quad \|\mathbf{R}_{m+2;\varepsilon}\|_{0,\Omega} + \|\operatorname{curl} \mathbf{R}_{m+2;\varepsilon}\|_{0,\Omega} \leq C\varepsilon^{m+1}.$$

But we have the formula

$$(4.22) \quad \mathbf{R}_{m;\varepsilon} = \sum_{j=m+1}^{m+2} \varepsilon^j \mathbf{H}_j + \mathbf{R}_{m+2;\varepsilon},$$

and using (4.2), we have also for any $j \in \mathbb{N}$

$$(4.23) \quad \|\mathbf{H}_j^+\|_{\mathbf{H}(\operatorname{curl},\Omega_+)} + \varepsilon^{-\frac{1}{2}} \|\mathbf{H}_j^-\|_{0,\Omega_-} + \varepsilon^{\frac{1}{2}} \|\operatorname{curl} \mathbf{H}_j^-\|_{0,\Omega_-} \leq C.$$

We finally deduce the wanted estimate (4.12) from (4.21) to (4.23). \square

5. CONCLUSION AND PERSPECTIVES

In this work we tackled a transmission problem in materials with high contrast in magnetic permeabilities. We proved uniform a priori estimates for the electromagnetic field solution of the time-harmonic Maxwell equations as the relative magnetic permeability μ_r between a magnetic conductor and a non-magnetic material tends to infinity, and when the interface between the subdomains is Lipschitz. Assuming smoothness for the interface between the subdomains, we derived a multiscale expansion for the magnetic field in powers of $\frac{1}{\sqrt{\mu_r}}$ with profiles inside the magnetic conductor, and we proved error estimates at any order.

One perspective would be to carry out numerical experiments in axisymmetric domains [8, 16, 24, 10]. This would make it possible in particular to illustrate our theoretical results. The multiscale expansion provides also an a priori knowledge on the magnetic field useful for the design of meshes in view of a quality finite element approximation. Thus, the mesh should be matched to the boundary of the magnetic conductor and can be coarse far from its boundary in the magnetic conductor, as with the *skin effect* in non-magnetic conductive materials. It would be useful also to derive impedance boundary conditions (see, e.g., [7, 20, 15] for scattering problems from absorbing obstacles) on the interface (magnetic conductor / non-magnetic material) in order to reduce the computational costs.

APPENDIX A. ELEMENTS OF PROOF FOR THE MULTISCALE EXPANSION

In the framework of Section 4, we assume that the surfaces Σ and Γ are smooth (cf. Assumption 4.1). Recall that there exists $\varepsilon_* > 0$ and such that for all $\varepsilon \in (0, \varepsilon_*)$, the problem (2.1)-(2.3) has a unique solution $(\mathbf{E}_{(\varepsilon)}, \mathbf{H}_{(\varepsilon)})$ which is denoted by $(\mathbf{E}_{(\varepsilon)}^+, \mathbf{H}_{(\varepsilon)}^+)$ in Ω_+ , and by $(\mathbf{E}_{(\varepsilon)}^-, \mathbf{H}_{(\varepsilon)}^-)$ in Ω_- . Furthermore, we assume that the right hand side \mathbf{j} is smooth and its support does not the domain Ω_- .

Recall that (y_α, y_3) is a *normal coordinate system* to the surface Σ in \mathcal{U}_- , see Figure 2. The function $\mathbf{y} \mapsto \chi(y_3)$ is a smooth cut-off with support in $\bar{\mathcal{U}}_-$ and equal to 1 in a smaller tubular neighborhood of Σ .

Theorem A.1. *Under the above assumptions, the magnetic field $\mathbf{H}_{(\varepsilon)}$ possesses the asymptotic expansion (see Theorem 4.2 in Section 4.2 for precise estimates):*

$$(A.1) \quad \mathbf{H}_{(\varepsilon)}^+(\mathbf{x}) \approx \sum_{j \geq 0} \varepsilon^j \mathbf{H}_j^+(\mathbf{x}),$$

$$(A.2) \quad \mathbf{H}_{(\varepsilon)}^-(\mathbf{x}) \approx \sum_{j \geq 0} \varepsilon^j \mathbf{H}_j^-(\mathbf{x}; \varepsilon) \quad \text{with} \quad \mathbf{H}_j^-(\mathbf{x}; \varepsilon) = \chi(y_3) \underline{\mathfrak{H}}_j(y_\alpha, \frac{y_3}{\varepsilon}),$$

where $\underline{\mathfrak{H}}_j(y_\alpha, Y_3) \rightarrow 0$ when $Y_3 \rightarrow \infty$. Moreover, for any $j \in \mathbb{N}$, we have

$$(A.3) \quad \mathbf{H}_j^+ \in \mathbf{H}(\text{curl}, \Omega_+) \quad \text{and} \quad \underline{\mathfrak{H}}_j \in \mathbf{H}(\text{curl}, \Sigma \times \mathbb{R}_+).$$

Hereafter, we present elements of proof of this theorem and details about the terms in asymptotics (A.1)–(A.2). In §A.1, we expand the “magnetic” Maxwell operators in power series of ε inside the boundary layer \mathcal{U}_- . We deduce in §A.2 the equations satisfied by the magnetic profiles, and we derive explicitly the first ones in §A.3.

A.1. Expansion of the Maxwell operators. Integrating by parts in the variational formulation (2.8), we find the following Maxwell transmission problem for the magnetic field (we recall $\kappa_+ = \omega \sqrt{\varepsilon_0 \mu_+}$, cf. (2.9))

$$(A.4) \quad \begin{cases} \text{curl curl } \mathbf{H}_{(\varepsilon)}^+ - \kappa_+^2 (1 + \frac{i}{\delta_+^2}) \mathbf{H}_{(\varepsilon)}^+ = \text{curl } \mathbf{j} & \text{in } \Omega_+ \\ \varepsilon^2 \text{curl curl } \mathbf{H}_{(\varepsilon)}^- - \kappa_+^2 (1 + \frac{i}{\delta_-^2}) \mathbf{H}_{(\varepsilon)}^- = 0 & \text{in } \Omega_- \\ (1 + \frac{i}{\delta_+^2})^{-1} \text{curl } \mathbf{H}_{(\varepsilon)}^+ \times \mathbf{n} = (1 + \frac{i}{\delta_-^2})^{-1} \text{curl } \mathbf{H}_{(\varepsilon)}^- \times \mathbf{n} & \text{on } \Sigma \\ \mathbf{H}_{(\varepsilon)}^+ \times \mathbf{n} = \mathbf{H}_{(\varepsilon)}^- \times \mathbf{n} & \text{on } \Sigma \\ \mathbf{H}_{(\varepsilon)}^+ \times \mathbf{n} = 0 & \text{on } \Gamma. \end{cases}$$

As a consequence of the Maxwell transmission problem, we have also

$$\text{div } \underline{\mu} \mathbf{H}_{(\varepsilon)} = 0 \quad \text{in } \Omega.$$

Therefore, we have in particular $\underline{\mu} \mathbf{H}_{(\varepsilon)} \in \mathbf{H}(\text{div}, \Omega)$, where

$$\mathbf{H}(\text{div}, \Omega) = \{ \mathbf{u} \in \mathbf{L}^2(\Omega) \mid \text{div } \mathbf{u} \in L^2(\Omega) \},$$

and from which we deduce the extra transmission condition

$$(A.5) \quad \varepsilon^2 \mathbf{H}_{(\varepsilon)}^+ \cdot \mathbf{n} = \mathbf{H}_{(\varepsilon)}^- \cdot \mathbf{n} \quad \text{on } \Sigma.$$

In that follows we denote by $\mathbf{L}(y_\alpha, h; D_\alpha, \partial_3^h)$ the second order Maxwell operator $\varepsilon^2 \text{curl curl} - \kappa_+^2 (1 + \frac{i}{\delta_-^2}) \mathbb{I}$ set in \mathcal{U}_- in a *normal coordinate system* (y_α, h) . Here D_α is the covariant derivative on the interface Σ , and ∂_3^h is the partial derivative with respect to the normal coordinate $y_3 = h$. Then after the scaling $Y_3 = \varepsilon^{-1}h$, the operator which is written $\mathbf{L}[\varepsilon]$, expands in power series of ε with coefficients intrinsic operators :

$$\mathbf{L}[\varepsilon] = \sum_{n=0}^{\infty} \varepsilon^n \mathbf{L}^n.$$

We denote by L_α^n the surface components of \mathbf{L}^n . Using a splitting a three-dimensional vector field $\underline{\mathfrak{H}}$ into its normal component \mathfrak{h} and its tangential component \mathfrak{H} that can be viewed as a one-form field \mathfrak{H}_α , and using the summation convention of repeated two dimensional indices (represented by greek letters), we have

$$(A.6) \quad L_\alpha^0(\underline{\mathfrak{H}}) = -\partial_3^2 \mathfrak{H}_\alpha - \kappa_+^2 (1 + \frac{i}{\delta_-^2}) \mathfrak{H}_\alpha \quad \text{and} \quad L_\alpha^1(\underline{\mathfrak{H}}) = -2b_\alpha^\beta \partial_3 \mathfrak{H}_\beta + \partial_3 \partial_\alpha \mathfrak{h} + b_\beta^\beta \partial_3 \mathfrak{H}_\alpha,$$

where ∂_3 is the partial derivative with respect to Y_3 and $b_\alpha^\beta = a^{\beta\gamma} b_{\gamma\alpha}$. Here $a^{\beta\gamma}$ is the inverse of the metric tensor $a_{\beta\gamma}$ in Σ , and $b_{\gamma\alpha}$ is the curvature tensor in Σ . Then we denote by L_3^n the transverse components of \mathbf{L}^n . We have

$$(A.7) \quad L_3^0(\underline{\mathfrak{H}}) = -\kappa_+^2 (1 + \frac{i}{\delta_-^2}) \mathfrak{h} \quad \text{and} \quad L_3^1(\underline{\mathfrak{H}}) = \gamma_\alpha^\alpha (\partial_3 \mathfrak{H}) + b_\beta^\beta \partial_3 \mathfrak{h}.$$

Here $\gamma_{\alpha\beta}(\underline{\mathfrak{H}}) = \frac{1}{2}(D_\alpha \mathfrak{H}_\beta + D_\beta \mathfrak{H}_\alpha) - b_{\alpha\beta} \mathfrak{h}$ is the change of metric tensor.

Then we denote by $\mathbf{B}(y_\alpha, h; \partial_\alpha, \partial_3^h)$ the tangent trace operator $\text{curl} \cdot \times \mathbf{n}$ on Σ in a *normal coordinate system*. If $\underline{\mathfrak{H}} = (\mathfrak{H}_\alpha, \mathfrak{h})$, we have (cf. [9, Sec. A.4])

$$(\mathbf{B}(y_\alpha, h; \partial_\alpha, \partial_3^h) \underline{\mathfrak{H}})_\alpha = \partial_3^h \mathfrak{H}_\alpha - \partial_\alpha \mathfrak{h}.$$

Then if $\mathbf{B}[\varepsilon]$ is the operator obtained from \mathbf{B} in \mathcal{U}_- after the scaling $Y_3 = \varepsilon^{-1}h$, we have

$$\mathbf{B}[\varepsilon] = \varepsilon^{-1} \mathbf{B}^0 + \mathbf{B}^1.$$

Finally we denote by B_α^n the surface components of \mathbf{B}^n . We have

$$(A.8) \quad B_\alpha^0(\underline{\mathfrak{H}}) = \partial_3 \mathfrak{H}_\alpha \quad \text{and} \quad B_\alpha^1(\underline{\mathfrak{H}}) = -\partial_\alpha \mathfrak{h}.$$

A.2. Equations for the coefficients of the magnetic field. According to the second and third equations in system (A.4), the profiles $\underline{\mathfrak{H}}_j$ and the coefficients \mathbf{H}_j^+ of the magnetic field satisfy the following system

$$(A.9) \quad \mathbf{L}[\varepsilon] \sum_{j \geq 0} \varepsilon^j \underline{\mathfrak{H}}_j(y_\alpha, Y_3) = 0 \quad \text{in } \Sigma \times I,$$

$$(A.10) \quad \mathbf{B}[\varepsilon] \sum_{j \geq 0} \varepsilon^j \underline{\mathfrak{H}}_j(y_\alpha, 0) = \left(1 + \frac{i}{\delta_-^2}\right) \left(1 + \frac{i}{\delta_+^2}\right)^{-1} \sum_{j \geq 0} \varepsilon^j \operatorname{curl} \mathbf{H}_j^+ \times \mathbf{n} \quad \text{on } \Sigma,$$

with $I = (0, +\infty)$.

Then we perform in (A.9)-(A.10) the identification of terms with the same power in ε . The components of the equation (A.9) are the collections of equations

$$(A.11) \quad \mathbf{L}^0(\underline{\mathfrak{H}}_0) = 0, \quad \mathbf{L}^0(\underline{\mathfrak{H}}_1) + \mathbf{L}^1(\underline{\mathfrak{H}}_0) = 0, \quad \text{and} \quad \sum_{l=0}^n \mathbf{L}^{n-l}(\underline{\mathfrak{H}}_l) = 0,$$

for all $n \geq 2$. The surface components of the equation (A.10) write

$$(A.12) \quad B_\alpha^0(\underline{\mathfrak{H}}_0) = 0 \quad \text{and} \quad B_\alpha^0(\underline{\mathfrak{H}}_n) + B_\alpha^1(\underline{\mathfrak{H}}_{n-1}) = \left(1 + \frac{i}{\delta_-^2}\right) \left(1 + \frac{i}{\delta_+^2}\right)^{-1} (\operatorname{curl} \mathbf{H}_{n-1}^+ \times \mathbf{n})_\alpha,$$

for all $n \geq 1$.

Using the components of the operator \mathbf{L}^0 (A.6)-(A.7), and the surface components of \mathbf{B}_0 and \mathbf{B}_1 (A.8), and expanding $\mathbf{H}_{(\varepsilon)}^+$ in Ω_+ , we thus see that, according to the system (A.4), the profiles $\underline{\mathfrak{H}}_n = (\mathfrak{H}_n, \mathfrak{h}_n)$ and the terms \mathbf{H}_n^+ have to satisfy, for all $n \geq 0$,

$$(A.13) \quad \left\{ \begin{array}{ll} (i) & \partial_3^2 \mathfrak{H}_{n,\alpha} - \lambda^2 \mathfrak{H}_{n,\alpha} = \sum_{j=0}^{n-1} L_\alpha^{n-j}(\underline{\mathfrak{H}}_j) & \text{in } \Sigma \times I \\ (ii) & \partial_3 \mathfrak{H}_{n,\alpha} = \partial_\alpha \mathfrak{h}_{n-1} + \left(1 + \frac{i}{\delta_-^2}\right) \left(1 + \frac{i}{\delta_+^2}\right)^{-1} (\operatorname{curl} \mathbf{H}_{n-1}^+ \times \mathbf{n})_\alpha & \text{on } \Sigma \\ (iii) & -\lambda^2 \mathfrak{h}_n = \sum_{j=0}^{n-1} L_3^{n-j}(\underline{\mathfrak{H}}_j) & \text{in } \Sigma \times I \\ (iv) & \operatorname{curl} \operatorname{curl} \mathbf{H}_n^+ - \kappa_+^2 \left(1 + \frac{i}{\delta_+^2}\right) \mathbf{H}_n^+ = \delta_n^0 \operatorname{curl} \mathbf{j} & \text{in } \Omega_+ \\ (v) & \mathbf{n} \times \mathbf{H}_n^+ \times \mathbf{n} = \mathfrak{H}_n & \text{on } \Sigma \\ (vi) & \mathbf{H}_n^+ \times \mathbf{n} = 0 & \text{on } \Gamma. \end{array} \right.$$

Here we have

$$\lambda = \kappa_+ \sqrt[4]{1 + \frac{1}{\delta_-^4}} e^{i \frac{\theta(\delta_-) - \pi}{2}} \quad \text{with} \quad \theta(\delta_-) = \arctan \frac{1}{\delta_-^2},$$

so that: $-\lambda^2 = \kappa_+^2 \left(1 + \frac{i}{\delta_-^2}\right)$ and $\operatorname{Re} \lambda > 0$.

Note that the transmission condition (A.5) implies the extra condition

$$(A.14) \quad \mathfrak{h}_n = \mathbf{H}_{n-2}^+ \cdot \mathbf{n} \quad \text{on } \Sigma.$$

The set of equations (A.13) allows to determine $\underline{\mathfrak{H}}_n$ and \mathbf{H}_n^+ by induction.

A.3. First terms of the magnetic field asymptotics. According to equations (i)-(ii) in system (A.13) for $n = 0$, \mathfrak{H}_0 satisfies the following ODE

$$(A.15) \quad \begin{cases} \partial_3^2 \mathfrak{H}_0(\cdot, Y_3) - \lambda^2 \mathfrak{H}_0(\cdot, Y_3) = 0 & \text{for } Y_3 \in I, \\ \partial_3 \mathfrak{H}_0(\cdot, 0) = 0. \end{cases}$$

The unique solution of (A.15) such that $\mathfrak{H}_0 \rightarrow 0$ when $Y_3 \rightarrow \infty$, is $\mathfrak{H}_0 = 0$. From equation (iii) in system (A.13) for $n = 0$, there holds $\mathfrak{h}_0 = 0$. We infer

$$(A.16) \quad \underline{\mathfrak{H}}_0(y_\beta, Y_3) = 0.$$

From equations (iv)-(vi) in (A.13) for $n = 0$, and from (A.16), the first asymptotic of the magnetic field in the non-magnetic region solves the following problem

$$\begin{cases} \operatorname{curl} \operatorname{curl} \mathbf{H}_0^+ - \kappa_+^2 (1 + \frac{i}{\delta_+^2}) \mathbf{H}_0^+ = \operatorname{curl} \mathbf{j} & \text{in } \Omega_+ \\ \mathbf{H}_0^+ \times \mathbf{n} = 0 & \text{on } \Sigma \cup \Gamma. \end{cases}$$

The next term determined in the asymptotic expansion is \mathfrak{H}_1 . From equations (i)-(ii) in (A.13) for $n = 1$, \mathfrak{H}_1 satisfies for $Y_3 \in I$

$$\begin{cases} \partial_3^2 \mathfrak{H}_1(\cdot, Y_3) - \lambda^2 \mathfrak{H}_1(\cdot, Y_3) = 0, \\ \partial_3 \mathfrak{H}_1(\cdot, 0) = (1 + \frac{i}{\delta_-^2})(1 + \frac{i}{\delta_+^2})^{-1} (\operatorname{curl} \mathbf{H}_0^+ \times \mathbf{n})(\cdot, 0). \end{cases}$$

We denote by $\mathbf{j}_k(y_\beta) = \lambda^{-1}(1 + \frac{i}{\delta_-^2})(1 + \frac{i}{\delta_+^2})^{-1} (\operatorname{curl} \mathbf{H}_k^+ \times \mathbf{n})(y_\beta, 0)$ for $k = 0, 1$. Hence,

$$(A.17) \quad \mathfrak{H}_1(y_\beta, Y_3) = -\mathbf{j}_0(y_\beta) e^{-\lambda Y_3}.$$

From equation (iii) in (A.13) for $n = 1$, and from (A.16), we obtain $\mathfrak{h}_1 = 0$. From equations (iv)-(vi) in (A.13) for $n = 1$, and from (A.17), the asymptotic of order 1 for the magnetic field in the non-magnetic part solves :

$$\begin{cases} \operatorname{curl} \operatorname{curl} \mathbf{H}_1^+ - \kappa_+^2 (1 + \frac{i}{\delta_+^2}) \mathbf{H}_1^+ = 0 & \text{in } \Omega_+ \\ \mathbf{H}_1^+ \times \mathbf{n} = -\mathbf{j}_0 \times \mathbf{n} & \text{on } \Sigma \\ \mathbf{H}_1^+ \times \mathbf{n} = 0 & \text{on } \Gamma. \end{cases}$$

Then, from equations (i)-(ii) in (A.13) for $n = 2$, $\mathfrak{H}_{2,\alpha}$ solves the ODE for $Y_3 \in I$:

$$\begin{cases} \partial_3^2 \mathfrak{H}_{2,\alpha}(\cdot, Y_3) - \lambda^2 \mathfrak{H}_{2,\alpha}(\cdot, Y_3) = -2b_\alpha^\sigma \partial_3 \mathfrak{H}_{1,\sigma}(\cdot, Y_3) + b_\beta^\beta \partial_3 \mathfrak{H}_{1,\alpha}(\cdot, Y_3) \\ \partial_3 \mathfrak{H}_{2,\alpha}(\cdot, 0) = (1 + \frac{i}{\delta_-^2})(1 + \frac{i}{\delta_+^2})^{-1} (\operatorname{curl} \mathbf{H}_1^+ \times \mathbf{n})_\alpha(\cdot, 0). \end{cases}$$

We denote by $\mathbf{j}_{k,\alpha}$ the surface components of \mathbf{j}_k , for $k = 0, 1$. We obtain

$$(A.18) \quad \mathfrak{H}_{2,\alpha}(y_\beta, Y_3) = \left[-\mathbf{j}_{1,\alpha} + (\lambda^{-1} + Y_3)(b_\alpha^\sigma \mathbf{j}_{0,\sigma} - \mathcal{H} \mathbf{j}_{0,\alpha}) \right] (y_\beta) e^{-\lambda Y_3}.$$

From equation (iii) in (A.13) for $n = 2$, we infer

$$\mathfrak{h}_2(y_\beta, Y_3) = -\lambda^{-1} D_\alpha \mathbf{j}_0^\alpha(y_\beta) e^{-\lambda Y_3}.$$

From equations (iv)-(vi) in (A.13) for $n = 2$, and from (A.18), the asymptotic of order 2 for the magnetic field in the non-magnetic part solves :

$$\begin{cases} \operatorname{curl} \operatorname{curl} \mathbf{H}_2^+ - \kappa_+^2 (1 + \frac{i}{\delta_+^2}) \mathbf{H}_2^+ = 0 & \text{in } \Omega_+ \\ \mathbf{n} \times \mathbf{H}_2^+ \times \mathbf{n} = -\mathbf{j}_1 + \lambda^{-1} (\mathcal{C} - \mathcal{H}) \mathbf{j}_0 & \text{on } \Sigma \\ \mathbf{H}_2^+ \times \mathbf{n} = 0 & \text{on } \Gamma. \end{cases}$$

REFERENCES

- [1] D. Abou El Nasser El Yafi and V. Péron. Efficient asymptotic models for axisymmetric eddy current problems in linear ferromagnetic materials. *Computers & Mathematics with Applications*, 151:335–345, 2023.
- [2] D. Abou El Nasser El Yafi, V. Péron, R. Perrussel, and L. Krähenbühl. Numerical study of the magnetic skin effect: Efficient parameterization of 2d surface-impedance solutions for linear ferromagnetic materials. *International Journal of Numerical Modelling: Electronic Networks, Devices and Fields*, 36(3):e3051, 2023.
- [3] H. Ammari, B. Li, and J. Zou. Superresolution in recovering embedded electromagnetic sources in high contrast media. *SIAM Journal on Imaging Sciences*, 13(3):1467–1510, 2020.
- [4] H. Ammari, B. Li, and J. Zou. Mathematical analysis of electromagnetic scattering by dielectric nanoparticles with high refractive indices. *Transactions of the American Mathematical Society*, 376(01):39–90, 2023.
- [5] C. Amrouche, C. Bernardi, M. Dauge, and V. Girault. Vector potentials in three-dimensional non-smooth domains. *Math. Methods Appl. Sci.*, 21(9):823–864, 1998.
- [6] M. Artola and M. Cessenat. Diffraction d’une onde électromagnétique par une couche composite mince accolée à un corps conducteur épais. ii, cas d’inclusions (ou d’une couche) de forte permittivité et de forte perméabilité. *Comptes rendus de l’Académie des sciences. Série 1, Mathématique*, 313(6):381–385, 1991.
- [7] M. Artola and M. Cessenat. Diffraction d’une onde électromagnétique par un obstacle borné à permittivité et perméabilité élevées. *Comptes rendus de l’Académie des sciences. Série 1, Mathématique*, 314(5):349–354, 1992.
- [8] C. Bernardi, M. Dauge, and Y. Maday. *Spectral methods for axisymmetric domains*, volume 3 of *Series in Applied Mathematics (Paris)*. Gauthier-Villars, Éditions Scientifiques et Médicales Elsevier, Paris, 1999. Numerical algorithms and tests due to Mejdı Azaïez.
- [9] G. Caloz, M. Dauge, E. Faou, and V. Péron. On the influence of the geometry on skin effect in electromagnetism. *Computer Methods in Applied Mechanics and Engineering*, 200(9-12):1053–1068, 2011.
- [10] G. Caloz, M. Dauge, and V. Péron. Uniform estimates for transmission problems with high contrast in heat conduction and electromagnetism. *Journal of Mathematical Analysis and Applications*, 370(2):555–572, 2010.
- [11] M. Costabel and M. Dauge. Singularities of electromagnetic fields in polyhedral domains. *Arch. Ration. Mech. Anal.*, 151(3):221–276, 2000.
- [12] M. Costabel, M. Dauge, and S. Nicaise. Singularities of Maxwell interface problems. *M2AN Math. Model. Numer. Anal.*, 33(3):627–649, 1999.
- [13] M. Dauge, E. Faou, and V. Péron. Comportement asymptotique à haute conductivité de l’épaisseur de peau en électromagnétisme. *C. R. Acad. Sci. Paris Sér. I Math.*, 348(7-8):385–390, 2010.
- [14] L. Eskola, R. Puranen, and H. Soininen. Measurement of magnetic properties of steel sheets. *Geophysical prospecting*, 47(4):593–602, 2001.
- [15] H. Haddar, P. Joly, and H.-M. Nguyen. Generalized impedance boundary conditions for scattering problems from strongly absorbing obstacles: the case of Maxwell’s equations. *Math. Models Methods Appl. Sci.*, 18(10):1787–1827, 2008.
- [16] R. Hiptmair and P. D. Ledger. Computation of resonant modes for axisymmetric Maxwell cavities using *hp*-version edge finite elements. *Internat. J. Numer. Methods Engrg.*, 62(12):1652–1676, 2005.
- [17] J. Huang and J. Zou. Uniform a priori estimates for elliptic and static maxwell interface problems. *Discrete and Continuous Dynamical Systems Series B*, 7(1):145, 2007.
- [18] M. Issa, J.-R. Poirier, R. Perrussel, O. Chadebec, and V. Péron. Boundary element method for 3d conductive thin layer in eddy current problems. *COMPEL-The international journal for computation and mathematics in electrical and electronic engineering*, 38(2):502–521, 2019.

- [19] H. Kawaguchi, T. Muramatsu, M. Katoh, M. Hosaka, and Y. Takashima. Time-domain eddy current and wake fields analysis of pulsed multipole magnet beam injector in synchrotron radiation ring. *IEICE Transactions on Electronics*, 105(4):146–153, 2022.
- [20] O. Lafitte and G. Lebeau. Équations de maxwell et opérateur d’impédance sur le bord d’un obstacle convexe absorbant. *Comptes rendus de l’Académie des sciences. Série 1, Mathématique*, 316(11):1177–1182, 1993.
- [21] R. C. MacCamy and E. Stephan. Solution procedures for three-dimensional eddy current problems. *J. Math. Anal. Appl.*, 101(2):348–379, 1984.
- [22] R. C. MacCamy and E. Stephan. A skin effect approximation for eddy current problems. *Arch. Rational Mech. Anal.*, 90(1):87–98, 1985.
- [23] S. Mukherjee and M. E. Everett. 3d controlled-source electromagnetic edge-based finite element modeling of conductive and permeable heterogeneities. *Geophysics*, 76(4):F215–F226, 2011.
- [24] B. Nkemzi. On the solution of Maxwell’s equations in axisymmetric domains with edges. *ZAMM Z. Angew. Math. Mech.*, 85(8):571–592, 2005.
- [25] K. Noh, S. Oh, S. J. Seol, K. H. Lee, and J. Byun. Analysis of anomalous electrical conductivity and magnetic permeability effects using a frequency domain controlled-source electromagnetic method. *Geophysical Journal International*, 204(3):1550–1564, 2016.
- [26] D. Pardo, L. Demkowicz, C. Torres-Verdín, and M. Paszynski. Two-dimensional high-accuracy simulation of resistivity logging-while-drilling (LWD) measurements using a self-adaptive goal-oriented hp finite element method. *SIAM J. Appl. Math.*, 66(6):2085–2106, 2006.
- [27] V. Péron. Asymptotic models and impedance conditions for highly conductive sheets in the time-harmonic eddy current model. *SIAM Journal on Applied Mathematics*, 79(6):2242–2264, 2019.
- [28] V. Péron and C. Poignard. On a magnetic skin effect in eddy current problems: the magnetic potential in magnetically soft materials. *Zeitschrift für Angewandte Mathematik und Physik*, 72(4), 2021.
- [29] X. Ren, R. Corcolle, and L. Daniel. A homogenization technique to calculate eddy current losses in soft magnetic composites using a complex magnetic permeability. *IEEE Transactions on Magnetics*, 52(12):1–9, 2016.
- [30] E. Stephan. Solution procedures for interface problems in acoustics and electromagnetics. In *Theoretical acoustics and numerical techniques*, volume 277 of *CISM Courses and Lectures*, pages 291–348. Springer, Vienna, 1983.

LABORATOIRE DE MATHÉMATIQUES ET DE LEURS APPLICATIONS DE PAU (UMR CNRS 5142),
E2S UPPA, CNRS, UNIVERSITÉ DE PAU ET DES PAYS DE L’ADOUR, 64013 PAU CEDEX, FRANCE,
VICTOR.PERON@UNIV-PAU.FR