



**HAL**  
open science

# Wastewater treatment plant efficiency and contaminant levels in a Mediterranean coastal area: a comprehensive inventory and assessment

P. Merdy, Floriane Delpy, Gilles Baratto, E. Plessis, Y. Lucas

## ► To cite this version:

P. Merdy, Floriane Delpy, Gilles Baratto, E. Plessis, Y. Lucas. Wastewater treatment plant efficiency and contaminant levels in a Mediterranean coastal area: a comprehensive inventory and assessment. International Journal of Environmental Science and Technology, In press, 10.1007/s13762-024-05801-7. hal-04622028

**HAL Id: hal-04622028**

**<https://hal.science/hal-04622028v1>**

Submitted on 24 Jun 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **Wastewater treatment plant efficiency and contaminant levels in a Mediterranean coastal area: a comprehensive inventory and assessment**

This Accepted Manuscript (AM) is a PDF file of the manuscript accepted for publication after peer review, when applicable, but does not reflect post-acceptance improvements, or any corrections. Use of this AM is subject to the publisher's embargo period and AM terms of use. Under no circumstances may this AM be shared or distributed under a Creative Commons or other form of open access license, nor may it be reformatted or enhanced, whether by the Author or third parties. By using this AM (for example, by accessing or downloading) you agree to abide by Springer Nature's terms of use for AM versions of subscription articles: <https://www.springernature.com/gp/open-research/policies/accepted-manuscript-terms>

The Version of Record (VOR) of this article, as published and maintained by the publisher, is available online at: <https://doi.org/10.1007/s13762-024-05801-7>. The VOR is the version of the article after copy-editing and typesetting, and connected to open research data, open protocols, and open code where available. Any supplementary information can be found on the journal website, connected to the VOR.

For research integrity purposes it is best practice to cite the published Version of Record (VOR), where available (for example, see ICMJE's guidelines on overlapping publications). Where users do not have access to the VOR, any citation must clearly indicate that the reference is to an Accepted Manuscript (AM) version.

# Wastewater treatment plant efficiency and contaminant levels in a Mediterranean coastal area: a comprehensive inventory and assessment.

P. Merdy<sup>1\*</sup>, F. Delpy<sup>1</sup>, G. Baratto<sup>2</sup>, E. Plessis<sup>2</sup>, Y. Lucas<sup>1</sup>

(1) Université de Toulon, Aix Marseille Univ., CNRS, IM2NP, 83041 TOULON CEDEX 9, France

(2) Véolia Eau, - Var Provence Méditerranée- Rue des Oliviers - Le Pouverel, La GARDE (83), France

\* Corresponding author: merdy@univ-tln.fr

## ORCID numbers

P. Merdy: 0000-0002-4540-6464

Y. Lucas: 0000-0003-3073-7065

## Acknowledgement

Financial support was obtained from the Water Agency (Agence de l'Eau Rhône Méditerranée Corse) and the VEOLIA Foundation.

## CRediT authorship contribution statement

**P. Merdy:** Conceptualization, Supervision, Funding acquisition, Project administration, Investigation, Methodology, Formal analysis, Writing – original draft. **F. Delpy:** Investigation, Writing – review & editing. **G. Baratto:** Conceptualization, Funding acquisition. **E. Plessis:** Conceptualization, Funding acquisition. **Y. Lucas:** Investigation, Formal analysis, Writing – review & editing.

# Wastewater treatment plant efficiency and contaminant levels in a Mediterranean coastal area: a comprehensive inventory and assessment.

## Abstract

The contamination of estuarine and coastal waters raises the question of risk assessment, management and remediation, all of which require a good knowledge of the contamination present. Most studies focus on specific species in samples from a given site, but few examine the presence of multiple contaminants at once. Here we considered 7 nonmetallic inorganics, 32 metals and metalloids, 4 chlorobenzenes, 16 PAHs, 8 PCBs, 34 organochloride pesticides and 239 other pesticides or pesticide metabolites, 12 pharmaceuticals and 2 others, in sea water, sediments, and in urban wastewater before and after treatment. Our study sites included a reference station, a harbor, a coastal bay and a treated wastewater outlet plume. Our results provide insight into the current contamination levels and the efficiency of the wastewater treatment plant for a wide range of contaminants. Although some organic contaminants remained at concentrations above the limit of reporting in the treated water, the dilution effect in the plume was sufficient to bring most of the organic contaminants studied below detectable levels, except for 17 $\alpha$ -ethinylestradiol, which was the most commonly detected contaminants. This finding **underscores** the insufficient reduction of endocrine disruptors by current wastewater treatment.

### **Keywords**

Persistent compounds; Trace metals; Pharmaceuticals; Coastal waters; WWTP effluents; Mass load

## Introduction

Coastal marine environments face significant impact from human activities, as they receive and concentrate numerous anthropogenic pollutants. The coastal zone, acting as a buffer between the land and the ocean, plays a critical role in providing various ecosystem services (Lu et al., 2018). Consequently, conserving this zone becomes imperative to ensure the sustainable utilization of marine resources (ICSU-ISSC, 2015). Contaminants found in the coastal zone can originate either directly from the marine environment or more commonly, they are transported from the continental environment (Riechers et al., 2021), particularly from large rivers and the surrounding coastal areas. Depending on the specific type of contaminant, local sources may exert greater influence, primarily due to the high population density and urban development in coastal regions (Creel, 2003; Nicolau et al., 2012; Merkens et al., 2016). The contaminants that tend to accumulate or transit in coastal regions encompass chemicals, metal, metalloids, nutrients, oil, microplastics, and plastics (Riechers et al., 2021). These contaminants can emanate from diffuse, non-point sources or localized, point sources, and their contamination can manifest as either sporadic or long-term occurrences.

Within the realm of chemical micropollutants, pesticides and pharmaceuticals stand out as major concerns (Fent et al., 2006). Notably, pesticides like glyphosate, a non-selective herbicide extensively employed by farmers and gardeners, pose a significant risk off runoff into watersheds and groundwater through soil infiltration (Daouk et al., 2015; Dollinger, 2016). The concentrations of glyphosate in surface waters can, in the most severe cases, reach the  $\text{mg L}^{-1}$  range, while they typically reside in the  $\mu\text{g L}^{-1}$  range in rivers and  $\text{ng L}^{-1}$  in seawater (Annett et al., 2014; Bento et al., 2021; Feltraco et al., 2022). However, the prevalence of glyphosate in aquatic environments is frequently underestimated due to its degradation into **aminomethylphosphonic acid (AMPA)**, which can reach concentrations surpassing those of glyphosate by more than fourfold (Bento et al., 2021).

Pharmaceuticals compounds, including antibiotics, analgesics, and nonsteroidal anti-inflammatory drugs, have been identified in seawater by dedicated studies (Gaw et al., 2014; Puckowski et al., 2016; Madikizela et al., 2020). Despite growing awareness of the potential impact of these drugs on aquatic organisms, current regulations do not account for their release into the environment, whether directly or through wastewater treatment plant (WWTP) effluents. In Europe, the regulatory framework for medicinal products is outlined in Directive 2001/83/EC, which lacks provisions addressing the discharge of these substances into the environment. In contrast, guidelines specifying permissible limits exist for most potentially toxic metals and metalloids. While veterinary pharmaceuticals can be directly released into the environment, the majority of human medicine products pass through WWTPs. However, the effectiveness of WWTPs in removing these pharmaceuticals varies significantly depending on the specific compound (Strenn et al., 2004).

In Europe, the main approach to reducing the presence of micropollutants in the environment is to implement reductions, or bans, on their use upstream of rivers or **WWTPs**. The Water Framework

Directive (CEC, 2000) aims to achieve a "good" ecological and chemical status for water bodies. The chemical status assessment is based on the analysis of 45 priority substances, encompassing 24 pesticides and only three pharmaceuticals, while the ecological status assessment involves 177 molecules (EU-CDI, 2013). However, this list, established nearly a decade ago, falls short of reflection the advancements in understanding micropollutants and their evolving sources.

Chemicals also fall **within** the purview of the European REACH regulation, designed to establish a harmonized system for the registration, evaluation, and authorization of chemical substances at the European level (Williams et al., 2009). Currently, this framework does not encompass pharmaceutical substances, although their inclusion is under investigation.

Scientists engage in research efforts aimed at informing and shaping regulatory frameworks, often with the support and encouragement of government agencies. In 2009, the NORMAN's Bioassay Working Group was established with the mission of fostering information exchange, method and tool validation, and project coordination related to the assessment of emerging environmental compounds (INERIS, 2018). These substances fall outside the scope of routine monitoring programs in Europe, and there remains limited understanding of their behavior, fate, and (eco)toxicological impacts. To address these concerns, the OSPAR convention binds its signatory nations to identify environmental threats and collaboratively undertake effective measures against the most hazardous substances in marine environments. **The European Parliament voted on 5/10/2023 for a proposal for a Directive specifically addressing urban wastewater treatment.**

It is therefore crucial to enhance our comprehension of contaminants in the environment and to translate this knowledge into regulatory actions. While individual studies on specific contaminants are essential to decipher their behavior and impacts, a holistic understanding of all potential contaminants within a given area becomes crucial, especially when assessing the cumulative effects of multiple contaminants on aquatic ecosystems (Relyea, 2009; Saaristo et al., 2018; Petersen et al., 2014). To this end, we conducted an inventory of potential contaminants in the coastal waters and sediments of a Mediterranean coastal agglomeration characterized by a wide array of activities. Our analysis **targeted a large number of potential contaminants on the same site, which is innovative:** 10 nonmetallic inorganic parameters, 32 dissolved metals, 5 chlorobenzenes, 19 PAHs, 10 PCBs, 40 organochlorine pesticides, 242 non-organochlorine pesticides, 1 PBBs, and 12 pharmaceuticals. To evaluate the influence of wastewater treatment plants (WWTPs) on the coastal environment, we conducted sampling at both **the WWTP entrance and exit**, as well as in the sea near the WWTP outlet. Furthermore, we conducted sediment analysis in addition to surface water analysis at three distinct sampling stations, recognizing the tendency of certain organic compounds to adsorb to solid particles.

## Material and methods

## Sample site description

Located on the French coast of the Mediterranean Sea, the Toulon Bay is bordered by a metropolis of about 450,000 inhabitants. The site has been subjected to various human activities (e.g., intense maritime traffic, commercial, military port and marina, urban, industrial and agricultural discharges) resulting in high contamination of organometallic compounds (mono-, di- and tributyltin), organic (PAHs, PCBs, pesticides, pharmaceuticals, etc.) and metals (mercury, copper, lead, cadmium, etc.) whether in the water column or in sediments. Six sampling stations were chosen, distributed inside and outside the Toulon Bay (Figure 1 and Table 1), representative of the main environmental situations observed in the area. At each station, water was sampled at 1 m depth. The REF station is a point of reference, situated upstream of Toulon with regard to the liguro-provençal current, therefore **the closest point** relatively protected from the anthropogenic impacts that affect the Toulon **area**. The Plume1, Plume2 and Plume3 stations are located in the treated water discharge plume of the Amphitria WWTP, which has a capacity of  $500 \cdot 10^3$  Population Equivalent (PE) for a reference flow of  $86 \cdot 10^3 \text{ m}^3 \text{ d}^{-1}$ . The **Plume1** station is vertical to the outlet located at a depth of 6 m, the **Plume2** and **Plume3** stations are located 150 m and 300 m, respectively, downstream in the plume. Two other WWTPs with a capacity of  $106 \cdot 10^3$  and  $122 \cdot 10^3$  PE treat the wastewater of the metropolis (Fig. 1). The Bay station is located in the middle of the Toulon Bay. The Ha station (**Harbour station**) is located within the protected port area, shielded from the waves by a dyke which limits water exchanges with the outer part of the bay. The port area has a depth of less than 20 m and a multi-contaminated sediment (Tessier et al., 2011, Pougnet et al., 2014).




Figure 1. Location of the sampling points. Each colored area **refers** to a different wastewater treatment plant. 1: Amphitria WWTP.

## Sampling

The samples at sea were taken 1 m below the surface using a NISKIN-type sampling bottle. Temperature, salinity, pH, dissolved oxygen, redox potential, turbidity and chlorophyll of seawater were measured in situ at sampling time using a multiparametric probe (Hydrolab DS5). Various procedures were carried out immediately after sampling, depending on the type of analysis required. For dissolved organic carbon (DOC), 10 mL of water was filtered through a 0.45  $\mu\text{m}$  nitrocellulose syringe filter pre-rinsed with 10%  $\text{HNO}_3$  nitric acid, then stored at 10°C in a 15 mL centrifuge tube, to which 100  $\mu\text{L}$  of 1 M  $\text{NaN}_3$  (sodium azide) was added to prevent microbial activity until analysis. For ions, 10 mL of water was filtered through the same type of syringe filter, then collected in a 15 mL centrifuge tube and stored at -20°C until analysis. For metals, 40 mL of water was filtered through the same type of syringe filter, then collected in a 50 mL centrifuge tube, acidified with 200  $\mu\text{L}$  of 70%  $\text{HNO}_3$  nitric acid and stored at 10°C until analysis. For organic molecules, 2 L of water was taken in a glass bottle, then stored at 10°C until analysis.

Wastewater was sampled at the Amphitria WWPT (station 1 on Fig. 1). Raw sewage (RS) and treated wastewater (TWW) were sampled upstream and downstream, respectively, of the WWTP to assess the % of abatement for the targeted chemicals. RS and TWW were sampled twice, once in July 2017 and again in February 2018, to investigate possible differences between summer, with high tourist load, and winter. The July 2017 sampling took place during a dry period, with more than 30 days without rain, whereas the February 2018 sampling occurred after approximately 50 mm fell over the area in the preceding 10 days. Annual species fluxes to the sea were calculated using the average TWW concentrations from the two sampling, with a concentration of 0 assigned when a species concentration was below the Limit of Reporting (LOR). The average annual volume of TWW was 18.6  $10^6 \text{ m}^3$ .

Sediment samples were collected using a Van Walt stainless steel weighted grab. After collection, they were oven dried at 45°C then sieved at 2 mm.

## Chemical analysis

Phosphates were analyzed by spectrophotometry (SHIMADZU®, UV-1800) according to the method of Murphy and Riley (1962), with the modifications of Strickland and Parson (1968) (dl 0.02  $\mu\text{M}$ ). All other analysis were performed in the certified laboratory ALS Czech Republic, Prague, using the following methods:  $\text{NH}_3$  and  $\text{NH}_4^+$  by discrete spectrophotometry;  $\text{F}^-$ ,  $\text{NO}_3^-$ ,  $\text{NO}_2^-$  and  $\text{SO}_4^{2-}$  by ion liquid chromatography; major cations Ca, P, K and Na by atomic emission spectrometry with inductively coupled plasma; other dissolved metals (excepting Hg) and metalloids by mass spectrometry with inductively coupled plasma; Hg by atomic fluorescence spectrometry; clarithromycin and fluoxetine by high-performance liquid chromatography; tamoxifen, 17 $\beta$  oestradiol and 17 $\alpha$ -ethinyloestradiol by gas chromatography-mass spectrometry; octachlorostyrene, chlorobenzenes, PAHs, PCBs, organochlorine pesticides, trifluralin, PBB163 by gas chromatography-tandem mass spectrometry; all other pesticides by liquid chromatography-tandem mass spectrometry with prior derivatization for amitrole, AMPA, glufosinate ammonium and glyphosate. Sediment total analysis was performed on the <2 mm fraction. Organics in the sediments were analyzed on a water extract at the solid/liquid ratio of 1/10 of the pulverized sample. The analyzed species are listed on Table 1.



**Table 1.** List of elements or compounds analyzed. For each of them is given the limit of reporting (LOR). The elements or compounds marked in blue had a concentration higher than the LOR for at least one of the samples analyzed.

Non metallic species					
Ammonia and ammonium ions as N (PHO)	0.040 mg/L	Ammonia and ammonium ions as N (SPC)	0.040 mg/L	Fluoride	0.200 mg/L
Nitrates	2.00 mg/L	Nitrites	0.040 mg/L	Orthophosphate as P	0.010 mg/L
Sulphate as SO <sub>4</sub> <sup>2-</sup>	5.00 mg/L				
Dissolved metals					
Ag 1.0 µg/L	Bo 10.0 µg/L	Fe 2.0 µg/L	Mo 1.0 µg/L	Se 1.0 µg/L	V 5.0 µg/L
Al 5.0 µg/L	Ca 0.005 mg/L	Hg 0.01 µg/L	Na 0.03 mg/L	Sn 1.0 µg/L	Zn 2.0 µg/L
As 1.0 µg/L	Cd 0.5 µg/L	K 0.015 mg/L	Ni 3.0 µg/L	Sr 1.0 µg/L	
Ba 1.0 µg/L	Co 0.5 µg/L	Li 1.0 µg/L	P 0.05 mg/L	Te 5.0 µg/L	
Be 0.2 µg/L	Cr 5.0 µg/L	Mg 10 µg/L	Pb 1.0 µg/L	Th 0.5 µg/L	
Bi 1.0 µg/L	Cu 1.0 µg/L	Mn 0.5 µg/L	Sb 1.0 µg/L	Ti 5.0 µg/L	
Chlorobenzenes					
1,2,3,5- & 1,2,4,5-Tetrachlorobenzene	0.004 µg/L	1,2,3,4-Tetrachlorobenzene	0.002 µg/L		
Hexachlorobenzene (HCB)	0.002 µg/L	Pentachlorobenzene	0.002 µg/L		
PAHs					
Naphthalene 0.005 µg/L	Fluoranthene 0.001 µg/L	Benzo(k)fluoranthene 0.001 µg/L			
Acenaphthylene 0.001 µg/L	Pyrene 0.001 µg/L	Benzo(a)pyrene 0.001 µg/L			
Acenaphthene 0.001 µg/L	Benz(a)anthracene 0.001 µg/L	Indeno(1.2.3.cd)pyrene 0.001 µg/L			
Fluorene 0.001 µg/L	Chrysene 0.001 µg/L	Dibenz(a,h)anthracene 0.001 µg/L			
Phenanthrene 0.001 µg/L	Benzo(b)fluoranthene 0.001 µg/L	Benzo(g,h,i)perylene 0.001 µg/L			
Anthracene 0.001 µg/L					
PCBs					
PCB 28 0.001 µg/L	PCB 101 0.001 µg/L	PCB 138 0.001 µg/L	PCB 180 0.001 µg/L		
PCB 52 0.001 µg/L	PCB 118 0.001 µg/L	PCB 153 0.001 µg/L	PCB 194 0.001 µg/L		
Organochlorine pesticides					
2,4-DDD 0.002 µg/L	Dichlobenil 0.002 µg/L	Hexachlorobutadiene 0.002 µg/L			
2,4-DDE 0.002 µg/L	Dieldrin 0.002 µg/L	Hexachloroethane 0.002 µg/L			
2,4-DDT 0.002 µg/L	Endosulfan sulfate 0.005 µg/L	Isodrin 0.002 µg/L			
4,4'-DDD 0.002 µg/L	Endrin 0.002 µg/L	Methoxychlor 0.005 µg/L			
4,4'-DDE 0.002 µg/L	Heptachlor 0.002 µg/L	Mirex 0.002 µg/L			
4,4'-DDT 0.002 µg/L	Heptachloroepoxide-cis 0.005 µg/L	Oxychlorane 0.005 µg/L			
Alachlor 0.005 µg/L	Heptachloroepoxide-trans 0.005 µg/L	Pentachlorotoluene 0.002 µg/L			
Aldrin 0.002 µg/L	Hexachlorocyclohexane Alpha 0.002 µg/L	Telodrin 0.002 µg/L			
alpha-Endosulfan 0.002 µg/L	Hexachlorocyclohexane Beta 0.002 µg/L	trans-Chlordane 0.003 µg/L			
beta-Endosulfan 0.002 µg/L	Hexachlorocyclohexane Delta 0.002 µg/L	Trans-Nonachlor 0.003 µg/L			
cis-Chlordane 0.005 µg/L	Hexachlorocyclohexane Epsilon 0.002 µg/L				
cis-Nonachlor 0.002 µg/L	Hexachlorocyclohexane Gamma 0.002 µg/L				
Pesticides					
1-(3,4-Dichlorophenyl)urea (DCPU) 0.05 µg/L	2-amino-N-(isopropyl)benzamide 0.05 µg/L	2-Chloro-2,6-diethylacetanilide 0.05 µg/L	Chlorotoluron-desmethyl 0.05 µg/L		
Acetamidid 0.05 µg/L	Azoxystrobin 0.05 µg/L	Carfentrazone-ethyl 0.05 µg/L	Cybutryne (Irgarol) 0.05 µg/L		
Acetochlor 0.05 µg/L	BAM 0.05 µg/L	Chlorbromuron 0.05 µg/L	Cymoxanil 0.05 µg/L		
Acibenzolar-S-methyl 0.05 µg/L	BDMC 0.05 µg/L	Chlorfenvinphos 0.05 µg/L	Cyprazine 0.05 µg/L		
Aclonifen 0.05 µg/L	Benalaxyl 0.05 µg/L	Chloridazon 0.05 µg/L	Cyproconazole 0.05 µg/L		
Alachlor 0.05 µg/L	Bendiocarb 0.05 µg/L	Chloridazon-desphenyl 0.05 µg/L	Cyprodinil 0.05 µg/L		
Aldicarb 0.05 µg/L	Bentazone methyl 0.05 µg/L	Chlorotoluron 0.05 µg/L	Cyromazine 0.05 µg/L		
Aldicarb sulfone 0.05 µg/L	Bifenox 0.05 µg/L	Chloroxuron 0.05 µg/L	DEET 0.05 µg/L		
Ametryn 0.05 µg/L	Bitertanol 0.05 µg/L	Chlorpropham 0.05 µg/L	Desmetryn 0.05 µg/L		
Amidosulfuron 0.05 µg/L	Boscalid 0.05 µg/L	Chlorpyrifos 0.05 µg/L	Diazinon 0.05 µg/L		
Amitrole 0.1 µg/L	Bromacil 0.05 µg/L	Chlorpyrifos-methyl 0.05 µg/L	Dichlofenthion 0.05 µg/L		
AMPA 0.05 µg/L	Bromophos-ethyl 0.05 µg/L	Chlorsulfuron 0.05 µg/L	Dichlormid 0.05 µg/L		
Atraton 0.05 µg/L	Cadusafos 0.05 µg/L	Clodinafop 0.05 µg/L	Dichlorvos 0.05 µg/L		
Atrazine 0.05 µg/L	Carbaryl 0.05 µg/L	Clomazone 0.05 µg/L	Dicrotophos 0.05 µg/L		
Atrazine-2-hydroxy 0.05 µg/L	Carbendazim 0.05 µg/L	Clomeprop 0.05 µg/L	Diethofencarb 0.05 µg/L		
Atrazine-desethyl 0.05 µg/L	Carbetamide 0.05 µg/L	Clothianidin 0.05 µg/L	Difenacoum 0.05 µg/L		
Atrazine-desisopropyl 0.05 µg/L	Carbofuran 0.05 µg/L	Coumaphos 0.05 µg/L	Difenoconazole 0.05 µg/L		
Azinphos-ethyl 0.05 µg/L	Carbofuran-3-hydroxy 0.05 µg/L	Crimidine 0.05 µg/L	Difenoxuron 0.05 µg/L		
Azinphos-methyl 0.05 µg/L	Carboxin 0.05 µg/L	Cyanazine 0.05 µg/L	Diflubenzuron 0.05 µg/L		

Pesticides							
Diflubenzuron	0.05 µg/L	Imazamox	0.05 µg/L	Neburon	0.05 µg/L	Pyribenzoxim	0.05 µg/L
Diflufenican	0.05 µg/L	Imazethapyr	0.05 µg/L	Nicosulfuron	0.05 µg/L	Pyrimethanil	0.05 µg/L
Dimefuron	0.05 µg/L	<b>Imidacloprid</b>	<b>0.05 µg/L</b>	Nuarimol	0.05 µg/L	Pyriproxifen	0.05 µg/L
Dimethachlor	0.05 µg/L	Indoxacarb	0.05 µg/L	Omethoate	0.05 µg/L	Quinclorac	0.05 µg/L
Dimethenamid	0.05 µg/L	Iprodione	0.05 µg/L	Oxadixyl	0.05 µg/L	Quinmerac	0.05 µg/L
Dimethoate	0.05 µg/L	Iprovalicarb	0.05 µg/L	Oxamyl	0.05 µg/L	Quinoxifen	0.05 µg/L
Dimethomorph	0.05 µg/L	Isoproturon	0.05 µg/L	Paclobotrazol	0.05 µg/L	Quizalofop	0.05 µg/L
Diuron	0.05 µg/L	Isoproturon-desmethyl	0.05 µg/L	Paraoxon-ethyl	0.05 µg/L	Rimsulfuron	0.05 µg/L
Diuron desmethyl (DCPMU)	0.05 µg/L	Isoproturon-monodesmethyl	0.05 µg/L	Paraoxon-methyl	0.05 µg/L	Sebuthylazine	0.05 µg/L
Epoiconazole	0.05 µg/L	Isopyrazam	0.05 µg/L	Parathion-ethyl	0.05 µg/L	Secbumeton	0.05 µg/L
EPTC	0.05 µg/L	Kresoxim-methyl	0.05 µg/L	Penconazole	0.05 µg/L	Sethoxydim	0.05 µg/L
Ethiofencarb	0.05 µg/L	Lenacil	0.05 µg/L	Pencycuron	0.05 µg/L	Simazine	0.05 µg/L
Ethion	0.05 µg/L	Linuron	0.05 µg/L	Pendimethalin	0.05 µg/L	Simazine-2-hydroxy	0.05 µg/L
Ethofumesate	0.05 µg/L	Malaoxon	0.05 µg/L	Phorate	0.05 µg/L	Simetryn	0.05 µg/L
Ethoprophos	0.05 µg/L	Malathion	0.05 µg/L	Phosalone	0.05 µg/L	Spiroxamine	0.05 µg/L
Fenamiphos	0.05 µg/L	Mandipropamid	0.05 µg/L	Phosmet	0.05 µg/L	Sulfosulfuron	0.05 µg/L
Fenarimol	0.05 µg/L	Mecarbam	0.05 µg/L	Phosphamidon	0.05 µg/L	Tebuconazole	0.05 µg/L
Fenhexamid	0.05 µg/L	Mefenpyr-diethyl	0.05 µg/L	Picloram	0.05 µg/L	Tebuthiuron	0.05 µg/L
Fenoxaprop	0.05 µg/L	Mesosulfuron-methyl	0.05 µg/L	Picoxystrobin	0.05 µg/L	Teflubenzuron	0.05 µg/L
Fenoxycarb	0.05 µg/L	Mesotrione	0.05 µg/L	Pirimicarb	0.05 µg/L	Terbuthylazine	0.05 µg/L
Fenpropidin	0.05 µg/L	Metalaxyl (isomers)	0.05 µg/L	Pirimiphos-ethyl	0.05 µg/L	Terbuthylazine-desethyl	0.05 µg/L
Fenpropimorph	0.05 µg/L	Metamitron	0.05 µg/L	Pirimiphos-methyl	0.05 µg/L	Terbuthylazine-desethyl-2-hydroxy	0.05 µg/L
Fensulfotion	0.05 µg/L	Metazachlor	0.05 µg/L	Pretilachlor	0.05 µg/L	Terbuthylazine-hydroxy	0.05 µg/L
Fenuron	0.05 µg/L	Metconazole	0.05 µg/L	Primisulfuron-methyl	0.05 µg/L	Terbutryn	0.05 µg/L
Fipronil	0.05 µg/L	Methabenzthiazuron	0.05 µg/L	Prochloraz	0.05 µg/L	Thiabendazole	0.05 µg/L
Florasulam	0.05 µg/L	Methamidophos	0.05 µg/L	Prodiamine	0.05 µg/L	Thiamethoxam	0.05 µg/L
Fluazifop	0.05 µg/L	Methidathion	0.05 µg/L	Profenofos	0.05 µg/L	Thifensulfuron-methyl	0.05 µg/L
Fluazifop-butyl (isomers)	0.05 µg/L	Methiocarb	0.05 µg/L	Promecarb	0.05 µg/L	Thiobencarb	0.05 µg/L
Flusilazole	0.05 µg/L	Methomyl	0.05 µg/L	Prometon	0.05 µg/L	Thiophanate-methyl	0.05 µg/L
Flutolanil	0.05 µg/L	Methoxyfenozide	0.05 µg/L	Prometryn	0.05 µg/L	Triadimefon	0.05 µg/L
Fonofos	0.05 µg/L	Metobromuron	0.05 µg/L	Propachlor	0.05 µg/L	Triadimenol	0.05 µg/L
Foramsulfuron	0.05 µg/L	Metolachlor (isomers)	0.05 µg/L	Propamocarb	0.05 µg/L	Tri-allate	0.05 µg/L
Furathiocarb	0.05 µg/L	Metoxuron	0.05 µg/L	Propanil	0.05 µg/L	Triasulfuron	0.05 µg/L
Glufosinate ammonium	0.1 µg/L	Metribuzin	0.05 µg/L	Propaquizafop	0.05 µg/L	Triazophos	0.05 µg/L
<b>Glyphosate</b>	<b>0.05 µg/L</b>	Metribuzin-desamino	0.05 µg/L	Propazine	0.05 µg/L	Tribenuron-methyl	0.05 µg/L
Haloxifop	0.05 µg/L	Metsulfuron-methyl	0.05 µg/L	Propham	0.05 µg/L	Tricyclazole	0.05 µg/L
Haloxifop-methyl (isomers)	0.05 µg/L	Molinate	0.05 µg/L	Propiconazole	0.05 µg/L	Trifloxysulfuron-sodium	0.05 µg/L
Hexaconazole	0.05 µg/L	Monocrotophos	0.05 µg/L	Propoxur	0.05 µg/L	Trifluralin	0.005 µg/L
Hexazinone	0.05 µg/L	Monolinuron	0.05 µg/L	Propoxycarbazone-sodium	0.05 µg/L	Triflusulfuron-methyl	0.05 µg/L
Hexythiazox	0.05 µg/L	Monuron	0.05 µg/L	Propyzamide	0.05 µg/L	Triforine	0.05 µg/L
Imazalil	0.05 µg/L	Napropamide	0.05 µg/L	Prosulfocarb	0.05 µg/L	Triticonazole	0.05 µg/L
Imazamethabenz-methyl	0.05 µg/L	Naptalam	0.05 µg/L	Prothioconazole	0.05 µg/L		

## Others

PBB 153	0.002 µg/L	Octachlorostyrene	0.003 µg/L
---------	------------	-------------------	------------

## Pharmaceuticals

<b>Carbamazepine</b>	0.01	mg/L	<b>17β-oestradiol (E2)</b>	0.0003	µg/L
Clorofibric acid	0.01	mg/L	<b>17α-ethinyloestradiol (EE2)</b>	0.00003	µg/L
<b>Clarithromycin</b>	0.001	µg/L	<b>Ibuprofen</b>	0.01	mg/L
<b>Diclofenac</b>	0.01	mg/L	Paracetamol	0.01	mg/L
<b>Fluoxetin</b>	0.0002	µg/L	<b>Propanolol</b>	0.01	mg/L
<b>Gemfibrosil</b>	0.01	mg/L	Tamoxifen	0.005	µg/L

## Results and discussion

This article discusses the concentration data for substances with concentrations above the limit of reporting (LOR) for at least one of the samples. The set of results can be consulted in the supplementary data.

### From wastewater to WWTP plume at sea

Inorganic species did not show significant differences between the two sampling periods (see Table 2). Nitrates and nitrites were absent in the WWTP inlet but present at the outlet due to nitrification of the ammonia produced by organic matter mineralization in the treatment process. Sulfates levels may also increase as a result of the oxidation of sulfides present in the raw sewage during the treatment process. Both N and P compounds were dispersed in the plume, but their concentrations were below the limit of reporting at the second point of the plume, indicating the absence of eutrophic conditions. Metals and metalloids fell within the expected range at both inlet and outlet of the WWTP. However, higher concentrations of Fe, Ca, and Mn in TWW compared to RS could be attributed to the use of flocculants during the treatment process. In the plume, concentrations of Fe and Zn increased, indicating concentrations in the seawater higher than the typical values of approximately 0.03 and 0.4  $\mu\text{g L}^{-1}$ , respectively (Bruland et al., 2014). Beryllium was not detected in sewage, TWW, or the first sample of the plume (Plume1), but it was found in the Plume2 and Plume3 samples at a concentration higher than the predicted no effect concentration (PNEC), which was evaluated at 0.08  $\mu\text{g L}^{-1}$  (Assoumain and Salomon, 2020). This non-radioactive metal is highly toxic and is classified as a category 1 carcinogen by the European Union. It can be generated by rock weathering or coal burning and can be found in various products such as cosmetics, electronic products, alloys, and ceramics use by various industries, including nuclear weapons, aeronautics, and coal mining. It can be found in manufacture goods like electronic compounds, alloys or ceramics, but beyond specialized industrial sites its occurrence in the environment is primarily associated with the alteration of rocks or the combustion of coal. However, further studies are required to fully interpret these observations.

Table 2. Nonmetallic inorganics and metals in the raw sewage (RS), the treated wastewater (TWW) and in the TWW plume at sea. LOR, limit of reporting, nd: non defined

Chemical species	Feb. 2018 sampling			Jul. 2017 sampling			Plume 1	Plume 2	Plume 3
	RS	TWW	Abatement	RS	TWW	Abatement			
<b>Nonmetallic inorganics</b>									
NH <sub>3</sub> and NH <sub>4</sub> <sup>+</sup> as N (mg L <sup>-1</sup> )	65.3	50.5	23 %	64.5	5	92 %	0.301	<LOR	<LOR
Nitrates (mg L <sup>-1</sup> )	<LOR	10.2	nd	<LOR	163	nd	<LOR	<LOR	<LOR
Nitrites (mg L <sup>-1</sup> )	<LOR	3.2	nd	<LOR	2.17	nd	<LOR	<LOR	<LOR
Orthophosphate as P (mg L <sup>-1</sup> )	6.21	1.57	75 %	5.68	2.05	64 %	0.018	<LOR	<LOR
Sulfate (mg L <sup>-1</sup> )	287	345	-20 %	379	363	4 %	2780	2830	2810
<b>Metals</b>									
Al (µg L <sup>-1</sup> )	53.5	<LOR	nd	51	<LOR	nd	<LOR	<LOR	<LOR
Ba (µg L <sup>-1</sup> )	32	14	56 %	32	14.9	53 %	<LOR	<LOR	<LOR
Be (µg L <sup>-1</sup> )	< LOR	< LOR	nd	<LOR	<LOR	nd	<LOR	4.02	4.72
B (µg L <sup>-1</sup> )	586	638	-9 %	682	596	13 %	3960	3950	4160
Ca (mg L <sup>-1</sup> )	116	140	-21 %	134	145	-8 %	419	461	413
Cu (µg L <sup>-1</sup> )	< LOR	3	nd	<LOR	<LOR	nd	47.3	<LOR	<LOR
Fe (µg L <sup>-1</sup> )	55.3	156	-182 %	208	37.6	82 %	146	77.6	131
Hg (µg L <sup>-1</sup> )	< LOR	< LOR	nd	<LOR	0.113	nd	<LOR	<LOR	<LOR
K (mg L <sup>-1</sup> )	69.3	64.6	7 %	77.2	74.7	3 %	485	508	496
Li (µg L <sup>-1</sup> )	26.7	25.6	4 %	30.9	29.5	5 %	164	162	185
Mg (mg L <sup>-1</sup> )	142	146	-3 %	162	167	-3 %	1430	1360	1350
Mn (µg L <sup>-1</sup> )	41.1	76.6	-86 %	607	421	31 %	21.8	19.6	21.6
Na (g L <sup>-1</sup> )	1.1	1.0	3 %	1.4	1.4	-5 %	11.9	11.4	11.8
P (mg L <sup>-1</sup> )	6.1	1.73	72 %	5.52	1.9	66 %	<LOR	<LOR	<LOR
Se (µg L <sup>-1</sup> )	13.1	< LOR	nd	< LOR	< LOR	nd	< LOR	< LOR	< LOR
Sr (µg L <sup>-1</sup> )	1190	1200	-1 %	1430	1430	0 %	9120	8580	8670
Zn (µg L <sup>-1</sup> )	11.5	11.5	0 %	<LOR	17	nd	92.2	89	162

Table 3 presents the concentrations of polycyclic aromatic hydrocarbons (PAHs) and polychlorinated biphenyls (PCBs). Out of the 16 PAHs examined, 15 were detected above the LOR. High concentrations of phenanthrene, fluoranthene, pyrene and chrysene were found in the samples from July, with values exceeding 0.02 µg L<sup>-1</sup>. PAH concentrations in untreated wastewater exhibited a substantial increase in July 2017 compared to February 2018. Although forest fires occurred in the Toulon region before the sampling period in July 2017, it is challenging to ascertain how the generated PAHs could have entered the sewer system. A more plausible hypothesis involves a significant surge in backyard barbecues during the summer in the Toulon region, notably driven by the substantial rise in population associated with tourism. These barbecues emit PAHs, which may find their way into the sewer system through the washing of dishes and utensils. The WWTP demonstrated varying degrees of effectiveness in reducing PAH concentrations, with an abatement ranging from 32% (acenaphthene) to 78% (phenanthrene). In February, the raw sewage exhibited considerably lower concentrations compared to July, except for naphthalene and acenaphthylene. This discrepancy may account for the observed 100% abatement for all PAHs in February. As the plume extended, all PAH concentrations became diluted, dropping below the LOR at the very first point in the plume.

Table 3. PAHS and PCBs in the raw sewage (RS), the treated wastewater (TWW) and in the TWW plume at sea. LOR, limit of reporting, nd: non defined

Chemical species	Feb. 2018 sampling			Jul. 2017 sampling			Plume 1	Plume 2	Plume 3
	RS	TWW	Abatement	RS	TWW	Abatement			
<b>PAHs (ng L<sup>-1</sup>)</b>									
Naphtalene	34	<LOR	100 %	<LOR	<LOR	nd	<LOR	<LOR	<LOR
Acenaphthylene	1.6	<LOR	100 %	<LOR	<LOR	nd	<LOR	<LOR	<LOR
Acenaphthene	<LOR	<LOR	nd	3.4	2.3	32 %	<LOR	<LOR	<LOR
Fluorene	5.6	<LOR	100 %	7.1	3.3	54 %	<LOR	<LOR	<LOR
Phenanthrene	15.5	<LOR	100 %	32.6	7.3	78 %	<LOR	<LOR	<LOR
Anthracene	<LOR	<LOR	nd	5.5	2.1	62 %	<LOR	<LOR	<LOR
Fluoranthene	1.2	<LOR	100 %	37.3	14.9	60 %	<LOR	<LOR	<LOR
Pyrene	13.9	<LOR	100 %	35.0	13.2	62 %	<LOR	<LOR	<LOR
Benz(a)anthracene	2.6	<LOR	100 %	16.4	8.4	49 %	<LOR	<LOR	<LOR
Chrysene	6.9	<LOR	100 %	24.3	15.3	37 %	<LOR	<LOR	<LOR
Benzo(b)fluoranthene	3.9	<LOR	100 %	15.2	7.0	54 %	<LOR	<LOR	<LOR
Benzo(k)fluoranthene	1.8	<LOR	100 %	10.0	4.3	57 %	<LOR	<LOR	<LOR
Benzo(a)pyrene	2.6	<LOR	100 %	9.2	3.3	64 %	<LOR	<LOR	<LOR
Indeno(1.2.3.cd)pyrene	1.4	<LOR	100 %	6.6	2.37	59 %	<LOR	<LOR	<LOR
Benzo(g,h,i)perylene	2.5	<LOR	100 %	7.7	2.7	65 %	<LOR	<LOR	<LOR
<b>PCBs (ng L<sup>-1</sup>)</b>									
PCB 28	<LOR	<LOR	nd	14	1	90 %	<LOR	<LOR	<LOR
PCB 138	1.8	<LOR	100 %	<LOR	<LOR	nd	<LOR	<LOR	<LOR
PCB 153	1.7	<LOR	100 %	<LOR	<LOR	nd	<LOR	<LOR	<LOR

Out of the eight polychlorinated biphenyls (PCBs) analyzed, three were found at concentrations above the LOR but at low levels: PCB 28 was detected in July, while PCB 138 and 153 were detected in February (Table 3). PCB 28 exhibited a 90% **abatement** during wastewater treatment, while the other two were completely eliminated. At all plume sampling points, concentrations of all PCBs were below the LOR.

Regarding organochlorines and other pesticides, out of the 279 compounds analyzed, four exceeded the LOR in July 2017, and 15 exceeded it in February 2018 (Table 4). The higher number of compounds detected in February 2018 is likely due to heavy rainfall in the days preceding the sampling, resulting in the overflow of rainwater from agricultural areas into the sewer network. **Pesticides have been detected in wastewater (Sutton et al., 2019; Budd et al., 2023), yet the mechanisms facilitating their entry into sewers remain inadequately characterized (Knight et al., 2023). In the context of the July 2017 and February 2018 samplings, a notable distinction lies in the heavy rainfall observed in the days leading up to the latter sampling. This precipitation may have triggered runoff from agricultural areas, causing an overflow into the sewer system and introducing professional herbicides like imazamox, propyzamide, or terbuthryn.** While most of these substances were present at low concentrations, three stood out: glyphosate, AMPA, and DEET. Glyphosate is the active herbicide ingredient in Roundup®, commonly used by both professionals and individuals. AMPA (aminomethylphosphonic acid) is one of its degradation residues. In raw sewage, concentrations of glyphosate and AMPA were higher in July 2017

compared to February 2018. This can be attributed to the intensive use of the herbicide during the period of vegetation growth, especially in areas where the separation between the stormwater network and the sewer network is not perfect. DEET (N,N-diethyl-3-methylbenzamide) is a widely used mosquito repellent. The higher concentration of DEET in raw sewage in July 2018 can be explained by its increased usage during the summer months.

The **abatement** of organochlorines and other pesticides ranged from 8% for acetamiprid (an insecticide) to 47% for propamocarb (a fungicide). The few negative **abatements** can be attributed to low concentrations near the detection limit, resulting in higher relative uncertainties. In the plume samples, all pesticide concentrations were below the LOR, except for AMPA in the Plume 1 sample.

Table 4. Organochlorine and other pesticides in the raw sewage (RS), the treated wastewater (TWW) and in the TWW plume at sea. PNEC, Predicted No Effect Concentration; LOR, limit of reporting; nd: non defined

Chemical species	Feb. 2018 sampling			Jul. 2017 sampling			Plume 1	Plume 2	Plume 3
	RS	TWW	Abatement	RS	TWW	Abatement			
<b>Organochlorine pesticides (<math>\mu\text{g L}^{-1}</math>)</b>									
Heptachloroepoxide-cis	0.164	<LOR	nd	<LOR	<LOR	nd	<LOR	<LOR	<LOR
Heptachloroepoxide- $\gamma$	0.0056	0.0046	18 %	<LOR	<LOR	nd	<LOR	<LOR	<LOR
<b>Pesticides (<math>\mu\text{g L}^{-1}</math>)</b>									
Acetamiprid	0.013	0.012	8 %	<LOR	<LOR	nd	<LOR	<LOR	<LOR
AMPA	3.66	3.77	-3 %	12.9	9.28	28 %	0.077	<LOR	<LOR
Boscalid	0.013	0.011	15 %	<LOR	<LOR	nd	<LOR	<LOR	<LOR
DEET	1.05	0.795	24 %	3.77	2.31	39 %	<LOR	<LOR	<LOR
Diuron	0.012	0.014	-8 %	<LOR	<LOR	nd	<LOR	<LOR	<LOR
Glyphosate	0.50	0.40	20 %	1.19	0.7	41 %	<LOR	<LOR	<LOR
Imazamox	0.033	<LOR	100 %	<LOR	<LOR	nd	<LOR	<LOR	<LOR
Imidacloprid	<LOR	0.054	nd	0.09	0.13	-46 %	<LOR	<LOR	<LOR
Propamocarb	0.099	0.053	47 %	<LOR	<LOR	nd	<LOR	<LOR	<LOR
Propyzamid	0.014	0.020	-15 %	<LOR	<LOR	nd	<LOR	<LOR	<LOR
Terbutryn	0.051	0.056	10 %	<LOR	<LOR	nd	<LOR	<LOR	<LOR

Among the 12 pharmaceutical substances analyzed, 10 were detected at concentrations above the detection threshold, exhibiting variable **% of abatements** between July 2017 and February 2018 (Table 5). Negative **abatements** can be attributed to low concentrations or fluctuations in the input of raw sewage into the WWTP. No definitive conclusion can be drawn for gemfibrozil because its PNEC is lower than the LOR.

Concentrations of clarithromycin, fluoxetine,  $17\beta$ -estradiol, diclofenac, and carbamazepine in TWW exceeded their respective PNECs. However, in the plume, the dilution effect reduced the concentrations below the LOR and, consequently, below the PNEC for all of these substances, except for  $17\alpha$ -ethinylestradiol, which remained above the PNEC in all plume samples. The efficient removal of ibuprofen and paracetamol in the WWTP aligns with findings reported in the literature (Beausse, 2004). The high concentration of paracetamol observed in the February 2018 sample may be attributed to a

peak in influenza activity during that period (Bernard-Stoecklin, 2018). The limited % of abatement of diclofenac and carbamazepine in the WWTP is consistent with previous studies (Zhang et al., 2008).

Table 5. Pharmaceuticals in the raw sewage (RS), the treated wastewater (TWW) and in the TWW plume at sea. PNEC, Predicted No Effect Concentration; LOR, limit of reporting; nd: non defined. <sup>(1)</sup>: Bengtsson-Palme and Larsson 2016; <sup>(2)</sup>: Petrie et al. 2016; <sup>(3)</sup>: EU-CID 2018, Klaic and Jirsa 2022; <sup>(4)</sup>: Gamarra et al. 2015 ; <sup>(5)</sup>: Heye et al. 2019; <sup>(6)</sup>: Calopulo et al. 2018; <sup>(7)</sup>: Ferrari et al. 2003; <sup>(8)</sup>: Orias et al. 2015.

Chemical species	PNEC	Feb. 2018 sampling			Jul. 2017 sampling			Plume 1	Plume 2	Plume 3
		RS	TWW	Abatement	RS	TWW	Abatement			
<b>Pharmaceuticals (<math>\mu\text{g L}^{-1}</math>)</b>										
Clarithromycin	0.13 <sup>(1)</sup>	0.373	0.381	-2 %	0.095	0.145	-53 %	<LOR	<LOR	<LOR
Fluoxetine	0.012 <sup>(2)</sup>	0.0606	0.0404	33 %	0.0041	0.0129	-215 %	<LOR	<LOR	<LOR
17 $\beta$ -estradiol	0.0004 <sup>(3)</sup>	0.0177	0.0001	99 %	0.0166	0.0019	89 %	<LOR	<LOR	<LOR
17 $\alpha$ -ethinylestradiol	35 10 <sup>-6(3)</sup>	383 10 <sup>-6</sup>	256 10 <sup>-6</sup>	32 %	82 10 <sup>-6</sup>	116 10 <sup>-6</sup>	-41 %	196 10 <sup>-6</sup>	98 10 <sup>-6</sup>	88 10 <sup>-6</sup>
Diclofenac	10 <sup>(4)</sup>	2.04	2.06	-1 %	1.104	1.261	-14 %	<LOR	<LOR	<LOR
Carbamazepine	8 <sup>(5)</sup>	nd	nd	nd	0.376	0.745	-98 %	<LOR	<LOR	<LOR
Propranolol	0.0383 <sup>(6)</sup>	0.362	0.355	2 %	0.127	0.1841	-45 %	<LOR	<LOR	<LOR
Gemfibrozil	0.0043 <sup>(6)</sup>	<LOR	<LOR	nd	0.08	<LOR	100 %	<LOR	<LOR	<LOR
Ibuprofen	7.1 <sup>(4)</sup>	8.38	1.32	84 %	1.34	<LOR	100 %	<LOR	<LOR	<LOR
Paracetamol	13 <sup>(3)</sup>	983.3	<LOR	100 %	<LOR	<LOR	nd	<LOR	<LOR	<LOR
Clofibric acid	21 <sup>(7)</sup>	<LOR	<LOR	nd	<LOR	<LOR	nd	<LOR	<LOR	<LOR
Tamoxifen	0.081 <sup>(8)</sup>	<LOR	<LOR	nd	<LOR	<LOR	nd	<LOR	<LOR	<LOR

Table 6 presents the annual fluxes of species from the WWTP to the sea. While all potentially contaminating species, except for 17 $\alpha$ -ethinylestradiol, were below the LOR in the WWTP plume, the annual fluxes carried by the TWW effluent could still have environmental implications. For instance, the fluxes of Cu and Zn from the WWTP are of comparable magnitude (27.9 and 265 kg per year, respectively) to those contributed by a small coastal river in the Bay of Toulon (7.5 and 42.3 kg per year as dissolved species and 88 and 401 kg per year as particulate species, respectively) (Nicolau et al., 2012). This highlights the potential impact of these metals in the marine environment.

Regarding organic micropollutants, their potential impact on human health can be attributed to both a high characterization factor and a substantial emitted mass (Aemig et al., 2021). Our findings address the lack of data regarding the latter parameter, i.e., the mass released into the environment, which is crucial for accurate machine learning-based modeling of environmental impacts (Servien et al., 2022).

Table 6.

Chemical species	kg y <sup>-1</sup>	Chemical species	kg y <sup>-1</sup>	Chemical species	kg y <sup>-1</sup>
<b>Nonmetallic inorganics</b>		<b>PAHs</b>		<b>Pesticides</b>	
NH <sub>3</sub> and NH <sub>4</sub> <sup>+</sup> as N	516 10 <sup>3</sup>	Acenaphthene	0,02139	Acetamiprid	0,112
Nitrates	1611 10 <sup>3</sup>	Fluorene	0,03069	AMPA	121
Nitrites	49.9 10 <sup>3</sup>	Phenanthrene	0,06789	Boscalid	0,102
Orthophosphate as P	33.6 10 <sup>3</sup>	Anthracene	0,01953	DEET	28,9
Sulfate	6584 10 <sup>3</sup>	Fluoranthene	0,13857	Diuron	0,130
<b>Metals</b>		Pyrene	0,12276	Glyphosate	10,2
Ba	269	Benz(a)anthracene	0,07812	Hexachlorocyclohexane-γ	0,043
B	11,5 10 <sup>3</sup>	Chrysene	0,14229	Imidacloprid	1,71
Ca	2651 10 <sup>3</sup>	Benzo(b)fluoranthene	0,06510	Propamocarb	0,493
Cu	27,9	Benzo(k)fluoranthene	0,03999	Propyzamid	0,186
Fe	1.80 10 <sup>3</sup>	Benzo(a)pyrene	0,03069	Terbutryn	0,521
Hg	1,05	Indeno(1.2.3.cd)pyrene	0,02204	<b>Pharmaceuticals</b>	
K	1295 10 <sup>3</sup>	Benzo(g,h,i)perylene	0,02511	Clarithromycin	4,89
Li	512	<b>PCBs</b>		Fluoxetine	0,496
Mg	2911 10 <sup>3</sup>	PCB 28	0,0093	17β-estradiol	0,017
Mn	4.63 10 <sup>3</sup>	<b>Organochlorine pesticides</b>		17α-ethinylestradiol	0,00346
Na	22.32 10 <sup>6</sup>	Heptachloroepoxide-γ	0,04278	Diclofenac	30,9
P	33.8 10 <sup>3</sup>			Carbamazepine	6,93
Sr	24.5 10 <sup>3</sup>			Propranolol	5,01
Zn	265			Ibuprofen	12,3

## Seawater

Table 6 presents the results obtained from the Harbor (Ha), Toulon Bay (Bay), and reference (REF) stations, with a reminder for comparison of the most diluted WWTP plume station (PLUME3).

For nonmetallic inorganics and most metals (Table 6), the concentrations fell within the expected range for seawater. However, beryllium was observed at a high concentration in the Bay sample, although it remained below the limit of reporting (LOR) in the REF and Ha samples. This underscores the need for additional dedicated studies to better understand its occurrence. Iron and zinc concentrations were also significantly higher than their respective seawater solubility values, typically in the nanomolar range. These observations exceed what would be expected from colloidal aerosol deposition (Aguilar-Islas et al., 2010), highlighting the necessity for further investigations regarding these elements.

Among all the organic compounds analyzed (PAHs, PCBs, pesticides, and pharmaceuticals), only 17α-ethinylestradiol was found at levels exceeding the LOR. This was observed across all sea stations, including the reference station. This indicates that the studied area is relatively less impacted by organic contaminants compared to other heavily influenced coastal environments (Togola, 2018; Kang et al., 2022; Wang et al., 2022). However, it is important to note that 17α-ethinylestradiol concentrations exceeded the PNEC of  $35 \times 10^{-3} \text{ ng L}^{-1}$  at all sea stations. The presence of elevated concentrations at the reference station can be attributed to the transportation of WWTP effluents from small cities located east of Toulon by the liguro-provençal current, which flows from east to west.. Although the recorded



concentrations are in the lower range of those found in contaminated coastal areas like estuaries, lagoons, or semi-enclosed seas (Almeida et al., 2020), the observed values underscore the widespread contamination of coastal environments by this estrogen, which can potentially impact reproductive function and energy metabolism.

Table 6. Nonmetallic inorganics, metals and organic species with at least one value above the LOR in sea stations and in the last TWW plume (Plume3), Jul. 2017 sampling. LOR, limit of reporting, nd: non defined

Chemical species	LOR	Ha	Bay	REF	Plume3
<b>Nonmetallic inorganics</b>					
Sulfate (mg L <sup>-1</sup> )	5.00	2800	2840	2640	2810
<b>Metals</b>					
Be (µg L <sup>-1</sup> )	0.20	<LOR	6.09	<LOR	4.72
B (µg L <sup>-1</sup> )	10.0	4980	4130	3800	4160
Ca (mg L <sup>-1</sup> )	0.005	411	414	436	413
Fe (µg L <sup>-1</sup> )	2.00	<LOR	191	153	131
K (mg L <sup>-1</sup> )	0.015	489	501	509	496
Li (µg L <sup>-1</sup> )	1.00	166	179	152	185
Mg (mg L <sup>-1</sup> )	10.0	1420	1500	1390	1350
Mn (µg L <sup>-1</sup> )	0.5	18.2	17.7	20.6	21.6
Na (g L <sup>-1</sup> )	0.03	11.7	11.5	11.8	11.8
Sr (µg L <sup>-1</sup> )	1.00	8200	9530	8900	8670
Zn (µg L <sup>-1</sup> )	2.00	44.8	117	49.9	162
<b>Organics (ng L<sup>-1</sup>)</b>					
17α-ethinylestradiol	30 10 <sup>-3</sup>	69 10 <sup>-3</sup>	97 10 <sup>-3</sup>	443 10 <sup>-3</sup>	88 10 <sup>-3</sup>

## Sediments

Table 7 provides the concentrations of the species of interest in sediment samples that exceeded the limit of reporting (LOR), excluding iron which was not analyzed. **The sediment samples collected beneath the WWTP plume exhibited comparable contents of elements (metals and metalloids). Only two elements, arsenic (As) and nickel (Ni), slightly exceeded the threshold effect level (TEL). The high manganese (Mn) content can be attributed to elevated concentrations in the plume. In comparison to the plume sample, the Ha sample had higher contents of chromium (Cr), copper (Cu), mercury (Hg), and lead (Pb), with Hg and Pb concentrations surpassing the probable effect level (PEL) and TEL, respectively. This indicates historical contamination of harbor sediments due to military activities since the Second World War (Tessier et al., 2011).**

Table 7. Content of species of interest with at least one value above the LOR in the sediments. Fe was not analyzed. TEL: threshold effect level; PEL: probable effect level (Buchman, 2008).

Chemical species	Plume1	Plume2	Ha	TEL	PEL
<b>Metals (mg kg<sup>-1</sup>)</b>					
Al	>17300	>17300	5570	/	/
As	14	14	8	7.2	41.6
Cr	22	22	12	52.3	160
Cu	13	10	33	18.7	108
Hg	0.023	nd	0.894	0.13	0.70
Li	48	56	9	/	
Mn	945	654	67	/	
Ni	24	26	9	15.9	42.8
Pb	11	7	32	30.2	112
Sr	69	198	521	/	
Ti	467	163	56	/	
V	22	23	16	/	
Zn	70	70	51	124	271

None of the organic compounds of interest were identified at concentrations above the LOR in the studied stations. This suggests that inputs from the WWTP, coastal rivers, and runoff are sufficiently dispersed or degraded to prevent accumulation in the sediments. Acidic compounds (such as aspirin, ibuprofen, ketoprofen, naproxen, diclofenac, indometacin) with pKa values ranging from 4.9 to 4.1, as well as clofibric acid and gemfibrozil, are expected to be primarily present in the water column before undergoing gradual degradation at their own rates. More hydrophobic compounds like 17 $\beta$ -estradiol are prone to adsorb onto mineral surfaces, but their detection was not observed in this study. This could be attributed to the limitations of the water extraction method chosen for assessing compound bioavailability, necessitating solvent extraction for further investigations.

## Conclusion

The species of interest in this study, i.e. the species detected in at least one of the samples, included 7 nonmetallic inorganics, 32 metals and metalloids, 4 chlorobenzenes, 16 PAHs, 8 PCBs, 34 organochloride pesticides and 239 other pesticides or pesticide metabolites, 12 pharmaceuticals, and 2 other compounds.

The concentrations of nonmetallic inorganics, metals, and metalloids fell within the expected range in both raw sewage at the WWTP inlet and treated wastewater at the WWTP outlet. Among the organic compounds, 13 PAHs, 3 PCBs, 14 pesticides and pesticide metabolites (including AMPA), and 9 pharmaceuticals were detected in raw sewage. Most of these species were also detected in treated wastewater. The **abatement efficiencies** were favorable for PCBs, ranging from 32% to 78% for PAHs, but less effective for pesticides and pharmaceuticals. However, in the treated wastewater plume at sea, the dilution effect led to concentrations of all organic compounds falling below the limit of reporting (LOR), and thus, in the case of pharmaceuticals, below their predicted no effect concentration (PNEC),

except for 17 $\alpha$ -ethinylestradiol. At the three offshore stations located far from the WWTP outlets, 17 $\alpha$ -ethinylestradiol was the only organic compound of interest detected, and its concentration exceeded the PNEC. This finding underscores the insufficient reduction of endocrine disruptors by current wastewater treatment.

Furthermore, while the concentrations of contaminants were found to be low in the treated wastewater plume, the annual masses emitted by the WWTP could still have an environmental impact. In addition to pharmaceuticals, significant annual emissions were observed for AMPA, DEET, and glyphosate (121 kg y<sup>-1</sup>, 30 kg y<sup>-1</sup>, and 10 kg y<sup>-1</sup>, respectively). Sediment analysis did not reveal any contamination related to the treatment plants, but the sediment in the port area exhibited contamination with Hg and Pb, likely associated with long-term military activities in the port.

## Conflict of interest

The authors declare that they have no conflict of interest.

## References

- Aemig Q, Hélias A, Patureau D (2021) Impact assessment of a large panel of organic and inorganic micropollutants released by wastewater treatment plants at the scale of France. *Wat Res* 188:116524. <https://doi.org/10.1016/j.watres.2020.116524>
- Aguilar-Islas AM, Wu J, Rember R, Johansen AM, Shank LM (2010) Dissolution of aerosol-derived iron in seawater: Leach solution chemistry, aerosol type, and colloidal iron fraction. *Mar Chem* 120:25-33. <https://doi.org/10.1016/j.marchem.2009.01.011>
- Almeida Â, Silva MG, Soares AM, Freitas R (2020) Concentrations levels and effects of 17 $\alpha$ -Ethinylestradiol in freshwater and marine waters and bivalves: A review. *Env Res* 185:109316. <https://doi.org/10.1016/j.envres.2020.109316>
- Annett R, Habibi HR, Hontela A (2014) Impact of glyphosate and glyphosate-based herbicides on the freshwater environment. *J Appl Toxicol* 34:458-479. <https://doi.org/10.1002/jat.2997>
- Assoumain A, Salomon M (2021) Substances pertinentes à surveiller (SPAS) dans les eaux de surface. Bilan des données de surveillance acquises de 2016 à 2018 pour l'eau et le sédiment. INERIS 181881-v3.0, Verneuil-en-Halatte.
- Beausse J (2004) Selected drugs in solid matrices: a review of environmental determination, occurrence and properties of principal substances. *Trends Anal Chem* 23:753-761. <https://doi.org/10.1016/j.trac.2004.08.005>

- Bengtsson-Palme J, Larsson DGJ (2016) Concentrations of antibiotics predicted to select for resistant bacteria: Proposed limits for environmental regulation. *Env Internat* 86:140-149. <https://doi.org/10.1016/j.envint.2015.10.015>
- Bento CPM, Naumann T, Wittmann A, Tang J, Zhen X, Liu L, Ebinghaus R (2021) River-Sea Systems: Spatial and temporal occurrence of Neonicotinoids, Glyphosate and related transformation products in the Chinese Bohai Sea and 36 surrounding Rivers. EGU21-13296 Copernicus Meetings. <https://doi.org/10.5194/egusphere-egu21-13296>
- Bernard-Stoecklin S, Campèse C, Savitch Y et al (2018) Surveillance de la grippe en France, saison 2017-2018. *Bulletin Épidémiologique Hebdomadaire-BEH* 34:664-674.
- Bruland KW, Middag R and Lohan, MC (2013) Controls of Trace Metals in Seawater. In: Mottl MJ, Elderfield H (eds) *Treatise on Geochemistry: Second Edition*. Elsevier, Philadelphia, pp 19-51. <https://doi.org/10.1016/B978-0-08-095975-7.00602-1>
- Budd R, Teerlink J, Alaimo C, Wong L, Young TM (2023) Sub-sewershed Monitoring to Elucidate Down-the-Drain Pesticide Sources. *Env Sci Technol* 57:5404-5413. <https://doi.org/10.1021/acs.est.2c07443>**
- Buchman MF (2008) NOAA Screening Quick Reference Tables (SQuiRTs). NOAA OR&R Report 08-1. [https://repository.library.noaa.gov/view/noaa/9327/noaa\\_9327\\_DS1.pdf](https://repository.library.noaa.gov/view/noaa/9327/noaa_9327_DS1.pdf). Accessed 11 May 2023.
- Capolupo M, Díaz-Garduño B, Martín-Díaz ML (2018) The impact of propranolol, 17 $\alpha$ -ethinylestradiol, and gemfibrozil on early life stages of marine organisms: effects and risk assessment. *Env Sci Poll Res* 25:32196-32209. <https://doi.org/10.1007/s11356-018-3185-6>
- CEC (2000) Directive of the European Parliament and of the Council establishing a framework for Community action in the field of water policy: Joint text approved by the Conciliation Committee. Commission of the European Communities, 1997/0067(COD) C5-0347/00.
- Creel L (2003) Ripple effects: population and coastal regions. Population reference bureau, Washington DC, pp 1-7.
- Daouk S, Frege C, Blanc N, Mounier S, Redon R, Merdy P, Lucas Y, Pfeifer HR (2015) Fluorescence spectroscopy to study dissolved organic matter interactions with agrochemicals applied in Swiss vineyards. *Env Sci Poll Res* 22:9284-9292. <https://doi.org/10.1007/s11356-015-4086-6>
- Dollinger J, Dagès C, Negro S, Bailly JS, Voltz M (2016) Variability of glyphosate and diuron sorption capacities of ditch beds determined using new indicator-based methods. *Sci Tot Env* 573:716-726. <https://doi.org/10.1016/j.scitotenv.2016.08.168>
- EU-CDI (2013) Directive 2013/39/EU of the European Parliament and of the Council of 12 August 2013 amending Directives 2000/60/EC and 2008/105/EC as regards priority substances in the field of water policy. *Off J Eur Union* L226, pp. 1-17.
- EU-CDI (2018) Commission implementing decision (EU) 2018/840 of 5 June 2018 establishing a watch list of substances for Union-wide monitoring in the field of water policy pursuant to Directive

- 2008/105/EC of the European Parliament and of the Council and repealing Commission Implementing Decision (EU) 2015/495, European Parliament.
- Feltracco M, Barbaro E, Morabito E, Zangrando R, Piazza R, Barbante C, Gambaro A (2021) Assessing glyphosate in water, marine particulate matter, and sediments in the Lagoon of Venice. *Env Sci Poll Res* 29:16383–16391. <https://doi.org/10.1007/s11356-021-16957-x>
- Fent K Weston A, Caminada D (2006) Ecotoxicology of Human Pharmaceuticals. *Aquat Toxicol* 76:122-159. <https://doi.org/10.1016/j.aquatox.2005.09.009>
- Ferrari B, Paxéus N, Giudice RL, Pollio A, Garric J (2003) Ecotoxicological impact of pharmaceuticals found in treated wastewaters: study of carbamazepine, clofibric acid, and diclofenac. *Ecotoxicol Env Safety* 55:359-370. [https://doi.org/10.1016/S0147-6513\(02\)00082-9](https://doi.org/10.1016/S0147-6513(02)00082-9)
- Gamarra Jr JS, Godoi AFL, de Vasconcelos EC, de Souza KMT, de Oliveira CMR (2015) Environmental Risk Assessment (ERA) of diclofenac and ibuprofen: A public health perspective. *Chemosphere* 120:462-469. <https://doi.org/10.1016/j.chemosphere.2014.08.020>
- Gaw S Thomas K, Hutchinson T (2014) Sources, impacts and trends of pharmaceuticals in the marine and coastal environment. *Phil Trans Roy Soc B: Biol Sci.* 369:20130572. [10.1098/rstb.2013.0572](https://doi.org/10.1098/rstb.2013.0572)
- Heye K, Wiebusch J, Becker J, Rongstock L, Bröder K, Wick A, Schulte-Oehlmann U, Oehlmann J (2019) Ecotoxicological characterization of the antiepileptic drug carbamazepine using eight aquatic species: baseline study for future higher tier tests. *J Env Sci Health Part A* 54:441-451. <https://doi.org/10.1080/10934529.2018.1562819>
- ICSU-ISSC (2015) Review of the sustainable development goals: the science perspective. International Council for Science (ICSU), Paris.
- INERIS (2018) NORMAN: un bilan après 10 ans d'existence, DRC-18-158374-04175A, INERIS, Verneuil-en-Halatte
- JO (2017) Arrêté du 24/08/2017 modifiant dans une série d'arrêtés ministériels les dispositions relatives aux rejets de substances dangereuses dans l'eau en provenance des installations classées pour la protection de l'environnement. *JO Rep Fra* 234. <https://aida.ineris.fr/reglementation/arrete-240817-modifiant-serie-darretes-ministeriels-dispositions-relatives-rejets>. Accessed 11 October 2022
- Kang Y, Zhang R, Yu K, Han M, Wang Y, Huang X, Wang R, Liu F (2022) First report of organochlorine pesticides (OCPs) in coral tissues and the surrounding air-seawater system from the South China sea: distribution, source, and environmental fate. *Chemosphere* 286:131711. <https://doi.org/10.1016/j.chemosphere.2021.131711>
- Klaic M, Jirsa F (2022) 17 $\alpha$ -Ethinylestradiol (EE2): concentrations in the environment and methods for wastewater treatment – an update. *RSC Adv* 12:12794-12805. <https://doi.org/10.1039/D2RA00915C>
- Knights ER, Verhagen R, Mueller JF, Tschärke BJ (2023) Spatial and temporal trends of 64 pesticides and their removal from Australian wastewater. *Sci Tot Env* 905:166816. <https://doi.org/10.1016/j.scitotenv.2023.166816>**

- Lu Y, Yuan J, Lu X, Su C, Zhang Y, Wang C, Cao X, Li Q, Su J, Ittekkot V, Garbutt RAA, Bush S, Fletcher S, Wagey T, Kachur A, Sweijd N (2018) Major threats of pollution and climate change to global coastal ecosystems and enhanced management for sustainability. *Env Poll* 239:670-680. <https://doi.org/10.1016/j.envpol.2018.04.016>
- Madikizela LM, Ncube S, Tutu H, Richards H, Newman B, Ndungu K, Chimuka L (2020) Pharmaceuticals and their metabolites in the marine environment: sources, analytical methods and occurrence. *Trends Env Anal Chem* 28:e00104. <https://doi.org/10.1016/j.teac.2020.e00104>
- Merkens JL, Reimann L, Hinkel J, Vafeidis AT (2016) Gridded population projections for the coastal zone under the Shared Socioeconomic Pathways. *Global Planet Change* 145:57-66. <https://doi.org/10.1016/j.gloplacha.2016.08.009>
- Murphy JAMES, Riley JP (1962) A modified single solution method for the determination of phosphate in natural waters. *Anal Chim Acta* 27:31-36. [https://doi.org/10.1016/S0003-2670\(00\)88444-5](https://doi.org/10.1016/S0003-2670(00)88444-5)
- Nicolau R, Lucas Y, Merdy P, Raynaud M (2012) Base flow and stormwater net fluxes of carbon and trace metals to the Mediterranean Sea by an urbanized small river. *Wat Res* 46:6625-6637. <https://doi.org/10.1016/j.watres.2012.01.031>
- Orias F, Bony S, Devaux A, Durrieu C, Aubrat M, Hombert T, Wigh A, Perrodin Y (2015) Tamoxifen ecotoxicity and resulting risks for aquatic ecosystems. *Chemosphere* 128:79-84. <https://doi.org/10.1016/j.chemosphere.2015.01.002>
- Petersen K, Heiaas HH, Tollefsen KE (2014) Combined effects of pharmaceuticals, personal care products, biocides and organic contaminants on the growth of *Skeletonema pseudocostatum*. *Aquat Toxicol* 150:45–54. <https://doi.org/10.1016/j.aquatox.2014.02.013>
- Petrie B, Youdan J, Barden R, Kasprzyk-Hordern B (2016) New framework to diagnose the direct disposal of prescribed drugs in wastewater—a case study of the antidepressant fluoxetine. *Env Sci Technol* 50:3781-3789. <https://doi.org/10.1021/acs.est.6b00291>
- Pougnat F, Schäfer J, Dutruch L, Garnier C, Tessier E, Dang DH, Lancelot L, Mullot JU, Lenoble V, Blanc G (2014) Sources and historical record of tin and butyl-tin species in a Mediterranean bay (Toulon Bay, France). *Env Sci Poll Res* 21:6640–6651. <https://doi.org/10.1007/s11356-014-2576-6>
- Puckowski A, Mioduszevska K, Łukaszewicz P, Borecka M, Caban M, Maszkowska J, Stepnowski P (2016) Bioaccumulation and analytics of pharmaceutical residues in the environment: A review. *J Pharm Biomed Anal* 127:232-255. <https://doi.org/10.1016/j.jpba.2016.02.049>
- Relyea RA (2009) A cocktail of contaminants: how mixtures of pesticides at low concentrations affect aquatic communities. *Oecologia* 159:363-376. <https://doi.org/10.1007/s00442-008-1213-9>
- Riechers M, Brunner B P, Dajka JC, Duşu IA, Lübker HM, Manlosa AO, Sala JE, Schaal T, Weidlich S (2021) Leverage points for addressing marine and coastal pollution: A review. *Mar Poll Bull* 167:112263. <https://doi.org/10.1016/j.marpolbul.2021.112263>
- Saaristo M, Brodin T, Balshine S, Bertram MG, Brooks BW, Ehlman SM, McCallum ES, Sih A, Sundin J, Wong BBM, Arnold KE (2018) Direct and indirect effects of chemical contaminants on

- the behaviour, ecology and evolution of wildlife. *Proc Royal Soc B* 285, 1885:20181297. <https://doi.org/10.1098/rspb.2018.1297>
- Servien R, Latrille E, Patureau D, Hélias A (2022) Machine learning models based on molecular descriptors to predict human and environmental toxicological factors in continental freshwater. *Peer Community J* 2:e15. <https://doi.org/10.24072/pcjournal.90>
- Strenn, B Clara, M Gans O, Kreuzinger N (2004) Carbamazepine, diclofenac, ibuprofen and bezafibrate - investigations on the behaviour of selected pharmaceuticals during wastewater treatment. *Wat Sci Technol* 50:269–276. <https://doi.org/10.2166/wst.2004.0337>
- Strickland JDH, Parsons TR (1972) *A Practical Handbook of Seawater Analysis*. 2nd edition. Bull Fisheries Res Board Canada 167, Ottawa. <http://dx.doi.org/10.25607/OBP-1791>
- Sutton R., Xie Y, Moran KD, Teerlink J (2019). Occurrence and sources of pesticides to urban wastewater and the environment. In: Goh K, Gan J, Young DF (eds) *Pesticides in surface water: Monitoring, modeling, risk assessment, and management*. American Chemical Society, Washington, pp 63-88. <http://dx.doi.org/10.1021/bk-2019-1308.ch005>.
- Tessier E, Garnier C, Mullot JU, Lenoble V, Arnaud M, Raynaud M, Mounier S (2011) Study of the spatial and historical distribution of sediment inorganic contamination in the Toulon bay (France). *Mar Poll Bull* 62:2075-2086. <https://doi.org/10.1016/j.marpolbul.2011.07.022>
- Togola A (2018) Etat de la contamination des eaux usées, eaux superficielles et eaux souterraines par les substances pharmaceutiques. *Environment Risque Sante* 17:7-14. <https://doi.org/10.1684/ers.2017.1075>
- Wang L, Zheng M, Xu H, Hua Y, Liu A, Li Y, Fang L, Chen X (2022) Fate and ecological risks of current-use pesticides in seawater and sediment of the Yellow Sea and East China Sea. *Env Res* 207:112673. <https://doi.org/10.1016/j.envres.2021.112673>
- Williams ES, Panko J, Paustenbach D J (2009) The European Union's REACH regulation: a review of its history and requirements. *Critical Rev Toxicol* 39:553-575. <https://doi.org/10.1080/10408440903036056>
- Zhang Y, Geißen S-U, Gal C (2008) Carbamazepine and diclofenac: Removal in wastewater treatment plants and occurrence in water bodies. *Chemosphere* 73:1151–1161. <https://doi.org/10.1016/j.chemosphere.2008.07.086>