



HAL
open science

Physiological and behavioural responses of bottlenose dolphin *Tursiops truncatus* to gustatory, olfactory and trigeminal chemical stimuli

Pablo Pezzino, Bertrand Bouchard, Julieta Arenarez, Alba Rincon Monne, Andreas Fahlman, Sylvia Campagna, Angelo Torrente, Aurélie Célrier

► To cite this version:

Pablo Pezzino, Bertrand Bouchard, Julieta Arenarez, Alba Rincon Monne, Andreas Fahlman, et al.. Physiological and behavioural responses of bottlenose dolphin *Tursiops truncatus* to gustatory, olfactory and trigeminal chemical stimuli. 35th ECS Conference, Apr 2024, Catania, Italy. hal-04622407

HAL Id: hal-04622407

<https://hal.science/hal-04622407>

Submitted on 25 Jun 2024


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Physiological and behavioural responses of Bottlenose Dolphin *Tursiops truncatus* to gustatory chemical stimuli

Pablo Pezzino¹ and Bertrand Bouchard¹, Julieta Arenarez³, Alba Rincon Monne³, Andreas Fahlman³, Sylvia Campagna¹, Angelo G. Torrente² and Aurélie Célérier¹



¹CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France ²Institut de Génétique Fonctionnelle, Université de Montpellier, CNRS, INSERM, Montpellier, France ³Fundación Oceanogràfic de la Comunitat Valenciana, Gran Vía Marqués del Turia 19, 46005, Valencia, Spain



Introduction

Chemical perception is the **first modality** that evolved in animals to sense the surrounding world. In terrestrials' mammals, chemical perception plays a crucial role in feeding, orientation, communication and much more. Conversely, information regarding this aspect is **notably scarce in marine mammals**, especially in cetaceans. The limited studies available so far indicate a **partial regression** of both anatomical structures and key genes families involved in olfaction and gustation. However, different behavioural observations suggest that chemical perception could be functional. In this study, we aimed to shed light on the chemical perception of Bottlenose Dolphins, focusing one of the chemical facet : **gustatory sense**. Gustation is characterized by the detection of molecules through taste receptors on the surface of papillae. We expected to unveil conserved processes and signalling pathways, through physiological and behavioural measurements. These findings would indicate that chemical senses could have kept a biological utility through odontocetes evolution.

Experimental design



Physiological variables


Logger developed to record underwater electrocardiograms (ECG) in dolphins

Dolphins ECGs


Behavioural variables

- Mouth openings and closing
- Vocalisations
- Breathing rate
- Avoiding behaviours (turning back, recoil)


Main results



We observed a **significant decrease in heart rate** when the Phase 2 of the session was either **salt or lemon** compared to the Phase 1 (water). This effect was not observed when the Phase 2 was Water, Quinine or fish hydrolysate.



The animals significantly **closed less their mouth** only when **lemon** was presented.



The animals **vocalized more** during the Phase 2 when it was only Water. That did not happen with any actual taste.

Discussion & Conclusion

- ❖ Significant response for **Salt and Lemon** ➡ trigger behavioural and physiological response ➡ **functional gustatory perception** | **emotional response** by autonomic nervous system ?
- ❖ Perception through **channels** (in opposition to GPCR for quinine/fish taste) ➡ some **molecular pathways** still operational in cetacean and may play a role in the sensory ecology : ➡ detection of **salinity gradient** near estuaries or in depths, perception of the **acid compounds** that may be clues of organic activity (through dissolved CO₂)
- ❖ **Vocal response** detected only when Water in Phase 2 ("no taste" session) ➡ **frustration** reaction from the animal due to the perceived absence of taste stimuli ?