



HAL
open science

Sex-related impairment of synaptic plasticity in the hippocampus of hypergravity-exposed mice: new insights for cognition in space

Wullen Mathilde, Thomas Freret, Valentine Bouet, Jean-Marie Billard

► To cite this version:

Wullen Mathilde, Thomas Freret, Valentine Bouet, Jean-Marie Billard. Sex-related impairment of synaptic plasticity in the hippocampus of hypergravity-exposed mice: new insights for cognition in space. FENS Forum 2024, FENS, Jun 2024, Wien, Austria. hal-04624901

HAL Id: hal-04624901

<https://hal.science/hal-04624901v1>

Submitted on 25 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


Wullen M, Bouet V, Freret T and Billard J-M.
Normandie Univ, UNICAEN, INSERM, COMETE, 14000 Caen, France

Introduction

One of the difficulties for space human missions is the exposure to repeated gravitational fluctuations during flight, which particularly affect the vestibular system. Given the **vestibulo-hippocampal connections** (Hitier, et al. 2014), hippocampal activity could be altered and would affect **cognitive performances** of the crew (Clement, et al. 2013).

This preclinical study aimed to investigate functional synaptic changes in hippocampal neuronal networks after **2G hypergravity** (HG) for different durations.

Materials & methods




Functional assessments

1. Long Term Potentiation (LTP) of synaptic strength was induced by high-frequency stimulation (HFS) and theta burst stimulation (TBS). LTP magnitude was measured over the last 15min relative to baseline.


2. Activation of NMDA receptors is essential for LTP. fEPSP/PFV ratio of **isolated NMDA-mediated fEPSPs** were constructed in a low magnesium medium supplemented with the non-NMDAR antagonist NBQX (10µM).

3. LTP impairment and altered NMDAR activation were pharmacologically characterized with **exogenous D-serine** (100µM) to assess the role of the NMDA glycine binding site.




Recordings in males

1. LTP HFS




HFS-induced LTP magnitude was depressed in mice exposed to **24h** of HG. No impairment was found for TBS-LTP.

2. NMDAR activation



As for LTP, NMDAR activation was depressed in **24h** exposed mice.

3. D-serine pharmacological supplementation




Exogenous D-serine partially **reversed LTP deficit** exhibited by **24h** exposed mice.

Exogenous D-serine **reversed NMDAR activation deficit** in **24h** exposed mice.




Recordings in females

1. LTP TBS




TBS-induced LTP was depressed in **15d** exposed mice. In contrast to males, HFS-induced LTP was not affected.

2. NMDAR activation



As for LTP, NMDAR activation was depressed in **15d** exposed mice.

3. D-serine pharmacological supplementation



Exogenous D-serine **did not reverse LTP deficit** exhibited by **15d** exposed mice.

Exogenous D-serine **did not reverse NMDAR activation deficit** in **15d** exposed mice.

This study reveals changes in hippocampus functional plasticity of mice exposed to hypergravity. Evidence is provided for sex differences that rely not only on HG duration but also on the type of functional plasticity and on the underlying synaptic mechanisms. Whether these changes in functionality may trigger significant cognitive deficits is currently under investigation.

