



HAL
open science

Quality control of the predatory mite *Euseius scutalis* (Acari: Phytoseiidae) against *Tetranychus turkestanii* (Acari: Tetranychidae) over 30 generations of rearing on cattail pollen

Zahra Gorji, Parviz Shishehbor, Farhan Kocheili, Elham Riahi

► To cite this version:

Zahra Gorji, Parviz Shishehbor, Farhan Kocheili, Elham Riahi. Quality control of the predatory mite *Euseius scutalis* (Acari: Phytoseiidae) against *Tetranychus turkestanii* (Acari: Tetranychidae) over 30 generations of rearing on cattail pollen. *Acarologia*, 2024, 64 (3), pp.819-832. 10.24349/gz0z-vqgh . hal-04625613

HAL Id: hal-04625613

<https://hal.science/hal-04625613>

Submitted on 26 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Quality control of the predatory mite *Euseius scutalis* (Acari: Phytoseiidae) against *Tetranychus turkestanii* (Acari: Tetranychidae) over 30 generations of rearing on cattail pollen

Zahra Gorji ^a, Parviz Shishehbor ^a, Farhan Kocheili ^a, Elham Riahi ^b

^a Department of Plant Protection, Faculty of Agriculture, Shahid Chamran University of Ahvaz, Ahvaz, Iran.

^b Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, P.O. Box 14115-336, Tehran, Iran.

Original research

ABSTRACT

In this study life table parameters and predation activity of *E. scutalis* on a diet of *T. turkestanii* were studied before the supply of *Typha* pollen (G0) and, after 10 (G10), 20 (G20), and 30 (G30) generations of being reared on cattail pollen (*Typha latifolia* L.). Our results showed that the longer the rearing of *E. scutalis* on cattail pollen before switching to *T. turkestanii*, the most impact it had on the developmental duration of the predatory mite. The female adult longevity was the highest in G0. In the older generations (G20 and G30), adult pre-oviposition period (APOP) was shorter than in younger generations (G10). The adult females of G0 had the longest oviposition days. The intrinsic rate of increase (r) and finite rate of increase (λ) of *E. scutalis* were highest in G20. There was no significant difference in the parameters R_0 , r , λ , and T (mean generation time) before and after thirty generations of rearing on cattail pollen when exposed to *T. turkestanii*. The proportion of male progeny produced by this predator was higher than that of females during both the early and late ages of the oviposition period. The predation rate of *E. scutalis* was not affected by the time of rearing on cattail pollen before switching to *T. turkestanii*. *Euseius scutalis* consumed an average of 435.93 *T. turkestanii* over its lifetime before the predator was reared on cattail pollen. This predation rate did not differ significantly after ten and thirty generations passed on a diet of *Typha* pollen. Overall, the findings of the current study indicated that *E. scutalis* long-term reared on cattail pollen could be a good candidate as a profitable predator on plants infested with *T. turkestanii*.

Received 21 November 2023

Accepted 13 June 2024

Published 20 June 2024

Corresponding author

Elham Riahi 

elham.riahi@modares.ac.ir

Academic editor

Kreiter, Serge

<https://doi.org/10.24349/gz0z-vqgh>

ISSN 0044-586X (print)

ISSN 2107-7207 (electronic)



Gorji Z. *et al.*

Licensed under

Creative Commons CC-BY 4.0



Keywords biological control; long-term rearing; growth rate; predation; pollen

Introduction

Tetranychus turkestanii Ugarov and Nikolski (*Tetranychus atlanticus* McGregor), known commonly as the strawberry spider mite, has become an important pest worldwide (Jeppson *et al.* 1975; Bolland *et al.* 1998). The species is polyphagous and is known to feed on more than 25 plant species belonging to more than 15 plant families (Jeppson *et al.* 1975; Kamali *et al.* 2001). It is found in the United States, Europe, Russia, Japan, China, the Near East, and the Middle East (Jeppson *et al.* 1975; Helle and Sabelis 1985; Zhang, 2003). The pest is frequent in Khuzestan province, southwest of Iran attacking various plant species from Solanaceae, Cucurbitaceae and Fabaceae (Mossadegh & Kocheili 2003).

How to cite this article Gorji Z. *et al.* (2024), Quality control of the predatory mite *Euseius scutalis* (Acari: Phytoseiidae) against *Tetranychus turkestanii* (Acari: Tetranychidae) over 30 generations of rearing on cattail pollen. *Acarologia* 64(3): 819-832. <https://doi.org/10.24349/gz0z-vqgh>

Tetranychus turkestanii produces extensive webbing and its feeding activity causes white chlorotic spots on the upper side of leaves. Severe infestations can result in the dropping of leaves and a reduction in flowering. In favorable conditions, the mite develops fast thanks to its short developmental period and high reproductive capacity. Heavy infestations lead to a rapid decrease in crop yields and cause serious economic losses (Carey & Bradley 1982; Tomczyk & Kropczynska 1985; Sohrabi & Shishchbor 2008).

Predaceous mites in the Phytoseiidae family play a crucial role in protecting different crops against many phytophagous mites and other pests (Bounfour & McMurtry 1987; McMurtry & Croft 1997). Therefore, they can be used in biocontrol programmes against pest mites and some insects (Gerson *et al.* 2003; Fouly *et al.* 2011). The members of genus *Euseius* are potential predators. They can prey on other food sources like fungal spores (Kostiainen & Hoy 1994; Schausberger 1997), and pollen (Fouly & Hassan 1991; Abou-Awad *et al.* 1992; Abdel halim *et al.* 2000). They have a major impact on low spider mite population densities, and could play a preventative role to limit the spread and the growth of the infestations (McMurtry 1992).

The first *Euseius scutalis* were described by Athias-Henriot (1958) in Algeria as *Typhlodromus scutalis*. In Middle East countries (Lebanon, Iran, Egypt, Israel, Jordan) and North Africa, *E. scutalis* is widely reported on a variety of host plants (Bounfour & McMurtry 1987; Momen & Abdel-Khalek 2008). It is one of the most common and widely distributed phytoseiid mites in Iran. *Euseius scutalis* is classified by McMurtry *et al.* (2013) as a specialized pollen feeder and generalist predator that is part of the type IV life style. The species is known to prey on spider mites, but also consumes eggs and immature stages of whiteflies, as demonstrated by previous studies (Swirski *et al.* 1967; Meyerdirk & Coudriet 1986; Nomikou *et al.* 2001).

Biological data and predation rates are important to evaluate the performance and potential of natural enemies on pest. They are useful in many studies, such as pest management, pesticide resistance, predator-prey relationships, biological control, mass rearing and harvesting (Chi *et al.* 2020). As the developmental stage and sex of most arthropods have a significant impact on their morphology, physiology, survival, reproduction, and predation rate, it is necessary to accurately describe stage differentiation and include both sexes in the experiments. They enable the achievement of realistic and precise life table analyses and accurately assess the overall population's fitness (Chi *et al.* 2020). Both sexes and stage differentiation can be included in data analysis, description, interpretation, and practical applications, thanks to the age-stage, two-sex life table. This method has been extensively used in entomological research for the past twenty years. Achieving effective sustainable management of a prey population using phytoseiid mites requires determining their full effects. There are numerous studies that utilize the age-stage, two-sex life table theory to evaluate the effects of diet on phytoseiid mites (Ansari-Shiri *et al.* 2022; Gravandian *et al.* 2022; Hadadi *et al.* 2022; Yazdanpanah *et al.* 2023a,b).

Among the various types of high-quality plant pollen, there is a commercially available product called Nutrimite™ (Biobest N.V., Westerlo, Belgium). This product is derived from the narrow-leaved cattail (*Typha angustifolia* L.) and has been recommended as an alternative food source for supporting generalist phytoseiid populations in crop fields and greenhouses (Samaras *et al.* 2015). Both *T. angustifolia* and *Typha latifolia* L. pollen have been shown to enhance phytoseiid performance (Goleva & Zebitz 2013; Vangansbeke *et al.* 2014). Compared to other plant pollen species, which are primarily insect-pollinated, *Typha* species have reduced allergenic properties and are more cost-effective and less labor-intensive to collect (Goleva & Zebitz 2013). However, there is a lack of published information on the long-term mass production of different phytoseiid species using this pollen, despite the potential of *Typha* species for mass rearing generalist phytoseiids.

Diets can affect mites' performance over many generations and not only in one generation (Nguyen *et al.* 2014). Any nutrient imbalances can cause a decline in the life table parameters of predatory mites over several generations (De Clercq *et al.* 2005). It is thus vital to investigate *E. scutalis* biological parameters, and ability to catch and feed on natural prey after being long-term reared on a non-prey food diet. The present work aimed to study with an age-stage, two-sex life

table theory, the biology, life table and predation rate of *E. scutalis* on a diet *T. turkestanii*, after being reared during different periods on cattail pollen. The growth rate and predation activity of four different generations of the species (0, 10th, 20th, 30rd generation) reared on cattail pollen was evaluated after being switched to a diet *T. turkestanii*. It seems that a predatory mite that can feed on both pollen and arthropod prey will likely not be significantly affected by a switch between these two diets. We hypothesize that since *E. scutalis* is specialized in feeding on pollen, rearing it exclusively on pollen for 30 generations will not affect its performance.

Material and methods

Prey and predator cultures

Tetranychus turkestanii was collected from heavily infested leaves of field bindweed plants (*Convolvus arvensis* L.) grown on the campus of the Faculty of Agriculture, Shahid Chamran University of Ahvaz, Ahvaz, Iran. The spider mites were then reared on cowpea seedlings (*Vigna unguiculata* (L.) Walp.) in laboratory conditions at a temperature of 25 °C (\pm 1 °C), 60 \pm 10% RH, and 16:8 h (L: D) photoperiod. Wood-frame culture cages (120 \times 60 \times 60 cm) with 210 μ m nylon mesh were used to contain the infested plants, with new plants added as needed.

A colony of *E. scutalis* population was obtained in February 2022 from a laboratory-grown culture in the Faculty of Agriculture, Shahid Chamran University of Ahvaz, Ahvaz, Iran. This population had initially been collected from marshmallow plants (*Althea officinalis* L.) infested with *T. turkestanii* on the campus of the Faculty of Agriculture, Shahid Chamran University of Ahvaz, in February 2021. A stereo-microscope (Olympus SZ51 Stereo Microscope) was used to examine the leaf samples and a thin-tipped brush was used to transfer the predatory mite females to green plastic sheet discs that had been soaked in water in Petri dishes. A mixture of *T. turkestanii* and date palm pollen (*Phoenix dactylifera* L.) was provided to the plastic discs as a food source for the predators. The date palm pollen was collected from Magtu Village, Khuzestan Province, and stored in a freezer throughout the experiments. Cotton threads were placed on the plastic sheets to provide egg-laying sites and shelter for the predators. To guarantee population continuity, predatory mite's eggs were collected at 1–2-day intervals and new mite colonies were established by transferring them to clean plastic discs. *Euseius scutalis* individuals were reared on date palm pollen for one month before starting the experiments. All the mite species were reared under laboratory conditions of 25 \pm 1 °C, 60 \pm 10% RH and photoperiod of 16: 8 (L: D) conditions throughout the experiments.

Experimental unit

All experiments were conducted in arenas similar to the rearing arenas used for predator rearing. A green plastic sheet (3 \times 3 \times 0.1 cm) was positioned over a sponge in a Petri dish (6 \times 1 cm, diam \times height) containing water. A small amount of sewing threads was added to the green plastic sheet to create shelter and oviposition spots. The rearing arena was provided daily with some cattail pollen, prior to and after *E. scutalis* adults appeared.

How *Euseius scutalis* performs before and after long-term rearing on cattail pollen

A random selection of around 150 *E. scutalis* females was taken from the stock colony and then moved to an experimental unit for oviposition. When the oviposited eggs hatched, defrost cattail pollen (*Typha latifolia* L.) collected from Dorud County, Lorestan Province, Iran, was offered to mites until they reached adulthood (G1). The diet of female progeny from the first generation was kept unchanged for as long as 30 generations (G30). After 10 (G10), 20 (G20), and 30 (G30) generations, the life table parameters of *E. scutalis* were estimated when switched to *T. turkestanii*. The trial was set-up with four treatments: individuals from the stock colony

(G0), individuals reared on cattail pollen for 10 generations (G10), individuals reared on cattail pollen for 20 generations (G20), and those raised on cattail pollen for 30 generations (G30) were compared. About 100 females were first randomly chosen from the related cultures and put in experimental arenas for 24 hours. The newly laid eggs were transferred individually to the experimental units. After the eggs hatched, *T. turkestanii* was added to the units and the number of consumed prey was replaced daily by new ones. The number of prey provided to each experimental unit was increased by changing the predator stage. Specifically, 20, 25, 30, and 50 immature stages of *T. turkestanii* were offered as food at the larval, protonymphal, deutonymphal, and adult stages, respectively. Each day, experimental arenas were checked and the duration and survival of each life stage were documented. The males and females of each treatment were paired immediately after the adults appeared. When there were more females than males or if a male who was paired with a female died, additional males of the same age from the laboratory colony were paired with them; however, the longevity and number of consumed prey were not documented. There were 51, 41, 43, and 45 replicate experimental arenas used, respectively for the treatments mentioned earlier. The daily data recording was conducted to determine the pre-oviposition period, oviposition days, adult longevity, survival, and fertility. The observations were carried on until all adults had died. The daily sex ratio was estimated by collecting eggs during the oviposition period daily and kept in the same conditions as their female parents until adulthood.

The predation ability of *E. scutalis* was assessed in G0, G10, G20, and G30 populations in life table experiments as well. Daily recording of the number of dead *T. turkestanii* was done until all predators died. To distinguish between the number of *T. turkestanii* preyed by male and female of *E. scutalis*, the predation rate of 20 males was examined under the same conditions. The female daily consumption rate was determined by subtracting the average male consumption from the consumption of pairs (Farhadi *et al.* 2011). All the experimental units were placed in a growth chamber at 25 ± 1 °C, $60 \pm 10\%$ RH, and a photoperiod of 16L:8D h.

Statistical analysis

Using TWSEX-MSChart, the age-stage, two-sex life table method was applied to analyze the life history data of *E. scutalis* (Chi & Liu 1985; Chi 1988; Chi 2023a). The TWSEX-MS Chart program was used to estimate age-stage-specific survival rate (s_{xj}), age-stage-specific fecundity (f_{xj}), age-specific fecundity (m_x), age-specific survival rate (l_x), and age-specific fecundity (m_x), as well as population growth parameters for each treatment (Chi 2023a). By using this program, the bootstrap method with 100000 re-sampling was used to estimate the means and standard errors of different stage durations, total pre-oviposition period (TPOP), adult pre-oviposition period (APOP), fecundity, oviposition days, and adult longevity, and the life table parameters [intrinsic rate of increase (r), finite rate of increase (λ), gross reproductive rate (GRR), net reproductive rate (R_0), and mean generation time (T)] (Chi 2023a). The mean of all parameters was compared between different treatments using the paired bootstrap test.

The computer program CONSUME-MS Chart (Chi 2023b) was used to analyze the daily consumption of all individuals. To compute the age-stage predation rate (c_{xj}), the daily consumption of each individual was taken into account. The mean number of *T. turkestanii* preyed by an individual *E. scutalis* during its life span (C_0), the number of *T. turkestanii* required for the production of an offspring from a *E. scutalis* (Qp), the entire consumption capacity of a stable population whose maximum value is one (ψ), and the predation potential of *E. scutalis* population by combining its growth rate (λ), age-stage predation rate (c_{xj}), and stable age-stage structure (a_{xj}) (ω) were computed based on Chi and Yang (2003). The bootstrap resampling method was used to estimate the variations and standard errors of predation parameters. Different stages' predation rates were compared using the Tukey test ($P < 0.05$) (SPSS 21.0). Using the TWSEX-MS Chart program, the comparison of C_0 , Qp , ψ , and ω was made based on the paired bootstrap test (with 100,000 resampling) (Chi 2023a).

Results

Life table parameters

Euseius scutalis developed on *T. turkestanii* even after being reared for thirty generations on pollen (Table 1). The developmental duration of *E. scutalis* was significantly affected by increasing the duration of rearing on cattail pollen before switching to *T. turkestanii*. The female protonymphal period was significantly longer in G20 than in G0, G10, and G30; this period was longer in G20 than in G10 in the case of males. The female pre-adult duration in G20 was significantly longer than other treatments. In G10, developmental durations of *E. scutalis* males were the shortest, followed by those in G30 (Table 1). The female and male adult longevities of *E. scutalis* were significantly different between various treatments. The longest female adult longevity was obtained in G0 (20.60 days), while there was no significant difference between G10 (14.64 days), G20 (14.97 days), and G30 (14.71 days). The same trend was observed for female total life span (Table 1). The male adult longevity of *E. scutalis* was not significantly different between G20 and G30, both were significantly shorter than that in G0 and G10. In G0, the male total life span was longest, while there was no significant difference between others (Table 1).

The adult pre-oviposition period (APOP), oviposition days, fecundity and sex ratio were significantly affected by the time of rearing on cattail pollen (Table 2). TPOP (total pre-oviposition period) did not significantly differ between treatments. In older generations (G20 and G30), APOP was shorter than in younger generations (G10). The individuals of G0 had the longest oviposition days (Table 2). There was no significant difference in oviposition days of this predator between G10, G20, and G30. The number of eggs produced by *E. scutalis* females in G0 and G20 was significantly higher than those laid by the females in G10 and G30 (Table 2). The sex ratio of the predatory mite in G20 (74% female adults/total individuals) was significantly higher than others, while not significantly different to G30. The lowest sex ratio (49%) was observed when *E. scutalis* fed *T. turkestanii* before rearing on cattail pollen (Table 2).

Figure 1 illustrates the age-stage survival rate (s_{xj}) of *E. scutalis* before and after different periods of rearing on cattail pollen with obvious overlap between stages as a consequence of differences in development rates among various individuals. Eggs and nymphs experienced

Table 1 Mean (\pm SE) duration of different life stages, developmental time, adult longevity, and total life span of female and male individuals of *Euseius scutalis* fed *Tetranychus turkestanii* before (G0) and after (G10–G30) long-term rearing on cattail pollen.

Parameter	G0	G10	G20	G30
Female				
Egg	1.84 \pm 0.07 a	1.72 \pm 0.09 ab	1.38 \pm 0.09 b	1.79 \pm 0.08 c
Larva	1.08 \pm 0.06 a	1.04 \pm 0.04 a	1.16 \pm 0.07 a	1.07 \pm 0.05 a
Protonymph	1.12 \pm 0.07 b	1.08 \pm 0.06 b	1.41 \pm 0.10 a	1.11 \pm 0.06 b
Deutonymph	1.00 \pm 0.00 b	1.20 \pm 0.10 ab	1.44 \pm 0.11 a	1.21 \pm 0.08 a
Pre-adult	5.04 \pm 0.04 b	5.04 \pm 0.11 b	5.38 \pm 0.11 a	5.18 \pm 0.12 ab
Adult longevity	20.60 \pm 1.32 a	14.64 \pm 1.44 b	14.97 \pm 1.02 b	14.71 \pm 1.47 b
Total life span	25.64 \pm 1.31 a	19.68 \pm 1.42 b	20.34 \pm 1.01 b	19.89 \pm 1.46 b
Male				
Egg	1.94 \pm 0.06 a	1.80 \pm 0.11 a	1.82 \pm 0.12 a	1.88 \pm 0.08 a
Larva	1.00 \pm 0.00 b	1.00 \pm 0.00 b	1.18 \pm 0.12 a	1.00 \pm 0.00 b
Protonymph	1.17 \pm 0.09 ab	1.00 \pm 0.00 b	1.36 \pm 0.15 a	1.18 \pm 0.10 ab
Deutonymph	1.28 \pm 0.14 a	1.13 \pm 0.09 a	1.09 \pm 0.09 a	1.12 \pm 0.08 a
Pre-adult	5.39 \pm 0.14 a	4.93 \pm 0.15 b	5.45 \pm 0.16 a	5.18 \pm 0.10 ab
Adult longevity	22.28 \pm 1.24 a	13.93 \pm 1.28 b	12.09 \pm 1.47 c	10.00 \pm 1.52 c
Total life span	27.67 \pm 1.27 a	18.87 \pm 1.22 b	17.55 \pm 1.46 b	15.18 \pm 1.54 b

The SE were estimated using the bootstrap method with 100,000 samples. Mean values followed by different letters in the same row are significantly different ($P < 0.05$), based on paired bootstrap test.




Figure 1 The age-stage survival rate (s_{xy}) of *Euseius scutalis* fed *Tetranychus turkestanii* before (G0) and after (G10–G30) long-term rearing on cattail pollen.

generation-based variations in survival rates; rates were higher in G30 and lower in G20. Furthermore, the survival rates of adults in G20 and G30 were higher than G0 and G10 (Figure 1). The female age-specific fecundity (f_x) first raised quickly, peaked, and then decreased. The peak f_x of *E. scutalis* was 3.41 eggs/female in G20, which was higher than that in G10 (3.39 eggs/female), G30 (3.30 eggs/female), and G0 (3.28 eggs/female) (Figure 2). The age-specific survivorship (l_x) for immature stages was 1 in all generations tested except for G0 (0.92). The age specific fecundity (m_x) was highest in the mentioned treatments at the age of 11 (1.91 eggs), 10 (2.11 eggs), 17 (2.92 eggs), and 23 (2.64 eggs) days, respectively (Figure 2).

As shown in Table 3, the population growth parameters of *E. scutalis* were affected by the time of rearing on cattail pollen when switched to *T. turkestanii* (Table 3). The gross reproductive rate (*GRR*) in the older generations was significantly higher than that in younger

Table 2 Mean (\pm SE) duration of adult pre-oviposition period (APOP), total pre-oviposition period (TPOP), oviposition days, fecundity, and sex ratio of *Euseius scutalis* fed *Tetranychus turkestanii* before (G0) and after (G10–G30) long-term rearing on cattail pollen.

Parameter	G0	G10	G20	G30
APOP	1.48 \pm 0.10 a	1.62 \pm 0.13 a	1.00 \pm 0.06 c	1.21 \pm 0.08 b
TPOP	6.52 \pm 0.10 a	6.67 \pm 0.19 a	6.38 \pm 0.11 a	6.39 \pm 0.12 a
Oviposition days	15.08 \pm 0.85 a	10.42 \pm 1.02 b	12.09 \pm 0.90 b	11.54 \pm 1.32 b
Fecundity (eggs/F)	42.68 \pm 2.58 a	27.92 \pm 3.05 b	36.81 \pm 2.98 a	27.07 \pm 3.66 b
Sex ratio (Nf/N)	0.49 \pm 0.07 c	0.61 \pm 0.08 b	0.74 \pm 0.07 a	0.62 \pm 0.07 ab

Means within rows followed by different letters are significantly different ($P < 0.05$). Differences between parameters were examined using paired bootstrap test.




Figure 2 The age-specific survivorship (l_x), age-stage specific fecundity of females (f_{xj}) (eggs) and age-specific fecundity (m_x) of *Euseius scutalis* fed *Tetranychus turkestanii* before (G0) and after (G10–G30) long-term rearing on cattail pollen.

generations. The net reproductive rate (R_0) of *E. scutalis* in G20 was not significantly different from that in G0, but the R_0 values in both generations were significantly higher than in G10 and G30 (Table 3). The intrinsic rate of increase (r) and finite rate of increase (λ) of *E. scutalis* were highest in G20 ($r = 0.2793 \text{ day}^{-1}$, $\lambda = 1.3222 \text{ day}^{-1}$) but did not differ significantly from those of in G10 ($r = 0.2469 \text{ day}^{-1}$, $\lambda = 1.2801 \text{ day}^{-1}$) (Table 3). The generation time (T) in G30 (13.178 days) was longer than those in G10 (11.48 days) and G20 (11.85 days), while there was no significant difference between G30 and G0 (Table 3).

Table 3 Mean (\pm SE) population growth parameters of *Euseius scutalis* fed *Tetranychus turkestanii* before (G0) and after (G10–G30) long-term rearing on cattail pollen.

Parameter	G0	G10	G20	G30
GRR (eggs/individual)	26.87 \pm 3.86 b	25.73 \pm 4.57 b	39.05 \pm 3.88 a	38.88 \pm 4.95 a
R_0 (eggs/individual)	20.92 \pm 3.23 ab	17.02 \pm 2.79 b	27.39 \pm 3.28 a	16.84 \pm 2.97 b
r (day^{-1})	0.2391 \pm 0.0137 b	0.2469 \pm 0.0147 ab	0.2793 \pm 0.0103 a	0.2143 \pm 0.0126 b
λ (day^{-1})	1.2701 \pm 0.0174 b	1.2801 \pm 0.0187 ab	1.3222 \pm 0.0136 a	1.2390 \pm 0.0156 b
T (day)	12.72 \pm 0.23 a	11.48 \pm 0.23 b	11.85 \pm 0.257 b	13.178 \pm 0.412 a

The SE were estimated by using the bootstrap technique with 100,000 samples. Mean values followed by different letters in the same row are significantly different ($P < 0.05$), based on paired bootstrap test. increase, λ : finite rate of increase, and T mean generation time.




Figure 3 Offspring sex ratio of females of *Euseius scutalis* fed *Tetranychus turkestanii* before (G0) and after (G10–G30) long-term rearing on cattail pollen. At each sampling date, white and black bars indicate the percentages of female and male offspring, respectively.

Sex ratio

Figure 3 shows how the female predatory mite, *E. scutalis*, changes the sex ratio of its progeny in response to the female age. The results demonstrate that the predator’s age influences the sex ratio in their offspring. In all treatments tested, lifetime sex ratios were consistently female biased. On the first day of oviposition, the sex ratio was male biased in all generations tested, then the percentage of females increased when the age of the predator increased. The three and two last days of oviposition period were male biased in G0 and G30, respectively, while the percentage of males increased in the last day of oviposition in G10 and G20. In other words, during the early and late ages of the oviposition period, the proportion of males produced in this predator increased. Generation does not affect this pattern.

Consumption rate

The predation rate of *E. scutalis* remained constant as the generation increased within the tested generations range (Table 4). The predation rate of the predator increased as stage changed from larva to protonymph, protonymph to deutonymph, and deutonymph to adult. For female and male protonymphs, the predation rate showed no significant difference between G10 (9.60 and 9.67 prey/predator, respectively) and G20 (11.50 and 10.18 prey/predator, respectively), but these values were significantly higher than that in G0 (7.12 and 6.11 prey/predator, respectively) and G30 (6.61 and 7.12 prey/predator, respectively). The total numbers of *T. turkestanii* consumed during the pre-adult stage were 25.44, 30.12, 30.28, and 24.67 per predator for females and 26.33, 28.07, 23.36, and 23.71 per predator for males, in G0, G10,

G20, and G30, respectively. The number of prey consumed during female pre-adult duration in G20 was significantly higher than other generations, while the male pre-adult consumption rate was not significantly changed between different generations. The highest female adult (594.42 prey) and male predation rates (338.21 prey) were in G0, and there were no significant differences between G10, G20, and G30 (Table 4).

The age-specific consumption rates (k_x), age-specific net consumption rates (q_x), and the cumulative consumption rate (C_x) of *E. scutalis* are shown in Figure 4. The value of k_x and q_x remained the same until the first death occurred, as indicated by Figure 4. The first peaks of these parameters of *E. scutalis* fed on *T. turkestanii* occurred at the age of 9, 7, 8, and 8 days in G0, G10, G20, and G30, respectively.

The net predation rate (C_0) changed significantly among generations tested (Table 4). Each *E. scutalis* consumed an average of 435.93 *T. turkestanii* over its lifetime when exposed to it before long-term rearing on cattail pollen, which did not differ significantly from the number of prey fed after ten (344.79) and thirty generations (341.25) on this pollen. The finite predation rate (ω) in G20 was lower than that in G30 and G10, while there was no significant difference between other three generations. The transformation rate (Q_p), the average number of prey needed to produce one predator egg, in G20 was 12.15 prey, which was significantly lower than that in G0 (20.84 prey), G10 (20.25 prey), and G30 (20.26 prey) (Table 4). Stable predation rate (ψ) was lowest in G20, and highest in G30, which the letter was not significantly different from that in G10.

Discussion

One of the most significant concerns in the biological control industry today is the quality of the natural enemies produced (Chambers 1977; Morrison & King 1977; Dicke *et al.* 1989; van Lenteren *et al.* 1993; Kim *et al.* 2001). Natural enemies may lose wild-insect traits during production due to genetic variability loss (Paspati *et al.* 2019). Therefore, monitoring the production, the process, and the product over successive laboratory generations is a major step in mass-rearing of natural enemies for biological control (Parra 2008). The overall quality of a laboratory population is assessed based on its effectiveness against the targeted prey after

Table 4 Mean (\pm SE) predation rates of *Euseius scutalis* fed *Tetranychus turkestanii* before (G0) and after (G10– G30) long-term rearing on cattail pollen.

Stage and statistic	Predation rate (prey/predator)				F (df), P-value
	G0	G10	G20	G30	
Larva (Female)	3.56 \pm 0.47 a	2.68 \pm 0.40 a	1.50 \pm 0.25 b	1.11 \pm 0.08 b	12.09 (3,106), < 0.0001
Protonymph (Female)	7.12 \pm 0.70 b	9.60 \pm 0.65 a	11.50 \pm 1.10 a	6.61 \pm 0.67 b	7.714 (3,106), < 0.0001
Deutonymph (Female)	14.76 \pm 1.01 a	17.84 \pm 1.75 a	17.28 \pm 1.30 a	5.53 \pm 1.05 a	0.989 (3,106), = 0.401
Pre-adult (Female)	25.44 \pm 1.18 ab	30.12 \pm 1.97 b	30.28 \pm 1.47 a	24.67 \pm 1.15 b	4.163 (3,106), = 0.008
Adult (Female)	594.42 \pm 38.67 a	395.66 \pm 43.80 b	357.13 \pm 28.52 b	417.79 \pm 47.35 b	6.81 (3,106), < 0.0001
Larva (Male)	4.78 \pm 0.65 a	2.33 \pm 0.57 b	1.91 \pm 0.51 b	1.00 \pm 0.00 b	11.399 (3,57), < 0.0001
Protonymph (Male)	6.11 \pm 0.89 b	9.67 \pm 0.76 a	10.18 \pm 1.36 a	7.12 \pm 0.83 b	4.318 (3,57), = 0.008
Deutonymph (Male)	15.44 \pm 1.99 a	16.07 \pm 1.90 a	11.27 \pm 1.19 a	15.59 \pm 1.12 a	1.375 (3,57), = 0.260
Pre-adult (Male)	26.33 \pm 2.07 a	28.07 \pm 2.50 a	23.36 \pm 1.94 a	23.71 \pm 1.57 a	1.108 (3,57), = 0.354
Adult (Male)	338.21 \pm 15.20 a	202.87 \pm 19.41 b	151.21 \pm 19.13 b	150.84 \pm 25.07 b	20.486 (3,57), < 0.0001
Net Predation rate (C_0)	435.93 \pm 35.53 a	344.79 \pm 31.61 ab	332.96 \pm 25.71 b	341.25 \pm 36.02 ab	—
Stable predation rate (ψ)	7.96 \pm 0.36 b	8.90 \pm 0.32 ab	6.84 \pm 0.25 c	9.00 \pm 0.34 a	—
Finite predation rate (ω)	10.11 \pm 0.58 ab	11.39 \pm 0.56 a	9.04 \pm 0.41 b	11.15 \pm 0.55 a	—
Transformation rate (Q_p)	20.84 \pm 2.23 a	20.25 \pm 2.13 a	12.15 \pm 0.84 b	20.26 \pm 1.96 a	—

Means followed by different small letters in the same row are significantly different using the paired bootstrap test at a 5% significance level. Means followed by different capital letters in the same row are significantly different ($P < 0.05$, Tukey).




Figure 4 Age-specific predation rate (k_x), age-specific net predation rate (q_x) and cumulative predation rate (C_x) of *Euseius scutalis* fed *Tetranychus turkestani* before (G0) and after (G10 –G30) long-term rearing on cattail pollen.

release (Huettel 1976). Pollen is a common food source for rearing and sustaining predatory mite populations.

Our results indicated that the development, reproduction and population growth rate of *E. scutalis* on *T. turkestani* were not impacted by the different rearing periods on cattail pollen, ranging from 10 to 30 generations. The duration of rearing on this pollen did not have any significant effects on female and male pre-adult period, female adult longevity, female and male total lifespan, and oviposition days. Furthermore, despite fluctuations in G20, no changes were observed between G10 and G30 in terms of fecundity, sex ratio, intrinsic rate of increase, and finite rate of increase. It can be concluded that the predator's performance remained constant during 30 generations of rearing on cattail pollen. Similar to our findings, duration of rearing (ranging from 10 to 30 generations) on almond pollen did not significantly impact the finite rate of increase (λ) and intrinsic rate of increase (r) of *N. cucumeris* after switching to *T. urticae* (Yazdanpanah *et al.* 2023b). Another study demonstrated that although a longer period of rearing on cattail pollen (20 generations) before switching to *T. urticae* resulted in shorter adult longevity, fewer oviposition days, reduced fecundity, GRR , and R_0 of *A. swirskii* compared to a shorter period (10 generations) of feeding on this pollen, the intrinsic and finite rates of increase remained unchanged (Hadadi *et al.* 2022). In contrast to our results, Gravandian *et al.* (2022) found that individuals of *N. cucumeris*, which was reared on cattail pollen for a longer period before being switched to *Tetranychus urticae*, exhibited shorter adult longevity and oviposition days, along with a higher population growth rate compared to those fed on the pollen for a shorter duration before the switch to *T. urticae*.

As mentioned previously, evaluating the effectiveness of the rearing process and the quality of the of reared organisms is necessary to assess their quality over time (Huettel 1976; Parra 2008). To ensure the success of a biological control program, it is crucial to monitor the performance of a predator that has been reared on a non-prey diet for a long period when it switches to a prey diet. Therefore, in this study we compared the life table parameters of *E. scutalis* before (G0) and after long-term rearing (G30) on cattail pollen when exposed to *T.*

turkestani. Our results revealed that although the female and male adult longevity, female and male total life span, APOP, oviposition days, and fecundity of individuals of G30 was shorter or lower than those of G0, pre-adult duration, TPOP, R_0 , r , λ , and T remained consistent. Given the significance of intrinsic rate of increase (r) as the most important parameter summarizing all demographic and biological parameters of a population (Birch 1948), it can be concluded that long-term rearing on cattail pollen did not negatively affect the performance and success rate of this beneficial biocontrol agent. Both populations (G0 and G30) were fed exclusively on date palm pollen for at least one month before experiments and this might explain why the G0 population was not significantly different from the G30 population. In other words, the biological parameters and population growth of the population that was fed date palm pollen and then transferred to prey did not differ much from those fed cattail pollen after initially being fed date palm and then transferred to prey. From this, we can conclude that there is not a significant difference in nutritional value between date pollen and cattail pollen for this predator. Similarly, stable performance of other phytoseiid mite, *A. swirskii*, before and after twenty generations of rearing on saffron pollen (Pourbahram *et al.* 2022), and ten generations of rearing on bird-of-paradise pollen (Yazdanpanah *et al.* 2023a) has been previously reported. In another study, a higher population growth rate and net reproductive rate of *A. swirskii* were estimated after forty-five generations of rearing on almond pollen when switched to *T. urticae* compared to the initial population (Ansari-Shiri *et al.* 2022).

After long-term rearing on alternative food, predators should still be capable of locating, seizing, and killing target prey (Grenier & De Clercq 2003). Our results indicated that *E. scutalis* can still catch and kill its natural prey (*T. turkestani*) even after thirty generations of rearing on cattail pollen. Similar results were observed for *N. cucumeris* against *T. urticae* after twenty generations (Gravandian *et al.* 2022) and thirty generations (Yazdanpanah *et al.* 2023b). *Amblyseius swirskii* also remained capable of preying on *T. urticae* after being reared on almond and maize pollen for six generations (Nemati & Riahi 2020), bird-of-paradise pollen for ten generations (Yazdanpanah *et al.* 2023a), saffron pollen for twenty generations (Pourbahram *et al.* 2022), and almond pollen for forty-five generations (Ansari-Shiri *et al.* 2022).

The potential predation ability of natural enemies can be evaluated using the finite predation rate (ω) (Chi *et al.* 2011), which remained constant even as the length of the rearing period on cattail pollen increased (from ten to thirty generations) in the current study. This parameter serves as a standard measure by establishing a connection between the finite rate, stable age-stage distribution, and age-stage specific predation rate (Chi *et al.* 2011). The predation rate of both immature and adult stages, as well as the net predation rate of *E. scutalis* when exposed to *T. turkestani*, remained stable regardless of the rearing period on pollen. In addition, individuals from G30, preyed on the same number of *T. turkestani* as individuals from G10, with no significant difference in stable predation rate, finite predation rate, and transformation rate between these populations. Similar to our results, the duration of rearing on pollen before switching to natural prey did not notably affect the stable predation rate and finite predation rate of *A. swirskii* in other studies (Hadadi *et al.* 2022; Pourbahram *et al.* 2022). In contrast, Gravandian *et al.* (2022) discovered that a longer period of rearing on cattail pollen led to a more stable and finite predation rates of *N. cucumeris* on *T. urticae* compared to a shorter period. Yazdanpanah *et al.* (2023b) reported opposite results. By comparing the finite predation rate of *E. scutalis* between the G30 and the initial population (G0), we found that the predation potential of this predatory mite against *T. turkestani* remained unaffected after 30 generations of rearing on cattail pollen, aligning with trends observed in previous studies (Ansari-Shiri *et al.* 2022; Hadadi *et al.* 2022; Pourbahram *et al.* 2022).

Phenotypic plasticity refers to the ability of animals to adjust their developmental, physiological, or behavioral characteristics in response to changes in their environment (Stephen *et al.* 2011). Various environmental factors can induce plastic responses in organisms (Westneat *et al.* 2019). Research on insect nutrition has demonstrated that the composition of the diet can greatly influence plasticity (Koyama *et al.* 2013). Insects that fail to adapt to new food sources in a timely manner may enter diapause or even die (Gu *et al.* 2021). Based on our

findings, *E. scutalis* appears to possess the ability to maintain population stability due to its high plasticity in response to environmental changes. In other words, this predatory mite shows no difficulty in adapting to different diets, whether it is pollen or prey. This is consistent with the study of McMurtry *et al.* (2013) that mentioned this predator's ability to be a specialized pollen feeder and a generalist predator. This characteristic allows *E. scutalis* to easily adjust to various diets within a short period of time, without experiencing a significant decline in fitness. In contrast, *A. swirskii* took some time to adjust to a new food source after switching its diet (Hadadi *et al.* 2022). This finding confirms that *A. swirskii* is a Type III predator, rather than a specialist pollen feeder (Type IV). This suggests that although *A. swirskii* can consume pollen, it is not as highly specialized as Type IV predators. Consequently, its ability to adapt to diet changes may differ from that of *E. scutalis*. There is a need to explore field applications of mass-reared predatory mites and the nutritional changes that occur when diets are altered in further researches.

In conclusion, our results showed that *E. scutalis*, considered a promising biological control agent against certain mites and insect pests, maintained successful predation on *T. turkestanii* even after a long-term rearing on cattail pollen. Moreover, the biological and predation parameters of this predator did not suffer any adverse effects due to the long-term feeding history on cattail pollen. The growth and consumption rates of the predators remained consistent when they switched from cattail pollen to prey, comparable to the population that was not reared on cattail pollen at all. Consequently, *E. scutalis* specimens that have been reared for an extended period on cattail pollen may represent excellent candidates for establishing an effective predator on plants infested with *T. turkestanii*. However, further research is necessary to assess the effectiveness of these mass-reared predators in greenhouses or fields.

Acknowledgements

Financial support (Grant no. SCU.AP1401.400) provided by the research deputy of Shahid Chamran University of Ahvaz, Ahvaz, Iran is gratefully acknowledged.

ORCID

Zahra Gorji  <https://orcid.org/0000-0001-8069-3684>

Parviz Shishehbor  <https://orcid.org/0000-0001-7843-4317>

Farhan Kocheili  <https://orcid.org/0000-0003-2572-4913>

Elham Riahi  <https://orcid.org/0000-0002-0429-4224>

References

- Abdel Halim S.M., Hanna M.A., Abdella M.M., Ramadan M.F. 2000. Evaluation of food type, availability of food and competition as factors affecting mass rearing of the predaceous mite *Euseius scutalis* (A.-H.) (Acari: Phytoseiidae). *Egyptian J. Biol. Pest Control.*, 10(1): 33-38.
- Abou-Awad B.A., Reda A.S., Elsawi S.A. 1992. Effects of artificial and natural diets on the development and reproduction of two phytoseiid mites *Amblyseius gossipi* and *Amblyseius swirskii* (Acari: Phytoseiidae). *Insect Sci. Appl.*, 13: 441-445. <https://doi.org/10.1017/S1742758400013746>
- Ansari-Shiri H., Fathipour Y., Hajiqaanbar H., Riahi E., Riddick E.W. 2022. Quality control of the predatory mite *Amblyseius swirskii* during long-term rearing on almond *Prunus amygdalis* pollen. *Arthropod-Plant Inte.*, 16: 645-655. <https://doi.org/10.1007/s11829-022-09929-6>
- Athias-Henriot C. 1958. Contribution à la connaissance du genre *Typhlodromus* Scheuten. (Acariens Parasitiformes, Phytoseiidae). Description de deux espèces nouvelles d'Algérie et clé des espèces du groupe *finlandicus*. *Rev. Patol. Veg. Entom. Agric. France*, 37(2): 181-186.
- Birch L.C. 1948. The intrinsic rate of natural increase of an insect population. *J. Anim. Ecol.*, 17: 15-26. <https://doi.org/10.2307/1605>
- Bolland H.R., Gutierrez J., Flechtmann C.H.W. 1998. World catalogue of the spider mite family (Acari: Tetranychidae), with reference to taxonomy, synonymy, host plants and distribution. Leiden, The Netherlands: Brill academic Publisher. 392 pp.
- Bounfour M., McMurtry J.A. 1987. Biology and ecology of *Euseius scutalis* (Athias-Henriot) (Acarina: Phytoseiidae). *Hilgardia*, 55: 1-23. <https://doi.org/10.3733/hilg.v55n05p023>

- Carey J.R., Bradley J.W. 1982. Developmental rate, vital schedules, sex ratios, and life tables of *Tetranychus urticae*, *T. turkestanii* and *T. pacificus* (Acarina: Tetranychidae) on cotton. *Acarologia*, 23(4): 333-345.
- Chambers D. L. 1977. Quality control in mass rearing. *Annu. Rev. Entomol.*, 22(1): 289-308. <https://doi.org/10.1146/annurev.en.22.010177.001445>
- Chi H. 1988. Life-table analysis incorporating both sexes and variable development rates among individuals. *Environ. Entomol.*, 17: 26-34. <https://doi.org/10.1093/ee/17.1.26>
- Chi H. 2023a. CONSUME-MSChart: computer program for consumption rate analysis based on the age stage, two-sex life table. 2019.02.01 ed. <http://140.120.197.173/Ecology/prod02.htm>. Accessed 19 Feb 2023
- Chi H. 2023b. TWSEX-MSChart: a computer program for the age stage, two-sex life table analysis. 2019.01.09 Edition. <http://140.120.197.173/Ecology/prod02.htm>. Accessed 19 Feb 2023
- Chi H., Liu H. 1985. Two new methods for the study of insect population ecology. *Bull. Inst. Zool. Acad. Sin.*, 24: 225-240.
- Chi H., Yang T. 2003. Two-sex life table and predation rate of *Propylaea japonica* Thunberg (Coleoptera: Coccinellidae) fed on *Myzus persicae* (Sulzer). *Environ. Entomol.*, 32: 327-333. <https://doi.org/10.1603/0046-225X-32.2.327>
- Chi H., Huang Y.B., Allahyari H., Yu J.Z., Mou D.F., Yang T.C., Farhadi R., Gholizadeh M. 2011. Finite predation rate: a novel parameter for the quantitative measurement of predation potential of predator at population level. *Nat. Prec.* <https://doi.org/10.1038/npre.2011.6651.1>
- Chi H., You M., Atlihan R., Smith C.L., Kavousi A., Özgökçe M.S., et al. 2020. Age-stage, two-sex life table: an introduction to theory, data analysis, and application. *Entomol. Gen.*, 40: 103-124. <https://doi.org/10.1127/entomologia/2020/0936>
- De Clercq P., Arijis Y., van Meir T., van Stappen G., Sorgeloos P., Dewettinck K., Rey M., Grenier S., Febvay G. 2005. Nutritional value of brine shrimp cysts as a factitious food for *Orius laevisgatus* (Heteroptera: Anthocoridae). *Biocontrol Sci. Technol.*, 15: 467-479. <https://doi.org/10.1080/09583150500086706>
- Dicke M., de Jong M., Alers M.P.T., Stelder F.C.T., Wunderink R., Post J., 1989. Quality control of mass-reared arthropods: Nutritional effects on performance of predatory mites 1. *J. Appl. Entomol.*, 108(1-5): 462-475. <https://doi.org/10.1111/j.1439-0418.1989.tb00480.x>
- Fouly A.H., Al-Daghairi M.A., Abdel Baky N.F. 2011. Biological aspects and life tables of *Typhlodromips swirskii* (Acari: Phytoseiidae) fed *Bemisia tabaci* (Homoptera: Aleyrodidae). *J. Entomol.*, 8(1): 52-62. <https://doi.org/10.3923/je.2011.52.62>
- Fouly A.H., Hassan M.F. 1991. Effect of crowding and food level on the predaceous phytoseiid mite *Amblyseius gossipi* (El-Badry) fed on white fly *Bemisia tabaci* (Gennadius). *Bull. Zool. Soc. Egypt*, 40: 141-146.
- Gerson U., Smiley R.L., Ochoa R. 2003. Mites (Acari) for Pest Control. UK: Blackwell Science, Oxford, UK. 539 pp. <https://doi.org/10.1002/9780470750995>
- Goleva I., Zebitz C.P.W. 2013. Suitability of different pollen as alternative food for the predatory mite *Amblyseius swirskii* (Acari, Phytoseiidae). *Exp. Appl. Acarol.*, 61: 259-283. <https://doi.org/10.1007/s10493-013-9700-z>
- Gravandian M., Fathipour Y., Hajiqaanbar H., Riahi E., Riddick E.W. 2022. Long-term effects of cattail *Typha latifolia* pollen on development, reproduction, and predation capacity of *Neoseiulus cucumeris*, a predator of *Tetranychus urticae*. *BioControl*, 67: 149-160. <https://doi.org/10.1007/s10526-021-10116-4>
- Grenier S., De Clercq P. 2003. Comparison of artificially vs. naturally reared natural enemies and their potential for use in biological control. In: van Lenteren J. (Ed). *Quality control and production of biological control agents. Theory and testing procedures*. Wallingford, UK: CABI Publishing. p. 115-131. <https://doi.org/10.1079/9780851996882.0115>
- Gu H.J., Dai C.C., Yang Y.Z., Lu Y.H. 2021. Effects of different preys on development and fecundity of *Hippodamia variegata* (Goeze). *J. of Plant Prot.*, 48(05): 1187-1188.
- Hadadi A., Fathipour Y., Hajiqaanbar H., Riahi E. 2022. Long-term effects of cattail pollen on development, population growth potential, and predation capacity of *Amblyseius swirskii* (Acari: Phytoseiidae). *Biocontrol Sci. Technol.*, 32: 1403-1416. <https://doi.org/10.1080/09583157.2022.2134555>
- Helle W., Sabelis M.W. 1985. Spider mites: their biology, natural enemies and control. Volume 1A. Amsterdam, The Netherlands: Elsevier Science Publisher. 405 pp.
- Huettel, M.D. 1976. Monitoring the quality of laboratory-reared insects: A biological and behavioral perspective. *Environ. Entomol.*, 5: 807-14. <https://doi.org/10.1093/ee/5.5.807>
- Jeppson L.R., Keifer H.H., Baker E.W. 1975. Mites injurious to economic plants. Berkeley, USA: University of California Press. 614 pp. <https://doi.org/10.1525/9780520335431>
- Kamali K., Ostovan H., Atamehr A. 2001. A catalogue of mites and ticks (Acari) of Iran. Tehran, Iran: Islamic Azad University Scientific Publication Centre. pp. 192.
- Kim J.H., Broadbent A.B., Lee S.G. 2001. Quality control of the mass-reared predatory mite, *Amblyseius cucumeris* (Acarina: Phytoseiidae). *J. Asia Pac. Entomol*, 4(2): 175-179. [https://doi.org/10.1016/S1226-8615\(08\)60120-X](https://doi.org/10.1016/S1226-8615(08)60120-X)
- Kostiainen T., Hoy M.A. 1994. Egg-harvesting allows large scale rearing of *Amblyseius finlandicus* (Acari: Phytoseiidae) in the laboratory. *Exp. Appl. Acarol.*, 18: 155-165. <https://doi.org/10.1007/BF02353683>
- Koyama T., Mendes C.C., Mirth C.K. 2013. Mechanisms regulating nutrition-dependent developmental plasticity through organ-specific effects in insects. *Front. Physiol.*, 4: 263. <https://doi.org/10.3389/fphys.2013.00263>
- McMurtry J.A. Dynamics and potential impact of 'generalist' phytoseiids in agroecosystems and possibilities for establishment of exotic species. *Exp. Appl. Acarol.*, 14: 371-382. <https://doi.org/10.1007/BF01200574>

- McMurtry J.A., Croft B.A. 1997. Life-styles of Phytoseiid mites and their roles in biological control. *Annu. Rev. Entomol.*, 42: 291-321. <https://doi.org/10.1146/annurev.ento.42.1.291>
- McMurtry J.A., De Moraes G.J., Sourassou N.F. 2013. Revision of the lifestyles of phytoseiid mites (Acari: Phytoseiidae) and implications for biological control strategies. *Syst. Appl. Acarol.*, 18: 297-320. <https://doi.org/10.11158/saa.18.4.1>
- Meyerdirk D.E., Coudriet D.L. 1986. Evaluation of two biotypes of *Euseius scutalis* (Acarina: Phytoseiidae) as predators of *Bemisia tabaci* (Homoptera: Aleyrodidae). *J. Econ. Entomol.*, 79(3): 659-663. <https://doi.org/10.1093/jee/79.3.659>
- Momen F.M., Abdel-Khalek A. 2008. Influence of diet on biology and life-table parameters of the predacious mite *Euseius scutalis* (A.-H.) (Acari: Phytoseiidae). *Acta Phytopathol. Entomol. Hungarica*, 41(6): 418430. <https://doi.org/10.1080/03235400600813508>
- Morrison R.K., King E.G. 1977. Mass production of natural enemies. In: Ridgway R.L., Vinson S.B. (Eds). *Biological control by augmentation of natural enemies*. New York: Plenum: 183-217. https://doi.org/10.1007/978-1-4684-2871-1_6
- Mossadegh M.S., Kocheili F. 2003. A semi descriptive checklist of identified species of arthropods (agricultural, medical) and other pests from Khuzestan, Iran. Shahid Chamran U., Ahvaz Press, Ahvaz, Iran.
- Nemati A., Riahi E. 2020. Does feeding on pollen grains affect the performance of *Amblyseius swirskii* (Acari: Phytoseiidae) during subsequent generations? *Bull. Entomol. Res.*, 110: 449-456. <https://doi.org/10.1017/S0007485319000804>
- Nguyen D.T., Vangansbeke D., De Clercq P. 2014. Artificial and factitious foods support the development and reproduction of the predatory mite *Amblyseius swirskii*. *Exp. Appl. Acarol.*, 62: 181-194. <https://doi.org/10.1007/s10493-013-9749-8>
- Nomikou M., Janssen, A., Schraag R., Sabelis M.W. 2001. Phytoseiid predators as potential biological control agents for *Bemisia tabaci*. *Exp. Appl. Acarol.*, 25: 271 - 291. <https://doi.org/10.1023/A:1017976725685>
- Parra J.R.P. 2008. Mass rearing of natural enemies. In: Capinera J.L. (Eds). *Encyclopedia of entomology*. Dordrecht: Springer. https://doi.org/10.1007/978-1-4020-6359-6_1741
- Paspati A., Ferguson K.B., Verhulst E.C., Urbaneja A., González-Cabrera J., Pannebakker B.A. 2019. Effect of mass rearing on the genetic diversity of the predatory mite *Amblyseius swirskii*. *Entomol Exp Appl*, 167(7): 670-681. <https://doi.org/10.1111/eea.12811>
- Pourbahram P., Hajiqaanbar H., Yazdanpanah S., Fathipour Y. 2022. Generation-dependent demography and predation capacity of *Amblyseius swirskii* (Acari: Phytoseiidae) fed on saffron pollen. *Int. J. Acarol.*, 48: 669-678. <https://doi.org/10.1080/01647954.2023.2171481>
- Samaras K., Pappas M.L., Fytas E., Broufas G.D. 2015. Pollen suitability for the development and reproduction of *Amblydromalus limonicus* (Acari: Phytoseiidae). *BioControl*, 60: 773-782. <https://doi.org/10.1007/s10526-015-9680-5>
- Schausberger P. 1997. Inter- and intraspecific predation on immatures by adult females in *Euseius finlandicus*, *Typhlodromus pyri* and *Kampimodromus aberrans* (Acari: Phytoseiidae). *Exp. Appl. Acarol.*, 21: 131-150.
- Sohrabi F., Shishehbor P. 2008. Effect of host plant and temperature on growth and reproduction of the strawberry spider mite *Tetranychus turkestanii* Ugarov and Nikolski (Acari: Tetranychidae). *Syst. Appl. Acarol.*, 13: 26-32. <https://doi.org/10.11158/saa.13.1.2>
- Stephen J.S., Gregory A.S., Nathan L. 2011. Polyphenism in insects. *Curr. Biol.*, 21(18): 738-749. <https://doi.org/10.1016/j.cub.2011.06.006>
- Swirski E., Amitai S., Dorzia N. 1967. Laboratory studies on the feeding, development and reproduction of the predaceous mites *Amblyseius rubini* Swirski and Amitai and *Amblyseius swirskii* Athias-Henriot (Acarina: Phytoseiidae) on various kinds of food substances. *Israel J. Agric. Res.*, 17: 101-112.
- Tomczyk A., Kropczynska D. 1985. Effects of host plants. In: Helle W., Sabelis M.W. (Eds). *Spider mites: their biology, natural enemies and control*. Volume 1A. Amsterdam, The Netherlands: Elsevier Science Publisher: 317-330.
- Vangansbeke D., Nguyen D.T., Audenaert J., Verhoeven R., Deforce K., Gobin B., Tirry L., De Clercq P. 2014. Diet dependent cannibalism in the omnivorous phytoseiid mite *Amblydromalus limonicus*. *Biol. Control.*, 74: 30-35. <https://doi.org/10.1016/j.biocontrol.2014.03.015>
- van Lenteren J.C., Bigler F., Waddington C. 1993. Quality control guidelines for natural enemies. *Proc. 7th workshop global IOBC working group quality control of mass reared arthropods*: 222-230.
- Westneat D.F., Potts L.J., Sasser K.L., Shaffer J.D. 2019. Causes and consequences of phenotypic plasticity in complex environments. *Trends Ecol. Evol.*, 34(6): 555-568. <https://doi.org/10.1016/j.tree.2019.02.010>
- Yazdanpanah Sh., Pourbahram P., Fathipour Y. 2023a. Bird-of-paradise, thorn apple and maize pollens are promising alternative diets for the long-term rearing of predatory mite *Amblyseius swirskii* (Acari: Phytoseiidae). *Syst. Appl. Acarol.*, 28(10): 1678-1690. <https://doi.org/10.11158/saa.28.10.8>
- Yazdanpanah Sh., Fathipour Y., Riahi E., Zalucki M.P. 2023b. Effects of diet switching from almond pollen to natural prey on predation capacity of *Neoseiulus cucumeris*. *J. Appl. Entomol.*, 147: 72-84. <https://doi.org/10.1111/jen.13081>
- Zhang Z.Q. 2003. *Mites of greenhouses: identification, biology and control*. London, UK: CABI Publishing., pp. 244. <https://doi.org/10.1079/9780851995908.0000>