



HAL
open science

Element of a pluri-disciplinary project on karstic reservoir characterization : groundwater modelling of the Dévoluy Massif, France.

Nathan Rispal, Bruno Arfib, Philippe Audra, Pierre Henry, Ludovic Mocochain, Benoît Viguiier, Alexandre Zappelli, Helene Miche, Christine Vallet-Coulomb

► To cite this version:

Nathan Rispal, Bruno Arfib, Philippe Audra, Pierre Henry, Ludovic Mocochain, et al.. Element of a pluri-disciplinary project on karstic reservoir characterization : groundwater modelling of the Dévoluy Massif, France.. Eurokarst, Jun 2024, Roma Sapienza Università, Italy. 2024. hal-04628392v2

HAL Id: hal-04628392

<https://hal.science/hal-04628392v2>

Submitted on 27 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - ShareAlike 4.0 International License

Element of a pluri-disciplinary project on karstic reservoir characterization : groundwater modelling of the Dévoluy Massif, France.

Nathan RISPAL ⁽¹⁾⁽²⁾, Bruno ARFIB ⁽²⁾, Philippe AUDRA ⁽¹⁾, Pierre HENRY ⁽²⁾, Ludovic Mocochain ⁽⁴⁾, Benoît VIGUIER ⁽³⁾, Alexandre ZAPPELLI ⁽²⁾, Hélène MICHE ⁽²⁾, Christine VALLET-COULOMB ⁽²⁾

(1) Université Côte d'Azur, Polytech'Lab - UPR 7498, Sophia-Antipolis, France. (2) Aix Marseille Univ, CNRS, IRD, INRAE, CEREGE, Aix-en-Provence, France. (3) Géoazur, Université Côte d'Azur - UCA - France. (4) Independent researcher.

Objectives : the aim of this study is to understand the behavior of a mountainous karstic reservoir. Exploration of hydrodynamic and meteorological data coupled to simulation on KarstMod platform with a snow routine module helps to decipher if discharge variations at the springs are related wether to recharge processes or properties of storage and flow in the reservoir.

How the snow routine works in KarstMod

Snow routine parameters in KarstMod

T _s	0.0	to	0.0	°C
MF	5.0	to	6.0	mm/°C
CFR	0.3	to	0.8	
CWH	0.05	to	0.25	

Subcatchments parametrisation

N°	Proportion	Temperature shift (°C)
1	0.35	1.92
2	0.33	-1.05
3	0.32	-4.65

Model

Modified from Pianosi and Wagener (2016)

Temperature and precipitation

temperature: TS
precipitation: snowfall, rainfall

solid component ↔ snowmelt ↔ liquid component

refreezing ↔ snowfall

Degree day factor :
 $M = MF \times TDD$

M (mm/d) : water from melted snow per day
MF = DDF : melting factor or degree-day factor (mm / °C . Day)
TDD = Active melting temperature (°C)
TDD = Air Temp. - Ts
(Ts = threshold temperature, default = 0°C)

CFR (refreezing factor)
CWH (water holding capacity)

Case study : the Dévoluy karstic reservoir

Modified from Gidon et al., 1980

Reservoir Cover

Oligocene
Eocene
Maastrichtian
Campanian
Santonian
Coniacian
Turonian, C2
"Marnes bleues"
Aptian-Albian
Cenomanian n°1-3
Barrémian-Beudantic
Hauteriviain n°1
Berriassien-Vauxin
Triassien n°8-9
Kimmeridgien n°6-7
Bathonian
Oxfordian
"Terres Noires" (> 1000 m)
n°1-5

Basement

400 m

Discharge :
- Max. 29m³/s
- Min. 200l/s
- Mean 5m³/s

Data :
- EDF gauge station : Q, T°C, P
- Météo France SAFRAN models : P, T°C, PET

Workflow : modelling a snow-influenced karstic aquifer

Field observations and data visualisation

Model 1 :
Classical configuration with two reservoir and no snow routine

Model 2 :
Adding snow routine

Model 3 :
Adding subcatchments

Model 4 :
Refining model with third reservoir

key elements

Step 1 : identifying the role of snow in the recharge

During spring and summer, discharge increases at the Gillardes spring while there is no significant Q rise in la Souloise river.

Modelling without snowroutine shows that part of the discharge is lost during spring and summer.

Step 2 : Implementing snow routine in the model

Snow routine helps to better match observed discharge during spring and summer.

Subcatchments in snow routine enhance the recharge simulation (rainfall, snow and snowmelt) over the catchment area.

Adding third reservoir allows to better reproduce fast variations in discharge.

Discussion

How to use the model once it is well calibrated ?

1/ Highlights the variations in flow rate in relation with the conditions of reservoir recharge. Possible causes of variation in recharge in Dévoluy: artificial snow at the ski resort, P and/or T°C variations due to climate change.

2/ Provides a better understanding of the hydrodynamic of the system in relation with groundwater management : Architecture of the system, contribution of each reservoir to the discharge at spring, water resources.

Conclusion and way-forward

Importance of recharge mechanisms : work is needed on recharge modeling to improve rainfall-discharge modeling with conceptual models («reservoir or compartments models» like KarstMod). Good model structure is not enough.

Snow routine module is efficient : the degree-day method used by KarstMod is effective when applied to a karstic spring.

Way-forward : threshold temperature or temperature shift in the subcatchments could be used to simulate futur climate change with KarstMod.

References and acknowledgments :

Thanks to EDF DTG for P, T°C and Q data and Météo France for meteorological data from SAFRAN models.
Gidon, M. et al. (1980) «Carte Géologique de la France, BRGM, feuille de Saint Bonnet XXXIII 37, 1:50000.» BRGM.
Michard, A. et al. (2010) «Cretaceous folding in the Dévoluy mountains (Subalpine Chains, France) gravity-driven detachment at the European paleomargin versus compressional event.» Bulletin de la Société Géologique de France, 181(6), pp. 565-581.
Pianosi, F., and Wagener, T. (2016) Understanding the time-varying importance of different uncertainty sources in hydrological modelling using global sensitivity analysis. Hydrol. Process., 30, 3991-4003. doi: 10.1002/hyp.10968
Siville, V., Cinkus, G., Mazzilli, N., Labat, D., Arfio, B., Massei, N., Cousquer, Y., Bertin, D., and Jourde, H. Improvement of the KarstMod modeling platform for a better assessment of karst groundwater resources. Hydrol. Earth Syst. Sci. Discuss. [preprint]. https://doi.org/10.5194/hess-2023-17, in review, 2023.
Zappelli, A., Belleville, A. and Jagercikova, M. (2018) Hydrology of Dévoluy karstic system: New insights with dye tracing and rainfall-discharge model. Eurokarst, Besançon.

KarstMod is free to use and can be downloaded via : www.sokarst.org