

Adult non-parents can recognize babies from their pain cries

Siloé Corvin, Camille Fauchon, Alexis Koutseff, David Reby, Nicolas Mathevon, Roland Peyron

▶ To cite this version:

Siloé Corvin, Camille Fauchon, Alexis Koutseff, David Reby, Nicolas Mathevon, et al.. Adult non-parents can recognize babies from their pain cries. IASP 2021 Virtual World Congress on Pain, Jun 2021, Online, France. hal-04628993

HAL Id: hal-04628993 https://hal.science/hal-04628993v1

Submitted on 28 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Adult non-parents can recognize babies from their pain cries






Experience (n=21)

Siloé Corvin^{1,2}, Camille Fauchon^{1,3}, Alexis Koutseff², David Reby², Nicolas Mathevon² and Roland Peyron¹


1 Central Integration of Pain (NeuroPain) Lab - Lyon Neuroscience Research Center, INSERM U1028, CNRS UMR5292, Université Claude Bernard, Bron, F-69677, France; 2 Equipe de Neuro-Ethologie Sensorielle ENES/CRNL, University of Lyon/ Saint-Etienne, INSERM U1028, CNRS UMR5292, Saint Etienne 42023, France; 3 Division of Brain, Imaging, and Behaviour, Krembil Brain Institute, University Health Network, Toronto, ON, Canada Contact: Siloé Corvin, siloe.corvin@inserm.fr; Camille Fauchon, camille.fauchon@univ-st-etienne.fr N° 1033631

Rationale: information in baby's cries

• Acoustic roughness is specific to pain cries




- Parents can distinguish between milddiscomfort and pain cries (Koutseff et al., 2017, **Bioacoustics**)
- Individual recognition from milddiscomfort (bath) cries (Gustafsson et al., 2013, Nat. Commun.; Bouchet et al., 2020, Proc. Biol. Sci.)



Methods

22 babies recorded during bath and vaccine



Acoustic similarity index per baby: median of spectrogram correlation bath/vaccine

Playback experiment:

42 non-parent adult listeners were assigned to one baby

Training phase

J1: 5 bath cries of the "assigned" baby

J2: 5 other bath cries of the "assigned" baby

Test phase

J2: 20 pain cries of the "assigned" baby (4 cries) and 4 unknown babies (4x4cries)

Which one is "my" baby crying from pain?

Multiple logistic regression

Results





Fig.1. Assigned babies were successfully

effects of listener's sex nor age were found.

Fig.2. Non-parents with no experience identified more than stranger babies. No at caring for babies seemed to be more sensitive to acoustic similarities



Conclusion

After listening few non-painful cries, adults are able to recognize a specific baby from its pain cries. Thus, there is an individual vocal signature present in the baby's discomfort and pain cries. Human without prior experience are able to learn quickly these important signals suggesting that it requires cognitive and perceptual abilities common to all adults rather than specific to mothers or fathers.

These findings will help to understand how distress is encoded in babies' cries to better assess their pain in pediatric care services.