



HAL
open science

Adult non-parents can recognize babies from their pain cries

Siloé Corvin, Camille Fauchon, Alexis Koutseff, David Reby, Nicolas Mathevon, Roland Peyron

► **To cite this version:**

Siloé Corvin, Camille Fauchon, Alexis Koutseff, David Reby, Nicolas Mathevon, et al.. Adult non-parents can recognize babies from their pain cries. IASP 2021 Virtual World Congress on Pain, Jun 2021, Online, France. hal-04628993

HAL Id: hal-04628993

<https://hal.science/hal-04628993>


Submitted on 28 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Rationale: information in baby's cries


- Acoustic roughness is specific to pain cries



- Parents can distinguish between mild-discomfort and pain cries (Koutseff et al., 2017, *Bioacoustics*)
- Individual recognition from mild-discomfort (bath) cries (Gustafsson et al., 2013, *Nat. Commun.*; Bouchet et al., 2020, *Proc. Biol. Sci.*)

Is it my baby?

- ✓ Mothers
- ✓ Fathers
- ✓ Non-parents




Is this ability transferred to pain (vaccine) cries?



Methods

- 22 babies recorded during bath and vaccine



Acoustic similarity index per baby: median of spectrogram correlation bath/vaccine

- Playback experiment: 42 non-parent adult listeners were assigned to one baby

Training phase

- J1: 5 bath cries of the "assigned" baby
- J2: 5 other bath cries of the "assigned" baby



Test phase

- J2: 20 pain cries of the "assigned" baby (4 cries) and 4 unknown babies (4x4cries)



Which one is "my" baby crying from pain?

- Multiple logistic regression

Results




Fig.1. Assigned babies were **successfully identified** more than stranger babies. No effects of listener's sex nor age were found.




Fig.2. Non-parents with no experience at caring for babies seemed to be more sensitive to acoustic similarities

Conclusion

After listening few non-painful cries, **adults are able to recognize a specific baby** from its pain cries. Thus, there is an **individual vocal signature** present in the baby's discomfort and pain cries. Human without prior experience are able to learn quickly these important signals suggesting that it requires **cognitive and perceptual abilities common to all adults** rather than specific to mothers or fathers.

These findings will help to **understand how distress is encoded in babies' cries** to better assess their pain in pediatric care services.