

## Has behavioural thermoregulation evolved solely to stay alive in insects, nothing more?

Sylvain Pincebourde

#### ► To cite this version:

Sylvain Pincebourde. Has behavioural thermoregulation evolved solely to stay alive in insects, nothing more?. Functional Ecology, 2024, 38 (6), pp.1342-1343. 10.1111/1365-2435.14574 . hal-04630107

### HAL Id: hal-04630107 https://hal.science/hal-04630107v1

Submitted on 9 Oct 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

- 1 FE Spotlights [1002 words]
- 2 Commentary on the article FE-2023-01052: Thermoregulation enhances survival but not

3 reproduction in a plant-feeding insect, by Noah T. Leith, Em A. Miller and Kasey D. Fowler-

- 4 Finn.
- 5
- 6 Has behavioural thermoregulation evolved solely to stay alive in
- 7 insects, nothing more?
- 8 Sylvain Pincebourde
- 9 Institut de Recherche sur la Biologie de l'Insecte, UMR 7261, CNRS Université de Tours,
- 10 37200 Tours, France
- 11 Contact : sylvain.pincebourde@univ-tours.fr
- 12 Conflict of interest: none
- 13

# Has behavioural thermoregulation evolved solely to stay alive ininsects, nothing more?

16 Temperature is probably the most influential abiotic variable as it drives nearly all 17 physiological rates. Temperature is both easy to measure, especially with our technological level, and complex to apprehend, because it varies widely across temporal and spatial scales. 18 The mere question of what temperature a given ectotherm experiences at the level of its cells 19 20 and enzymes (ie, body temperature) has generated tons of excellent works since a century 21 (Gates, 1980; Angilletta, 2009; Clarke, 2017). The process of behavioural thermoregulation is 22 a key tenet in these studies because it bridges organismal performance and temperature heterogeneity. In its broad definition, behavioural thermoregulation is the use of locomotion 23 or behavioural adjustments to meet permissive temperatures in the environment (Lahondère, 24 2024). For thermal ecologists, behavioural thermoregulation is a stimulating topic as it 25 involves several disciplines including behaviour, physiology, biometeorology and biophysics 26 (Gates, 1980; Helmuth, 1998; Kearney & Porter, 2004; Pincebourde & Woods, 2012). 27 28 Behavioural thermoregulation occurs in various ectotherm taxa. Amazingly, studies on reptiles' thermoregulation largely focused on their ability to find the optimal temperature for 29 their performance (eg, locomotion, activity window) while works on insects mostly 30 investigated their capacity to avoid overheating and improve survival under extreme heat. 31 32 However, in both cases, the picture is only partial, and only few studies so far have analysed 33 behavioural choices in the context of thermoregulation to both avoid lethal temperatures and 34 maximise performance by selecting the optimal temperature. This is precisely the aim of the study by Leith et al. (2024). In a small herbivore insect, Leith and colleagues assess if 35 36 behavioural thermoregulation both improves survival and maximise reproduction performance within the mosaic of thermal microenvironments of the host plant. 37

In an open-air mesocosm, Leith et al. (2024) surveyed body and operative temperatures of 38 39 treehoppers across different plant structures using infrared imaging. The operative temperature (ie, the body temperature at a given position without any thermoregulatory effect) 40 was inferred using judicious 3D printed models with colour, size and shape matching the 41 42 treehopper body. Operative temperatures are used to describe the available microclimates. Among the most astonishing results, Leith et al. reports high heterogeneity of operative 43 temperatures within and across plant individuals at any point in time, by up to almost 20°C. 44 The variability in actual body temperatures is lower, suggesting that the insect actively 45 thermoregulate to some extent, especially to avoid the most stressful temperatures above 46 47  $\sim$ 36°C that elicit the rapid heat escape behaviour. However, the thermal preference range 48 remains wide, and the insect is unlikely to select directly its microclimate within this range to optimize mating activity. Indeed, the thermal preference of the treehopper does not vary with 49 50 sex and mating status (as inferred by playing playbacks of acoustic courtship primers during thermal preference assays), suggesting that mating behaviours do not modify thermal biology 51 52 metrics.

Finally, the core of this study consists in analysing the effects of thermal quality (ie, if lethal 53 temperatures are present within the plant) and variability (ie, the temperature range present 54 55 across the plant) on the thermoregulation accuracy. The relationship between thermoregulation metrics unambiguously demonstrates that the cost-benefit conceptual model 56 applies in this system: the insect actively select body temperature within its thermal 57 preference range mostly when lethal temperature are present somewhere in the plant, and 58 thermoregulation is even more accurate in a highly heterogeneous thermal environment. 59 60 Indeed, thermoregulation was mostly effective under high ambient air temperature and when the insect is on a leaf, which are the conditions with higher probability to meet lethal 61 temperatures. Otherwise, the insect mostly thermoconforms suggesting that the treehopper 62

does not thermoregulate behaviourally to meet the narrow range of body temperatures thatmaximise mating activity.

The study of Leith et al. is highly significant for the field of thermal ecology. Conceptually, 65 66 studies that allow to partition between the cost-benefit and the inhibited-movement models of thermoregulation remain exceptional. Indeed, the thermal heterogeneity at fine scale is such 67 68 that it may become unpredictable for the insect which should spend a huge amount of energy to search and exploit body temperatures near optimum for mating – in other words the cost 69 outweighs the benefits of being precise. This cost-benefit model certainly applies to numerous 70 arthropod species that display similar heat escape behaviours such as aphids (Ma et al., 2018). 71 72 The rapid heat escape behavior to ovoid exposure to lethal temperature is not without 73 consequences, however. Insects that fall on the ground to escape overheating at the leaf 74 surface are suddenly exposed to soil predators and are at risk of starvation before they relocate themselves on the plant (Ma et al., 2018). An important trade-off should exist therefore 75 between heat avoidance, predation exposure, and feeding constraint which complexifies the 76 picture. Comparative analyses across species differing in behaviour are necessary to better 77 comprehend the drivers of this compromise. 78

Although survival is improved by behavioural thermoregulation, the population level 79 performance may still be challenged under elevated atmospheric temperatures because most 80 individuals remain at suboptimal temperatures for reproduction. Thus, this study provides 81 mechanistic understanding of recent works reporting that the thermal sensitivity of 82 reproduction more accurately predicts species distributions and vulnerability to climate 83 change (Parratt et al., 2021; van Heerwaarden & Sgrò, 2021). Since a decade or so, we have 84 85 seen a resurgence of macroecological studies considering microclimates (air temperature) as a potential buffer of species vulnerability to climate change (Zellweger et al., 2020) - this is not 86 enough as neglecting body temperatures within fine scales misses the actual level of exposure 87

to limiting temperatures. This is particularly true in ecosystems where organisms are exposed 88 89 to solar radiation which generate strong levels of thermal heterogeneity within fine scales (Saudreau et al., 2017; Pincebourde & Suppo, 2016), by contrast to the understorey of forests 90 which are shielded from radiation and display homogeneous thermal environments (Zellweger 91 et al., 2020). Training on heat transfer processes certainly is key to comprehend the 92 mechanisms generating thermal heterogeneity and to adjust appropriate designs (Briscoe et 93 94 al., 2023). Ecologists working in this area should develop their own "biophysical intuition" to better anticipate the amplitude of body temperature gradients within fine spatial scales. 95

96

#### 97 **References**

- 98 Angilletta, M. J. J. (2009). *Thermal Adaptation A theoretical and empirical synthesis*.
- 99 Oxford: Oxford University Press.
- 100 Clarke, A. (2017). *Principles of thermal ecology Temperature, energy and life*. Oxford:
- 101 Oxford University Press.
- 102 Lahondère, C. (2023). Recent advances in insect thermoregulation. Journal of Experimental
- 103 *Biology*, 226: jeb245751. https://doi.org/10.1242/jeb.245751
- 104 Gates, D. M. (1980). *Biophysical Ecology*. New-York: Springer-Verlag.
- Helmuth, B. (1998). Interdidal mussel microclimates: predicting the body temperature of a
  sessile invertebrate. *Ecological Monographs*, 68, 51-74.
- 107 Kearney, M., & Porter, W. P. (2004). Mapping the fundamental niche: physiology, climate,
- and the distribution of a nocturnal lizard. *Ecology*, *85*, 3119-3131.
- 109 doi:https://doi.org/10.1890/03-0820
- 110 Pincebourde, S., & Woods, H. A. (2012). Climate uncertainty on leaf surfaces: the biophysics
- 111 of leaf microclimates and their consequences for leaf-dwelling organisms. *Functional*
- 112 *Ecology*, *26*, 844-853. doi:10.1111/j.1365-2435.2012.02013.x

- 113 Leith, N. T., Miller, E. A., & Fowler-Finn, K. D. (2024). Thermoregulation enhances survival
- but not reproduction in a plant-feeding insect. *Functional Ecology*, 00, 1–13.
- 115 https://doi.org/10.1111/1365-2435.14546
- 116 Parratt, S. R., Walsh, B. S., Metelmann, S., White, N., Manser, A., Bretman, A. J., Hoffmann,
- 117 A. A., Snook, R. R., & Price, T. A. R. (2021). Temperatures that sterilize males better
- 118 match global species distributions than lethal temperatures. *Nature Climate Change*, *11*(6),

119 481–484. https://doi.org/10.1038/s41558-021-01047-0

- 120 van Heerwaarden, B., & Sgrò, C. M. (2021). Male fertility thermal limits predict vulnerability
- to climate warming. *Nature Communications*, *12*, 2214. https://doi.org/10.1038/s41467-
- 122 690 021-22546-w
- 123 Saudreau, M., Ezanic, A., Adam, B., Caillon, R., Walser, P., & Pincebourde, S. (2017).
- 124 Temperature heterogeneity over leaf surfaces: the contribution of the lamina
- microtopography. *Plant Cell and Environment*, 40, 2174-2188. 10.1111/pce.13026
- 126 Pincebourde, S., & Suppo, C. (2016). The vulnerability of tropical ectotherms to warming is
- 127 modulated by the microclimatic heterogeneity. *Integrative and Comparative Biology*, 56,
- 128 85-97. 10.1093/icb/icw014
- 129 Briscoe, N. J., Morris, S. D., Mathewson, P. D., Buckley, L. B., Jusup, M., Levy, O.,
- 130 Maclean, I. M. D., Pincebourde, S., Riddell, E. A., Roberts, J. A., Schouten, R., Sears, M.
- 131 W., & Kearney, M. R. (2023). Mechanistic forecasts of species responses to climate
- 132 change: The promise of biophysical ecology. *Global Change Biology*, 29, 1451–1470.
- 133 https://doi.org/10.1111/gcb.16557
- 134 Ma, G., Bai, C.-M., Wang, X.-J., Majeed, M. Z., & Ma, C.-S. (2018). Behavioural
- thermoregulation alters microhabitat utilization and demographic rates in ectothermic
- invertebrates. *Animal Behaviour*, *142*, 49-57.

- 138 Zellweger, F., De Frenne, P., Lenoir, J., Vangansbeke, P., Verheyen, K., Bernhardt-
- 139 Römermann, M., ... Coomes, D. (2020). Forest microclimate dynamics drive plant
- responses to warming. *Science*, *368*, 772-775.