



HAL
open science

X-ray optical activity in crystals of coordination complexes

Elen Duverger-Nédellec, Elisabeth A. Hillard, Patrick Rosa, Ph. Saintavit,
James Ablett, Fabrice Wilhelm, Andrei Rogalev

► **To cite this version:**

Elen Duverger-Nédellec, Elisabeth A. Hillard, Patrick Rosa, Ph. Saintavit, James Ablett, et al.. X-ray optical activity in crystals of coordination complexes. 33rd International Symposium on Chirality - CHIRALITY 2023, Sapienza Università di Roma; Università di Pisa, Jul 2023, Rome, Italy. hal-04631452

HAL Id: hal-04631452

<https://hal.science/hal-04631452v1>

Submitted on 2 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

X-ray optical activity in crystals of coordination complexes

Elen Duverger-Nédellec,¹ Elizabeth A. Hillard,¹ Patrick Rosa,¹ Philippe Saintavit,² James Ablett,³ Fabrice Wilhelm,⁴ Andrei Rogalev⁴

¹Univ. Bordeaux, CNRS, Bordeaux INP, ICMCB, UMR 5026, F-33600 Pessac, France

²IMPMC, CNRS, Sorbonne Université, Muséum National d'Histoire Naturelle, UMR 7590, 75252 Paris, France

³Synchrotron SOLEIL, L'Orme des Merisiers, BP 48, 91192 Saint Aubin, France

⁴European Synchrotron Radiation Facility (ESRF), 38043 Grenoble, France

E-mail: elizabeth.hillard@icmcb.cnrs.fr

Natural Circular Dichroism (NCD), the differential absorption of right- and left-circularly polarized light, is a workhorse technique for the stereochemical characterization of molecules and macromolecules. While this technique is most often applied in the UV-visible energy range, core-level optical activity using X-rays demonstrates some unique features such as element specificity and orbital selectivity, but is limited to single crystal measurements.¹ The majority of X-ray Natural Circular Dichroism (XNCD) studies have been performed using synchrotron sources, especially the ESRF ID12 at the K-edge of light transition metals² or at the L_{2,3}-edge of heavier atoms, such as lanthanides.³

Our work is devoted to extending X-ray optical activity to new techniques and molecular crystals, especially those based on coordination complexes. To this end, we have recently observed the first example of Resonant Inelastic X-ray Scattering – NCD (RIXS-NCD) at the GALAXIES beamline at the SOLEIL synchrotron. We have also recently investigated at the ID12 optically active but non-chiral systems and fully characterized the polar and azimuthal dependence of a tetragonal crystal system, thus supporting theoretical predictions.⁴

These new results will be presented in conjunction with a presentation of our approach to the conception and synthesis of model systems.




Figure a) Angular (azimuthal) dependence of the XNCD intensity of an achiral iron coordination complex; b) RIXS-NCD of enantiomers of a chiral cobalt complex.

- (1) Goulon, J.; Rogalev, A.; Brouder, C. X-Ray Detected Optical Activity. In *Comprehensive Chiroptical Spectroscopy*; John Wiley & Sons, Ltd, 2011; pp 457–491. <https://doi.org/10.1002/9781118120187.ch17>.
- (2) Srinivasan, A.; Cortijo, M.; Bulicanu, V.; Naim, A.; Clerac, R.; Saintavit, P.; Rogalev, A.; Wilhelm, F.; Rosa, P.; Hillard, E. A. Enantiomeric Resolution and X-Ray Optical Activity of a Tricobalt Extended Metal Atom Chain. *Chem. Sci.* **2018**, *9* (5), 1136–1143. <https://doi.org/10.1039/c7sc04131d>.
- (3) Mitcov, D.; Platonov, M.; Buch, C. D.; Reinholdt, A.; Døssing, A. R.; Wilhelm, F.; Rogalev, A.; Piliigkos, S. Hard X-Ray Magnetochiral Dichroism in a Paramagnetic Molecular 4f Complex. *Chem. Sci.* **2020**, *11* (31), 8306–8311. <https://doi.org/10.1039/D0SC02709J>.
- (4) Natoli, C. R.; Brouder, C.; Saintavit, Ph.; Goulon, J.; Goulon-Ginet, C.; Rogalev, A. Calculation of X-Ray Natural Circular Dichroism. *Eur. Phys. J. B* **1998**, *4* (1), 1–11. <https://doi.org/10.1007/s100510050344>.