

#### Rapid selection of multiple-resistance to eprinomectin and benzimidazole in a dairy goat farm

Léa Bordes, Nicolas Dumont, Anne Lespine, Elise Souil, Jean-Francois Sutra, Françoise Prévot, Christelle Grisez, Lola Romanos, Aurélie Dailledouze, Philippe Jacquiet

#### ▶ To cite this version:

Léa Bordes, Nicolas Dumont, Anne Lespine, Elise Souil, Jean-Francois Sutra, et al.. Rapid selection of multiple-resistance to eprinomectin and benzimidazole in a dairy goat farm. 27th Conference of the World Association for the Advancement of Veterinary Parasitology, Jul 2019, Madison, WI, USA, United States. hal-04632464

HAL Id: hal-04632464

https://hal.science/hal-04632464

Submitted on 2 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.






## PS02-74

27th Conference of the World Association for the Advancement of Veterinary Parasitology (WAAVP 2019), Madison, WI, USA





# Rapid selection of multiple-resistance to eprinomectin and benzimidazole in a dairy goat farm

Bordes L.a, Dumont N.b, Lespine A.c, Souil E.a, Sutra J.F.c, Prévot F.a, Grisez C.a, Romanos L.a, Dailledouze A.a, Jacquiet P.a\*


a: UMR INRA/ENVT IHAP, UMT Santé des Troupeaux de Petits Ruminants, Université de Toulouse, Ecole Nationale Vétérinaire de Toulouse, France b: Vétérinaires GARAZI, 64220 Saint Jean le Vieux, France

c: INTHERES, Université de Toulouse, INRA, ENVT, Toulouse, France, \* corresponding author E-mail address: p.jacquiet@envt.fr (P. Jacquiet)

### Observations and objectives

In a French dairy goat farm with high mortality rates and severe anemia, a vet practitioner noted a poor efficacy of eprinomectin treatments. This flock includes 70 lactating goats from the Alpine breed and was firstly constituted in 2012 with young kids. They are allowed to graze from March to September on 5ha, with complementation food and are regularly treated with eprinomectin each year in June and in August. The aims of this study were i) to explore this suspected case of eprinomectin resistance with a Fecal Egg Count Reduction Test completed with the measurements of eprinomectin concentrations in serum and ii) in case of resistance, to identify the eprinomectin-resistant gastrointestinal nematodes species.

#### **Materials and Methods**



Individual Fecal Egg Counts

- before treatment
- 14 days after treatment

Identification of L3 larvae species



After composite larval culture (D0 and D14) morphological identifications (Van Wyk and Mayhew, 2013) and Real-Time PCR (Milhes et *al.*, 2017)

Fecal Egg Counts Reduction test

(FECRT); guidelines of WAAVP

(Coles et *al.*, 1992)

Blood sampling: 2 and 5 days post-treatment



Eprinomectin concentrations in plasma with HPLC (Sutra et *al.*, 1998)

#### Result of Fecal Egg Count Reduction Test





Fig. 1: FECRT showed a low efficacy of fenbendazole and of eprinomectin whatever the route of administration (IR or TR).


Moxidectin was totally effective in this farm.

## Identification of GIN Larvae L3 species after bulk cultures

Morphological and molecular analyzes showed similar results: *Haemonchus contortus* is the major species (99%) found after treatments of the two groups Eprinomectin IR and TR and in the Fenbendazole group.



#### Serum concentrations of eprinomectin in Eprinomectin IR and Eprinomectin TR groups






Fig 2. A & B: At day 2, eprinomectin concentrations were higher than 2ng/ml (the minimal active concentration) whatever the goats and the treated groups. The lack of efficacy was not due to a suboptimal exposition of nematodes to the eprinomectin but to a resistant population.

This is the first report of multiple-resistance to eprinomectin and benzimidazole in a *Haemonchus contortus* population in France.