

Which domestic refrigerator temperatures in Europe? Focus on shelf-life studies regarding Listeria monocytogenes (Lm) in ready-to-eat (RTE) foods

Ludivine Bonanno, Hélène Bergis, Nathalie Gnanou-Besse, Adrien Asséré, Corinne Danan

▶ To cite this version:

Ludivine Bonanno, Hélène Bergis, Nathalie Gnanou-Besse, Adrien Asséré, Corinne Danan. Which domestic refrigerator temperatures in Europe? - Focus on shelf-life studies regarding Listeria monocytogenes (Lm) in ready-to-eat (RTE) foods. Food Microbiology, 2024, 123, pp.104595. 10.1016/j.fm.2024.104595. hal-04633366

HAL Id: hal-04633366 https://hal.science/hal-04633366v1

Submitted on 16 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

Which domestic refrigerator temperatures in Europe?

- Focus on shelf-life studies regarding *Listeria monocytogenes (Lm)* in ready-to-eat
- 3 (RTE) foods

1

4

10

12

17

- 5 Ludivine BONANNO, Hélène BERGIS, Nathalie GNANOU-BESSE, Adrien ASSÉRÉ,
- 6 Corinne DANAN
- 7 Unit Salmonella & Listeria, Agence Nationale de Sécurité Sanitaire (Anses), French Agency
- 8 for Food, Environmental and Occupational Health Safety. Laboratory for Food Safety, 14 rue
- 9 Pierre et Marie Curie, 94702 Maisons Alfort cedex, France
- 11 Corresponding author: Ludivine BONANNO, ludivine.bonanno@anses.fr

13 ABBREVIATIONS

- 14 Lm, Listeria monocytogenes; RTEF, Ready-To-Eat Food; FBO, Food Business Operator;
- 15 TGD, Technical Guidance Document; EURL, European Union Reference Laboratory; NRL,
- 16 National Reference Laboratory

18 ABSTRACT

- 19 Listeria monocytogenes (Lm) is a pathogenic bacteria able to grow at refrigerated temperatures,
- widely distributed in the environment. This bacteria is susceptible to contaminate various food
- 21 products of which refrigerated ready-to-eat foods (RTEF) may pose a risk for public health. In
- Europe, food business operators (FBOs) shall ensure that foodstuffs comply with the relevant
- 23 microbiological criteria set out in the Regulation (EC) N°2073/2005. Food safety criteria for
- 24 Lm are defined in RTEF throughout their shelf-life. FBOs should implement studies to
- demonstrate that the concentration of Lm does not exceed 100 CFU/g at the end of the shelf-

life, taking into account foreseeable conditions of distributions, storage and use, including the use by consumers. However, this last part of the cold chain for food products is the most difficult to capture and control.

For this purpose, the European Union Reference Laboratory for Lm (EURL Lm) launched an inquiry to its National Reference Laboratory network and reviewed the scientific literature from 2002 to 2020. The outcomes were integrated in the technical guidance document of the EURL Lm to assess shelf-life of RTEF which resulted in the recommendation to use 10°C as the reference temperature to simulate the reasonably foreseen storage conditions in domestic

KEYWORDS

refrigerators.

- Listeria monocytogenes, ready-to-eat foods, shelf-life studies, temperature, domestic
- 38 refrigerators, food safety

1. INTRODUCTION

Listeriosis is one of the most severe foodborne diseases under European Union surveillance in terms of fatality case in high-risk groups of immunocompromised population (EFSA and ECDC, 2023). This disease is mainly caused by *Listeria monocytogenes* (*Lm*), a ubiquitous pathogenic bacterium that can contaminate and grow in a large variety of environments. Due to its tolerance to various physico-chemical conditions and its ability to grow at low temperature, *Lm* is considered as a major microbial hazard for the food safety management from farm to fork, especially in refrigerated ready-to-eat foods (RTEF). In this frame, the European regulation lays down general food safety requirements, according to which food must not be placed on the market if it is unsafe. Especially, specific microbiological criteria

for Lm are defined in the Regulation (EC) N°2073/2005¹ for different categories of RTEF. Pursuant to Regulation (EC) No 2073/2005 Lm is not to be detected in 25g of RTEF intended for infants and for special medical purposes, and in 25g of other RTEF able to support the growth of Lm before they have left the immediate control of the producing food business operator. For all other situations, the food safety criterion sets a limit of 100 CFU/g throughout

When the limit is 100 CFU/g, food business operators (FBOs) shall conduct, as necessary, studies to evaluate the growth of Lm that may be present in the product during the shelf-life (Reg. (EC) N°2073/2005: article 3 and annex II). These studies should take into account reasonably foreseeable storage conditions susceptible to impact bacterial growth and their fluctuations along the cold chain (from production to consumption). As temperature has a major effect on bacterial growth, challenge tests assessing Lm growth potential, and durability studies for chilled RTEF shall mimic realistic time-temperature profiles at all steps of the food chain (at manufacturer, retail and consumer level). Furthermore, storage periods are clearly identified as a factor of tremendous influence on the risk of listeriosis associated with RTEF that support Lm growth (FAO and WHO, 2007). Indeed, the relative high and homogenous growth of Lm population at refrigeration temperatures has been shown by Rosa Rodrigues de Souza et al. (2023) for a collection of 39 strains. Moreover, quantitative modelling suggests that more than 90% of invasive listeriosis are caused by ingestion of refrigerated RTEF containing Lm at concentrations higher than 2,000 CFU/g (EFSA BIOHAZ Panel, 2018).

The knowledge of the temperature of domestic fridges leads to assess the growth of *Lm* in potential contaminated foods and consumer exposure. This plays therefore a very important role to assess the risk and contribute to manage food safety.

the shelf-life.

¹ Commission Regulation (EC) No 2073/2005 of 15 November 2005 on microbiological criteria for foodstuffs OJ L 338,

^{22.12.2005,} pp. 1–26.

However, this part of the cold chain remains a critical point for harmonized shelf-life studies. At international level, current recommendations of the Codex Alimentarius lead to adjust the temperature of household refrigerators so that the temperature of products does not exceed 6°C (preferably 2°C - 4°C) (FAO and WHO, 2007). In Europe, EURL Lm Technical Guidance Document (TGD) gives specifications to be considered by laboratories assessing the shelf-life of RTEF related to Lm (EURL Lm, 2021). The EURL Lm TGD is specific for Lm and is considered complementary to standard EN ISO 20976-1 for conducting challenge tests studies especially for the growth potential and the maximum growth rate (EN ISO 20976-1, 2019). This TGD has also a part dedicated to durability studies. An important point is the use of a time temperature profile to cover each cold chain (at manufacture, retail and consumer level) when performing a challenge test allowing to determine a shelf-life of a product. The TGD recommends that temperatures have to be properly justified and documented by FBOs. It is indicated that "when the FBO has its own data, the use of this information is preferred. In this case, use the 95th percentile of the FBO's data observation. If no data is available, default temperatures are defined in EURL Lm TGD on shelf-life studies. For the last step of the cold chain (consumer storage), a default reference temperature of 12°C has been recommended in the previous version of the EURL Lm TGD on shelf-life studies (2008), on the basis of expert opinion. Since then, numerous studies had been carried out in Europe on domestic storage temperatures. Some studies have already reviewed temperature of household refrigerators on a world

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

Some studies have already reviewed temperature of household refrigerators on a world scale (James et al., 2008; James et al., 2017; Dumitrașcu *et al.*, 2020), but this is the first study to summarize domestic refrigerators temperatures at European level and to assess the impact of these temperatures on shelf-life studies and European application guidelines. Therefore, the goal of this study was to account and gather the recent available data on storage temperatures

at consumer level in order to provide FBOs a harmonized approach in setting up shelf-life of
 prepacked refrigerated products.

2. MATERIAL AND METHODS

2.1. Data collection

To collect the highest number of available data on the temperature storage of foodstuff in household refrigerators, EURL *Lm* used two information channels:

the network of 42 National Reference Laboratories (NRLs) for *Lm*. EURL *Lm* launched an enquiry (Sphinx IQ2 software) in July 2019 among the NRLs *Lm* network in order to identify any surveys that may have been or conducted on the temperature of domestic refrigerator from 2000 to 2020, and to collect data from these surveys. In case of publications of such surveys, the corresponding author was contacted to collect the raw temperature data.

- in parallel, a review on domestic refrigerator temperatures was performed (PubMed) from the scientific literature published between 2000 to 2020 and referring to the keywords: refrigerator, temperature, food and household. Only scientific publications providing data on the temperature of European household refrigerators were selected.

2.2. Temperature distribution and determination of percentiles.

- Based on data collected from 10 national surveys and 17 scientific publications, the 75th and 95th percentiles of refrigerator temperature distribution were:
 - (1) Directly provided from the survey or the scientific publication (Method 1 M1),
- (2) Calculated from raw temperature data (R software version 1.4.1106, "quantile" 121 function) (Method 2 – M2),

122 -	(3) Estimated from graphical temperature distribution reported in the survey report or
123	the publication (Method 3 - M3) or
124 -	(4) Estimated from the mean temperature and standard deviation of N measurements
125using "rno	orm" and "matrix" functions in R (Method 4 - M4) The function "rnorm"
126	generates a random simulation of temperatures following a normal distribution. The
127	function "matrix" repeats the simulation (1000 iterations were performed in this
128	study). In order to take into account the uncertainty related to the number of refrigerators
129	investigated, the 75 th and 95 th percentiles were determined by bootstrap approach from

" and the mean were calculated with the

132

131

133 2.3. Statistical analysis

130each simulation by the function "rowQuantiles"

associated 95% confidence intervals (CI).

134Three kind of statistical analyses were performed in this study. First, a Student's test was used 135 to evaluate the determination of percentiles (Method M4) in comparison to the other methods. another Student's test was used to evaluate the difference between refrigerator 136Next. 137 temperatures measured in air or in food. 138 Finally, countries were grouped according to the Köppen-Geiger climate classification and the 139 weighted average refrigerator temperatures was calculated to take into account the number of refrigerators investigated. A Kruskal-Wallis's test was used to compare the weighted average 140 141refrigerator temperatures according to this classification following by a Dunn's test to compare 142 Geoclimatic groups by pair.

143

144

3. RESULTS AND DISCUSSION

145 3.1. Data from national surveys

146 Twenty-nine out of 42 enquired NRLs (69%) replied to the EURL *Lm* questionnaire and among 147 them, seven (24%), have conducted an inquiry in their country. Table 1 presents the general 148 information from each national survey. 149 The number of monitored refrigerators were comprised between 40 and 5 428, mainly 150 investigated in the whole randomly sampled population or by a stratified sampling strategy for 151 two studies (Anses, 2017; Daminelli, 2019). One survey used a selective strategy, targeting 152 elderly people (Denmark survey, 2015). Temperature measurements were performed mostly by 153 a home investigator who installed data loggers inside the refrigerators for a duration varying 154 from 30 min to 7 days. For some studies, the monitoring was performed in the refrigerator (air 155 measurement) at different areas (top, middle and bottom). For one study, loggers were placed 156 between packages of RTEF (Denmark survey, 2015). For two studies, the data loggers were 157 directly put into the food product purchased by consumers, allowing monitoring during several days (Derens et al., 2006; Derens-Bertheau et al., 2015). 158

159 Table 1. General information from national surveys

Country	Year	Scope	Title of the survey	N refrigerators	Target population	Sampling strategy	Data reporting	Monitoring tool	Duration	Air (area of fridge) or Food
Austria	2011	National	Food safety and hygiene in the private household. (Austria survey 2011)	40	Whole population	Random	Interview by phone	TF	?	Air
Belgium	2004	National	Belgian food consumption survey 1 – 2004. (Devriese et al. 2004)	3001	Whole population	Random	Home investigator	?	?	Air
Denmark	2015	District	Investigate the temperature of elderly people refrigerators. (Denmark survey 2015, <i>Unpublished results</i>)	77	Elderly people	Selective	Health Care workers	DL	30 min	Air
France	1999	National	Study of domestic refrigerator temperature and analysis of factors affecting temperature: a French study. (Laguerre et al. 2002)	118	Whole population	Random	Home investigator	DL	7 days	Air (T; M; B)
France	2006	National	The cold chain of chilled products under supervision in France. (Derens et al. 2006)	251	Whole population covered French territory	Random	Recorder inside food returning by the consumer	DL	4 days	Food
France	2013	National	Cold chain of chilled food in France. (Derens-Bertheau et al. 2015)	75	Whole population	Random	Recorder inside food returning by the consumer	DL	6 days	Food
France	2014 - 2015	National	Estimate of food consumption & nutritional intake of people living in France. (Anses, 2017)	5428	Whole population	Stratified multi-stage design	Home investigator	TH	30 – 45 min	Air (T; M; B)
Ireland	2005	National	Food safety knowledge of consumers and the microbiological and temperature status of their refrigerators. (Kennedy et al., 2005)	100	Whole population	Random	Home investigator	DL	72 h	Air (M)
Italy	2019 - 2020	National	National survey of domestic refrigerators temperatures in Italy (preliminary results) (Daminelli , <i>Unpublished results</i>)	494	Whole population	Stratified sampling plan	Home investigator	DL	?	Air
The Netherlands	2012	National	Food waste. (The Netherland survey 2012, <i>Unpublished results</i>)	326	?	?	Self-reporting via a questionnaire	ТН	12 h	Air

DL: Data Logger; TH: thermometer; TF: Temperature display on the fridge; T: Top; M: Middle; B: Bottom;?: unknown;

160

3.2. Data from scientific literature

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

The general information from the 17 selected publications is presented in Table 2. These studies were conducted in 11 European countries. The number of monitored refrigerators varied from 30 and 250 and were mainly investigated in the whole population. Regarding the sampling strategy, five studies used a random approach. For two studies, the sampling strategy was performed by a selective approach by targeting specific population like students (Galvão et al., 2017) or listeria-sensitive population (elderly, family with children and/or pregnant women (Dumitrascu et al., 2020)). Two studies used a snowball approach which is used when researchers have difficulties finding participants for their studies (Rossvoll et al., 2014; Ovca et al., 2020). This is a non-probabilistic sampling method where currently enrolled participants help to recruit new participants to form part of the sample. One study was performed by a passive approach based on volunteer families (Azevedo et al., 2005). Finally, the sampling strategy was unknown for five studies. As previously observed, temperature was monitored mainly by data loggers placed inside the refrigerators (air measurement) at different area (top, middle, bottom and door) excepted for two studies where they were placed in water (Garrido et al. 2010) or foods (Marklinder et al., 2004). The duration of the recording varied between 40 min to 14 days. For one study (Garrido et al. 2010), only three measures were performed with a calibrated probe.

Table 2. General information from scientific literature

180

181

Country	Year	Title of the publication	Target population	Sampling strategy	Monitoring tool	Duration	Air (area of fridge) / Food
Germany; Spain; France; United Kingdom	2002	Cold storage in private households: recommendations and consumer real life behaviour. (Geppert et al. 2010)	?	? DL 11 days		11 days	Air (M)
Sweden	2004	Home storage temperatures and consumer handling of refrigerated foods in Sweden. (Marklinder et al. 2004)	Whole population (Uppsala)	Random	DL	40 min	Food
Portugal	2005	Incidence of <i>Listeria</i> spp. in domestic refrigerators in Portugal. (Azevedo et al. 2005)	North of Portugal	Passive	TH	?	Air
Greece	2005	Modelling the effect of house hold chilled storage condition on the risk distribution of meat product. (Taoukis et al. 2005)	Whole population (Athens metropolitan)	?	DL	7 days	Air
Spain	2007	Survey of Temperature and Consumption Patterns of Fresh-Cut Leafy Green Salads: Risk Factors for Listeriosis. (Carrasco et al. 2007)	Whole population (Cordoba)	Random	DL	24 h	Air
Spain	2010	Temperature distribution in Spanish domestic refrigerators and its effect on <i>Listeria</i> monocytogenes growth in sliced ready-to-eat ham. (Garrido et al. 2010)	Whole population (Pamplona)	?	sntativa frida		Air (T; M; B) Water (T; M; B)
United Kingdom	2010	Reducing food waste through the chill chain. (George et al. 2010)	Whole population	Representative fridge type	DL	4 days	Air (T; M; B)
Greece	2010	Probabilistic Model for <i>Listeria monocytogenes</i> growth during distribution, retail storage, and domestic storage of pasteurized milk. (Koutsoumanis et al. 2010)	?	?	DL 24 h		Air (T; M; B; D)
Italy	2013	Survey conducted in Italy on the consumer refrigeration temperatures and their impact on food safety illustrated with Salmonella. (Roccato, 2013)	Volunteers population	Random	DL 6 days		Air (T; B; D)
Norway	2014	Toxin production and growth of pathogens subjected to temperature fluctuations simulating consumer handling of cold cuts. (Rossvoll et al. 2014)	Specific population (Oslo and Akershus)	Snowball	DL 11 days		Air
Italy	2014	The management of the domestic refrigeration: microbiological status and temperature. (Vegara et al. 2014)	Whole population (Central and North Italy)	?	TH	24 h	Air (M)
United Kingdom	2016	Time-temperature profiling of United Kingdom consumers' domestic refrigerators. (Evans and Redmond, 2016)	Whole population (urban community, South Wales)	Random	P	6 days	Air (M; D)
Portugal	2017	Thermal performance, usage behaviour and food waste of domestic refrigerators in a university student community: findings towards cities sustainability (Galvão et al. 2017)	Student community University (Beira Interior)	Selective	DL	8 days	Air (M)
Spain	2019	Domestic refrigerator temperatures in Spain: Assessment of its impact on the safety and shelf-life of cooked meat products. (Jofre et al. 2019)	Whole population	Random	DL	24 h	Air (M; D)
Greece	2019	Field survey on temperature distribution and hygienic status of domestic refrigerators in Lemnos, Greece. (Andritsos et al. 2021)	Island of Lemnos	Random	DL	24 h	Air (T; B; D)
Romania; France; Norway; Portugal; United Kingdom	2020	Time-temperature profiles and <i>Listeria monocytogenes</i> presence in refrigerators from households with vulnerable consumers. (Dumitrașcu et al. 2020)	One city per country	Selective	DL	14 days	Air
Slovenia	2020	Temperatures and storage conditions in domestic refrigerators - Slovenian scenario (Ovca et al. 2020)	?	Snowball	DL	7 days	Air (M)

DL: Data Logger; TH: thermometer; P: Calibrated probe; T: Top; M: Middle; B: Bottom; D: Door; ?: Unknown

3.3. Combined data and temperature analysis

The refrigerator data were collected from 16 European countries: 14 out of the 27 EU member states (MSs) and two non-MSs (Norway and the United Kingdom) (Figure 1). Published storage temperatures at consumer level may present a large scale at European level, depending on multi-criteria factors linked to, the area of the refrigerator where the temperature was registered, individual knowledge of proper management of domestic refrigerator, consumer awareness on their role in food safety, etc. A total 10 national surveys and 17 scientific publications were collected for this study to have a general overview of refrigerator temperature at European level. Most of the studies/surveys targeted the whole population of a defined geographic area. Few of them focused on a specific population such as students, elderly, families with infants and/or pregnant women and young single men. The temperature measurements were mainly carried out using data loggers at different places inside the refrigerator, including the top, middle, bottom and the door or inside a food, mainly RTEF. Measurement using the data logger method allows the true operating conditions of the refrigerator to be represented. However, some limitations can be pointed out. In some surveys/studies the sampling was not randomly performed and the number of refrigerators investigated was lower than 100, which may lead to a bias and not reflect the overall population. For a few studies, the measurement method was performed with a thermometer and did not allow continuous measurements to be taken over a given period of time, as with data loggers, in order to establish a precise mean temperature value. Finally, there was an underrepresentation of Northern Europe (i.e. Denmark, Estonia, Finland, Latvia and Lithuania). It would have been preferable to have data from a European survey that covered all European countries and use the same methodology to measure the temperature of domestic refrigerators as the method described by Derens-Bertheau et al. 2015.

204

205

206

207

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

2031

In total, from the 27 studies, temperature measures were performed on 9,926 refrigerators. The mean temperature and the percentile (75th and 95th) were determined from each of them (Table 3). The 75th and 95th percentiles were obtained for 25 out of the 27 studies. For one study, no raw data or graphic

representation of the data was available to perform the percentile determination (Rossvoll et al. 2014). For 13 of the 26 studies (50%), the 75th and 95th percentiles were calculated using method M2. For the remaining 13 studies, percentiles were estimated using methods M4 (8), M1 (3) and M3 (2). Regarding the eight studies (30%) for which the determination of percentiles was estimated based on a simulation (Method M4) from available data (number of refrigerators, the average of refrigerator temperature and the associated standard deviation), this method is conditioned by a normal distribution assumption of the data (Pouillot et al. 2007). The normality of the data sets was verified with statistical test within three studies: Kennedy et al., 2005 (by Roccato et al., 2017), Taoukis et al., 2005, Carrasco et al., 2007. For three other studies (Koutsoumanis et al., 2010; Roccato, 2013; Evans and Redmond, 2016), it was assumed that the data follow a normal distribution based on the visual appearance of the temperature distribution curves presented in the publications. Finally, for two last studies, this information was unknown (Austria survey, 2011; The Netherland survey, 2012). This approach allowed to take these studies into account, and not reject them (due to the absence of raw data), in order to cover as many European countries as possible, such as Austria, Republic of Ireland and the Netherlands. The robustness of this approach was also assessed by comparing percentiles obtained from studies with M2 method (based on raw data). For these studies, percentiles were calculated with M4 method. Results obtained with M2 method were included in the confidence interval obtained from M4 method. The 75th percentiles average and the 95th percentiles were not significantly different by the use of the four different methods (p-value = 0.48 and 0.79, respectively).

227!

228

229

230

231

232

233

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

The refrigerator temperatures measured either in the air or in food, ranged respectively between 4.1 to 8.7°C and 5.7 to 8.0°C. The average of these temperatures from air (6.4°C±1.0°C) and food (6.6°C±0.6°C) were not significantly different (p-value=0.44, Student test). Based on these results, the analysis was conducted combining air and food data. Figure 2 presents a graph of the refrigerator temperature mean (A), the 75th percentile (B) and the 95th percentile (C) per European country analyzed in this study. The refrigerator temperatures in European countries ranging from 4.1 to 8.7°C, with a

234 mean of 6.4°C±0.9°C (dotted line in figure 2A). This mean is close to the temperature that products 235 should not exceed according to the Codex guidelines (i.e. 6°C). Moreover, this value is similar to the 236 global mean air temperatures (6.1°C) measured in the 35 worldwide surveys reviewed by James et al., 2017. The 75th percentiles are comprised between 5.0 and 10.4°C, with a mean of 7.9°C±1.1°C (dotted 237 238 line in Figure 2B). The 95th percentiles are comprised between 8.0 and 13.9°C, with a mean of 239 10.3°C±1.4°C (dotted line in Figure 2C). 240 All these results indicated that, variability of temperature depends on the refrigerator age, the internal 241 thermostat that allows temperature adjustment, the frequency and the duration of door opening and the refrigerator's loading rate. 242

243

244

245

246

247

248

249

250

252

253

254

255

256

257

258

The 16 European countries were classified according to the Köppen-Geiger climate classification (Beck et al., 2018) (Figure 1). The data used for this classification correspond to the period considered in our study. The aim was to compare the temperature of domestic refrigerators according to geoclimatic zones. Our results showed that the weighted average refrigerator temperatures of the three geoclimatic groups was 6.9°C, 6.3°C and 6.6°C respectively for temperate climate with dry summer (Cs), temperate climate without dry season (Cf) and cold climate without dry season (Df) (Table 4). The Kruskal-Wallis's test showed that there was a significant difference (p-value = 0.0311) in the 251251 mean refrigerator temperature mean within the three categories. The Dunn's test showed that the mean refrigerator temperature of the Cf group countries was significantly lower than that of the Cs group countries (p-value = 0.0283). On the other hand, there was no significant difference between the Cf group and the Df group and between the Df group and the Cs group. However, in the case of Df group, the low number of studies and refrigerators investigated could explain the lack of significant results. Moreover, these results only represent between 0.6 to 10.5% of the households in European countries surveyed (Table 4). Thus, it would be interesting to recommend an update of these data according to or on the basis of more representative population data and climate change.



Figure 1. European countries where data on temperature of domestic refrigerators (from national surveys and scientific literature) were collected clustered according the Köppen-Geiger climate classification (described by Beck et al., 2018)

- Legend of figure 1: Dark grey, Cold climate without dry season (Df); Medium grey, Temperate climate without dry season (Cf); Light grey, Temperate climate with dry summer (Cs)
- 265 (Map created from https://www.mapchart.net/europe.html)

- Table 3. Collected data from national surveys and scientific literature and determination of the
- **75**th and **95**th percentile of refrigerator temperatures


Country	Reference	N refrigerator	Monitoring tool	Recorder position (Air / Food)	T° Mean ± SD (°C)	T° min (°C)	T° max (°C)	75 th percentile (°C)	95 th percentile (°C)	Percentile method
AT	Austria survey 2011	40	TF	Air	7.0 ± 2.5	2.0	13.0	$8.6 [7.6, 9.7]^a$	10.8 [9.4, 12.4] ^a	M4
BE	Devriese et al., 2004	3001	?	Air	7.0 (±?)	?	?	9.0	?	M1
DE	Geppert et al,. 2010	100	DL	Air	6.2 (±?)	0.0	11.0	7.0	10.0	M3
DK	Denmark survey 2015	77	DL	Air	6.1 ± 2.1	0.0	10.0	7.2	9.5	M2
	Carrasco et al,. 2007	30	DL	Air	6.6 ± 2.6	?	?	$8.3 [7.0, 9.6]^a$	10.5 [8.9, 12.3] ^a	M4
	Garrido et al. 2010	33	Р	Air	8.3 ± 2.8	3.9	14.6	9.3	13.5	M2
ES	Garrido et al. 2010		_	Food	8.0 ± 2.7	3.9	14.4	8.9	12.6	M2
	Geppert et al. 2010	100	DL	Air	4.1 (±?)	0.0	10.0	6.0	9.0	M3
	Jofre et al. 2019	106	DL	Air	5.4 ± 2.1	0.6	12.7	6.9	9.2	M1
	Anses 2017	3767	TH	Air	5.9 ± 2.9	-3.0	15.0	8.0	11.0	M2
				Food 1	5.7 ± 3.2	-4.8	13.8	7.4	11.1	M2
	Derens et al. 2006	251	DL	Food 2	6.1 ± 2.9	-2.0	12.5	7.9	11.7	M2
FR				Food 3	6.0 ± 2.2	1.1	10.8	7.5	9.7	M2
ľK	Derens-Bertheau et al., 2015	75	DL	Food	6.3 ± 2.2	1.1	10.7	8.1	9.9	M2
	Dumitrașcu et al., 2020	15	DL	Air	6.4 ± 1.4	4.0	8.3	7.3	8.2	M2
	Geppert et al., 2010	100	DL	Air	6.7 (±?)	1.0	12.0	9.0	12.0	M3
	Laguerre et al. 2002	118	DL	Air	6.6 ± 2.2	0.9	11.4	8.2	10.2	M2
				Air (Top area)	7.6 ± 3.0	-1.8	14.5	9.6 [8.8, 10.4] ^a	12.4 [11.3, 13.6] ^a	M4
	Koutsoumanis et al., 2010	100	DL	Air (Middle area)	6.3 ± 2.7	-0.7	13.0	$8.1 [7.4, 8.8]^a$	10.6 [9.6, 11.7] ^a	M4
GR				Air (Bottom area)	6.7 ± 2.3	-2.7	18.1	$8.2 [7.6, 8.8]^a$	10.4 [9.5, 11.3] ^a	M4
	Andritsos et al. 2021	70	DL	Air	8.7 ± 3.2	-0.8	17.5	10.4	13.9	M2
	Taoukis et al., 2005	250	DL	Air	6.3 ± 2.7	?	?	$8.1 [7.7, 8.6]^a$	$10.7 [10.0, 11.4]^a$	M4
ΙE	Kennedy et al., 2005; Roccato et al.2017	100	DL	Air	5.9 ± 2.5	-1.7	11.8	$7.6 [6.9, 8.2]^a$	9.9 [8.9, 10.9] ^a	M4
	Daminelli 2019	494	DL	Air	7.5 ± 3.1	-1.2	16.4	9.5	12.5	M1
IT	Roccato 2013	106	DL	Air	7.2 ± 2.5	-3.5	15.2	$8.9 [8.2, 9.5]^a$	11.2 [10.3, 12.2] ^a	M4
	Vegara et al. 2014	84	TH	Air	8.1 (±?)	2.5	15.9	?	10.1	M3
NL	The Netherland survey 2012	326	TH	Air	5.3 ± 1.7	?	?	$6.4 [6.2, 6.7]^a$	8.1 [7.7, 8.5] ^a	M4
NO	Dumitrașcu et al., 2020	13	DL	Air	5.4 ± 2.1	1.2	8.8	7.0	8.3	M2
NO	Rossvoll et al., 2014	46	DL	Air	6.2 (±?)	2.9	12.0	?	?	-
	Azevedo et al., 2005	86	TH	Air	7.3 ± 2.6	0.9	15.2	9.0	10.9	M2
PT	Dumitrașcu et al., 2020	15	DL	Air	5.5 ± 1.9	3.2	9.1	6.3	8.9	M2
	Galvão et al., 2017	51	DL	Air	5.5 ± 2.7	-1.8	11.9	6.9	10.5	M2
RO	Dumitrașcu et al., 2020	14	DL	Air	5.3 ± 2.2	1.7	9.5	6.4	8.7	M2
			l	Food 1	6.2 ± 2.3	0.8	11.3	7.3	10.1	M2
				Food 2	6.5 ± 2.5	0.2	12.8	7.9	10.6	M2
				Food 3	6.8 ± 2.0	2.4	13.6	8.1	9.9	M2
SE	Marklinder et al., 2004	102	DL	Food 4	6.9 ± 2.2	0.6	13.2	8.4	10.4	M2
				Food 5	7.2 ± 2.4	1.1	12.3	9.1	10.4	M2
				Food 6	7.1 ± 2.5	0.2	12.3	8.9	10.8	M2
				Food 7	6.9 ± 2.8	1.8	18.2	9.1	12.1	M2
SI	Ovca et al. 2020	50	DL	Air	5.9 ± 2.2	-0.3	10.7	7.2	9.6	M2
31				Food	6.0 ± 2.2	1.0	10.7	7.5	9.6	M2
	Evans and Redmond, 2016	43	P	Air	5.8 ± 1.9	1.8	11.4	$7.1 [6.3, 7.8]^a$	$8.7 [7.7, 9.9]^a$	M4
UK	Dumitrașcu et al., 2020	13	DL	Air	5.0 ± 2.7	-0.1	9.3	6.2	8.7	M2
UK	George et al., 2010	50	DL	Air	6.8 ± 2.3	1.5	11.6	8.4	10.2	M2
	Geppert et al. 2010	100	DL	Air	5.2 (±?)	3.0	10.0	5.0	8.0	M3

274

275

271

272






Figure 2. Graph of the refrigerator temperature mean (A), 75th percentile mean (B) and 95th percentile mean (C) and associated standard deviation per country.

276 Legend of Figure 2A: Underlined country, country for which a single study was collected.

279

Table 4: Weighted average refrigerator temperatures according to the Köppen-Geiger climate classification of European countries

Group	Number of studies	Number of refrigerators included in the studies	Weighted average refrigerator temperatures	Estimated number of households*	Proportion of households surveyed
cold climate without dry season (Df)	6	392	6.6°C	132 901 000	0.6%
temperate climate without dry season (Cf)	12	8009	6.3°C	170 590 000	10.5%
temperate climate with dry summer (Cs)	13	1525	6.9°C	127 633 000	2.8%

Legend of Table 4: *Global population number/number of people per household, sources: INSEE 2020

(for UK) and 2021 (others countries); Eurostat 2017 (for UK) and 2022 (others countries)

2821

283

280

2811

3.4. Taking these data into account for shelf-life studies

284 The results of this study were taken into account in order to revise the EURL Lm TGD for conducting 285 shelf-life studies on Listeria monocytogenes in ready-to-eat foods (EURL Lm 2019). This revision had 286 been conducted by the EURL Lm with a working group of representatives from seven EU NRLs Lm 287 (Belgium, Denmark, Ireland, Italy, Netherland, Portugal and Sweden) and one non-EU member 288 (United Kingdom). 289 This working group was also consulted in 2018 to revise the section "Storage conditions" of the 290 EURL Lm TGD for conducting shelf-life studies on Listeria monocytogenes in ready-to-eat foods 291 (EURL Lm, 2014) in order to refine the temperature at retail level. In this revision, the value of the temperature at retail level was based on the 95th percentile temperature distribution (EURL *Lm*, 2015) 292 293 and defined as consensus in the amendment 1 of version 3 of the EURL Lm TGD for conducting shelf-294 life studies on *Listeria monocytogenes* in ready-to-eat foods (EURL *Lm*, 2019). 295 To be consistent with the consensus value recommended for temperature at retail level of the EURL Lm 296 TGD v3 amendment 1 (EURL Lm, 2019), the working group decided to select the rounded value of the 95th percentile distribution as value for the temperature at consumer level. Therefore the value of 297

10°C was kept to propose the revision of the Table 4 of the EURL *Lm* TGD version 4 on challenge tests and durability studies for assessing shelf-life of ready-to-eat foods related to *Listeria monocytogenes* (EURL *Lm*, 2021).

4. CONCLUSIONS

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

determination of the shelf-life in a RTE foods.

For the first time, a study was carried out to collect data on the temperatures of household refrigerators in 16 European countries and to define a more realistic temperature to mimic the storage of RTEF at consumer stage, used in challenge tests and durability studies for shelf-life validation or verification. A temperature of 10°C, representing the 95th percentile of the studied dataset, was taken for the consumer level to revise the fourth version of the EURL Lm TGD on challenge tests and durability studies for assessing shelf-life of ready-to-eat foods related to Listeria monocytogenes (EURL Lm, 2021). The choice of a lower temperature for these tests may lead to increase shelf-life evaluations for RTEF, especially for those based on higher temperature at consumer level (e.g. 12°C as mentioned in the previous versions of the EURL Lm TDG). As RTEF shelf-life is not limited to the assessment of Lm growth, Food Business Operators may be limited by the growth of spoilage organisms and other quality criteria if they decide to increase the shelf-life of their products. This global approach should be included in a comprehensive food safety management system. These results might be also impacted by several factors, e.g. impact of climate change including overall increased temperatures and other climatic conditions at regional level, but also geopolitical situations which put pressure on energy prices, economic crisis associated to lack of financial resources particularly dedicated for keeping the cold chain. Thus, the situation of the status of home and retail temperatures should be regularly revised. Finally this should nevertheless not replace the need of continuous consumer education and awareness raising in a shared responsibility approach. These observations can be taken into account for the

324	We thank the NRL network for their collaboration and all authors who shared their raw data.
325	We also thank Laurent Guillier, Researcher in quantitative microbiology and risk assessment at
326	ANSES, for his helping regarding the statistical part.
327	
328	FOUNDING
329	EURL Listeria monocytogenes is co-funded by the European Union. Views and opinions expressed
330	are however those of the authors only and do not necessarily reflect those of the European Union or
331	the European Health and Digital Executive Agency (HaDEA). Neither the European Union nor
332	HaDEA can be held responsible for them.

ACKNOWLEDGEMENTS

333 **REFERENCES**

356

357

Andritsos N. D., Stasinou V., Tserolas D., Giaouris E. 2021. "Temperature distribution and hygienic 334 status of domestic refrigerators in Lemnos island, Greece". Food Control 127: 108121 https://doi.org/10.1016/j.foodcont.2021.108121. 335 Anses, 2017. Avis n° « 2014-SA-0234 » relatif à la troisième étude individuelle nationale des 336 3 *INCA 3)*. consommations alimentaires (Etude https://www.anses.fr/sites/default/files/NUT2014SA0234Ra.pdf 337 338 household." Austria 2011. "Food safety and hygiene in the private 339 https://www.ages.at/download/sdl-340 eyJ0eXAiOiJKV1QiLCJhbGciOiJIUzI1NiJ9.eyJpYXQiOjE2MDk0NTkyMDAsImV4cCI6N 341 DA3MDkwODgwMCwidXNlciI6MCwiZ3JvdXBzIjpbMCwtMV0sImZpbGUiOiJmaWxlY 342 WRtaW5cL0FHRVNfMjAyMlwvMl9NRU5TQ0hcL0Vyblx1MDBlNGhydW5nX0xlYmVuc 343 21pdHRlbFwvRXJuXHUwMGU0aHJ1bmdzZW1wZmVobHVuZ2VuXC9TaWNoZXJfa29i a GVuXC9MZWJlbnNtaXR0ZWxzaWNoZXJoZWl0X3VuZF9IeWdpZW5lX2ltX1ByaXZhd344 345 GhhdXNoYWx0XzEzXzEyXzIwMTMucGRmIiwicGFnZSI6MTM5N30.bbm1C4eyGJzqDZ 1U Rm0Hq-4t-346 347 toV0hmxb4bBx4hEhc/Lebensmittelsicherheit und Hygiene im Privathaushalt 13 12 2013 348 .pdf. 349 Azevedo, I., Regalo, M., Mena, C., Almeida, G., Carneiro, L., Teixeira, P., Hogg, T., Gibbs, P.A., 2005. 350 "Incidence of Listeria spp. in domestic refrigerators in Portugal." Food Control 16 (2): 121-351 124. https://doi.org/10.1016/j.foodcont.2003.12.006. 352 Beck, H. E., Zimmermann, N. E., McVicar, T.R., Vergopolan, N., Berg, A., Wood, E.F., 2018. "Present 353 and future Koppen-Geiger climate classification maps at 1-km resolution." Sci Data 5: 180214. 354 https://doi.org/10.1038/sdata.2018.214. 355 Carrasco, E., Perez-Rodriguez, F., Valero A., Garcia-Gimeno, R.M., Zurera, G., 2007. "Survey of

temperature and consumption patterns of fresh-cut leafy green salads: risk factors for

listeriosis." J Food Prot 70 (10): 2407-12. https://doi.org/10.4315/0362-028x-70.10.2407.

- 358 Food and Agriculture Organization of the United Nation (FAO) and World Health Organization 359 (WHO). 2007. Codex Alimentarius guidelines on the application of the general principles of 360 food hygiene to the control of Listeria monocytogenes in foods (CAC/GL 61 - 2007). 361 Daminelli, P.. 2019. National survey of domestic refrigerators temperatures in Italy. edited by Network 362 of the Italian Experimental Zooprophylactic Institutes (I.I.Z.Z.S.S.). Research Project IZS LER 363 2016/018 RC. Research funded by the Italian Ministry of Health. Unpublished results. 364 Denmark survey, 2015. Investigate the temperature of elderly people refrigerators. Unpublished 365 results. Derens-Bertheau, E., Osswald ,V., Laguerre, O., Alvarez, G., 2015. "Cold chain of chilled food in 366 367 France." International Journal of Refrigeration 52: 161-167. 368 https://doi.org/10.1016/j.ijrefrig.2014.06.012. Derens, E., Palagos B., Guilpart, J., 2006. "The cold chain of chilled products under supervision in 369 370 France." IUFoST World Congress, 13th World Congress of Food Science & Technology. 371 https://doi.org/10.1051/IUFoST:20060823. 372 Devriese, S., Huybrechts, I., Moreau, M., Van Oyen, H.. 2004. Enquête de consommation alimentaire *2004*. Institut scientifique 373 Belge de Santé publique. https://www.wiv-374 isp.be/Epidemio/epifr/foodfr/food04fr/food1fr.pdf. 375 Dumitrascu, L., Nicolau, A.I., Neagu, C., Didier, P., Maître, I., Nguyen-The, C., Skuland, S.E., Møretrø, T., Langsrud, S., Truninger, M., Teixeira, P., Ferreira, V., Martens, L., Borda, D., 376 2020. "Time-temperature profiles and Listeria monocytogenes presence in refrigerators from 377 378 households with vulnerable consumers." Food Control 111. 379 https://doi.org/10.1016/j.foodcont.2019.107078.
- EFSA BIOHAZ Panel (EFSA Panel on Biological Hazards), 2018. Scientific Opinion on the *Listeria*monocytogenes contamination of ready-to-eatfoods and the risk for human health in the EU.

 EFSA Journal 2018;16(1):5134, 173 pphttps://doi.org/10.2903/j.efsa.2018.5134.

383	European Food Safety Authority (EFSA), European Centre for Disease Prevention and Control
384	(ECDC). 2023. "The European Union One Health 2022 Zoonoses Report." EFSA Journal 21
385	(12): e8442. https://doi.org/10.2903/j.efsa.2023.8442.
386	EN ISO 20976-1. 2019. Microbiology of the Food Chain—Guidelines for Conducting Challenge Tests
387	of Food and Feed Products—Part 1: Challenge Tests to Study the Growth Potential, Lag Time
388	and the Maximum Growth Rate.
389	European Union Reference Laboratory for Listeria monocytogenes (EURL Lm). 2014. "EURL Lm
390	Technical Guidance Document for conducting shelf-life studies on Listeria monocytogenes in
391	ready-to-eat foods – version 3 of 6 July 2014." https://doi.org/10.5281/zenodo.10651272.
392	European Union Reference Laboratory for Listeria monocytogenes (EURL Lm). 2015. "Report on
393	temperature distribution of ready-to-eat food at retail level in Europe"
394	https://zenodo.org/records/11073001.
395	European Union Reference Laboratory for Listeria monocytogenes (EURL Lm). 2019. "EURL Lm
396	Technical Guidance Document for conducting shelf-life studies on Listeria monocytogenes in
397	ready-to-eat foods- version 3 amendement 1 of 21 February 2019"
398	https://doi.org/10.5281/zenodo.10651320.
399	European Union Reference Laboratory for Listeria monocytogenes (EURL Lm). 2021. "EURL Lm
400	Technical Guidance Document on challenge tests and durability studies for assessing shelf-life
401	of ready-to-eat foods related to Listeria monocytogenes - version 4 of 1 July 2021."
402	https://doi.org/10.5281/zenodo.7760700.
403	Evans, E. W., ad Redmond, E.C., 2016. "Time-Temperature Profiling of United Kingdom Consumers'
404	Domestic Refrigerators." J Food Prot 79 (12): 2119-2127. https://doi.org/10.4315/0362-
405	028X.JFP-16-270.
406	Galvão, D., Dinis Gaspar, P., Dinho Da Silva, P., Pires L., 2017. "Thermal Performance, Usage
407	Behaviour and Food Waste of Domestic Refrigerators in a University Student Community:

- Findings Towards Cities Sustainability." The Sustainable City XII., 223, 539-550.
- 409 http://dx.doi.org/10.2495/SC170471
- 410 Garrido, V., García-Jalón I., Vitas A.I., 2010. "Temperature distribution in Spanish domestic
- refrigerators and its effect on *Listeria monocytogenes* growth in sliced ready-to-eat ham." *Food*
- 412 *Control* 21 (6): 896-901. https://doi.org/10.1016/j.foodcont.2009.12.007.
- 413 George, R.M., Burgess, P.J., Thorn, R.D., 2010. Reducing Food Waste through the Chill Chain. An
- extensive study of consumer refrigerated food practices in the home and experimental
- measurements of refrigerated food temperatures during their transit from the retail store to the
- domestic fridge. Final report for WRAP funded project RSC007-003 Banbury, UK
- 417 Geppert, J., Reger, O., Bichler, S., Stamminger, R., 2010. "Cold storage in private households:
- recommendations and consumer real life behaviour." 4th Cold chain management workshop,
- 419 Bonn, DE..
- James, C., Bukola, A.O., Stephen J.J., 2017. "The Use and Performance of Household Refrigerators: A
- Review." Comprehensive Reviews in Food Science and Food Safety 16 (1): 160-179.
- 422 https://doi.org/10.1111/1541-4337.12242.
- James, S.J., Evans, J., James, C., 2008. "A review of the performance of domestic refrigerators."
- 424 *Journal of Food Engineering* 87 (1): 2-10. https://doi.org/10.1016/j.jfoodeng.2007.03.032.
- Jofre, A., Latorre-Moratalla, M.L., Garriga, M., Bover-Cid, S., 2019. "Domestic refrigerator
- temperatures in Spain: Assessment of its impact on the safety and shelf-life of cooked meat
- 427 products." *Food Res Int* 126: 108578. https://doi.org/10.1016/j.foodres.2019.108578.
- 428 Kennedy, J., Jackson, V., Blair, I.S., McDowell, D.A., Cowan, C., Bolton D.J., 2005. "Food safety
- knowledge of consumers and the microbiological and temperature status of their refrigerators."
- 430 J Food Prot 68 (7): 1421-30. https://doi.org/10.4315/0362-028x-68.7.1421.
- 431 Koutsoumanis, K., Pavlis, A., Nychas, G.J., Xanthiakos, K, 2010. "Probabilistic model for
- 432 Listeria monocytogenes growth during distribution, retail storage, and domestic storage of

- pasteurized milk." *Appl Environ Microbiol* 76 (7): 2181-91. https://doi.org/10.1128/AEM.02430-09.

 Laguerre, O., Derens, E., Palagos, B., 2002. "Study of domestic refrigerator temperature and analysis
- of factors affecting temperature: a French survey." I.J.R.,25,5,653:659.
- 437 https://doi.org/10.1016/S0140-7007(01)00047-0.
- 438 Marklinder, I. M., M. Lindblad, L. M. Eriksson, A. M. Finnson, and R. Lindqvist. 2004. "Home storage
- temperatures and consumer handling of refrigerated foods in Sweden." *J Food Prot* 67 (11):
- 440 2570-7. https://doi.org/10.4315/0362-028x-67.11.2570.
- 441 Ovca, A., Skufca, T., Jevsnik, M., 2020. "Temperatures and storage conditions in domestic
- refrigerators Slovenian scenario." Food Control: 107715.
- https://doi.org/10.1016/j.foodcont.2020.107715.
- 444 Pouillot, R., Miconnet, N., Afchain A.L., Delignette-Muller, M. L., Beaufort, A., Rosso, L., Denis, J.
- B., Cornu, M., 2007. "Quantitative risk assessment of Listeria monocytogenes in French cold-
- smoked salmon: I. Quantitative exposure assessment." Risk Anal 27 (3): 683-700.
- 447 https://doi.org/10.1111/j.1539-6924.2007.00921.x.
- Roccato, A. 2013. "Survey conducted in Italy on the consumer refrigeration temperatures and their
- impact on food safety illustrated with *Salmonella*." IAFP 15th-17th May 2013, Marseille, FR.
- Roccato, A., Uyttendaele, M., Membré, J. M., 2017. "Analysis of domestic refrigerator temperatures
- and home storage time distributions for shelf-life studies and food safety risk assessment."
- 452 Food Res Int 96: 171-181. https://doi.org/10.1016/j.foodres.2017.02.017.
- Rosa Rodrigues de Souza, C., Bergis, H., Ng, P., Guillier, L., Felix, B., Leclercq, A., Gnanou Besse
- N., 2023. "Assessment of the relationship between the MLST genetic diversity of Listeria
- 455 monocytogenes and growth under selective and non-selective conditions." Food Microbiol
- 456 114: 104303. https://doi.org/10.1016/j.fm.2023.104303.
- Rossvoll, E., Ronning, H.T., Granum, P.E., Moretro, T., Hjerpekjon, M.R., Langsrud S., 2014. "Toxin
- 458 production and growth of pathogens subjected to temperature fluctuations simulating consumer

459	handling	of	cold	cuts."	Int	J	Food	Microbiol	185:	82-92.			
460	https://doi.org/10.1016/j.ijfoodmicro.2014.05.020.												
461	Taoukis, P.S., Gian	nakou	rou, M.C	C., Koutso	umanis,	K., B	Bakalis, S.,	2005. "Mode	lling the	Effect of			
462	House Hold	l Chil	led Stora	age Condi	tions or	the	Risk Dist	ribution of M	eat Produ	icts." III			
463	Internationa	1 Sym	posium (on Applica	ations of	f Mod	elling as a	nn Innovative	Technolog	gy in the			
464	Agri-Food (Chain;	MODEL	-IT, 29 th N	May 200	5, Leu	iven, BE.						
465	The Netherland sur	vey. 20	012. Foo	d waste. U	npublis	ned res	sults.						
466	Vegara, A., Festino	, A.R.,	Di Cicci	o, P., Cost	anzo, C	, Penn	nisi, L., Ian	ieri, A., 2014.	"The man	agement			
467	of the dome	stic re	frigeration	on: microb	oiologica	ıl statı	us and ten	nperature." <i>Bri</i>	tish Food	l Journal			
470	116 (6): 104	7-105	7. https:/	/doi.org/10	0.1108/I	3FJ-05	5-2012-010	03.					
471	World Health Orga	nizatio	on., 2022	Listeria	топосу	togene	es in read	y-to-eat (RTE)	food: att	ribution,			
472	characteriza	ation a	ınd moni	toring: me	eeting re	port.	Food and	Agriculture O	rganizatio	on of the			
473	United Nation	ons.											