



**HAL**  
open science

# Theory of electromagnetic probes

Jacopo Ghiglieri

► **To cite this version:**

Jacopo Ghiglieri. Theory of electromagnetic probes. LHCb IFT workshop, Jul 2024, Santiago de Compostela, Spain. hal-04634560

**HAL Id: hal-04634560**

**<https://hal.science/hal-04634560v1>**

Submitted on 4 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Theory of electromagnetic probes



**NUCLÉAIRE  
& PARTICULES**




Jacopo Ghiglieri, SUBATECH, Nantes

LHCb IFT Workshop, July 1st 2024



# Relativistic Heavy-Ion Collisions

made by Chun Shen






# How EM probes are made

(and what they tell us)

- The hard partonic processes in the heavy ion collision produce quarks, gluons and *prompt photons and dileptons*,  $W$  and  $Z$  bosons. They can tell us about nPDFs, photon-tagged hadrons and jets
- At a later stage, quarks and gluons form a plasma
- Scatterings of (pre)thermal partons produce QGP *photons and dileptons*. thermalisation,  $T$ , hydro
- A jet traveling can radiate *jet-thermal photons*. Jet quenching
- Later on, hadronization. *hadron gas photons and dileptons*.  $T$ ,  $T_c$ , hydro
- (Some) hadrons decay into *decay photons and dileptons*



# In this talk



- Theoretical description: **convolution** of **microscopic rates** over the **macroscopic (hydro) evolution** of the medium
- In this talk
  - overview and recent results on the **microscopic rates**, mostly up to the *thermal phase*
  - Photons and dileptons in and before equilibrium from pQCD

# How to compute rates

- $\alpha \ll 1$  implies that photon production is a rare event and that rescatterings and back-reactions are negligible: medium is transparent to / not cooled by photons
- At leading order in QED and to all orders in QCD the **photon** and **dilepton** rates are given by

$$\frac{dN_{\gamma}}{d^4 X d^3 k} = \frac{d\Gamma_{\gamma}(k)}{d^3 k} = -\frac{\alpha}{4\pi^2 k} \int d^4 X e^{iK \cdot X} \text{Tr} \rho J^{\mu}(0) J_{\mu}(X)$$

$$\frac{d\Gamma_{l+l-}(k)}{dk^0 d^3 k} = -\frac{\alpha^2}{6\pi^3 K^2} \int d^4 X e^{iK \cdot X} \text{Tr} \rho J^{\mu}(0) J_{\mu}(X)$$


$\rho$  density operator,  $J$  electromagnetic current



# The basics of pQCD photons


$$\frac{d\Gamma_\gamma(k)}{d^3k} = -\frac{\alpha}{4\pi^2 k} \int d^4X e^{iK \cdot X} \text{Tr} \rho J^\mu(0) J_\mu(X) \quad J^\mu = \sum_{q=u,d,s} e_q \bar{q} \gamma^\mu q : \text{~}$$

- Real, hard photon:  $k^0 = k \gtrsim T$
- At one loop ( $\alpha_{\text{EM}} g^0$ ):




Kinematically forbidden. Need to kick one of the quarks off-shell. Works for dileptons

- Leading order photon is  $\alpha_{\text{EM}} g^2$
- Strength of the kick (virtuality) naturally divides the calculation in the distinct  $2 \leftrightarrow 2$  processes and collinear processes




# 2 ↔ 2 processes

- Cut two-loop diagrams ( $\alpha_{\text{EM}} g^2$ )



2 ↔ 2 processes (with crossings and interferences):




$$\int_{\text{ph. space}} f(p) f(p') (1 \pm f(k')) |\mathcal{M}|^2 \delta^4(P + P' - K - K')$$

- Equivalence with kinetic theory: **distributions** x **matrix elements**
- IR divergence (Compton) when  $t$  goes to zero




# $2 \leftrightarrow 2$ processes

- The IR divergence disappears when **Hard Thermal Loop** resummation is performed [Braaten Pisarski NPB337 \(1990\)](#)



# 2 ↔ 2 processes

- The IR divergence disappears when **Hard Thermal Loop** resummation is performed [Braaten Pisarski NPB337 \(1990\)](#)




- In the end one obtains the result

$$\left. \frac{d\Gamma_\gamma}{d^3k} \right|_{2\leftrightarrow 2} \propto e^2 g^2 \left[ \log \frac{T}{m_\infty} + C_{2\leftrightarrow 2} \left( \frac{k}{T} \right) \right]$$

[Kapusta Lichard Siebert PRD44 \(1991\)](#) [Baier Nakkagawa Niegawa Redlich ZPC53 \(1992\)](#)



# Collinear processes





- These diagrams contribute to LO if small ( $g$ ) angle radiation/annihilation [Aurenche Gelis Kobes Petitgirard Zaraket 1998-2000](#)
- Photon formation times is then of the same order of the soft scattering rate  $\Rightarrow$  interference: *LPM effect*
- Requires resummation of infinite number of ladder diagrams

$$\frac{d\Gamma_\gamma}{d^3k} \Big|_{\text{coll}} = \text{Re} \left( \left( \text{Ladder Diagram} \right)^* \left( \text{Ladder Diagram} \right) \right)$$

[AMY \(Arnold Moore Yaffe\) JHEP 0111, 0112, 0226 \(2001-02\)](#)

# QGP photons

Thermal photon rate,  $\alpha_s=0.2$



LO: AMY (2001-02) NLO: JG Hong Kurkela Lu Moore Teaney JHEP0503 (2013)


# Hadronic photons





# Hadronic photons

## Hadron Gas



+bremsstrahlung



6(15)









# Hadronic photons



Schäfer Garcia-Montero Paquet Elfner Gale 2111.13603

# From QGP photons to data

- The macroscopic expansion of the fireball gives rise to blueshift: the observed inverse slope of the direct photon spectrum at low  $p_T$  is not the real temperature





Shen Heinz Paquet Gale 1308.2440



# From QGP photons to data

- Importance of viscous corrections, i.e. perturbations around partonic equilibrium from shear flow




Shen Heinz Paquet Gale 1308.2440

# Bottom-up thermalisation

- Competition between **expansion** and **interaction**, **attractor** solution when they balance out

$$\left( \frac{\partial}{\partial t} - \frac{p_z}{\tau} \frac{\partial}{\partial p_z} \right) f(\mathbf{p}) = -C^{2 \leftrightarrow 2} - C^{1 \leftrightarrow 2}$$

- **Expansion** is driven by the specifics of the heavy-ion collision and the initial state, drives the system away from equilibrium. **Interaction** among the constituents tends to isotropize the system.



Baier Mueller Schiff Son (2001) Kurkela Moore (2011)

fig. Mazeliauskas





fig. Kurkela



- Initially
- strong isotropizing effect of interactions
- gluon-rich system
- Later, **bremstrahlung** in the cascade and mini-jet quench

Baier Mueller Schiff Son (2001) Kurkela Moore (2011)





# Pre-equilibrium photons

- Two competing effects:

↓ Quark number suppression and short duration of the phase

↑ Higher energy for the early quarks



Garcia-Montero Mazeliauskas Plaschke Schlichting **2308.09747**



# Pre-equilibrium photons

- Two competing effects:

↓ Quark number suppression and short duration of the phase


↑ Higher energy for the early quarks

See also

Churchill Yan Jeon Gale [2008.02902](#)


Garcia-Montero [1909.12294](#)

Garcia-Montero Mazeliauskas Plaschke Schlichting [2308.09747](#)



# pQCD dileptons

fig. Jackson





# pQCD dileptons

- Consider non-zero virtuality  $k^0 > k \geq 0$ .

$$\frac{d\Gamma_{l+l-}(k)}{dk^0 d^3k} = -\frac{\alpha^2}{6\pi^3 K^2} \int d^4X e^{iK \cdot X} \text{Tr} \rho J^\mu(0) J_\mu(X)$$



- Born contribution present, gets larger as **invariant mass  $M^2=K^2$**  grows



# pQCD dileptons

- Consider non-zero virtuality  $k^0 > k \geq 0$ .
- If  $K^2 \sim T^2$  loop corrections: real and virtual (with IR cancellations)

$$\frac{d\Gamma_{l+l-}(k)}{dk^0 d^3k} = -\frac{\alpha^2}{6\pi^3 K^2} \int d^4X e^{iK \cdot X} \text{Tr} \rho J^\mu(0) J_\mu(X)$$



NLO results [Laine JHEP1311 \(2013\)](#) extended to spacelike region in [Jackson PRD100 \(2019\)](#)




# pQCD dileptons

- Consider non-zero virtuality  $k^0 > k \geq 0$ .

$$\frac{d\Gamma_{l+l-}(k)}{dk^0 d^3k} = -\frac{\alpha^2}{6\pi^3 K^2} \int d^4X e^{iK \cdot X} \text{Tr} \rho J^\mu(0) J_\mu(X)$$



- If  $K^2 \ll T^2$  LPM and/or HTL resummations are again necessary, similar to  $K^2=0$   
 Braaten Pisarski Yuan **PRL64** (1990),  
 Aurenche Gelis Moore Zaraket **JHEP0212** (2002)  
 NLO results JG Moore **JHEP1412** (2014)
- Finite- $k$  rate available at NLO for all  $K^2 \geq 0$   
 Ghisoiu Laine **JHEP1410** (2014) JG Moore (2014)  
 JG Laine, **JHEP2201** (2021)



# pQCD dileptons

rate from a **static source**:

$$\frac{d\Gamma_{\ell\bar{\ell}}}{dM} = \int_{\mathbf{k}} \frac{M}{\sqrt{M^2 + k^2}} \frac{d\Gamma_{\ell\bar{\ell}}}{d\omega d^3\mathbf{k}}$$




Churchill Du Gale Jackson Jeon 2311.06951 2311.06675

# Connecting pQCD dilepton rates to data

$$M \gg T \quad \frac{d\Gamma_{\ell\bar{\ell}}}{dM} \propto (MT_{\text{eff}})^{3/2} \exp(-M/T_{\text{eff}})$$

- Look at intermediate masses
- Full stack to connect microscopic rates to data
- MC-Glauber IC
- MUSIC hydro
- Freeze-out and hadronic afterburner with UrQMD




Churchill Du Gale Jackson Jeon **2311.06951 2311.06675**

# Connecting pQCD dilepton rates to data

$$M \gg T \quad \frac{d\Gamma_{\ell\bar{\ell}}}{dM} \propto (MT_{\text{eff}})^{3/2} \exp(-M/T_{\text{eff}})$$

- Linear relation between highest temperature in the hydro phase and effective temperature in the intermediate mass regime
- Predicted in [Kajantie Miettinen \(1981\)](#)



Churchill Du Gale Jackson Jeon [2311.06951](#) [2311.06675](#)





# Pre-equilibrium dileptons

- Two competing effects:

↓ Quark number suppression and short duration of the phase

↑ Higher energy for the early quarks




# Pre-equilibrium dileptons

- Two competing effects:

↓ Quark number suppression and short duration of the phase

↑ Higher energy for the early quarks

$$K \text{ (wavy)} \rightarrow \text{circle} \leftarrow K \text{ (wavy)} = \left| \text{triangle} \right|^2$$





- Quadrupole moment of the angular distribution sensitive to change in polarization

# Pre-equilibrium dileptons

- Two competing effects:

↓ Quark number suppression and short duration of the phase

↑ Higher energy for the early quarks




# Conclusions

- Theoretical description: **convolution** of **microscopic rates** over the **macroscopic (hydro) evolution** of the medium
- Many theory advancements in connecting the two and in pre-equilibrium emission
  - Pre-eq. photon and dileptons overtake thermal radiation at some point
  - Linear relationship between highest temperature and effective temperature for dileptons
  - Angular structure of dileptons could shed light on their origin



# Backup





And the lattice?

# And the lattice?

- What is measured directly is the Euclidean correlator

$$G_E(\tau, k) = \int d^3x J_\mu(\tau, \mathbf{x}) J_\mu(0, 0) e^{i\mathbf{k}\cdot\mathbf{x}}$$

# And the lattice?

- What is measured directly is the Euclidean correlator

$$G_E(\tau, k) = \int d^3x J_\mu(\tau, \mathbf{x}) J_\mu(0, 0) e^{i\mathbf{k}\cdot\mathbf{x}}$$

- Analytical continuation  $G_E(\tau, k) = G^<(i\tau, k)$

$$G_E(\tau, k) = \int_0^\infty \frac{dk^0}{2\pi} \rho_V(k^0, k) \frac{\cosh(k^0(\tau - 1/2T))}{\sinh(\frac{k^0}{2T})} \quad W^<(K) = n_B(k^0) \rho_V(k^0, k)$$



# And the lattice?

- What is measured directly is the Euclidean correlator

$$G_E(\tau, k) = \int d^3x J_\mu(\tau, \mathbf{x}) J_\mu(0, 0) e^{i\mathbf{k} \cdot \mathbf{x}}$$

- Analytical continuation  $G_E(\tau, k) = G^<(i\tau, k)$

$$G_E(\tau, k) = \int_0^\infty \frac{dk^0}{2\pi} \rho_V(k^0, k) \frac{\cosh(k^0(\tau - 1/2T))}{\sinh(\frac{k^0}{2T})} \quad W^<(K) = n_B(k^0) \rho_V(k^0, k)$$

- It contains much more info (**full spectral function**), but hidden in the **convolution**. Inversion tricky, discrete dataset with errors

# And the lattice?

- What is measured directly is the Euclidean correlator


$$G_E(\tau, k) = \int d^3x J_\mu(\tau, \mathbf{x}) J_\mu(0, 0) e^{i\mathbf{k}\cdot\mathbf{x}}$$

- Analytical continuation


$$G_E(\tau, k) = G^<(i\tau, k)$$

$$G_E(\tau, k) = \int_0^\infty \frac{dk^0}{2\pi} \rho_V(k^0, k) \frac{\cosh(k^0(\tau - 1/2T))}{\sinh(\frac{k^0}{2T})}$$

$$W^<(K) = n_B(k^0) \rho_V(k^0, k)$$




- If  $k > 0$  *spf* describes **DIS** ( $k^0 < k$ ), photons ( $k^0 = k$ ) and **dileptons** ( $k^0 > k$ )



- Plots and spectral function from Jackson Laine **JHEP1109 (2019)**  
 Quenched lattice from JG Kaczmarek Laine F.Meyer **PRD94 (2016)**

# And the lattice?

- If  $k > 0$  *spf* describes **DIS** ( $k^0 < k$ ), photons ( $k^0 = k$ ) and **dileptons** ( $k^0 > k$ )






# Fitting to the lattice


- Main idea: assume a fitting form for the spf, relying on Ansätze
- Get the Euclidean correlator from this ansatz spf and fit the spf coeffs to the lattice data
- Two approaches so far
  - Quenched, continuum extrapolated lattice data, **standard vector spf**  $\rho_V = 2\rho_T + \rho_L$
  - Convolution dominated by (well-understood) vacuum physics at  $\omega \gg k$

JG Kaczmarek Laine F.Meyer **PRD94** (2016)




# Fitting to the lattice

- Main idea: assume a fitting form for the spf, relying on Ansätze
  - Get the Euclidean correlator from this ansatz spf and fit the spf coeffs to the lattice data
  - Two approaches so far
    - $N_f=2$  continuum extrapolated, **modified spf**  
 $\rho_{\text{Mainz}} = 2\rho_T - 2\rho_L$
    - Vacuum contribution vanishes identically (Lorentz invariance).  $\rho_{\text{Mainz}}(\omega=k) = \rho_V(\omega=k)$
- Brandt Francis Harris H.Meyer Steinberg  
 1710.07050 Cè Harris H.Meyer Steinberg Toniato  
 PRD102 (2020)




# Fitting to the lattice

- Define  $D_{\text{eff}}(k) \equiv \begin{cases} \frac{\rho_V(k, \mathbf{k})}{2\chi_q k} & , k > 0 \\ \lim_{\omega \rightarrow 0^+} \frac{\rho^{ii}(\omega, \mathbf{0})}{3\chi_q \omega} & , k = 0 \end{cases}$




- In the hydro limit  $k \ll T$   $D_{\text{eff}} \rightarrow D$   $\sigma = e^2 \sum_{f=1}^{N_f} Q_f^2 \chi_q D$




- If  $k > 0$  *spf* describes **DIS** ( $k^0 < k$ ), photons ( $k^0 = k$ ) and **dileptons** ( $k^0 > k$ )

$$T = 1.2T_c, N_f = 2$$



$$T = 1.2T_c, N_f = 2$$



- Plots and spectral function from Jackson Laine **JHEP1109 (2019)**  
Lattice from Cè Harris H.Meyer Steinberg Toniato **PRD102 (2020)**



## LO pQCD spf

- Backus-Gilbert method: linear map from the space of functions in the time domain,  $G$ , to the space of functions on the frequency domain,  $\rho_{\text{BG}}$
- It is exact for constant spfs and advantageous for a slowly varying spf
- The Mainz spf might indeed be slowly varying, or at least much slower than the vector one

