



HAL
open science

Theoretical analysis of a finite-volume scheme for a stochastic Allen-Cahn problem with constraint

Caroline Bauzet, Cédric Sultan, Guy Vallet, Aleksandra Zimmermann

► To cite this version:

Caroline Bauzet, Cédric Sultan, Guy Vallet, Aleksandra Zimmermann. Theoretical analysis of a finite-volume scheme for a stochastic Allen-Cahn problem with constraint. 2024. hal-04636670

HAL Id: hal-04636670

<https://hal.science/hal-04636670v1>

Preprint submitted on 5 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Theoretical analysis of a finite-volume scheme for a stochastic Allen-Cahn problem with constraint

Caroline Bauzet*, Cédric Sultan*, Guy Vallet ‡
Aleksandra Zimmermann§

July 5, 2024

Abstract

The aim of this contribution is to address the convergence study of a time and space approximation scheme for an Allen-Cahn problem with constraint and perturbed by a multiplicative noise of Itô type. The problem is set in a bounded domain of \mathbb{R}^d (with $d = 2$ or 3) and homogeneous Neumann boundary conditions are considered. The employed strategy consists in building a numerical scheme on a regularized version “à la Moreau-Yosida” of the constrained problem, and passing to the limit simultaneously with respect to the regularization parameter and the time and space steps, denoted respectively by ϵ , Δt and h . Combining a semi-implicit Euler-Maruyama time discretization with a Two-Point Flux Approximation (TPFA) scheme for the spatial variable, one is able to prove, under the assumption $\Delta t = \mathcal{O}(\epsilon^{2+\theta})$ for a positive θ , the convergence of such a “ $(\epsilon, \Delta t, h)$ ” scheme towards the unique weak solution of the initial problem, *a priori* strongly in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$ and *a posteriori* also strongly in $L^p(0, T; L^2(\Omega \times \Lambda))$ for any finite $p \geq 1$.

Keywords: Stochastic non-linear parabolic equation with constraint • Multiplicative Lipschitz noise • Finite-volume method • Variational approach • Convergence analysis • Multivoque maximal monotone operator • Differential inclusion • Lagrange multiplier.

Mathematics Subject Classification (2020): 60H15 • 35K05 • 65M08.

* Aix Marseille Univ, CNRS, Centrale Med, LMA, Marseille, France, caroline.bauzet@univ-amu.fr, cedric.sultan@univ-amu.fr

‡ Univ Pau & Pays Adour, LMAP, UMR CNRS 5142, IPRA, Pau, France, guy.vallet@univ-pau.fr

§ TU Clausthal, Institut für Mathematik, Clausthal-Zellerfeld, Germany, aleksandra.zimmermann@tu-clausthal.de

1 Introduction

We consider Λ a bounded, open, connected, and polygonal set of \mathbb{R}^d (with $d = 2$ or 3), and $(\Omega, \mathcal{A}, \mathbb{P})$ a probability space endowed with a right-continuous, complete filtration $(\mathcal{F}_t)_{t \geq 0}$. For $T > 0$, we are interested in finding a pair (u, ψ) with $\psi \in \partial I_{[0,1]}(u)$, satisfying the following time noise-driven Allen-Cahn equation:

$$\begin{aligned} du + (\psi - \Delta u) dt &= g(u) dW(t) + (\beta(u) + f) dt, & \text{in } \Omega \times (0, T) \times \Lambda; \\ u(0, \cdot) &= u_0, & \text{in } \Omega \times \Lambda; \\ \nabla u \cdot \mathbf{n} &= 0, & \text{on } \Omega \times (0, T) \times \partial\Lambda; \end{aligned} \tag{1.1}$$

where $(W(t))_{t \geq 0}$ is a standard, one-dimensional Brownian motion with respect to $(\mathcal{F}_t)_{t \geq 0}$ on $(\Omega, \mathcal{A}, \mathbb{P})$ and \mathbf{n} denotes the unit normal vector to $\partial\Lambda$ outward to Λ . As mentioned in [11], the sub-differential $\partial I_{[0,1]}$ represents a physical constraint on the solution of (1.1) forcing it to remain bounded in $(0, 1)$ by the presence of the Lagrange multiplier $\psi \in \partial I_{[0,1]}(u)$. More precisely, our equation in (1.1), can be written as a differential inclusion in the following manner:

$$\beta(u) + f - \partial_t \left(u - \int_0^\cdot g(u) dW \right) + \Delta u \in \partial I_{[0,1]}(u)$$

where the stochastic integral is understood in the sense of Itô and the sub-differential of the indicator function $I_{[0,1]} : \mathbb{R} \rightarrow \mathbb{R} \cup \{+\infty\}$ defined by

$$I_{[0,1]}(r) = \begin{cases} 0 & \text{if } r \in [0, 1] \\ +\infty & \text{else} \end{cases}$$

is the set-valued mapping $\partial I_{[0,1]} : [0, 1] \rightarrow \mathcal{P}(\mathbb{R})$ defined by

$$\partial I_{[0,1]}(r) = \begin{cases} \{0\} & \text{if } r \in (0, 1) \\ (-\infty, 0] & \text{if } r = 0 \\ [0, \infty) & \text{if } r = 1. \end{cases}$$

We consider the following assumptions on the data:

\mathcal{A}_1 : $u_0 \in L^2(\Omega; L^2(\Lambda))$ is \mathcal{F}_0 -measurable and verifies $0 \leq u_0(\omega, x) \leq 1$, for almost all $(\omega, x) \in \Omega \times \Lambda$.

\mathcal{A}_2 : $g : \mathbb{R} \rightarrow \mathbb{R}$ is a L_g -Lipschitz-continuous function (with $L_g \geq 0$), such that $\text{supp } g \subset [0, 1]$.

\mathcal{A}_3 : $\beta : \mathbb{R} \rightarrow \mathbb{R}$ is a L_β -Lipschitz-continuous function (with $L_\beta \geq 0$) such that for convenience $\beta(0) = 0$.

\mathcal{A}_4 : $f \in L^2_{\mathcal{P}_T}(\Omega \times (0, T); L^2(\Lambda))^\dagger$.

[†]For a given separable Banach space X , we denote by $L^2_{\mathcal{P}_T}(\Omega \times (0, T); X)$ the space of the predictable X -valued processes ([30] p.94 or [51] p.27). This space is the space $L^2(\Omega \times (0, T); X)$ for the product measure $d\mathbb{P} \otimes dt$ on the predictable σ -field \mathcal{P}_T (*i.e.* the σ -field generated by the sets $\mathcal{F}_0 \times \{0\}$ and the rectangles $A \times (s, t]$, for any $s, t \in [0, T]$ with $s \leq t$ and $A \in \mathcal{F}_s$).

1.1 Literature review

The equation in (1.1) is known in the literature as an Allen-Cahn type equation with constraint. It is applicable in modeling several physical phenomena, like phase transitions. In [11], a global existence and uniqueness result for this equation has been proposed to model the evolution of damage in continuum media. More precisely, it has been assumed that the solution u to (1.1) is a damage parameter, *i.e.* the local proportion of active cohesive bonds in the micro-structure of a material. Then, the function f on the right-hand side of (1.1) denotes an external source of damage (mechanical or chemical), while the nonlinear source term β is associated with the material's internal cohesion. A constraint was incorporated within the equation to restrict the values of u to the interval $[0, 1]$. This constraint has a physical meaning in the way that $u = 1$ signifies that the material is completely undamaged, $u = 0$ signifies that it is completely damaged while values of u in $(0, 1)$ represent varying degrees of intermediate damage. In [11], the physical constraint was ensured by the presence of a sub-differential graph, *i.e.* a multivalued maximal monotone operator. In addition, a stochastic force term given by an Itô integral has been added on the right-hand side of (1.1). Since its diffusion coefficient g depends on the damage parameter, the stochastic force is said to be multiplicative. From a physical point of view, the presence of this random force term reflects the fact that the phenomenon of damage is related to microscopic changes in the structure and configuration of the material lattice as a consequence of breaking bonds and the formation of cavities and voids. These changes are clearly related to stochastic processes occurring at a microscopic level (as introduced in Ising materials), which we aimed to take into account in the macroscopic description.

The literature on the deterministic Allen-Cahn equation is very rich, also including the presence of non-smooth (monotone) operators (see, among others, [1, 28, 29, 32, 60]), and the stochastic case has been addressed by a growing number of surveys. Some recent results are devoted to questions of existence and uniqueness of solutions for stochastic Allen-Cahn equation [56], in [19] well posedness for stochastic Allen-Cahn type equations with p -Laplacian as well as the random separation property are studied. Others are more interested in questions of existence and regularity of solution for stochastic Cahn-Hilliard/Allen-Cahn problems [4] and [33]. The study of degenerate Kolmogorov equations and questions of ergodicity for stochastic Allen-Cahn equations with logarithmic potential have been considered in [18, 57]. Other authors have studied a stochastic Allen-Cahn-Navier-Stokes system with inertial effects and multiplicative noise of jump type in a bounded domain [34, 46]. Let us mention that, in these last contributions, the sub-differential operator is replaced by smooth nonlinearities (possibly with prescribed growth conditions), as double-well potentials.

Furthermore, the study of stochastic partial differential inclusions in a rather general situation was carried out in [8, 16, 53], as well as questions concerning transition semigroup and invariant measures in [7], or even obstacle problems with Lewy-Stampacchia's inequality for a stochastic T -monotone obstacle problem (see [58]). According to [12], the convergence analysis of numerical schemes for stochastic PDEs of parabolic type has been

a very fashionable subject in recent decades and for this reason, an extensive literature on this topic is available (see [2], [31] and [49] for a general overview). In the past the use of finite-element methods was the preferred technique for the spatial discretisation of parabolic evolution equations (see [10], [24] for a state of the art on this subject), and this is particularly true for the numerical approximation of stochastic Allen-Cahn type equations without constraint (in the chronological order let us mention the contributions [40, 52, 27, 3, 45, 15, 25]).

The numerical analysis of differential inclusions was first carried out on multivalued differential equations in [59, 35, 5, 47], and new studies on the subject have continued to be published ever since [43, 9, 41, 21, 22, 55, 48]. In parallel, stochastic differential inclusions were studied in the early 2000's from a numerical point of view. Firstly, results of convergence analysis of time-discretization schemes have been derived in [50, 17, 44, 61], and secondly, convergence rate as long as error estimates have been investigated respectively in [62] and [38]. More recently, the time-space discretization of deterministic elliptic and parabolic partial differential inclusions was performed by combining Euler scheme with finite-element methods in [54, 23, 20].

To the best of our knowledge, the numerical analysis of stochastic partial differential inclusions is still an open topic. Our aim is then to fill the gap in the literature by addressing the convergence analysis of a time and space discretization scheme for our stochastic Allen-Cahn problem with constraint (1.1). Let us precise that the main originality of our approach consists in the use of a finite-volume method for the spatial discretization instead of a finite-element one.

1.2 Concept of solution and main result

Following our previous work [11], we are interested here in the following concept of solution for Problem (1.1):

Definition 1.1. *Any pair of stochastic processes $(u, \psi) \in (L^2_{\mathcal{P}_T}(\Omega \times (0, T); L^2(\Lambda)))^2$ with u belonging additionally to*

$$L^2(\Omega; \mathcal{C}([0, T]; L^2(\Lambda))) \cap L^2_{\mathcal{P}_T}(\Omega \times (0, T); H^1(\Lambda)),$$

is a solution to Problem (1.1) if almost everywhere in $(0, T) \times \Lambda$ and \mathbb{P} -a.s in Ω ,

$$0 \leq u \leq 1 \text{ and } \psi \in \partial I_{[0,1]}(u),$$

and if the pair (u, ψ) satisfies

$$u(t) = u_0 + \int_0^t (\Delta u(s) - \psi(s) + \beta(u(s)) + f(s)) ds + \int_0^t g(u(s)) dW(s),$$

in $L^2(\Lambda)$ and \mathbb{P} -a.s in Ω , where Δ denotes the Laplace operator on $H^1(\Lambda)$ associated with the formal Neumann boundary condition.

Remark 1.2. *A priori*, we have the predictability of u with values in $L^2(\Lambda)$. It is a direct consequence of, e.g., [42, Corollary 1.1.8] that we may *a posteriori* conclude that u belongs to $L^2_{\mathcal{P}_T}(\Omega \times (0, T); H^1(\Lambda))$.

Existence and uniqueness of a pair (u, ψ) solution of Problem (1.1) in the sense of Definition 1.1 has been proved in [11] for a given initial condition u_0 in $H^1(\Lambda)$ and under Assumptions \mathcal{A}_2 , \mathcal{A}_3 and \mathcal{A}_4 . To do so, we used a regularization procedure on the maximal monotone operator $\partial I_{[0,1]}$ by considering the following family of approximating problems depending on a parameter $\epsilon > 0$:

$$\begin{aligned} du_\epsilon + (\psi_\epsilon(u_\epsilon) - \Delta u_\epsilon) dt &= g(u_\epsilon) dW(t) + (\beta(u_\epsilon) + f) dt, & \text{in } \Omega \times (0, T) \times \Lambda; \\ u_\epsilon(0, \cdot) &= u_0, & \text{in } \Omega \times \Lambda; \\ \nabla u_\epsilon \cdot \mathbf{n} &= 0, & \text{on } \Omega \times (0, T) \times \partial\Lambda; \end{aligned} \quad (1.2)$$

where $\psi_\epsilon : \mathbb{R} \rightarrow \mathbb{R}$ denotes the Moreau-Yosida approximation of $\partial I_{[0,1]}$ (see e.g. [6, 26]), defined for all $v \in \mathbb{R}$ by

$$\psi_\epsilon(v) = -\frac{(v)^-}{\epsilon} + \frac{(v-1)^+}{\epsilon} = \begin{cases} v & \text{if } v \leq 0 \\ \frac{\epsilon}{0} & \text{if } v \in [0, 1] \\ \frac{v-1}{\epsilon} & \text{if } v \geq 1. \end{cases} \quad (1.3)$$

Firstly, we proved in [11], for fixed $\epsilon > 0$, existence and uniqueness of a solution u_ϵ for Problem (1.2) in the sense of Definition 1.3 below:

Definition 1.3. *A stochastic process $u_\epsilon \in L^2_{\mathcal{P}_T}(\Omega \times (0, T); H^1(\Lambda))$ element of*

$$L^\infty(0, T; L^2(\Omega; H^1(\Lambda))) \cap L^2(\Omega; \mathcal{C}(0, T; L^2(\Lambda)))$$

and such that $\partial_t(u_\epsilon - \int_0^\cdot g(u_\epsilon) dW)$ and Δu_ϵ belong to $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$, is a solution to the Problem (1.2) if almost everywhere in $(0, T)$ and \mathbb{P} -almost surely in Ω , the following variational formulation holds for any $v \in H^1(\Lambda)$

$$\int_\Lambda \partial_t \left(u_\epsilon - \int_0^\cdot g(u_\epsilon) dW(s) \right) v dx + \int_\Lambda \nabla u_\epsilon \cdot \nabla v dx + \int_\Lambda \psi_\epsilon(u_\epsilon) v dx = \int_\Lambda (\beta(u_\epsilon) + f) v dx,$$

with \mathbb{P} -a.s in Ω , $u_0 = \lim_{t \rightarrow 0} u_\epsilon(\cdot, t)$ in $L^2(\Lambda)$.

Secondly, the analysis of the sequences $(u_\epsilon)_{\epsilon > 0}$ and $(\psi_\epsilon(u_\epsilon))_{\epsilon > 0}$ allowed us (mainly thanks to monotonicity tools) to pass the limit with respect to the approximating parameter $\epsilon > 0$. We finally proved existence of a solution (u, ψ) of Problem (1.1) in the sense of Definition 1.1, as a weak limit of a subsequence of the pair $(u_\epsilon, \psi_\epsilon(u_\epsilon))_{\epsilon > 0}$. Then, we finished our study by proving the uniqueness of such a solution (u, ψ) .

The objective of the present paper is to propose a time and space approximation of the unique solution (u, ψ) of Problem (1.1) in the sense of Definition 1.1. To do so,

our idea consists in discretizing, for a given $\epsilon > 0$, Problem (1.2) by the way of finite-volume methods. Our main goal is to show that the resulting finite-volume approximation, depending on ϵ and on the time and space parameters (denoted respectively N and h in the sequel), can be bounded independently of these three parameters, with the idea of making them tend towards zero simultaneously. The aim of the game is then to find a relationship between ϵ , N and h that allows us to bound our finite-volume approximation and to pass to the limit in the numerical scheme, leading us to our main result stated hereafter:

Theorem 1.4. *Assume that hypotheses \mathcal{A}_1 to \mathcal{A}_4 hold and let (u, ψ) be the unique solution of Problem (1.1) in the sense of Definition 1.1. Let $(\mathcal{T}_m)_{m \in \mathbb{N}}$ be a sequence of admissible finite-volume meshes of Λ in the sense of Definition 2.1 such that the mesh size h_m tends to 0, let $(N_m)_{m \in \mathbb{N}} \subset \mathbb{N}^*$ be a sequence of positive integers which tends to infinity and let $(\epsilon_m)_{m \in \mathbb{N}} \subset \mathbb{R}_+^*$ be another sequence such that $\lim_{m \rightarrow +\infty} \epsilon_m = 0$. For a fixed $m \in \mathbb{N}$, let u_{h_m, N_m}^r and u_{h_m, N_m}^l be respectively the right and left in time finite-volume approximations defined by (2.2)-(3.1)-(3.2) with $\mathcal{T} = \mathcal{T}_m$, $N = N_m$ and $\epsilon = \epsilon_m$. If there exists $\theta > 0$ such that for any $m \in \mathbb{N}$, $\frac{T}{N_m} = \mathcal{O}((\epsilon_m)^{2+\theta})$, then the sequences $(u_{h_m, N_m}^r)_{m \in \mathbb{N}}$ and $(\psi_{\epsilon_m}(u_{h_m, N_m}^r))_{m \in \mathbb{N}}$ converge towards u and ψ , respectively strongly and weakly in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$. Moreover, the convergence of $(u_{h_m, N_m}^r)_{m \in \mathbb{N}}$ towards u also holds strongly in $L^p(0, T; L^2(\Omega; L^2(\Lambda)))$ for any finite $p \geq 1$.*

1.3 Outline

This contribution is organized as follows. In Section 2, the discretization framework is introduced: choice of the time step, definition of admissible finite-volume meshes of Λ , related notations and employed discrete norms. Then in Section 3, the semi-implicit TPFA scheme for the discretization of the regularized Problem (1.2) and its associated discrete solutions are defined, and the well-posedness of such a scheme is investigated. In Section 4, a clever relation between the regularization and the time and space discretization parameters (denoted ϵ , Δt and h , respectively), allows us to derive stability estimates satisfied by the discrete solutions, independently of these three parameters. Section 5 is then dedicated to the convergence analysis of our scheme by combining arguments we developed in [14] for the passage to the limit with respect to Δt and h , with the ones used in [11] to pass to the limit with respect to ϵ .

2 Discretization framework

Let us start this section by some general notations, then Subsections 2.2, 2.3, 2.4 contain all the definitions and notations related to temporal and spatial discretizations. Let us mention that they are the same as in our previous papers [12, 13, 14], but for a matter of self-containedness we choose to repeat them identically.

2.1 General notations

- The integral over Ω with respect to the probability measure \mathbb{P} is denoted by $\mathbb{E}[\cdot]$, and is called the expectation.
- For any x, y in \mathbb{R}^d , the euclidean norm of x is denoted by $|x|$, and the associated scalar product of x and y by $x \cdot y$.
- The d -dimensional Lebesgue measure of Λ is denoted by $|\Lambda|$, by overusing the euclidean norm notation.
- For $q \in \{1, d\}$, the $L^\infty(\mathbb{R}^q)$ norm is denoted by $\|\cdot\|_\infty$.

2.2 Uniform time step and admissible finite-volume meshes

With the aim of proposing a time and space approximation of the variational solution of Problem (1.2), a choice for the temporal and spatial discretizations must be made. The temporal one is achieved using a uniform subdivision: setting $N \in \mathbb{N}^*$, the fixed time step is defined by $\Delta t = \frac{T}{N}$ and the interval $[0, T]$ is decomposed in $0 = t_0 < t_1 < \dots < t_N = T$ equidistantly with $t_n = n\Delta t$ for all $n \in \{0, \dots, N-1\}$. For the spatial one, following [39, Definition 9.1], we consider admissible finite-volume meshes as defined hereafter:

Definition 2.1. (*Admissible finite-volume mesh*) An admissible finite-volume mesh of Λ , denoted by \mathcal{T} , is given by a family of “control volumes”, which are open polygonal convex subsets of Λ , a family of subsets of $\bar{\Lambda}$ contained in hyperplanes of \mathbb{R}^d , denoted by \mathcal{E} (these are the edges for $d = 2$ or sides for $d = 3$ of the control volumes), with strictly positive $(d-1)$ -dimensional Lebesgue measure, and a family of points of Λ denoted by \mathcal{P} satisfying the following properties*

- $\bar{\Lambda} = \bigcup_{K \in \mathcal{T}} \bar{K}$.
- For any $K \in \mathcal{T}$, there exists a subset \mathcal{E}_K of \mathcal{E} such that $\partial K = \bar{K} \setminus K = \bigcup_{\sigma \in \mathcal{E}_K} \bar{\sigma}$ and $\mathcal{E} = \bigcup_{K \in \mathcal{T}} \mathcal{E}_K$. \mathcal{E}_K is called the set of edges of K for $d = 2$ and sides for $d = 3$, respectively.
- For any $K, L \in \mathcal{T}$, with $K \neq L$ then either the $(d-1)$ Lebesgue measure of $\bar{K} \cap \bar{L}$ is 0 or $\bar{K} \cap \bar{L} = \bar{\sigma}$ for some $\sigma \in \mathcal{E}$, which will then be denoted by $K|L$ or $L|K$.
- The family $\mathcal{P} = (x_K)_{K \in \mathcal{T}}$ is such that $x_K \in \bar{K}$ for all $K \in \mathcal{T}$ and, if $K, L \in \mathcal{T}$ are two neighbouring control volumes, it is assumed that $x_K \neq x_L$, and that the straight line between x_K and x_L is orthogonal to $\sigma = K|L$.
- For any $\sigma \in \mathcal{E}$ such that $\sigma \subset \partial\Lambda$, let K be the control volume such that $\sigma \in \mathcal{E}_K$. If $x_K \notin \sigma$, the straight line going through x_K and orthogonal to σ has a nonempty intersection with σ .

*In fact, we shall denote, somewhat incorrectly, by \mathcal{T} the family of control volumes.




Figure 1: Notations of the mesh \mathcal{T} associated with $\Lambda \subset \mathbb{R}^2$

For a given admissible finite-volume mesh \mathcal{T} of Λ , the following associated notations will be used in the rest of the paper.

Notations.

- The mesh size is denoted by $h = \text{size}(\mathcal{T}) = \sup\{\text{diam}(K) : K \in \mathcal{T}\}$.
- The number of control volumes $K \in \mathcal{T}$ is denoted by $d_h \in \mathbb{N}$, where $h = \text{size}(\mathcal{T})$.
- The sets of interior and exterior interfaces are respectively denoted by $\mathcal{E}_{\text{int}} := \{\sigma \in \mathcal{E} : \sigma \not\subseteq \partial\Lambda\}$ and $\mathcal{E}_{\text{ext}} := \{\sigma \in \mathcal{E} : \sigma \subseteq \partial\Lambda\}$.
- For any $K \in \mathcal{T}$, the d -dimensional Lebesgue measure of K is denoted by m_K .
- For any $K \in \mathcal{T}$, the unit normal vector to ∂K outward to K is denoted by \mathbf{n}_K .
- For any $K \in \mathcal{T}$ and any $\sigma \in \mathcal{E}_K$, the unit vector on σ pointing out of K is denoted by $\mathbf{n}_{K,\sigma}$.
- For any $\sigma \in \mathcal{E}_{\text{int}}$, the $(d-1)$ -dimensional Lebesgue measure of σ is denoted by m_σ .
- For any neighboring control volumes $K, L \in \mathcal{T}$, the euclidean distance between x_K and x_L is denoted by $d_{K|L}$.
- The maximum of edges incident to any vertex of the mesh is denoted by \mathcal{N} .
- For any $K \in \mathcal{T}$ and any $\sigma \in \mathcal{E}_K$, the euclidean distance between x_K and σ is denoted by $d(x_K, \sigma)$.

The regularity of the mesh \mathcal{T} is measured by the following positive number

$$\text{reg}(\mathcal{T}) = \max \left(\mathcal{N}, \max_{\substack{K \in \mathcal{T} \\ \sigma \in \mathcal{E}_K}} \frac{\text{diam}(K)}{d(x_K, \sigma)} \right).$$

As in the deterministic setting, it is assumed that $\text{reg}(\mathcal{T})$ is uniformly bounded by a constant not depending on the mesh size h , which is one of the key point to prove the convergence of our finite-volume scheme. Indeed, the introduction of the number $\text{reg}(\mathcal{T})$ allows us particularly to derive the following inequality:

$$\forall K, L \in \mathcal{T}, \frac{h}{d_{K|L}} \leq \text{reg}(\mathcal{T}), \quad (2.1)$$

and such a uniform control of the ratio in the left-hand side of (2.1) will be essential in the proof of Lemma 5.11.

2.3 Discrete unknowns and piecewise constant functions

From here to the end of Section 4, let N be a positive integer, $\Delta t = \frac{T}{N}$ and \mathcal{T} be an admissible finite-volume mesh of Λ in the sense of Definition 2.1 with a mesh size $h > 0$. The ideology of a finite-volume method to approximate the variational solution of Problem (1.2) is to associate to each control volume $K \in \mathcal{T}$ and time $t_n \in \{0, \dots, t_N\}$ a discrete unknown value denoted by $u_K^n \in \mathbb{R}$, expected to be an approximation of $u_\epsilon(t_n, x_K)$.

For a given vector $(w_K^n)_{K \in \mathcal{T}} \in \mathbb{R}^{d_h}$, we introduce in what follows various associated functions. Firstly, we define the piecewise constant function in space $w_h^n : \Lambda \rightarrow \mathbb{R}$ by

$$w_h^n(x) := \sum_{K \in \mathcal{T}} w_K^n \mathbf{1}_K(x), \quad \forall x \in \Lambda.$$

Using the fact that the mesh \mathcal{T} is fixed, the continuous mapping defined from \mathbb{R}^{d_h} to $L^2(\Lambda)$ by

$$(w_K^n)_{K \in \mathcal{T}} \mapsto \sum_{K \in \mathcal{T}} \mathbf{1}_K w_K^n,$$

allows us to consider the space \mathbb{R}^{d_h} as a finite-dimensional subspace of $L^2(\Lambda)$ and to do the following natural identification between the function and the vector

$$w_h^n \equiv (w_K^n)_{K \in \mathcal{T}} \in \mathbb{R}^{d_h}.$$

Secondly, the knowledge for any $n \in \{0, \dots, N\}$ of the function w_h^n enables us to define the following right and left piecewise constant functions in time and space denoted respectively by $w_{h,N}^r$ and $w_{h,N}^l$, which are defined from $[0, T] \times \Lambda$ to \mathbb{R} by

$$\begin{aligned} w_{h,N}^r(t, x) &:= \sum_{n=0}^{N-1} w_h^{n+1}(x) \mathbf{1}_{[t_n, t_{n+1})}(t) \text{ if } t \in [0, T) \text{ and } w_{h,N}^r(T, x) := w_h^N(x), \\ w_{h,N}^l(t, x) &:= \sum_{n=0}^{N-1} w_h^n(x) \mathbf{1}_{[t_n, t_{n+1})}(t) \text{ if } t \in (0, T] \text{ and } w_{h,N}^l(0, x) := w_h^0(x). \end{aligned} \quad (2.2)$$

Since \mathcal{T} and N are fixed, reasoning as for the piecewise constant function in space above, the continuity of the mapping defined from $\mathbb{R}^{d_h \times N}$ to $L^2(0, T; L^2(\Lambda))$ by

$$(w_K^n)_{\substack{K \in \mathcal{T} \\ n \in \{0, \dots, N-1\}}} \mapsto \sum_{\substack{K \in \mathcal{T} \\ n \in \{0, \dots, N-1\}}} \mathbf{1}_K \mathbf{1}_{[t_n, t_{n+1})} w_K^n,$$

allows us to consider the space $\mathbb{R}^{d_h \times N}$ as a finite-dimensional subspace of $L^2(0, T; L^2(\Lambda))$ and to do naturally the identifications

$$\begin{aligned} w_{h,N}^l &\equiv (w_K^n)_{\substack{K \in \mathcal{T} \\ n \in \{0, \dots, N-1\}}} \in \mathbb{R}^{d_h \times N}, \\ w_{h,N}^r &\equiv (w_K^{n+1})_{\substack{K \in \mathcal{T} \\ n \in \{0, \dots, N-1\}}} \in \mathbb{R}^{d_h \times N}. \end{aligned}$$

Remark 2.2. In the following, when a time and space function $\phi : [0, T] \times \Lambda \rightarrow \mathbb{R}$ will be considered for fixed $x \in \Lambda$, the space variable will be omitted in the notations and $\phi(x)$ will be written instead of $\phi(\cdot, x)$. An analogous notation will apply for fixed $t \in [0, T]$, *i.e.*, we will write $\phi(t)$ for $\phi(t, \cdot)$.

2.4 Discrete norms and weak gradient

For the remainder of this subsection, let us set $n \in \{0, \dots, N-1\}$, consider an arbitrary vector $(w_K^n)_{K \in \mathcal{T}} \in \mathbb{R}^{d_h}$ and identify it with the piecewise constant function in space $w_h^n \equiv (w_K^n)_{K \in \mathcal{T}}$. Are introduced in what follows for w_h^n the definitions of its discrete $L^2(\Lambda)$ -norm, weak gradient and discrete $H^1(\Lambda)$ -semi-norm.

Definition 2.3 (Discrete $L^2(\Lambda)$ -norm). *The discrete $L^2(\Lambda)$ -norm of $w_h^n \in \mathbb{R}^{d_h}$ is defined by*

$$\|w_h^n\|_{L^2(\Lambda)} = \left(\sum_{K \in \mathcal{T}} m_K |w_K^n|^2 \right)^{\frac{1}{2}}.$$

Definition 2.4 (Weak gradient). *Let e_h be the number of elements in \mathcal{E} . The weak gradient operator $\nabla^h : \mathbb{R}^{d_h} \rightarrow (\mathbb{R}^d)^{e_h}$ maps any scalar fields $w_h^n \in \mathbb{R}^{d_h}$ into vector fields $\nabla^h w_h^n = (\nabla_\sigma^h w_h^n)_{\sigma \in \mathcal{E}} \in (\mathbb{R}^d)^{e_h}$, where for any $\sigma \in \mathcal{E}$, $\nabla_\sigma^h w_h^n \in \mathbb{R}^d$ is defined by*

$$\nabla_\sigma^h w_h^n := \begin{cases} d \frac{w_L^n - w_K^n}{d_{K|L}} \mathbf{n}_{K,\sigma}, & \text{if } \sigma = K|L \in \mathcal{E}_{\text{int}}; \\ 0, & \text{if } \sigma \in \mathcal{E}_{\text{ext}}. \end{cases}$$

Definition 2.5 (Discrete $H^1(\Lambda)$ -semi-norm). *The discrete $H^1(\Lambda)$ -semi-norm of $w_h^n \in \mathbb{R}^{d_h}$ is defined by*

$$|w_h^n|_{1,h} := \left(\sum_{\sigma=K|L \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} |w_K^n - w_L^n|^2 \right)^{\frac{1}{2}}.$$

Remark 2.6. We have the following relation between the discrete $(L^2(\Lambda))^d$ -norm of $\nabla^h w_h^n$ and the discrete $H^1(\Lambda)$ -semi-norm of w_h^n :

$$\|\nabla^h w_h^n\|_{(L^2(\Lambda))^d}^2 = \sum_{\sigma=K|L \in \mathcal{E}_{\text{int}}} \frac{d_{K|L} m_\sigma}{d} \left| d \frac{w_K^n - w_L^n}{d_{K|L}} \right|^2 = d |w_h^n|_{1,h}^2. \quad (2.3)$$

We end this subsection by recalling a classical trick of sum reordering, which will be used several times in the rest of the paper.

Remark 2.7 (Discrete partial integration). For any $\tilde{w}_h^n \equiv (\tilde{w}_K^n)_{K \in \mathcal{T}} \in \mathbb{R}^{d_h}$, the following rule of "discrete partial integration" holds:

$$\sum_{K \in \mathcal{T}} \sum_{\sigma=K|L \in \mathcal{E}_K \cap \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} (w_K^n - w_L^n) \tilde{w}_K^n = \sum_{\sigma=K|L \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} (w_K^n - w_L^n) (\tilde{w}_K^n - \tilde{w}_L^n). \quad (2.4)$$

Now, we have all the necessary definitions and notations to present the finite-volume scheme studied in this paper. This is the aim of the next section.

3 The semi-implicit TPFA scheme for Problem (1.2)

By the discretization of the initial condition u_0 of Problem (1.1) over each control volume:

$$u_K^0 := \frac{1}{m_K} \int_K u_0(x) dx, \quad \forall K \in \mathcal{T}, \quad (3.1)$$

we are firstly allowed to define the random vector $u_h^0 \equiv (u_K^0)_{K \in \mathcal{T}} \in \mathbb{R}^{d_h}$. Secondly, starting from this given initial \mathcal{F}_0 -measurable random vector $u_h^0 \in \mathbb{R}^{d_h}$, and fixing a parameter $\epsilon > 0$, we construct our semi-implicit TPFA scheme as follows:

For any $n \in \{0, \dots, N-1\}$, knowing $u_h^n \equiv (u_K^n)_{K \in \mathcal{T}} \in \mathbb{R}^{d_h}$, we search for $u_h^{n+1} \equiv (u_K^{n+1})_{K \in \mathcal{T}} \in \mathbb{R}^{d_h}$, solution of the following equations, \mathbb{P} -a.s in Ω :

$$\begin{aligned} & \frac{m_K}{\Delta t} (u_K^{n+1} - u_K^n) + \sum_{\sigma=K|L \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \frac{m_\sigma}{d_{K|L}} (u_K^{n+1} - u_L^{n+1}) + m_K \psi_\epsilon(u_K^{n+1}) \\ & = \frac{m_K}{\Delta t} g(u_K^n) (W^{n+1} - W^n) + m_K \beta(u_K^{n+1}) + m_K f_K^n, \quad \forall K \in \mathcal{T}, \end{aligned} \quad (3.2)$$

where $W^{n+1} - W^n$ denotes the increments of the Brownian motion between t_{n+1} and t_n :

$$W^{n+1} - W^n = W(t_{n+1}) - W(t_n) \text{ for } n \in \{0, \dots, N-1\},$$

and f_K^n is defined by

$$f_K^n = \frac{1}{\Delta t m_K} \int_{t_n}^{t_{n+1}} \int_K f(x, t) dx dt. \quad (3.3)$$

Remark 3.1. Although for any $n \in \{1, \dots, N\}$ and any $K \in \mathcal{T}$, the discrete unknowns u_K^n (and then the discrete solution u_h^n) depend on ϵ , we omit this dependency from the notation for the sake of clarity.

Proposition 3.2 (Well-posedness of the scheme). *Let \mathcal{T} be an admissible finite-volume mesh of Λ in the sense of Definition 2.1 with a mesh size h , let N be a positive integer and let $\epsilon \in \mathbb{R}_+^*$ be a given parameter. Then, under Assumptions \mathcal{A}_1 to \mathcal{A}_4 , there exists a unique solution $(u_h^n)_{1 \leq n \leq N} \in (\mathbb{R}^{d_h})^N$ to Problem (3.2) associated with the initial vector u_h^0 defined by (3.1). Moreover, for any $n \in \{0, \dots, N\}$, u_h^n is a \mathcal{F}_{t_n} -measurable random vector.*

Proof. It is a direct application of the main result of [13] in the particular case where the convection term is null and the source's one is equal to $u \mapsto \beta(u) - \psi_\epsilon(u) + f$. \square

The right and left finite-volume approximations $u_{h,N}^r$ and $u_{h,N}^l$ defined by (2.2) to approximate the solution u_ϵ of Problem (1.2) are then built from the discrete solution $(u_h^n)_{1 \leq n \leq N} \in (\mathbb{R}^{d_h})^N$ given by Proposition 3.2.

4 Stability estimates

In this section will be derived several stability estimates satisfied by the discrete solution $(u_h^n)_{1 \leq n \leq N} \in (\mathbb{R}^{d_h})^N$ given by Proposition 3.2, and also by the associated left and right finite-volume approximations $(u_{h,N}^l)_{h,N}$ and $(u_{h,N}^r)_{h,N}$ defined by (2.2). Let us start by bounding the discrete initial data:

Lemma 4.1. *Under Assumption \mathcal{A}_1 , the discrete initial data $u_h^0 \in \mathbb{R}^{d_h}$ associated to u_0 and defined by (3.1) satisfies the following inequality:*

$$\mathbb{E} \left[\|u_h^0\|_{L^2(\Lambda)}^2 \right] \leq \mathbb{E} \left[\|u_0\|_{L^2(\Lambda)}^2 \right].$$

Proof. It is a direct consequence of the definition of u_h^0 and Cauchy-Schwarz inequality. \square

This first lemma allows us to obtain the following first bounds on the discrete solutions:

Proposition 4.2 (Bounds on the discrete solutions). *There exists a constant $K_0 > 0$, depending only on u_0, L_g, L_β, f and T such that for any $\epsilon > 0$, any $N \in \mathbb{N}^*$ large enough (depending on L_β) and any $h \in \mathbb{R}_+^*$*

$$\mathbb{E} \left[\|u_h^n\|_{L^2(\Lambda)}^2 \right] + \sum_{k=0}^{n-1} \mathbb{E} \left[\|u_h^{k+1} - u_h^k\|_{L^2(\Lambda)}^2 \right] + \Delta t \sum_{k=0}^{n-1} \mathbb{E} [|u_h^{k+1}|_{1,h}^2] \leq K_0, \quad \forall n \in \{1, \dots, N\}.$$

Proof. Set $\epsilon > 0, N \in \mathbb{N}^*, h \in \mathbb{R}_+^*$ and fix $n \in \{1, \dots, N\}$. For any $k \in \{0, \dots, n-1\}$, we multiply the numerical scheme (3.2) with u_K^{k+1} , take the expectation, and sum over $K \in \mathcal{T}$ to obtain thanks to (2.4)

$$\begin{aligned} & \sum_{K \in \mathcal{T}} \frac{m_K}{\Delta t} \mathbb{E} [(u_K^{k+1} - u_K^k) u_K^{k+1}] + \sum_{\sigma=K|L \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} \mathbb{E} [|u_K^{k+1} - u_L^{k+1}|^2] \\ & + \sum_{K \in \mathcal{T}} m_K \mathbb{E} [\psi_\epsilon(u_K^{k+1}) u_K^{k+1}] \\ & = \sum_{K \in \mathcal{T}} \frac{m_K}{\Delta t} \mathbb{E} [g(u_K^k) u_K^{k+1} (W^{k+1} - W^k)] + \sum_{K \in \mathcal{T}} m_K \mathbb{E} [(\beta(u_K^{k+1}) + f_K^k) u_K^{k+1}]. \end{aligned} \quad (4.1)$$

We consider the terms of (4.1) separately. Firstly note that

$$\sum_{K \in \mathcal{T}} \frac{m_K}{\Delta t} \mathbb{E} [(u_K^{k+1} - u_K^k) u_K^{k+1}] = \frac{1}{2} \sum_{K \in \mathcal{T}} \frac{m_K}{\Delta t} \mathbb{E} [|u_K^{k+1}|^2 - |u_K^k|^2 + |u_K^{k+1} - u_K^k|^2]. \quad (4.2)$$

Secondly, since ψ_ϵ is monotone with $\psi_\epsilon(0) = 0$, one gets that

$$\sum_{K \in \mathcal{T}} m_K \mathbb{E} [\psi_\epsilon(u_K^{k+1}) u_K^{k+1}] \geq 0. \quad (4.3)$$

Thirdly, since u_K^k and $(W^{k+1} - W^k)$ are independent one obtains

$$\sum_{K \in \mathcal{T}} \frac{m_K}{\Delta t} \mathbb{E} [g(u_K^k) u_K^k (W^{k+1} - W^k)] = 0,$$

and so by applying Young's inequality and using the Itô isometry one arrives at

$$\begin{aligned} & \sum_{K \in \mathcal{T}} \frac{m_K}{\Delta t} \mathbb{E} [g(u_K^k) u_K^{k+1} (W^{k+1} - W^k)] \\ &= \sum_{K \in \mathcal{T}} \frac{m_K}{\Delta t} \mathbb{E} [g(u_K^k) (u_K^{k+1} - u_K^k) (W^{k+1} - W^k)] \\ &\leq \sum_{K \in \mathcal{T}} \frac{m_K}{\Delta t} \mathbb{E} [|g(u_K^k) (W^{k+1} - W^k)|^2] + \frac{1}{4} \sum_{K \in \mathcal{T}} \frac{m_K}{\Delta t} \mathbb{E} [|u_K^{k+1} - u_K^k|^2] \\ &\leq \Delta t L_g^2 \sum_{K \in \mathcal{T}} \frac{m_K}{\Delta t} \mathbb{E} [|u_K^k|^2] + \frac{1}{4} \sum_{K \in \mathcal{T}} \frac{m_K}{\Delta t} \mathbb{E} [|u_K^{k+1} - u_K^k|^2]. \end{aligned} \quad (4.4)$$

Fourthly, using the Lipschitz property of β with $\beta(0) = 0$, the following holds

$$\sum_{K \in \mathcal{T}} m_K \mathbb{E} [\beta(u_K^{k+1}) u_K^{k+1}] \leq L_\beta \sum_{K \in \mathcal{T}} m_K \mathbb{E} [|u_K^{k+1}|^2]. \quad (4.5)$$

Fifthly,

$$\begin{aligned} \sum_{K \in \mathcal{T}} m_K \mathbb{E} [f_K^k u_K^{k+1}] &\leq \frac{1}{2} \sum_{K \in \mathcal{T}} m_K \mathbb{E} [|f_K^k|^2] + \frac{1}{2} \sum_{K \in \mathcal{T}} m_K \mathbb{E} [|u_K^{k+1}|^2] \\ &\leq \frac{1}{2\Delta t} \sum_{K \in \mathcal{T}} \mathbb{E} \left[\int_{t_k}^{t_{k+1}} \int_K |f(x, t)|^2 dx dt \right] + \frac{1}{2} \sum_{K \in \mathcal{T}} m_K \mathbb{E} [|u_K^{k+1}|^2]. \end{aligned} \quad (4.6)$$

Combining (4.2)-(4.3)-(4.4)-(4.5) and (4.6) and multiplying the obtained inequality with $2\Delta t$, one gets

$$\begin{aligned} & \sum_{K \in \mathcal{T}} m_K \mathbb{E} [|u_K^{k+1}|^2 - |u_K^k|^2 + |u_K^{k+1} - u_K^k|^2] + 2\Delta t \sum_{\sigma=K|L \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} \mathbb{E} [|u_K^{k+1} - u_L^{k+1}|^2] \\ &\leq 2\Delta t L_g^2 \sum_{K \in \mathcal{T}} m_K \mathbb{E} [|u_K^k|^2] + \frac{1}{2} \sum_{K \in \mathcal{T}} m_K \mathbb{E} [|u_K^{k+1} - u_K^k|^2] \\ &\quad + \Delta t (2L_\beta + 1) \sum_{K \in \mathcal{T}} m_K \mathbb{E} [|u_K^{k+1}|^2] + \sum_{K \in \mathcal{T}} \mathbb{E} \left[\int_{t_k}^{t_{k+1}} \int_K |f(x, t)|^2 dx dt \right]. \end{aligned}$$

Then,

$$\begin{aligned}
& (1 - \Delta t(2L_\beta + 1)) \sum_{K \in \mathcal{T}} m_K \mathbb{E} [|u_K^{k+1}|^2 - |u_K^k|^2] + \frac{1}{2} \sum_{K \in \mathcal{T}} m_K \mathbb{E} [|u_K^{k+1} - u_K^k|^2] \\
& + 2\Delta t \sum_{\sigma=K|L \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} \mathbb{E} [|u_K^{k+1} - u_L^{k+1}|^2] \\
& \leq \Delta t(2L_g^2 + 2L_\beta + 1) \sum_{K \in \mathcal{T}} m_K \mathbb{E} [(u_K^k)^2] + \sum_{K \in \mathcal{T}} \mathbb{E} \left[\int_{t_k}^{t_{k+1}} \int_K |f(x, t)|^2 dx dt \right].
\end{aligned}$$

For Δt small enough so that $1 - \Delta t(2L_\beta + 1) \geq \frac{1}{4}$, after summing over $k \in \{0, \dots, n-1\}$, one arrives at

$$\begin{aligned}
& \frac{1}{4} \mathbb{E} \left[\|u_h^n\|_{L^2(\Lambda)}^2 - \|u_h^0\|_{L^2(\Lambda)}^2 \right] + \frac{1}{2} \sum_{k=0}^{n-1} \mathbb{E} \left[\|u_h^{k+1} - u_h^k\|_{L^2(\Lambda)}^2 \right] + 2\Delta t \sum_{k=0}^{n-1} \mathbb{E} [|u_h^{k+1}|_{1,h}^2] \\
& \leq \Delta t(2L_g^2 + 2L_\beta + 1) \sum_{k=0}^{n-1} \mathbb{E} \left[\|u_h^k\|_{L^2(\Lambda)}^2 \right] + \sum_{k=0}^{n-1} \sum_{K \in \mathcal{T}} \mathbb{E} \left[\int_{t_k}^{t_{k+1}} \int_K |f(x, t)|^2 dx dt \right].
\end{aligned} \tag{4.7}$$

Then, it follows that

$$\begin{aligned}
& \mathbb{E} \left[\|u_h^n\|_{L^2(\Lambda)}^2 \right] \\
& \leq \mathbb{E} \left[\|u_h^0\|_{L^2(\Lambda)}^2 \right] + 4\Delta t(2L_g^2 + 2L_\beta + 1) \sum_{k=0}^{n-1} \mathbb{E} \left[\|u_h^k\|_{L^2(\Lambda)}^2 \right] + 4\|f\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))}^2.
\end{aligned}$$

Applying the discrete Gronwall lemma yields

$$\mathbb{E} \left[\|u_h^n\|_{L^2(\Lambda)}^2 \right] \leq \left(\mathbb{E} \left[\|u_h^0\|_{L^2(\Lambda)}^2 \right] + 4\|f\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))}^2 \right) e^{4T(2L_g^2 + 2L_\beta + 1)}. \tag{4.8}$$

From (4.8) and Lemma 4.1 we may conclude that there exists a constant $\Upsilon > 0$ such that

$$\sup_{n \in \{1, \dots, N\}} \mathbb{E} \left[\|u_h^n\|_{L^2(\Lambda)}^2 \right] \leq \Upsilon. \tag{4.9}$$

Thanks to (4.9) one gets that for all $n \in \{1, \dots, N\}$

$$\Delta t \sum_{k=0}^{n-1} \mathbb{E} \left[\|g(u_h^k)\|_{L^2(\Lambda)}^2 \right] \leq L_g^2 \Delta t \sum_{k=0}^{n-1} \mathbb{E} \left[\|u_h^k\|_{L^2(\Lambda)}^2 \right] \leq L_g^2 T \Upsilon. \tag{4.10}$$

From (4.7), Lemma 4.1 and (4.9) it now follows that for all $n \in \{1, \dots, N\}$

$$\begin{aligned}
& \mathbb{E} \left[\|u_h^n\|_{L^2(\Lambda)}^2 \right] + 2 \sum_{k=0}^{n-1} \mathbb{E} \left[\|u_h^{k+1} - u_h^k\|_{L^2(\Lambda)}^2 \right] + 8\Delta t \sum_{k=0}^{n-1} \mathbb{E} [|u_h^{k+1}|_{1,h}^2] \\
& \leq \mathbb{E} \left[\|u_0\|_{L^2(\Lambda)}^2 \right] + 4\Upsilon T(2L_g^2 + 2L_\beta + 1) + 4\|f\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))}^2.
\end{aligned}$$

□

We are now interested in the bounds on the right and left finite-volume approximations defined by (2.2).

Lemma 4.3. *The sequences $(u_{h,N}^r)_{\epsilon,h,N}$ and $(u_{h,N}^l)_{\epsilon,h,N}$ are bounded in $L^\infty(0, T; L^2(\Omega; L^2(\Lambda)))$, independently of the regularization and discretization parameters $\epsilon \in \mathbb{R}_+^*$, $N \in \mathbb{N}^*$ and $h \in \mathbb{R}_+^*$. Additionally, $(u_{h,N}^l)_{\epsilon,h,N}$ is bounded in $L_{\mathcal{P}_T}^2(\Omega \times (0, T); L^2(\Lambda))$.*

Proof. We note that by (4.9)

$$\begin{aligned} \|u_{h,N}^r\|_{L^\infty(0,T;L^2(\Omega;L^2(\Lambda)))} + \|u_{h,N}^l\|_{L^\infty(0,T;L^2(\Omega;L^2(\Lambda)))} &\leq 2 \sup_{n \in \{0,1,\dots,N\}} \mathbb{E} \left[\|u_h^n\|_{L^2(\Lambda)}^2 \right] \\ &\leq 2\Upsilon + \mathbb{E} \left[\|u_0\|_{L^2(\Lambda)}^2 \right] \end{aligned}$$

The predictability of $(u_{h,N}^l)_{\epsilon,h,N}$ with values in $L^2(\Lambda)$ is a consequence of the \mathcal{F}_{t_n} -measurability of u_K^n for all $n \in \{0, \dots, N\}$ and all $K \in \mathcal{T}$. Indeed, by construction, $(u_{h,N}^l)_{\epsilon,h,N}$ is then an elementary process adapted to the filtration $(\mathcal{F}_t)_{t \geq 0}$ and so it is predictable. \square

Remark 4.4. Note that by Proposition 4.2, one gets the following useful estimate

$$\mathbb{E} \left[\|u_{h,N}^r - u_{h,N}^l\|_{L^2(0,T;L^2(\Lambda))}^2 \right] = \Delta t \mathbb{E} \left[\sum_{n=0}^{N-1} \|u_h^{n+1} - u_h^n\|_{L^2(\Lambda)}^2 \right] \leq K_0 \Delta t, \quad (4.11)$$

Thanks to Proposition 4.2 we can also obtain a $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$ -bound on the weak gradients of the finite-volume approximation $(u_{h,N}^r)_{\epsilon,h,N}$:

Lemma 4.5. *There exists a constant $K_1 \geq 0$ depending only on u_0 , L_g , L_β , f and T such that*

$$\int_0^T \mathbb{E} [|u_{h,N}^r(t)|_{1,h}^2] dt \leq K_1. \quad (4.12)$$

Proof. It is a direct consequence of Proposition 4.2. \square

Lemma 4.6. *The sequences $(g(u_{h,N}^r))_{\epsilon,h,N}$, $(g(u_{h,N}^l))_{\epsilon,h,N}$, $(\beta(u_{h,N}^r))_{\epsilon,h,N}$, and $(\beta(u_{h,N}^l))_{\epsilon,h,N}$ are bounded in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$ independently of the regularization and discretization parameters $\epsilon \in \mathbb{R}_+^*$, $N \in \mathbb{N}^*$ and $h \in \mathbb{R}_+^*$. Moreover, $(g(u_{h,N}^l))_{\epsilon,h,N}$ and $(\beta(u_{h,N}^l))_{\epsilon,h,N}$ are predictable processes with values in $L^2(\Lambda)$.*

Proof. It is a direct consequence of the boundedness of the sequences $(u_{h,N}^r)_{\epsilon,h,N}$ and $(u_{h,N}^l)_{\epsilon,h,N}$ in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$ given by Lemma 4.3 and of the Lipschitz nature of g and β . \square

Lemma 4.7. *There exists a constant $K_2 \geq 0$ depending only on u_0 , L_g , L_β , f and T such that*

$$\int_0^T \mathbb{E} [|g(u_{h,N}^r(t))|_{1,h}^2] dt \leq K_2. \quad (4.13)$$

Proof. After noticing that:

$$\int_0^T \mathbb{E} [|g(u_{h,N}^r(t))|_{1,h}^2] dt \leq L_g^2 \int_0^T \mathbb{E} [|u_{h,N}^r(t)|_{1,h}^2] dt,$$

the result is immediate thanks to Lemma 4.5. \square

Proposition 4.8. *If we assume that there exists $\theta > 0$ such that $\Delta t = \mathcal{O}(\epsilon^{2+\theta})$ and that $0 < \epsilon < 1$, then the sequences $(\psi_\epsilon(u_{h,N}^r))_{\epsilon,h,N}$ and $(\psi_\epsilon(u_{h,N}^l))_{\epsilon,h,N}$ are bounded respectively in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$ and $L^2_{\mathcal{P}_T}(\Omega \times (0, T); L^2(\Lambda))$, independently of the regularization and discretization parameters $\epsilon \in \mathbb{R}_+^*$, $N \in \mathbb{N}^*$ and $h \in \mathbb{R}_+^*$.*

Proof. Setting $0 < \epsilon < 1$, $K \in \mathcal{T}$, $N \in \mathbb{N}_+^*$ and $n \in \{0, \dots, N-1\}$, multiplying (3.2) by $\Delta t \psi_\epsilon(u_K^{n+1})$, taking the expectation, summing over $K \in \mathcal{T}$ and over $n \in \{0, \dots, N-1\}$ lead to

$$\begin{aligned} & \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} m_K \mathbb{E} [(u_K^{n+1} - u_K^n) \psi_\epsilon(u_K^{n+1})] + \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} m_K \Delta t \mathbb{E} [(\psi_\epsilon(u_K^{n+1}))^2] \\ & + \sum_{n=0}^{N-1} \Delta t \sum_{K \in \mathcal{T}} \mathbb{E} \left[\sum_{\sigma=K|L \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \frac{m_\sigma}{d_{K|L}} (u_K^{n+1} - u_L^{n+1}) \psi_\epsilon(u_K^{n+1}) \right] \\ & = \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} m_K \mathbb{E} [g(u_K^n) (W^{n+1} - W^n) \psi_\epsilon(u_K^{n+1})] \\ & + \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} \Delta t m_K \mathbb{E} [(\beta(u_K^{n+1}) + f_K^n) \psi_\epsilon(u_K^{n+1})]. \end{aligned} \tag{4.14}$$

Let us study separately each term of (4.14).

• For the study of the first term, we introduce the convex antiderivative of ψ_ϵ defined for any $v \in \mathbb{R}$ by

$$\phi_\epsilon(v) = \begin{cases} \frac{v^2}{2\epsilon} & \text{if } v \leq 0 \\ 0 & \text{if } v \in [0, 1] \\ \frac{(v-1)^2}{2\epsilon} & \text{if } v \geq 1. \end{cases} \tag{4.15}$$

Note that thanks to the convexity of ϕ_ϵ , the following holds

$$(u_K^{n+1} - u_K^n) \psi_\epsilon(u_K^{n+1}) = (u_K^{n+1} - u_K^n) \phi'_\epsilon(u_K^{n+1}) \geq \phi_\epsilon(u_K^{n+1}) - \phi_\epsilon(u_K^n),$$

and so

$$\sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} m_K \mathbb{E} [(u_K^{n+1} - u_K^n) \psi_\epsilon(u_K^{n+1})] \geq 0, \tag{4.16}$$

owing to the facts that $\sum_{K \in \mathcal{T}} m_K \mathbb{E} [\phi_\epsilon(u_K^N)] \geq 0$ and $\sum_{K \in \mathcal{T}} m_K \mathbb{E} [\phi_\epsilon(u_K^0)] = 0$ since from Assumption \mathcal{A}_1 , \mathbb{P} -a.s in Ω and for any $K \in \mathcal{T}$, $0 \leq u_K^0 \leq 1$.

• Using (2.4) and the monotonicity of ψ_ϵ , one proves that

$$\begin{aligned} & \sum_{n=0}^{N-1} \Delta t \sum_{K \in \mathcal{T}} \mathbb{E} \left[\sum_{\sigma=K|L \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \frac{m_\sigma}{d_{K|L}} (u_K^{n+1} - u_L^{n+1}) \psi_\epsilon(u_K^{n+1}) \right] \\ &= \sum_{n=0}^{N-1} \Delta t \mathbb{E} \left[\sum_{\sigma=K|L \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} (u_K^{n+1} - u_L^{n+1}) (\psi_\epsilon(u_K^{n+1}) - \psi_\epsilon(u_L^{n+1})) \right] \\ &\geq 0. \end{aligned} \quad (4.17)$$

• Firstly, by using the mean values theorem and the fact that for any $v \in \mathbb{R}$, $\psi_\epsilon(v)g(v) = 0$, we can prove that there exist several elements between u_K^n and u_K^{n+1} , all written in the form $\zeta_K^{n+1} = (1 - \lambda_K^{n+1})u_K^{n+1} + \lambda_K^{n+1}u_K^n$ (for some $\lambda_K^{n+1} \in [0, 1]$), and such that the following inequality holds true

$$\left| g(u_K^n) (\psi_\epsilon(u_K^{n+1}) - \psi_\epsilon(u_K^n)) \right| \leq \frac{1}{\epsilon} |g(u_K^n) - g(\zeta_K^{n+1})| |u_K^{n+1} - u_K^n|. \quad (4.18)$$

Indeed, this can be shown by separating the cases according to the position of u_K^n and u_K^{n+1} relative to 0 and 1 (by noting that u_K^n and u_K^{n+1} play a symmetrical role):

- If $u_K^n \notin (0, 1)$, by setting $\zeta_K^{n+1} = u_K^n$ we get

$$0 = \left| g(u_K^n) (\psi_\epsilon(u_K^{n+1}) - \psi_\epsilon(u_K^n)) \right| = \frac{1}{\epsilon} |g(u_K^n) - g(\zeta_K^{n+1})| |u_K^{n+1} - u_K^n|.$$

- If $u_K^n \in (0, 1)$ and $u_K^{n+1} \in [0, 1]$, by setting again $\zeta_K^{n+1} = u_K^n$,

$$0 = \left| g(u_K^n) (\psi_\epsilon(u_K^{n+1}) - \psi_\epsilon(u_K^n)) \right| = \frac{1}{\epsilon} |g(u_K^n) - g(\zeta_K^{n+1})| |u_K^{n+1} - u_K^n|.$$

- If $u_K^n \in (0, 1)$ and $u_K^{n+1} < 0$, there exists $\zeta_K^{n+1} \in (u_K^{n+1}, 0)$ such that

$$\begin{aligned} g(u_K^n) (\psi_\epsilon(u_K^{n+1}) - \psi_\epsilon(u_K^n)) &= g(u_K^n) \psi'_\epsilon(\zeta_K^{n+1}) u_K^{n+1} \\ &= (g(u_K^n) - g(\zeta_K^{n+1})) \psi'_\epsilon(\zeta_K^{n+1}) u_K^{n+1} \\ \text{and } \left| g(u_K^n) (\psi_\epsilon(u_K^{n+1}) - \psi_\epsilon(u_K^n)) \right| &\leq \frac{1}{\epsilon} |g(u_K^n) - g(\zeta_K^{n+1})| |u_K^{n+1} - u_K^n|. \end{aligned}$$

- If $u_K^n \in (0, 1)$ and $u_K^{n+1} > 1$, there exists $\zeta_K^{n+1} \in (1, u_K^{n+1})$ such that

$$\begin{aligned} g(u_K^n) (\psi_\epsilon(u_K^{n+1}) - \psi_\epsilon(u_K^n)) &= g(u_K^n) \psi'_\epsilon(\zeta_K^{n+1}) (u_K^{n+1} - 1) \\ &= (g(u_K^n) - g(\zeta_K^{n+1})) \psi'_\epsilon(\zeta_K^{n+1}) (u_K^{n+1} - 1) \\ \text{and } \left| g(u_K^n) (\psi_\epsilon(u_K^{n+1}) - \psi_\epsilon(u_K^n)) \right| &\leq \frac{1}{\epsilon} |g(u_K^n) - g(\zeta_K^{n+1})| |u_K^{n+1} - u_K^n|. \end{aligned}$$

Using the assumption that there exists $\theta > 0$ such that $\Delta t = \mathcal{O}(\epsilon^{2+\theta})$, then for any natural number p such that $p \geq 1 + \frac{2}{\theta}$, one has that (since $0 < \epsilon < 1$)

$$\frac{\Delta t^{p-1}}{\epsilon^{2p}} = \frac{\Delta t^{p-1}}{\epsilon^{(2+\theta)(p-1)}} \epsilon^{\theta(p-1)-2} \leq \left(\frac{\Delta t}{\epsilon^{2+\theta}} \right)^{p-1}. \quad (4.19)$$

Choosing $p \in \mathbb{N}$ according to (4.19), using successively the fact that $g(u_K^n) \psi_\epsilon(u_K^n) = 0$, Inequality (4.18), Young's inequality (with p and its conjugate $\frac{p}{p-1}$), the constant K_0 given by Proposition 4.2, the fact that

$$\mathbb{E} \left[(W^{n+1} - W^n)^{2p} \right] = \frac{(2p)!}{p!2^p} \Delta t^p = (2p-1)!! \Delta t^p$$

and Inequality (4.19), one gets the existence of a constant $C_p > 0$ only depending on p such that

$$\begin{aligned} & \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} m_K \mathbb{E} \left[g(u_K^n) (W^{n+1} - W^n) \psi_\epsilon(u_K^{n+1}) \right] \\ &= \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} m_K \mathbb{E} \left[g(u_K^n) (W^{n+1} - W^n) (\psi_\epsilon(u_K^{n+1}) - \psi_\epsilon(u_K^n)) \right] \\ &\leq \frac{1}{2} \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} m_K \mathbb{E} \left[\left(\frac{W^{n+1} - W^n}{\epsilon} \right)^2 (g(u_K^n) - g(\zeta_K^{n+1}))^2 \right] + \frac{1}{2} \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} m_K \mathbb{E} \left[(u_K^{n+1} - u_K^n)^2 \right] \\ &\leq \frac{1}{2p} \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} m_K \mathbb{E} \left[\left(\frac{W^{n+1} - W^n}{\epsilon} \right)^{2p} \right] + \frac{p-1}{2p} \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} m_K \mathbb{E} \left[(g(u_K^n) - g(\zeta_K^{n+1}))^{\frac{2p}{p-1}} \right] + K_0 \\ &= \frac{(2p-1)!!}{2p\epsilon^{2p}} \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} m_K \Delta t^p + \frac{p-1}{2p} L_g^2 (2\|g\|_\infty)^{\frac{2}{p-1}} \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} m_K \mathbb{E} \left[(u_K^n - \zeta_K^{n+1})^2 \right] + K_0 \\ &\leq C_p \frac{\Delta t^{p-1}}{\epsilon^{2p}} \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} m_K \Delta t + L_g^2 (2\|g\|_\infty)^{\frac{2}{p-1}} \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} m_K \mathbb{E} \left[(u_K^n - u_K^{n+1})^2 \right] + K_0 \\ &\leq C_p \left(\frac{\Delta t}{\epsilon^{2+\theta}} \right)^{p-1} |\Lambda| T + K_0 \left(L_g^2 (2\|g\|_\infty)^{\frac{2}{p-1}} + 1 \right). \end{aligned} \quad (4.20)$$

• The second right-hand side term of (4.14) can be handled in the following manner thanks to Young's inequality and the constant K_0 given by Proposition 4.2:

$$\begin{aligned} & \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} \Delta t m_K \mathbb{E} \left[(\beta(u_K^{n+1}) + f_K^n) \psi_\epsilon(u_K^{n+1}) \right] \\ &\leq \frac{1}{2} \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} \Delta t m_K \mathbb{E} \left[(\psi_\epsilon(u_K^{n+1}))^2 \right] + \frac{1}{2} \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} \Delta t m_K \mathbb{E} \left[(\beta(u_K^{n+1}) + f_K^n)^2 \right] \\ &\leq \frac{1}{2} \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} \Delta t m_K \mathbb{E} \left[(\psi_\epsilon(u_K^{n+1}))^2 \right] + L_\beta^2 T K_0 + \|f\|_{L^2(\Omega; L^2(0,T; L^2(\Lambda)))}^2. \end{aligned} \quad (4.21)$$

Finally, combining (4.16),(4.17),(4.20) and (4.21), we obtain that

$$\begin{aligned} & \frac{1}{2} \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} \Delta t m_K \mathbb{E} \left[(\psi_\epsilon(u_K^{n+1}))^2 \right] \\ & \leq C_p \left(\frac{\Delta t}{\epsilon^{2+\theta}} \right)^{p-1} |\Lambda| T + K_0 \left(L_g^2 (2 \|g\|_\infty)^{\frac{2}{p-1}} + 1 \right) + L_\beta^2 T K_0 + \|f\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))}^2 \end{aligned}$$

and the announced result holds since $\Delta t = \mathcal{O}(\epsilon^{2+\theta})$. \square

Remark 4.9. Note that using the constant $K_0 > 0$ given by Proposition 4.2, the following inequality holds directly

$$\|\psi_\epsilon(u_{h,N}^l) - \psi_\epsilon(u_{h,N}^r)\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))} \leq \frac{\Delta t}{\epsilon^2} K_0. \quad (4.22)$$

Lemma 4.10. *If we assume that there exists $\theta > 0$ such that $\Delta t = \mathcal{O}(\epsilon^{2+\theta})$ and that $0 < \epsilon < 1$, then the sequences $\left(\frac{(u_{h,N}^r)^-}{\epsilon}\right)_{\epsilon, h, N}$ and $\left(\frac{(u_{h,N}^r-1)^+}{\epsilon}\right)_{\epsilon, h, N}$ are bounded in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$, independently of the regularization and discretization parameters $\epsilon \in \mathbb{R}_+^*$, $N \in \mathbb{N}^*$ and $h \in \mathbb{R}_+^*$.*

Proof. Since $\psi_\epsilon(u_{h,N}^r) = -\frac{(u_{h,N}^r)^-}{\epsilon} + \frac{(u_{h,N}^r-1)^+}{\epsilon}$ and $(u_{h,N}^r)^- \times (u_{h,N}^r-1)^+ = 0$, one gets that

$$\left\| -\frac{(u_{h,N}^r)^-}{\epsilon} \right\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))}^2 + \left\| \frac{(u_{h,N}^r-1)^+}{\epsilon} \right\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))}^2 = \|\psi_\epsilon(u_{h,N}^r)\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))}^2,$$

and the result holds directly since the right-hand side is bounded by Proposition 4.8. \square

Using the same technique, one proves the following:

Lemma 4.11. *If we assume that there exists $\theta > 0$ such that $\Delta t = \mathcal{O}(\epsilon^{2+\theta})$ and that $0 < \epsilon < 1$, then the sequences $\left(\frac{(u_{h,N}^l)^-}{\epsilon}\right)_{\epsilon, h, N}$ and $\left(\frac{(u_{h,N}^l-1)^+}{\epsilon}\right)_{\epsilon, h, N}$ are bounded in $L^2_{\mathcal{P}_T}(\Omega \times (0, T); L^2(\Lambda))$, independently of the regularization and discretization parameters $\epsilon \in \mathbb{R}_+^*$, $N \in \mathbb{N}^*$ and $h \in \mathbb{R}_+^*$.*

Remark 4.12. Note that using Lemma 4.10, Lemma 4.11 and by expanding the square term of (4.22), one can prove that

$$\begin{aligned} & \left\| \frac{(u_{h,N}^r)^-}{\epsilon} - \frac{(u_{h,N}^l)^-}{\epsilon} \right\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))}^2 \leq \frac{\Delta t}{\epsilon^2} K_0 \\ \text{and} & \left\| \frac{(u_{h,N}^r-1)^+}{\epsilon} - \frac{(u_{h,N}^l-1)^+}{\epsilon} \right\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))}^2 \leq \frac{\Delta t}{\epsilon^2} K_0, \end{aligned}$$

which assures us that if the sequences $\left(\frac{(u_{h,N}^r)^-}{\epsilon}\right)_\epsilon$ and $\left(\frac{(u_{h,N}^l)^-}{\epsilon}\right)_\epsilon$ (respectively $\left(\frac{(u_{h,N}^r-1)^+}{\epsilon}\right)_\epsilon$ and $\left(\frac{(u_{h,N}^l-1)^+}{\epsilon}\right)_\epsilon$) converge, it is necessarily towards a common limit.

5 Convergence of the “ $(\epsilon, \Delta t, h)$ ” scheme

We have now all the necessary tools to pass to the limit in our “ $(\epsilon, \Delta t, h)$ ” scheme. In what follows, let $(\mathcal{T}_m)_{m \in \mathbb{N}}$ be a sequence of admissible meshes of Λ in the sense of Definition 2.1 such that the mesh size h_m tends to 0 when m tends to $+\infty$, let $(N_m)_{m \in \mathbb{N}} \subset \mathbb{N}^*$ be a sequence with $\lim_{m \rightarrow +\infty} N_m = +\infty$, set $\Delta t_m := \frac{T}{N_m}$, and let $(\epsilon_m)_{m \in \mathbb{N}} \subset]0, 1[$ be another sequence such that $\lim_{m \rightarrow +\infty} \epsilon_m = 0$, and assume that there exists $\theta > 0$ such that $\Delta t_m = \mathcal{O}(\epsilon_m^{2+\theta})$ for any $m \in \mathbb{N}$.

For the sake of simplicity, for $m \in \mathbb{N}$, we shall use the notations $\mathcal{T} = \mathcal{T}_m$, $h = \text{size}(\mathcal{T}_m)$, $\Delta t = \Delta t_m$, $N = N_m$ and $\epsilon = \epsilon_m$ when the m -dependency is not useful for the understanding of the reader.

5.1 Weak convergences of finite-volume approximations

First of all, owing to the bounds on the discrete solutions obtained in the previous section, we are able to derive the following weak convergences:

Proposition 5.1. *There exists a process $u \in L^2_{\mathcal{P}_T}(\Omega \times (0, T); H^1(\Lambda))$ such that, up to subsequences of $(u^r_{h,N})_m$ and $(u^l_{h,N})_m$ denoted in the same way,*

$$u^l_{h,N} \rightarrow u \text{ and } u^r_{h,N} \rightarrow u, \text{ both weakly in } L^2(\Omega; L^2(0, T; L^2(\Lambda))) \text{ as } m \rightarrow +\infty.$$

Proof. We refer to [14, Proposition 4.1], since the proof is exactly the same. \square

Lemma 5.2. *There exists a process ψ in $L^2_{\mathcal{P}_T}(\Omega \times (0, T); L^2(\Lambda))$ such that, up to subsequences of $(\psi_\epsilon(u^r_{h,N}))_m$ and $(\psi_\epsilon(u^l_{h,N}))_m$ denoted in the same way,*

$$\psi_\epsilon(u^r_{h,N}) \rightarrow \psi \text{ and } \psi_\epsilon(u^l_{h,N}) \rightarrow \psi, \text{ both weakly in } L^2(\Omega; L^2(0, T; L^2(\Lambda))) \text{ as } m \rightarrow +\infty.$$

Proof. This is a direct consequence of Proposition 4.8 and Remark 4.9. Let us mention that the predictability property of ψ with values in $L^2(\Lambda)$ is inherited from $(\psi_\epsilon(u^l_{h,N}))_m$ at the limit. \square

Lemma 5.3. *There exist not relabeled subsequences of $\left(-\frac{(u^r_{h,N})^-}{\epsilon}\right)_m$ and $\left(\frac{(u^r_{h,N-1})^+}{\epsilon}\right)_m$, and ψ_1, ψ_2 in $L^2_{\mathcal{P}_T}(\Omega \times (0, T); L^2(\Lambda))$ such that*

$$-\frac{(u^r_{h,N})^-}{\epsilon} \rightarrow \psi_1 \text{ and } \frac{(u^r_{h,N} - 1)^+}{\epsilon} \rightarrow \psi_2, \text{ both weakly in } L^2(\Omega; L^2(0, T; L^2(\Lambda))) \text{ as } m \rightarrow +\infty.$$

Proof. This is a direct consequence of Lemma 4.10, Lemma 4.11 and Remark 4.12. \square

Lemma 5.4. *Both strongly in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$, the following convergences hold:*

$$(u^r_{h,N})^- \rightarrow 0 \text{ and } (u^r_{h,N} - 1)^+ \rightarrow 0 \text{ as } m \rightarrow +\infty.$$

Proof. By Lemma 4.10, we have the existence of a constant $M > 0$ independent of the regularization and discretization parameters $\epsilon \in \mathbb{R}_+^*$, $N \in \mathbb{N}^*$ and $h \in \mathbb{R}_+^*$ such that

$$\|(u^r_{h,N})^-\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))}^2 + \|(u^r_{h,N} - 1)^+\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))}^2 \leq M\epsilon^2,$$

and the announced result holds. \square

Remark 5.5. [Additional informations about ψ_1 and ψ_2]

Firstly, note that since $-\frac{(u_{h,N}^r)^-}{\epsilon} \leq 0$ and $\frac{(u_{h,N}^r-1)^+}{\epsilon} \geq 0$, then $\psi_1 \leq 0$ and $\psi_2 \geq 0$.

Secondly, using the fact that $\psi_\epsilon(u_{h,N}^r) = -\frac{(u_{h,N}^r)^-}{\epsilon} + \frac{(u_{h,N}^r-1)^+}{\epsilon}$, one gets owing to Lemmas 5.2 and 5.3 that $\psi = \psi_1 + \psi_2$.

Thirdly, since

$$\psi_\epsilon(u_{h,N}^r)u_{h,N}^r = \frac{(u_{h,N}^r)^-}{\epsilon} \times (u_{h,N}^r)^- + \frac{(u_{h,N}^r-1)^+}{\epsilon} \times ((u_{h,N}^r-1)^+ + 1),$$

one obtains thanks to Lemmas 5.3 and 5.4 that

$$\mathbb{E} \left[\int_0^T \int_\Lambda \psi_\epsilon(u_{h,N}^r(t,x))u_{h,N}^r(t,x) dx dt \right] \rightarrow \mathbb{E} \left[\int_0^T \int_\Lambda \psi_2(t,x) dx dt \right],$$

as $m \rightarrow +\infty$.

Lemma 5.6. *There exists a process g_u in $L^2_{\mathcal{P}_T}(\Omega \times (0, T); H^1(\Lambda))$ such that, up to subsequences of $(g(u_{h,N}^r))_m$ and $(g(u_{h,N}^l))_m$ denoted in the same way,*

$$g(u_{h,N}^r) \rightarrow g_u \text{ and } g(u_{h,N}^l) \rightarrow g_u, \text{ both weakly in } L^2(\Omega; L^2(0, T; L^2(\Lambda))) \text{ as } m \rightarrow +\infty.$$

Proof. This is mainly due to Lemma 4.6, Lemma 4.7 and (4.11). A detailed proof can be found in [14]. \square

Lemma 5.7. *There exists a process β_u in $L^2_{\mathcal{P}_T}(\Omega \times (0, T); L^2(\Lambda))$ such that, up to subsequences of $(\beta(u_{h,N}^r))_m$ and $(\beta(u_{h,N}^l))_m$ denoted in the same way,*

$$\beta(u_{h,N}^r) \rightarrow \beta_u \text{ and } \beta(u_{h,N}^l) \rightarrow \beta_u, \text{ both weakly in } L^2(\Omega; L^2(0, T; L^2(\Lambda))) \text{ as } m \rightarrow +\infty.$$

Proof. This is a direct consequence of Lemma 4.6 and (4.11). \square

Lemma 5.8. *The sequence $(f_{h,N}^l)_m$ defined by (2.2) and (3.3) converges strongly towards f in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$ as $m \rightarrow +\infty$.*

Proof. There exists $\tilde{\Omega} \subset \mathcal{F}$ with $\mathbb{P}(\tilde{\Omega}) = 1$ such that, for all $\omega \in \tilde{\Omega}$, by a standard argument of Steklov average, it is well-known that $\lim_{m \rightarrow \infty} f_{h,N}^l(\omega) = f(\omega)$ in $L^2((0, T) \times \Lambda)$. Moreover, since $f \in L^2(\Omega; L^2((0, T) \times \Lambda))$, the mapping $\omega \mapsto \|f(\omega)\|_{L^2((0, T) \times \Lambda)}$ is an element of $L^2(\Omega)$, and using the fact that \mathbb{P} -a.s in Ω

$$\|f_{h,N}^l\|_{L^2((0, T) \times \Lambda)} \leq \|f\|_{L^2((0, T) \times \Lambda)},$$

one can conclude thanks to Lebesgue's dominated convergence theorem ([36, Theorem 1.3.3]) that $(f_{h,N}^l)_m$ converges strongly towards f in $L^2(\Omega; L^2((0, T) \times \Lambda))$, hence, by isometry, also in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$. \square

Proposition 5.9. *The weak limit u of our finite-volume scheme (3.1)-(3.2) introduced in Proposition 5.1 has \mathbb{P} -a.s. continuous paths with values in $L^2(\Lambda)$ and satisfies for all $t \in [0, T]$,*

$$u(t) = u_0 + \int_0^t (\Delta u(s) - \psi + \beta_u(s) + f(s)) ds + \int_0^t g_u(s) dW(s),$$

in $L^2(\Lambda)$ and \mathbb{P} -a.s. in Ω , where Δ denotes the Laplace operator on $H^1(D)$ associated with the formal Neumann boundary conditions, ψ , g_u and β_u respectively are given by Lemmas 5.2, 5.6 and 5.7.

Proof. Let $A \in \mathcal{A}$, $\xi \in \mathcal{D}(\mathbb{R})$ with $\xi(T) = 0$ and $\varphi \in \mathcal{D}(\mathbb{R}^2)$ with $\nabla \varphi \cdot \mathbf{n} = 0$ on $\partial\Lambda$, where we denote $\mathcal{D}(D) := \mathcal{C}_c^\infty(D)$ for any open subset $D \subseteq \mathbb{R}^d$, $d \in \mathbb{N}$. We introduce the discrete function $\varphi_h : \Lambda \rightarrow \mathbb{R}$ defined by $\varphi_h(x) = \sum_{K \in \mathcal{T}} \mathbf{1}_K(x) \varphi(x_K)$ for any $x \in \Lambda$.

For $K \in \mathcal{T}$, $n \in \{0, \dots, N-1\}$ and $t \in [t_n, t_{n+1})$ we multiply (3.2) with $\mathbf{1}_A \xi(t) \varphi(x_K)$ to obtain

$$\begin{aligned} & \mathbf{1}_A \xi(t) \frac{m_K}{\Delta t} \left(u_K^{n+1} - u_K^n - g(u_K^n) (W^{n+1} - W^n) \right) \varphi(x_K) \\ & + \mathbf{1}_A \xi(t) \sum_{\sigma=K|L \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \frac{m_\sigma}{d_{K|L}} (u_K^{n+1} - u_L^{n+1}) \varphi(x_K) \\ & + \mathbf{1}_A \xi(t) m_K \psi_\epsilon(u_K^{n+1}) \varphi(x_K) \\ & = \mathbf{1}_A \xi(t) m_K (\beta(u_K^{n+1}) + f_K^n) \varphi(x_K). \end{aligned} \quad (5.1)$$

Firstly, we sum (5.1) over each control volume $K \in \mathcal{T}$, we integrate over each time interval $[t_n, t_{n+1}]$, then we sum over $n = 0, \dots, N-1$, and finally we take the expectation to obtain

$$S_{1,m} + S_{2,m} + S_{3,m} + S_{4,m} = S_{5,m} + S_{6,m} \quad (5.2)$$

where

$$\begin{aligned} S_{1,m} &= \mathbb{E} \left[\sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \sum_{K \in \mathcal{T}} \mathbf{1}_A \xi(t) m_K \frac{u_K^{n+1} - u_K^n}{\Delta t} \varphi(x_K) dt \right] \\ S_{2,m} &= -\mathbb{E} \left[\sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \sum_{K \in \mathcal{T}} \mathbf{1}_A \xi(t) m_K g(u_K^n) \frac{W^{n+1} - W^n}{\Delta t} \varphi(x_K) dt \right] \\ S_{3,m} &= \mathbb{E} \left[\sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \sum_{K \in \mathcal{T}} \mathbf{1}_A \xi(t) \sum_{\sigma=K|L \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \frac{m_\sigma}{d_{K|L}} (u_K^{n+1} - u_L^{n+1}) \varphi(x_K) dt \right] \\ S_{4,m} &= \mathbb{E} \left[\sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \sum_{K \in \mathcal{T}} \mathbf{1}_A \xi(t) m_K \psi_\epsilon(u_K^{n+1}) \varphi(x_K) dt \right] \\ S_{5,m} &= \mathbb{E} \left[\sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \sum_{K \in \mathcal{T}} \mathbf{1}_A \xi(t) m_K \beta(u_K^{n+1}) \varphi(x_K) dt \right] \\ S_{6,m} &= \mathbb{E} \left[\sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \sum_{K \in \mathcal{T}} \mathbf{1}_A \xi(t) m_K f_K^n \varphi(x_K) dt \right]. \end{aligned}$$

Let us study separately the limit as m goes to $+\infty$ of $S_{1,m}$, $S_{2,m}$, $S_{3,m}$, $S_{4,m}$, $S_{5,m}$ and $S_{6,m}$.

- Study of $S_{1,m}$: Following [14, Proposition 4.5], one proves thanks to Proposition 5.1 and a discrete integration by parts formula, that up to a subsequence denoted in the same way

$$S_{1,m} \xrightarrow{m \rightarrow +\infty} -\mathbb{E} \left[\mathbf{1}_A \int_0^T \int_{\Lambda} u(t, x) \xi'(t) \varphi(x) dx dt \right] - \mathbb{E} \left[\mathbf{1}_A \int_{\Lambda} u_0(x) \xi(0) \varphi(x) dx \right].$$

- Study of $S_{2,m}$: Thanks to Lemma 5.6 and the properties of the stochastic integral, one shows that, up to a subsequence denoted in the same way (see [14, Proposition 4.5])

$$S_{2,m} \xrightarrow{m \rightarrow +\infty} \mathbb{E} \left[\mathbf{1}_A \int_0^T \int_{\Lambda} \int_0^t g_u(s, x) dW(s) \xi'(t) \varphi(x) dx dt \right].$$

- Study of $S_{3,m}$: Following the arguments we developed in [12, Proposition 4.16], one shows that

$$S_{3,m} \xrightarrow{m \rightarrow +\infty} -\mathbb{E} \left[\mathbf{1}_A \int_0^T \int_{\Lambda} \xi(t) \Delta \varphi(x) u(t, x) dx dt \right].$$

- Study of $S_{4,m}$: Using Lemma 5.2, one proves that

$$S_{4,m} \xrightarrow{m \rightarrow +\infty} \mathbb{E} \left[\mathbf{1}_A \int_0^T \int_{\Lambda} \psi(t, x) \varphi(x) \xi(t) dx dt \right].$$

To do so, we use the following decomposition for $S_{4,m} = S_{4,m} - \tilde{S}_{4,m} + \tilde{S}_{4,m}$, where

$$\tilde{S}_{4,m} = \mathbb{E} \left[\mathbf{1}_A \int_0^T \int_{\Lambda} \psi_{\epsilon}(u_{h,N}^r) \varphi(x) \xi(t) dx dt \right] \xrightarrow{m \rightarrow +\infty} \mathbb{E} \left[\mathbf{1}_A \int_0^T \int_{\Lambda} \psi(t, x) \varphi(x) \xi(t) dx dt \right].$$

Note that

$$\begin{aligned} |S_{4,m} - \tilde{S}_{4,m}| &= \left| \mathbb{E} \left[\mathbf{1}_A \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} \int_{t_n}^{t_{n+1}} \int_K \psi_{\epsilon}(u_K^{n+1}) (\varphi(x_K) - \varphi(x)) \xi(t) dx dt \right] \right| \\ &\leq h \|\xi\|_{\infty} \|\nabla \varphi\|_{\infty} \|\psi_{\epsilon}(u_{h,N}^r)\|_{L^1(\Omega; L^1(0, T; L^1(\Lambda)))} \xrightarrow{m \rightarrow +\infty} 0. \end{aligned}$$

- Study of $S_{5,m}$: Thanks to Lemma 5.7, one shows as for the study of $S_{4,m}$ that

$$S_{5,m} \xrightarrow{m \rightarrow +\infty} \mathbb{E} \left[\mathbf{1}_A \int_0^T \int_{\Lambda} \beta_u(t, x) \varphi(x) \xi(t) dx dt \right].$$

- Study of $S_{6,m}$: using Lemma 5.8 and the fact that f also belongs to $L^1(\Omega; L^1(0, T; L^1(\Lambda)))$, one proves that

$$S_{6,m} \xrightarrow{m \rightarrow +\infty} \mathbb{E} \left[\mathbf{1}_A \int_0^T \int_{\Lambda} f(t, x) \varphi(x) \xi(t) dx dt \right].$$

Indeed,

$$\begin{aligned}
& \left| S_{6,m} - \mathbb{E} \left[\mathbf{1}_A \int_0^T \int_{\Lambda} f(t,x) \varphi(x) \xi(t) dx dt \right] \right| \\
&= \left| \mathbb{E} \left[\sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \sum_{K \in \mathcal{T}} \mathbf{1}_A \xi(t) m_K f_K^n \varphi(x_K) dt \right] - \mathbb{E} \left[\mathbf{1}_A \int_0^T \int_{\Lambda} f(t,x) \varphi(x) \xi(t) dx dt \right] \right| \\
&= \left| \mathbb{E} \left[\mathbf{1}_A \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} \int_{t_n}^{t_{n+1}} \int_K \xi(t) \{ f_K^n \varphi(x_K) - f(t,x) \varphi(x) \} dx dt \right] \right| \\
&= \left| \mathbb{E} \left[\mathbf{1}_A \sum_{n=0}^{N-1} \sum_{K \in \mathcal{T}} \int_{t_n}^{t_{n+1}} \int_K \xi(t) \left\{ \varphi(x_K) (f_K^n - f(t,x)) + f(t,x) (\varphi(x_K) - \varphi(x)) \right\} dx dt \right] \right| \\
&\leq \|\xi\|_{\infty} \|\varphi\|_{\infty} \|f_{h,N}^1 - f\|_{L^1(\Omega; L^1((0,T) \times \Lambda))} + h \|\xi\|_{\infty} \|\nabla \varphi\|_{\infty} \|f\|_{L^1(\Omega; L^1(0,T; L^1(\Lambda)))} \xrightarrow{m \rightarrow +\infty} 0.
\end{aligned}$$

Gathering all the previous convergence results, we can pass to the limit in (5.2), and, by using the density of the set $\{\Psi \in \mathcal{D}(\mathbb{R}^2) \mid \nabla \Psi \cdot \mathbf{n} = 0 \text{ on } \partial\Lambda\}$ in $H^1(\Lambda)$ (given by [37, Theorem 1.1]), we get that \mathbb{P} -a.s. in Ω , for all $\xi \in \{\phi \in \mathcal{D}(\mathbb{R}) : \phi(T) = 0\}$ and all $\varphi \in H^1(\Lambda)$

$$\begin{aligned}
& - \int_0^T \int_{\Lambda} \left(u(t,x) - \int_0^t g_u(s,x) dW(s) \right) \xi'(t) \varphi(x) dx dt - \int_{\Lambda} u_0(x) \xi(0) \varphi(x) dx \\
&= - \int_0^T \int_{\Lambda} \nabla u(t,x) \cdot \nabla \varphi(x) \xi(t) dx dt - \int_0^T \int_{\Lambda} \psi(t,x) \varphi(x) \xi(t) dx dt \tag{5.3} \\
&+ \int_0^T \int_{\Lambda} \beta_u(t,x) \varphi(x) \xi(t) dx dt + \int_0^T \int_{\Lambda} f(t,x) \varphi(x) \xi(t) dx dt.
\end{aligned}$$

By identically repeating the arguments developed in the proof of [14, Proposition 4.5], we first obtain that $u \in L^2(\Omega; \mathcal{C}([0,T]; L^2(\Lambda)))$, and then that for any t in $[0,T]$

$$u(t) - u(0) - \int_0^t g_u(s) dW(s) + \int_0^t \psi(s) ds - \int_0^t \beta_u(s) ds - \int_0^t f(s) ds = \int_0^t \Delta u(s) ds,$$

in $H^1(\Lambda)^*$ and \mathbb{P} -a.s. in Ω . To conclude, let us mention that since the left-hand side of the above equality is in $L^2(\Lambda)$, it also holds in $L^2(\Lambda)$. \square

Lemma 5.10. (Stochastic energy equality) *For any $c > 0$, the stochastic process u introduced in Proposition 5.1 satisfies the following stochastic energy equality:*

$$\begin{aligned}
& e^{-ct} \mathbb{E} \left[\|u(t)\|_{L^2(\Lambda)}^2 \right] + 2 \int_0^t e^{-cs} \mathbb{E} \left[\|\nabla u(s)\|_{L^2(\Lambda)}^2 \right] ds \\
&= \mathbb{E} \left[\|u_0\|_{L^2(\Lambda)}^2 \right] - c \int_0^t e^{-cs} \mathbb{E} \left[\|u(s)\|_{L^2(\Lambda)}^2 \right] ds + \int_0^t e^{-cs} \mathbb{E} \left[\|g_u(s)\|_{L^2(\Lambda)}^2 \right] ds \tag{5.4} \\
&+ 2 \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} (\beta_u(s,x) + f(s,x) - \psi(s,x)) u(s,x) dx \right] ds, \quad \forall t \in [0,T].
\end{aligned}$$

Proof. It is a direct application of Itô formula to the stochastic process u and the functional $\mathcal{F} : (t,v) \mapsto e^{-ct} \|v\|_{L^2(\Lambda)}^2$ defined on $[0,T] \times L^2(\Lambda)$. \square

5.2 Identification of weak limits coming from the non-linear terms

We state here a result proved in [14, Lemma 4.7], which gives a lower bound for the inferior limit of the following quantity

$$\int_0^T \int_0^t e^{-cs} \mathbb{E}[|u_{h_m, N_m}^r(s)|_{1, h_m}^2] ds dt,$$

for any $c > 0$. This boundedness result will be one of the key points in identifying the weak limits ψ , g_u , β_u coming from the discretization of the non linear terms $\psi_\epsilon(u_{h, N}^r)$, $g(u_{h, N}^r)$ and $\beta(u_{h, N}^r)$, see Lemmas 5.2, 5.6, 5.7.

Lemma 5.11. *For any $c > 0$, the stochastic process u introduced in Proposition 5.1 satisfies the following inequality:*

$$\int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_\Lambda |\nabla u(x, s)|^2 dx \right] ds dt \leq \liminf_{m \rightarrow +\infty} \int_0^T \int_0^t e^{-cs} \mathbb{E}[|u_{h_m, N_m}^r(s)|_{1, h_m}^2] ds dt. \quad (5.5)$$

Now, we have all the necessary tools on the one hand for the identification of ψ , ψ_1 , ψ_2 , g_u and β_u , and on the other hand for completing the proof of Theorem 1.4.

Proposition 5.12. *The sequences $(u_{h, N}^r)_m$ and $(u_{h, N}^l)_m$ converge strongly in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$ to the unique variational solution of Problem (1.1) in the sense of Definition 1.1.*

Proof. Let us fix $n \in \{0, \dots, N-1\}$, $K \in \mathcal{T}$, and multiply (3.2) by $\Delta t u_K^{n+1}$, use the formula $a(a-b) = \frac{1}{2}(a^2 - b^2 + (a-b)^2)$ with $a = u_K^{n+1}$ and $b = u_K^n$, take the expectation, and proceed as for the obtention of (4.4) to arrive at

$$\begin{aligned} & \frac{m_K}{2} \mathbb{E} [(u_K^{n+1})^2 - (u_K^n)^2] + \frac{m_K}{2} \mathbb{E} [(u_K^{n+1} - u_K^n)^2] \\ & + \Delta t \sum_{\sigma=K|L \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \frac{m_\sigma}{d_{K|L}} \mathbb{E} [(u_K^{n+1} - u_L^{n+1})u_K^{n+1}] + \Delta t m_K \psi_\epsilon(u_K^{n+1})u_K^{n+1} \\ & \leq \frac{m_K}{2} \mathbb{E} [(u_K^{n+1} - u_K^n)^2] + \frac{m_K \Delta t}{2} \mathbb{E} [g^2(u_K^n)] + \Delta t m_K \mathbb{E} [(\beta(u_K^{n+1}) + f_K^n)u_K^{n+1}]. \end{aligned}$$

Now, we multiply the last inequality by e^{-ct_n} for arbitrary $c > 0$. Then, summing over $K \in \mathcal{T}$ and $n \in \{0, \dots, k\}$ for $k \in \{0, \dots, N-1\}$, using (2.4) and reasoning as in the proof of (4.3) one gets

$$\begin{aligned} & \frac{1}{2} \sum_{n=0}^k \sum_{K \in \mathcal{T}} m_K e^{-ct_n} \mathbb{E} [(u_K^{n+1})^2 - (u_K^n)^2] + \Delta t \sum_{n=0}^k e^{-ct_n} \sum_{\sigma=K|L \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} \mathbb{E} [|u_K^{n+1} - u_L^{n+1}|^2] \\ & + \Delta t \sum_{n=0}^k \sum_{K \in \mathcal{T}} m_K e^{-ct_n} \mathbb{E} [\psi_\epsilon(u_K^{n+1})u_K^{n+1}] \\ & \leq \frac{\Delta t}{2} \sum_{n=0}^k \sum_{K \in \mathcal{T}} m_K e^{-ct_n} \mathbb{E} [g^2(u_K^n)] + \Delta t \sum_{n=0}^k \sum_{K \in \mathcal{T}} m_K e^{-ct_n} \mathbb{E} [(\beta(u_K^{n+1}) + f_K^n)u_K^{n+1}]. \end{aligned}$$

Let us focus on each sum of this last inequality separately, by using the computations we developed in [14].

- Note that the general term of the first sum can be decomposed in the following way:

$$\begin{aligned} & e^{-ct_n} \mathbb{E} [(u_K^{n+1})^2 - (u_K^n)^2] \\ &= e^{-ct_n} \mathbb{E} [(u_K^{n+1})^2] - e^{-ct_{n-1}} \mathbb{E} [(u_K^n)^2] - \mathbb{E} [(u_K^n)^2] (e^{-ct_n} - e^{-ct_{n-1}}), \end{aligned}$$

where $t_{-1} := -\Delta t$. Firstly, we have

$$\begin{aligned} & \frac{1}{2} \sum_{n=0}^k \sum_{K \in \mathcal{T}} m_K (e^{-ct_n} \mathbb{E} [(u_K^{n+1})^2] - e^{-ct_{n-1}} \mathbb{E} [(u_K^n)^2]) \\ &= \frac{1}{2} \sum_{K \in \mathcal{T}} m_K e^{-ct_k} \mathbb{E} [(u_K^{k+1})^2] - \frac{1}{2} \sum_{K \in \mathcal{T}} m_K \mathbb{E} [(u_K^0)^2] e^{c\Delta t}. \end{aligned} \quad (5.6)$$

Then, using properties of the exponential function,

$$\begin{aligned} & -\frac{1}{2} \sum_{n=0}^k \sum_{K \in \mathcal{T}} m_K \mathbb{E} [(u_K^n)^2] (e^{-ct_n} - e^{-ct_{n-1}}) \\ &> -\frac{1}{2} \sum_{K \in \mathcal{T}} m_K \mathbb{E} [(u_K^0)^2] (1 - e^{c\Delta t}) + \frac{c}{2} e^{-c\Delta t} \int_0^{t_k} e^{-cs} \mathbb{E} [\|u_{h,N}^r(s)\|_{L^2(\Lambda)}^2] ds \\ &> \frac{c}{2} e^{-c\Delta t} \int_0^{t_k} e^{-cs} \mathbb{E} [\|u_{h,N}^r(s)\|_{L^2(\Lambda)}^2] ds. \end{aligned} \quad (5.7)$$

- The second sum can be handled similarly in the following manner

$$\begin{aligned} \Delta t \sum_{n=0}^k e^{-ct_n} \sum_{\sigma=K|L \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} \mathbb{E} [|u_K^{n+1} - u_L^{n+1}|^2] &= \Delta t \sum_{n=0}^k e^{-ct_n} \mathbb{E} [|u_h^{n+1}|_{1,h}^2] \\ &\geq \int_0^{t_{k+1}} e^{-cs} \mathbb{E} [|u_{h,N}^r(s)|_{1,h}^2] ds. \end{aligned} \quad (5.8)$$

- Since ψ_ϵ is non-decreasing and satisfies $\psi_\epsilon(0) = 0$, the third sum can be handled as follows:

$$\begin{aligned} & \int_0^{t_{k+1}} e^{-cs} \mathbb{E} \left[\int_\Lambda \psi_\epsilon(u_{h,N}^r(s, x)) u_{h,N}^r(s, x) dx \right] ds \\ &= \sum_{n=0}^k \sum_{K \in \mathcal{T}} \int_{t_n}^{t_{n+1}} e^{-cs} \mathbb{E} \left[\int_K \psi_\epsilon(u_K^{n+1}) u_K^{n+1} dx \right] ds \\ &\leq \Delta t \sum_{n=0}^k \sum_{K \in \mathcal{T}} m_K e^{-ct_n} \mathbb{E} [\psi_\epsilon(u_K^{n+1}) u_K^{n+1}]. \end{aligned} \quad (5.9)$$

- We have also the following majoration of the fourth sum:

$$\begin{aligned}
& \frac{\Delta t}{2} \sum_{n=0}^k \sum_{K \in \mathcal{T}} m_K e^{-ct_n} \mathbb{E} [g^2(u_K^n)] \\
& \leq \frac{\Delta t}{2} \sum_{K \in \mathcal{T}} m_K \mathbb{E} [g^2(u_K^0)] + \frac{1}{2} \int_0^{t_k} e^{-cs} \mathbb{E} \left[\|g(u_{h,N}^r)(s)\|_{L^2(\Lambda)}^2 \right] ds.
\end{aligned} \tag{5.10}$$

- Using the properties of the exponential function again, the last sum can be handled in the following manner:

$$\begin{aligned}
& \Delta t \sum_{n=0}^k \sum_{K \in \mathcal{T}} m_K e^{-ct_n} \mathbb{E} [(\beta(u_K^{n+1}) + f_K^n) u_K^{n+1}] \\
& \leq \int_0^{t_{k+1}} e^{-cs} \mathbb{E} \left[\int_{\Lambda} (\beta(u_{h,N}^r(s, x)) + f_{h,N}^l(s, x)) u_{h,N}^r(s, x) dx \right] ds \\
& \quad + c\Delta t \|u_{h,N}^r\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))} (L_{\beta} \|u_{h,N}^r\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))} + \|f\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))}) .
\end{aligned} \tag{5.11}$$

Combining (5.6), (5.7), (5.8), (5.9), (5.10) and (5.11), one gets

$$\begin{aligned}
& \sum_{K \in \mathcal{T}} m_K e^{-ct_k} \mathbb{E} [(u_K^{k+1})^2] + 2 \int_0^{t_{k+1}} e^{-cs} \mathbb{E} [|u_{h,N}^r(s)|_{1,h}^2] ds \\
& \quad + 2 \int_0^{t_{k+1}} e^{-cs} \mathbb{E} \left[\int_{\Lambda} \psi_{\epsilon}(u_{h,N}^r(s, x)) u_{h,N}^r(s, x) dx \right] ds \\
& \leq e^{c\Delta t} \sum_{K \in \mathcal{T}} m_K \mathbb{E} [(u_K^0)^2] + \Delta t \sum_{K \in \mathcal{T}} m_K \mathbb{E} [g^2(u_K^0)] + \int_0^{t_k} e^{-cs} \mathbb{E} \left[\|g(u_{h,N}^r)(s)\|_{L^2(\Lambda)}^2 \right] ds \\
& \quad + 2 \int_0^{t_{k+1}} e^{-cs} \mathbb{E} \left[\int_{\Lambda} (\beta(u_{h,N}^r(s, x)) + f_{h,N}^l(s, x)) u_{h,N}^r(s, x) dx \right] ds \\
& \quad - ce^{-c\Delta t} \int_0^{t_k} e^{-cs} \mathbb{E} \left[\|u_{h,N}^r(s)\|_{L^2(\Lambda)}^2 \right] ds \\
& \quad + 2c\Delta t \|u_{h,N}^r\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))} (L_{\beta} \|u_{h,N}^r\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))} + \|f\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))}) .
\end{aligned}$$

Moreover, for $t \in [t_k, t_{k+1})$ since $e^{-ct} \leq e^{-ct_k}$ and $(t - \Delta t)^+ \leq t_k$, one obtains that

$$\begin{aligned}
& e^{-ct} \mathbb{E} \left[\|u_{h,N}^r(t)\|_{L^2(\Lambda)}^2 \right] + 2 \int_0^t e^{-cs} \mathbb{E} [\|u_{h,N}^r(s)\|_{1,h}^2] ds \\
& + 2 \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} \psi_{\epsilon}(u_{h,N}^r(s, x)) u_{h,N}^r(s, x) dx \right] ds \\
\leq & e^{c\Delta t} \sum_{K \in \mathcal{T}} m_K \mathbb{E} [(u_K^0)^2] + \Delta t \sum_{K \in \mathcal{T}} m_K \mathbb{E} [g^2(u_K^0)] + \int_0^t e^{-cs} \mathbb{E} \left[\|g(u_{h,N}^r)(s)\|_{L^2(\Lambda)}^2 \right] ds \\
& + 2 \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} (\beta(u_{h,N}^r(s, x)) + f_{h,N}^l(s, x)) u_{h,N}^r(s, x) dx \right] ds \\
& + 2 \int_t^{t_{k+1}} e^{-cs} \mathbb{E} \left[\int_{\Lambda} (\beta(u_{h,N}^r(s, x)) + f_{h,N}^l(s, x)) u_{h,N}^r(s, x) dx \right] ds \\
& - ce^{-c\Delta t} \int_0^{(t-\Delta t)^+} e^{-cs} \mathbb{E} \left[\|u_{h,N}^r(s)\|_{L^2(\Lambda)}^2 \right] ds \\
& + 2c\Delta t \|u_{h,N}^r\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))} (L_{\beta} \|u_{h,N}^r\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))} + \|f\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))}) .
\end{aligned}$$

Using the constant $K_0 > 0$ given by Proposition 4.2, one gets the estimate

$$\begin{aligned}
& \int_{(t-\Delta t)^+}^t e^{-cs} \mathbb{E} \left[\|u_{h,N}^r(s)\|_{L^2(\Lambda)}^2 \right] ds \\
& + 2 \int_t^{t_{k+1}} e^{-cs} \mathbb{E} \left[\int_{\Lambda} (\beta(u_{h,N}^r(s, x)) + f_{h,N}^l(s, x)) u_{h,N}^r(s, x) dx \right] ds \\
& \leq \Delta t (1 + 2L_{\beta}) K_0 + 2\sqrt{\Delta t} \|f\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))} \sqrt{K_0},
\end{aligned}$$

and Chasles' relation $-\int_0^{(t-\Delta t)^+} = -\int_0^t + \int_{(t-\Delta t)^+}^t$ yields

$$\begin{aligned}
& e^{-ct} \mathbb{E} \left[\|u_{h,N}^r(t)\|_{L^2(\Lambda)}^2 \right] + 2 \int_0^t e^{-cs} \mathbb{E} [\|u_{h,N}^r(s)\|_{1,h}^2] ds \\
& + 2 \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} \psi_{\epsilon}(u_{h,N}^r(s, x)) u_{h,N}^r(s, x) dx \right] ds \\
\leq & e^{c\Delta t} \mathbb{E} [\|u_0\|_{L^2(\Lambda)}^2] + \Delta t L_g^2 \mathbb{E} [\|u_0\|_{L^2(\Lambda)}^2] + \int_0^t e^{-cs} \mathbb{E} \left[\|g(u_{h,N}^r)(s)\|_{L^2(\Lambda)}^2 \right] ds \quad (5.12) \\
& + 2 \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} (\beta(u_{h,N}^r(s, x)) + f_{h,N}^l(s, x)) u_{h,N}^r(s, x) dx \right] ds \\
& - ce^{-c\Delta t} \int_0^t e^{-cs} \mathbb{E} \left[\|u_{h,N}^r(s)\|_{L^2(\Lambda)}^2 \right] ds \\
& + c\Delta t (1 + 2L_{\beta}) K_0 + 2c\sqrt{\Delta t} \|f\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))} \sqrt{K_0} \\
& + 2c\Delta t \|u_{h,N}^r\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))} (L_{\beta} \|u_{h,N}^r\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))} + \|f\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))}) .
\end{aligned}$$

Furthermore,

$$\begin{aligned}
& \int_0^t e^{-cs} \mathbb{E} \left[\|g(u_{h,N}^r)(s)\|_{L^2(\Lambda)}^2 \right] ds \\
&= \int_0^t e^{-cs} \mathbb{E} \left[\|g(u_{h,N}^r)(s) - g(u)(s)\|_{L^2(\Lambda)}^2 \right] ds \\
& \quad + 2 \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} g(u_{h,N}^r)(s, x) g(u)(s, x) dx \right] ds - \int_0^t e^{-cs} \mathbb{E} \left[\|g(u)(s)\|_{L^2(\Lambda)}^2 \right] ds.
\end{aligned} \tag{5.13}$$

In the same manner,

$$\begin{aligned}
& - ce^{-c\Delta t} \int_0^t e^{-cs} \mathbb{E} \left[\|u_{h,N}^r(s)\|_{L^2(\Lambda)}^2 \right] ds \\
&= - ce^{-c\Delta t} \int_0^t e^{-cs} \mathbb{E} \left[\|u_{h,N}^r(s) - u(s)\|_{L^2(\Lambda)}^2 \right] ds \\
& \quad - 2ce^{-c\Delta t} \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} u_{h,N}^r(s, x) u(s, x) dx \right] ds + ce^{-c\Delta t} \int_0^t e^{-cs} \mathbb{E} \left[\|u(s)\|_{L^2(\Lambda)}^2 \right] ds.
\end{aligned} \tag{5.14}$$

And at last

$$\begin{aligned}
& \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} \beta(u_{h,N}^r(s, x)) u_{h,N}^r(s, x) dx \right] ds \\
&= \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} \beta(u_{h,N}^r(s, x)) u(s, x) dx \right] ds \\
& \quad + \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} (\beta(u_{h,N}^r(s, x)) - \beta(u(s, x))) (u_{h,N}^r(s, x) - u(s, x)) dx \right] ds \\
& \quad + \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} \beta(u(s, x)) (u_{h,N}^r(s, x) - u(s, x)) dx \right] ds.
\end{aligned} \tag{5.15}$$

Finally, by considering from now on a parameter $c > 0$ depending only on L_g and L_β such that for any N big enough

$$\begin{aligned}
& \int_0^t e^{-cs} \mathbb{E} \left[\|g(u_{h,N}^r(s)) - g(u(s))\|_{L^2(\Lambda)}^2 \right] ds - ce^{-c\Delta t} \int_0^t e^{-cs} \mathbb{E} \left[\|u_{h,N}^r(s) - u(s)\|_{L^2(\Lambda)}^2 \right] ds \\
& \quad + 2 \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} (\beta(u_{h,N}^r(s, x)) - \beta(u(s, x))) (u_{h,N}^r(s, x) - u(s, x)) dx \right] ds \leq 0,
\end{aligned}$$

we are able to prove that, after injecting (5.13), (5.14) and (5.15) in (5.12), and integrating

from 0 to T :

$$\begin{aligned}
& \int_0^T e^{-ct} \mathbb{E} \left[\|u_{h,N}^r(t)\|_{L^2(\Lambda)}^2 \right] dt + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} [|u_{h,N}^r(s)|_{1,h}^2] ds dt \\
& + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} \psi_{\epsilon}(u_{h,N}^r(s,x)) u_{h,N}^r(s,x) dx \right] ds dt \\
\leq & \int_0^T \mathbb{E} [\|u_0\|_{L^2(\Lambda)}^2] dt + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} g(u_{h,N}^r(s,x)) g(u)(s,x) dx \right] ds dt \\
& - \int_0^T \int_0^t e^{-cs} \mathbb{E} [\|g(u(s))\|_{L^2(\Lambda)}^2] ds dt - 2ce^{-c\Delta t} \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} u_{h,N}^r(s,x) u(s,x) dx \right] ds dt \\
& + ce^{-c\Delta t} \int_0^T \int_0^t e^{-cs} \mathbb{E} [\|u(s)\|_{L^2(\Lambda)}^2] ds dt + T (e^{c\Delta t} - 1) \mathbb{E} [\|u_0\|_{L^2(\Lambda)}^2] \\
& + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} \beta(u_{h,N}^r(s,x)) u(s,x) dx \right] ds dt \\
& + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} \beta(u(s,x)) (u_{h,N}^r(s,x) - u(s,x)) dx \right] ds dt \\
& + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} f_{h,N}^l(s,x) u_{h,N}^r(s,x) dx \right] ds dt \\
& + c\Delta t T (1 + 2L_{\beta}) K_0 + 2cT \sqrt{\Delta t} \|f\|_{L^2(\Omega; L^2(0,T; L^2(\Lambda)))} \sqrt{K_0} \\
& + 2c\Delta t T \|u_{h,N}^r\|_{L^2(\Omega; L^2(0,T; L^2(\Lambda)))} (L_{\beta} \|u_{h,N}^r\|_{L^2(\Omega; L^2(0,T; L^2(\Lambda)))} + \|f\|_{L^2(\Omega; L^2(0,T; L^2(\Lambda)))}) .
\end{aligned}$$

Firstly, by passing to the superior limit in this last inequality, Remark 5.5 and Lemma 5.8 allow us to state that:

$$\begin{aligned}
& \limsup_{m \rightarrow +\infty} \int_0^T e^{-ct} \mathbb{E} [\|u_{h,N}^r(t)\|_{L^2(\Lambda)}^2] dt + 2 \liminf_{m \rightarrow +\infty} \int_0^T \int_0^t e^{-cs} \mathbb{E} [|u_{h,N}^r(s)|_{1,h}^2] ds dt \\
& + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} \psi_2(s,x) dx \right] ds dt \\
\leq & \int_0^T \mathbb{E} [\|u_0\|_{L^2(\Lambda)}^2] dt + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} g_u(s,x) g(u)(s,x) dx \right] ds dt \\
& - \int_0^T \int_0^t e^{-cs} \mathbb{E} [\|g(u(s))\|_{L^2(\Lambda)}^2] ds dt - c \int_0^T \int_0^t e^{-cs} \mathbb{E} [\|u(s)\|_{L^2(\Lambda)}^2] ds dt \\
& + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} \beta_u(s,x) u(s,x) dx \right] ds dt + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} f(s,x) u(s,x) dx \right] ds dt.
\end{aligned}$$

Secondly, the stochastic energy equality (5.4) yields

$$\begin{aligned}
& \limsup_{m \rightarrow +\infty} \int_0^T e^{-ct} \mathbb{E} \left[\|u_{h,N}^r(t)\|_{L^2(\Lambda)}^2 \right] dt + 2 \liminf_{m \rightarrow +\infty} \int_0^T \int_0^t e^{-cs} \mathbb{E} [|u_{h,N}^r(s)|_{1,h}^2] ds dt \\
& + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} \psi_2(s, x) - \psi(s, x)u(s, x) dx \right] ds dt \\
& \leq \int_0^T e^{-ct} \mathbb{E} \left[\|u(t)\|_{L^2(\Lambda)}^2 \right] dt + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\|\nabla u(s)\|_{L^2(\Lambda)}^2 \right] ds dt \\
& \quad - \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\|g(u(s)) - g_u(s)\|_{L^2(\Lambda)}^2 \right] ds dt.
\end{aligned}$$

Thirdly, thanks to the key Inequality (5.5) given by Lemma 5.11, we arrive at

$$\begin{aligned}
& \limsup_{m \rightarrow +\infty} \int_0^T e^{-ct} \mathbb{E} \left[\|u_{h,N}^r(t)\|_{L^2(\Lambda)}^2 \right] dt + \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\|g(u(s)) - g_u(s)\|_{L^2(\Lambda)}^2 \right] ds dt \\
& + 2 \int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} \psi_2(s, x) - \psi(s, x)u(s, x) dx \right] ds dt \\
& \leq \int_0^T e^{-ct} \mathbb{E} \left[\|u(t)\|_{L^2(\Lambda)}^2 \right] dt. \tag{5.16}
\end{aligned}$$

Owing to the weak convergence of $(u_{h,N}^r)_m$ towards u in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$, we can affirm that the following inequality holds true

$$\int_0^T e^{-ct} \mathbb{E} \left[\|u(t)\|_{L^2(\Lambda)}^2 \right] dt \leq \liminf_{m \rightarrow +\infty} \int_0^T e^{-ct} \mathbb{E} \left[\|u_{h,N}^r(t)\|_{L^2(\Lambda)}^2 \right] dt,$$

so that

$$\int_0^T \int_0^t e^{-cs} \mathbb{E} \left[\int_{\Lambda} \psi_2(s, x) - \psi(s, x)u(s, x) + (g(u(s, x)) - g_u(s, x))^2 dx \right] ds dt \leq 0. \tag{5.17}$$

By Lemma 5.4 and Remark 5.5, we can affirm that \mathbb{P} -almost surely in Ω and almost everywhere in $(0, T) \times \Lambda$, $0 \leq u \leq 1$ and $\psi = \psi_1 + \psi_2$ with $\psi_1 \leq 0$ and $\psi_2 \geq 0$. Then $\psi_2 - \psi u = (1 - u)\psi_2 - \psi_1 u \geq 0$ and (5.17) allows us to say that $g_u = g(u)$ and that $\psi_2 - \psi u = 0$. In particular, we have since $\psi_2 - \psi u = (1 - u)\psi_2 - \psi_1 u$:

- In the set $\{u = 0\}$, then $\psi_2 - \psi u = 0$ implies that $\psi_2 = 0$ and so $\psi = \psi_1 \leq 0$.
- In the set $\{u = 1\}$, then $\psi_2 - \psi u = 0$ implies that $\psi_1 = 0$ and so $\psi = \psi_2 \geq 0$.
- In the set $\{0 < u < 1\}$, then $\psi_2 - \psi u = 0$ implies that $\psi_1 = \psi_2 = 0$ and so $\psi = 0$.

In this manner, $\psi \in \partial I_{[0,1]}(u)$. Going back to (5.16), we have

$$\limsup_{m \rightarrow +\infty} \int_0^T e^{-ct} \mathbb{E} \left[\|u_{h,N}^r(t)\|_{L^2(\Lambda)}^2 \right] dt \leq \int_0^T e^{-ct} \mathbb{E} \left[\|u(t)\|_{L^2(\Lambda)}^2 \right] dt, \tag{5.18}$$

and one can thus conclude from (5.18) and Proposition 5.1 that $(u_{h,N}^r)_m$ converges strongly to u in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$, that $\beta_u = \beta(u)$ and that (u, ψ) is the unique variational solution of Problem (1.1) in the sense of Definition 1.1. As a consequence of [42, Corollary 1.2.23, p.25] with $(S, \mathcal{A}, \mu) = (\Omega, \mathcal{F}, \mathbb{P})$, $(T, \mathcal{B}, \nu) = ((0, T), \mathcal{B}(0, T), \lambda)$, $X = L^2(\Lambda)$, $p = 2$, $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$ is isometrically isomorphic to $L^2(0, T; L^2(\Omega; L^2(\Lambda)))$, hence $(u_{h,N}^r)$ also converges strongly towards u in $L^2(0, T; L^2(\Omega; L^2(\Lambda)))$. Combining this last information with the boundedness of $(u_{h,N}^r)_m$ in $L^\infty(0, T; L^2(\Omega; L^2(\Lambda)))$ (see Lemma 4.3) allows us to conclude that $(u_{h,N}^r)_m$ converges strongly towards u in $L^p(0, T; L^2(\Omega; L^2(\Lambda)))$ for any finite $p \geq 1$ thanks to Vitali's theorem [36, Corollaire 1.3.3]. \square

Acknowledgments The authors would like to thank J. Droniou and T. Gallouët for their valuable suggestions. This work has been supported by the German Research Foundation project (ZI 1542/3-1) and various Procope programs: Project-Related Personal Exchange France-Germany (49368YE), Procope Mobility Program (DEU-22-0004 LG1) and Procope Plus Project (0185-DEU-22-001 LG 4).

References

- [1] S.M. Allen and J. W. Cahn. A microscopic theory for antiphase boundary motion and its application to antiphase domain coarsening. *Acta Metall.*, 27(6):1085–1095, 1979.
- [2] R. Anton, D. Cohen, and L. Quer-Sardanyons. A fully discrete approximation of the one-dimensional stochastic heat equation. *IMA J. Numer. Anal.*, 40(1):247–284, 2020.
- [3] D. C. Antonopoulou, L. Bañas, R. Nürnberg, and A. Prohl. Numerical approximation of the stochastic Cahn-Hilliard equation near the sharp interface limit. *Numerische Mathematik*, 147:1–47, 03 2021.
- [4] D. C. Antonopoulou, G. Karali, and A. Millet. Existence and regularity of solution for stochastic Cahn-Hilliard/Allen-Cahn equation with unbounded noise diffusion. *J. Differential Equations*, 260(3):2383–2417, 2016.
- [5] Z. Artstein. First-order approximations for differential inclusions. *Set-Valued Anal.*, 2:7–18, 1994.
- [6] V. Barbu. *Nonlinear Semigroups and Differential Equations in Banach Spaces*. Noordhoff, Leyden, 1976.
- [7] V. Barbu and G. Da Prato. Irreducibility of the transition semigroup associated with the stochastic obstacle problem. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, 8(3):397–406, 2005.
- [8] V. Barbu and A. Răscănu. Parabolic variational inequalities with singular inputs. *Differ. Integral Equ.*, 10(1):67–83, 1997.

- [9] J. Bastien and M. Schatzman. Numerical precision for differential inclusions with uniqueness. *ESAIM Math. Model. Numer. Anal.*, 36:427–460, 2002.
- [10] Ł. Bañas, Z. Brzezniak, M. Neklyudov, and A. Prohl. *Stochastic ferromagnetism*, volume 58 of *De Gruyter Studies in Mathematics*. De Gruyter, Berlin, 2014. Analysis and numerics.
- [11] C. Bauzet, E. Bonetti, G. Bonfanti, F. Lebon, and G. Vallet. A global existence and uniqueness result for a stochastic Allen-Cahn equation with constraint. *Math. Methods Appl. Sci.*, 40(14):5241–5261, September 2017.
- [12] C. Bauzet, F. Nabet, K. Schmitz, and A. Zimmermann. Convergence of a finite-volume scheme for a heat equation with a multiplicative lipschitz noise. *ESAIM Math. Model. Numer. Anal.*, 57(2):745–783, March 2023.
- [13] C. Bauzet, F. Nabet, K. Schmitz, and A. Zimmermann. Finite Volume Approximations for Non-Linear Parabolic Problems with Stochastic Forcing. In *Finite Volumes for Complex Applications X - Methods and Theoretical Aspects*. Springer, 2023.
- [14] C. Bauzet, K. Schmitz, and A. Zimmermann. On a finite-volume approximation of a diffusion-convection equation with a multiplicative stochastic force. Submitted, April 2023.
- [15] S. Becker, B. Gess, A. Jentzen, and P. E. Kloeden. Strong convergence rates for explicit space-time discrete numerical approximations of stochastic Allen-Cahn equations. *Stoch. Partial Differ. Equ.: Anal. Comput.*, 11:211–268, 2023.
- [16] A. Bensoussan and A. Rascanu. Stochastic variational inequalities in infinite dimensional spaces. *Numer. Funct. Anal. Optim.*, 18(1-2):19–54, 1997.
- [17] F. Bernardin. Multivalued stochastic differential equations : convergence of a numerical scheme. *Set-Valued Anal*, 11:393–415, 2003.
- [18] F. Bertacco. Stochastic Allen-Cahn equation with logarithmic potential. *Nonlinear Anal.*, 202:112122, 01 2021.
- [19] F. Bertacco, C. Orrieri, and L. Scarpa. Random separation property for stochastic Allen-Cahn-type equations. *Electron. J. Probab.*, 27, 01 2022.
- [20] W. J. Beyn, E. Emmrich, and J. Rieger. Semilinear parabolic differential inclusions with one-sided Lipschitz nonlinearities. *J. Evol. Equ.*, 18:1319–1339, 2018.
- [21] W. J. Beyn and J. Rieger. Numerical fixed grid methods for differential inclusions. *Computing*, 81:91–106, 2007.
- [22] W. J. Beyn and J. Rieger. The implicit Euler scheme for one-sided Lipschitz differential inclusions. *Discrete Contin. Dyn. Syst. Ser. B*, 14:409–428, 2010.

- [23] W. J. Beyn and J. Rieger. Galerkin finite element methods for semilinear elliptic differential inclusions. *Discrete Contin. Dyn. Syst. Ser. B*, 18:295–312, 2013.
- [24] D. Breit, M. Hofmanová, and S. Loisel. Space-time approximation of stochastic p -Laplace-type systems. *SIAM J. Numer. Anal.*, 59(4):2218–2236, 2021.
- [25] D. Breit and A. Prohl. Weak error analysis for the stochastic Allen–Cahn equation. *Stoch. Partial Differ. Equ.: Anal. Comput.*, 2024.
- [26] H. Brézis. *Opérateurs Maximaux Monotones et Semi-groupes de Contractions dans les Espaces de Hilbert*. Number 5 in North Holland Math. Studies, 1973.
- [27] C-E. Bréhier and L. Goudenège. Weak convergence rates of splitting schemes for the stochastic Allen–Cahn equation. *BIT Numer Math*, 60:543–582, 2020.
- [28] L. Cherfils, A. Miranville, and S. Zelik. The Cahn–Hilliard equation with logarithmic potentials. *Milan J. Math.*, 79(2):561–596, 2011.
- [29] P. Colli, G. Gilardi, P. Podio-Guidugli, and J. Sprekels. Global existence and uniqueness for a singular/degenerate Cahn–Hilliard system with viscosity. *J. Differ. Equ.*, 254(11):4217–4244, 2013.
- [30] G. Da Prato and J. Zabczyk. *Stochastic equations in infinite dimensions*, volume 152 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, second edition, 2014.
- [31] A. Debussche and J. Printems. Weak order for the discretization of the stochastic heat equation. *Math. Comp.*, 78(266):845–863, 2009.
- [32] M. Del Pino, M. Kowalczyk, F. Pacard, and J. Wei. Multiple-end solutions to the Allen–Cahn equation in \mathbb{R}^2 . *J. Funct. Anal.*, 258(2):458–503, 2010.
- [33] A. Di Primio, M. Grasselli, and L. Scarpa. Stochastic Cahn–Hilliard and conserved Allen–Cahn equations with logarithmic potential and conservative noise, 09 2023.
- [34] Andrea Di Primio, M. Grasselli, and Luca Scarpa. A stochastic Allen–Cahn–Navier–Stokes system with singular potential, 05 2022.
- [35] A. L. Dontchev and F. Lempio. Difference methods for differential inclusions: a survey. *SIAM Rev.*, 34:263–294, 1992.
- [36] J. Droniou. Intégration et Espaces de Sobolev à Valeurs Vectorielles. <https://hal.archives-ouvertes.fr/hal-01382368/document>, 2001.
- [37] J. Droniou. A density result in Sobolev spaces. *J. Math. Pures App.*, 81(7):697–714, 2002.
- [38] M. Eisenmann, M. Kovács, R. Kruse, and S. Larsson. Error estimates of the backward Euler–Maruyama method for multi-valued stochastic differential equations. *BIT Numer Math*, 62:803–848, 2021.

- [39] R. Eymard, T. Gallouët, and R. Herbin. Finite Volume Methods. In *Handbook of numerical analysis, Vol. VII*, Handb. Numer. Anal., VII, pages 713–1020. North-Holland, Amsterdam, 2000.
- [40] X. Feng and A. Prohl. Numerical analysis of the Allen-Cahn equation and approximation for mean curvature flows. *Numerische Mathematik*, 94:33–65, 03 2003.
- [41] G. Grammel. Towards fully discretized differential inclusions. *Set-Valued Anal*, pages 1–8, 2003.
- [42] T. Hytonen, J. van Neerven, M. Veraar, and L. Weis. *Analysis in Banach Spaces, Volume I: Martingales and Littlewood-Paley Theory*. 2016.
- [43] F. Lempio and V. Veliov. Discrete approximations of differential inclusions. *Bayreuth. Math. Schr.*, 54:149–232, 1998.
- [44] D. Leplingue and T. T. Nguyen. Approximating and simulating multivalued stochastic differential equations. *Monte Carlo Methods Appl.*, 10:129–152, 2004.
- [45] Z. Liu and Z. Qiao. Strong approximation of monotone stochastic partial differential equations driven by multiplicative noise. *Stoch. Partial Differ. Equ.: Anal. Comput.*, 9:559–602, 03 2021.
- [46] T. Medjo. A stochastic Allen-Cahn/Navier-Stokes model with inertial effects driven by multiplicative noise of jump type. *Math. Nachr.*, 296:4386–4428, 2023.
- [47] B. S. Mordukhovich. Discrete approximations and refined Euler-Lagrange conditions for differential inclusions. *SIAM J. Control Optim.*, 33:882–915, 1995.
- [48] B. S. Mordukhovich and Y. Tian. Implicit Euler approximation and optimization of one-sided Lipschitzian differential inclusions. *Contemp. Math.*, 659:165–188, 2016.
- [49] M. Ondrejat, A. Prohl, and N. Walkington. Numerical approximation of nonlinear SPDE’s. *Stoch. Partial Differ. Equ.: Anal. Comput.*, 2022.
- [50] R. Pettersson. Projection scheme for stochastic differential equations with convex constraints. *Stochastic Process. Appl.*, 88:125–134, 2000.
- [51] C. Prévôt and M. Röckner. *A concise course on stochastic partial differential equations*, volume 1905 of *Lecture Notes in Mathematics*. Springer, Berlin, 2007.
- [52] R. Qi and X. Wang. Optimal Error Estimates of Galerkin Finite Element Methods for Stochastic Allen–Cahn Equation with Additive Noise. *J. Sci. Comput.*, 80:1171–1194, 2019.
- [53] A. Rascanu. Deterministic and stochastic differential equations in Hilbert spaces involving multivalued maximal monotone operators. *PanAmer. Math. J.*, pages 83–119, 1996.

- [54] J. Rieger. Discretizations of linear elliptic partial differential inclusions. *Num. Funct. Anal. Opt.*, 32:904–925, 2011.
- [55] J. Rieger. Semi-implicit Euler schemes for ordinary differential inclusions. *SIAM J. Numer. Anal.*, 52:895–914, 2014.
- [56] M. D. Ryser, N. Nigam, and P. F. Tupper. On the well-posedness of the stochastic Allen-Cahn equation in two dimensions. *J. Comput. Phys.*, 231(6):2537–2550, 2012.
- [57] L. Scarpa and M. Zanella. Degenerate Kolmogorov equations and ergodicity for the stochastic Allen-Cahn equation with logarithmic potential. *Stoch. Partial Differ. Equ.: Anal. Comput.*, 12:1–45, 01 2023.
- [58] Y. Tahraoui and G. Vallet. Lewy-Stampacchia’s inequality for a stochastic T -monotone obstacle problem. *Stoch. Partial Differ. Equ. Anal. Comput.*, 10(1):90–125, 2022.
- [59] V. Veliov. Second order discrete approximations to strongly convex differential inclusions. *Syst. Control Lett.*, 13:263–269, 1989.
- [60] M. J. Ward. Metastable bubble solutions for the Allen-Cahn equation with mass conservation. *SIAM J. Appl. Math.*, 56(5):1247–1279, 1996.
- [61] J. Wu and H. Zhang. Penalization schemes for multi-valued stochastic differential equations. *Statist. Probab. Lett.*, 83:481–492, 2013.
- [62] H. Zhang. Strong convergence rate for multivalued stochastic differential equations via stochastic theta method. *STOCHASTICS*, 90:762–781, 2018.