



**HAL**  
open science

# Networks of sign-changing metamaterials: existence and spectral properties

Kaïs Ammari, Eric Bonnetier, Alessandro Duca

► **To cite this version:**

Kaïs Ammari, Eric Bonnetier, Alessandro Duca. Networks of sign-changing metamaterials: existence and spectral properties. 2024. hal-04639514v1

**HAL Id: hal-04639514**

**<https://hal.science/hal-04639514v1>**

Preprint submitted on 9 Jul 2024 (v1), last revised 29 Jul 2024 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# Superlinsing using sign-changing metamaterials on networks

Kaïs Ammari,<sup>\*</sup> Alessandro Duca,<sup>†</sup> Eric Bonnetier,<sup>‡</sup>

July 9, 2024

## Abstract

We study composite assemblages of dielectrics and metamaterials with respectively positive and negative material parameters. In the continuum case, for a scalar equation, such media may exhibit so-called plasmonic resonances for certain values of the (negative) conductivity in the metamaterial. This work investigates such resonances, and the associated eigenfunctions, in the case of composite conducting networks. Unlike the continuous media, we show a surprising specific dependence on the geometry of the network of the resonant values. We also study how the problem is affected by the choice of boundary conditions on the external nodes of the structure.

**AMS subject classifications:** 34B45, 47B25, 34L05.

**Keywords:** Metamaterials, networks, T-coercivity, spectral decomposition.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Star-graph networks with Dirichlet boundaries</b>	<b>4</b>
<b>3</b>	<b>Star-networks with Dirichlet and Neumann boundaries</b>	<b>10</b>
<b>4</b>	<b>Tadpole-shaped networks</b>	<b>13</b>
<b>5</b>	<b>Spectral representation for star-networks with Dirichlet boundaries</b>	<b>16</b>
<b>6</b>	<b>Spectral representation for star-networks with mixed boundaries</b>	<b>26</b>
<b>7</b>	<b>Spectral representation for two-phase tadpole networks</b>	<b>30</b>

---

<sup>\*</sup>LR Analysis and Control of PDEs, LR 22ES03, Department of Mathematics, Faculty of Sciences of Monastir, University of Monastir, Tunisia; e-mail: kais.ammari@fsm.rnu.tn

<sup>†</sup>Université de Lorraine, CNRS, INRIA, IECL, F-54000 Nancy, France; e-mail: alessandro.duca@inria.fr

<sup>‡</sup>Institut Fourier, Université Grenoble Alpes, 100 Rue des Mathématiques, 38610 Gières, France; e-mail: eric.bonnetier@univ-grenoble-alpes.fr

# 1 Introduction

Negative index materials possess spectacular properties of concentration and localization of light, which are very interesting for applications, in particular in optics and communication. The first works on negative index materials may date back to Veselago [32] in 1969. They have really become a subject of active research about two decades ago, when the development of lithographic techniques has made possible the manufacturing of metallic objects with dimensions smaller than the wavelengths in the range of visible light. Indeed, in this range of frequencies, objects of small dimensions made of metals such as silver, gold or aluminium, may have complex dielectric coefficients with a negative real part and a small imaginary part, and can exhibit concentration effects (hot spots) when interacting with an incident electromagnetic wave. Such features find useful applications in medical imaging or microscopy, such as for instance single molecule detectors.

The simplest model put forward to represent the associated resonance effect is the so-called plasmonic electrostatic model [21] for the voltage potential  $u$

$$\begin{cases} \operatorname{div}(a(x)\nabla u(x)) &= 0, & \text{in } \Omega \\ u(x) &= 0, & \text{in } \partial\Omega \end{cases} \quad (1)$$

where  $\Omega \subset \mathbb{R}^n$  is a domain that represents a composite medium made of a subset  $D$  of metamaterial with conductivity  $k < 0$ , embedded in a background dielectric material of conductivity 1, so that

$$\alpha(x) = 1 + (k - 1)1_D(x), \quad x \in \Omega.$$

Resonant states are defined as non-trivial solutions to (1), which, in view of the Lax Milgram Lemma, may only exist when  $k < 0$ .

There is a large body of work on the mathematical analysis of the well-posedness of (1) (see for instance [2, 3, 4, 10, 11, 18, 19, 21, 27, 28]), where techniques from potential theory or from variational analysis are used. The solution to (1) can indeed be represented as  $u = S_D\varphi$ , *i.e.* as the single layer potential of a function  $\varphi \in H^{-1/2}(\partial D)$ , and one is led to study an integral equation of the form  $(\lambda I - K^*)\varphi = 0$ , where  $\lambda = \frac{k+1}{2(k-1)}$  and  $K^*$  is the associated Neumann-Poincaré operator

$$K^*\varphi = \int_{\partial D} \frac{\nu(x) \cdot (x - y)}{|x - y|^2} \varphi(y) d\sigma(y).$$

When  $D$  is smooth,  $K^* : H^{-1/2}(\partial D) \rightarrow H^{-1/2}(\partial D)$  is a compact operator, and its spectrum is formed by a sequence of eigenvalues of finite multiplicity, that converge to 0. The latter value  $\lambda = 0$  corresponds to a conductivity

$$k = -1 \quad (2)$$

inside the inclusion  $D$ , which in this situation is the only value of the conductivity for which the operator  $Au = \operatorname{div}\left(1 + (k - 1)1_D(x)\nabla u\right)$  looses its Fredholm character [5, 6].

One could also investigate the well-posedness of second order scalar equations with sign-changing coefficient using the notion of T-coercivity [5, 6, 8, 16, 30], which relates to variational methods. A bilinear form  $a : H^1(\Omega) \times H^1(\Omega) \rightarrow \mathbb{R}$  is T-coercive if one can find an isomorphism

$$T : H^1(\Omega) \rightarrow H^1(\Omega)$$

such that  $a(T\cdot, \cdot)$  satisfies the hypotheses of the Lax-Milgram Lemma. In the framework of (1), T-coercivity may only provide necessary conditions on the values of  $k$  for which the associated operator is well-posed. However, contrarily to the integral equation approach, it may be applied to situations where the coefficients are not merely piecewise constant. The T-coercivity has been also adopted in homogenization problems with sign-changing materials in [12, 13, 14].

### Sign-changing scalar problems on networks and our results

In this work, we investigate the behaviour of networks made of mixtures of dielectric and metamaterials. We study the associated electrostatic plasmonic equation for star-shaped and tadpole networks, which takes the variational form : find  $u \in H$ , such that for all  $v \in H$ ,

$$\sum_{i=1}^N \int_{e_i} a_i(x) u'_i(x) v'_i(x) dx = \sum_{i=1}^N \int_{e_i} f_i(x) v_i(x) dx. \quad (3)$$

Here  $(e_i)_{i \leq N}$  denotes the  $N$  edge of the network,  $f = (f_1, \dots, f_N)$  is a source term, and the field  $u = (u_1, \dots, u_N)$  is sought in a subspace  $H$  of  $\prod_{i=1}^N H^1(e_i)$ , and satisfies transmission and boundary conditions. This problem can be studied much more directly than the case of continuous media, since the PDE can be reduced to a system of ODE's coupled via the transmission conditions between the edges of the network. In this work, we proceed as follows.

- We start by considering star-graph networks equipped with Dirichlet boundary conditions on the external vertices. We firstly study two-phase networks composed by  $N$  equal edges, where the conductivity of the  $D$  dielectric edges is 1 and the negative ones have conductivity  $k < 0$ . We ensure the well-posedness of the problem (3) via the T-coercivity when

$$k \neq -\frac{D}{N-D} \quad (4)$$

(see Theorem 2.1). Notice that the resonant value depends on the geometry of the structure and differs from the value  $-1$ , usually appearing for smooth continuous media (2). Secondly, we extend the result to the case of general star-graph networks equipped with Dirichlet boundary conditions (Corollary 2.2, Theorem 2.3 and Corollary 2.4). Also in the general case, the geometry of the graph plays an important role in the admissible conductivities for the well-posedness of (3).

- Afterwards, we consider star-shaped networks with Dirichlet and Neumann boundary conditions and composed by homogenous materials. We use a different technique from the T-coercivity: we study the transmission relation associated to the variational problem (3). Here, the loss of the Fredholm character occurs at some specific values, in general different from  $-1$ , again depending on the geometry of the structure. Also in this case, we start by considering two-phase networks (see Theorem 3.1), and later we extend the result to the general case (Corollary 3.4). An application of our general result covers the case of a star-network with edges of conductivities  $k_j$  and lengths  $L_j$ . In this framework, we denote  $I_d$  the edges equipped with Dirichlet boundary conditions and the well-posedness of (3) is ensured when

$$\sum_{l \in I_d} \frac{k_l}{L_l} \neq 0. \quad (5)$$

The identity (5) shows that the well-posedness of (3) is not affected by the presence of edges equipped with Neumann boundary conditions, but only by the "Dirichlet part" of the network. We conclude the first part of the work by applying the previous techniques to study the well-posedness for tadpole networks (Theorem 4.1, Theorem 4.3 and Corollary 4.4). Notice that this approach can be applied to any network: the PDE is reduced to a linear system of equation representing the transmission conditions between the edges of the network. Clearly, the more the structure is complex, the more the study of the well-posedness of the system becomes complicated.

- The second part of the work studies the spectral problem associated with (3) for the networks considered before (Section 5, Section 6 and Section 7). We seek for  $(u, \lambda)$  verifying

$$\forall v \in H, \quad \sum_{i=1}^N \int_{e_i} a_i(x) u'_i(x) v'_i(x) dx = \lambda \sum_{i=1}^N \int_{e_i} u_i(x) v_i(x) dx. \quad (6)$$

We characterize the spectral elements of the operator in different frameworks, which has both positive and negative eigenvalues. This implies that one cannot expect to construct a solution to a parabolic equation for a general initial datum, even when the corresponding stationary operator is T-coercive. One could, however, construct solutions for a Schrödinger-type equation

$$i \partial_t u(t, x) + \partial_x \left( a(x) \partial_x u(t, x) \right) = f(t, x).$$

Subsequently, we construct another 'natural' family of eigenfunctions, defined as the  $(u, \lambda)$ 's that satisfy

$$\forall v \in H, \quad \sum_{i=1}^N \int_{e_i} a_i(x) u'_i(x) v'_i(x) dx = \lambda \sum_{i=1}^N \int_{e_i} a_i(x) u_i(x) v_i(x) dx, \quad (7)$$

associated with the bilinear form  $(u, v)_a = \sum_{i=1}^N \int_{e_i} a_i(x) u_i(x) v_i(x) dx$ . This bilinear form is not a scalar product, as the  $a_i$ 's take positive and negative values. We show however that one can find a complete family of eigenfunctions of this form, which would allow one to define a solution to a 'pseudo-parabolic' equation of the form

$$a(x) \partial_t u(t, x) + \partial_x \left( a(x) \partial_x u(t, x) \right) = f(t, x).$$

## Acknowledgments

The authors acknowledge the financial support from VINCI Program 2020 of the "Université Franco Italienne" (UFI) for the project "Graphes quantiques de métamatériaux". The second author would like to thank the colleagues Karim Ramdani and Renata Bunoiu for the fruitful discussions on the behaviour of 'indefinite Laplacians' and on the well-posedness of sign-changing scalar problems.

## 2 Star-graph networks with Dirichlet boundaries

Let us consider a star-shaped network composed by  $N$  edges of lengths  $\{L_j\}_{1 \leq j \leq N}$  some composed by a material with positive conductivity and some with negative conductivity. We represent the network with a star-graph  $\mathcal{S}$  composed by  $N$  edges  $\{e_j\}_{1 \leq j \leq N}$ . We denote by  $v$  the internal vertex

connecting all the edges of  $\mathcal{S}$  and we parameterize each  $e_j$  with a coordinate going from 0 to its length  $L_j$  in  $v$ .



Figure 1: The figure shows the parametrization of a star-graph with 3 edges.

We define quantum states on  $\mathcal{S}$  as functions  $\psi = (\psi_1, \dots, \psi_N) \in L^2(\mathcal{S}) := \prod_{j=1}^N L^2(e_j, \mathbb{R})$ , where each  $\psi_j : e_j \rightarrow \mathbb{R}$ . The Hilbert space  $L^2(\mathcal{S})$  is equipped with the norm  $\|\cdot\|_{L^2}$  induced by the scalar product

$$(\psi, \varphi)_{L^2} := \sum_{1 \leq j \leq N} (\psi_j, \varphi_j)_{L^2(e_j, \mathbb{R})} = \sum_{1 \leq j \leq N} \int_0^{L_j} \psi_j(x) \varphi_j(x) dx, \quad \forall \psi, \varphi \in L^2(\mathcal{S}).$$

Let  $\widehat{H}^1(\mathcal{S}) := \prod_{j=1}^N H^1(e_j, \mathbb{R})$ . We denote by

$$H^1(\mathcal{S}) = \{\psi \in \widehat{H}^1(\mathcal{S}) : \psi_1(L_1) = \dots = \psi_N(L_N)\},$$

$$H_0^1(\mathcal{S}) = \{\psi \in H^1(\mathcal{S}) : \psi_1(0) = \dots = \psi_N(0) = 0\}.$$

By duality according to the scalar product of  $L^2(\mathcal{S})$ , we define the space  $H^{-1}(\mathcal{S})$ . For every  $f = (f_1, \dots, f_N) \in H^{-1}(\mathcal{S})$ , we investigate the existence of  $\psi = (\psi_1, \dots, \psi_N) \in H_0^1(\mathcal{S})$  solution of the following stationary problem in  $L^2(\mathcal{S})$

$$-\partial_x \left( k_j(x) \partial_x \psi_j(x) \right) = f_j(x), \quad x \in (0, L_j), \quad 1 \leq j \leq N, \quad (8)$$

where  $k_j : x \in [0, L_j] \rightarrow \mathbb{R}^*$  are the continuous functions representing the properties of the material composing the network and there exist  $I_j, S_j > 0$  such that

$$I_j \leq |k_j(x)| \leq S_j, \quad \forall x \in [0, L_j].$$

## 2.1 Two-phase networks with constant conductivities

Let us study the case of star graphs composed of  $N$  edges composed of two materials, one with positive conductivity and the other with negative conductivity. In the following theorem, we assume that all the conductivities are constants and that the star graph is equilateral.

**Theorem 2.1.** *Let us consider the problem (8) with  $L_1 = \dots = L_N = L \in \mathbb{R}_+^*$ . Assume  $(k_1, \dots, k_N)$  be such that  $k_j = 1$  for every  $1 \leq j \leq D$  and  $k_j = k^-$  with  $k^- < 0$  for every  $D + 1 \leq j \leq N$ . If*

$$|k^-| \neq \frac{D}{N - D},$$

*then for every  $f \in H^{-1}(\mathcal{S})$ , there exists a unique solution  $\psi \in H_0^1(\mathcal{S})$  solving (8).*

*Proof.* The proof is based on the T-coercivity approach [5, 6, 8, 16, 30]. We introduce the bilinear form  $a_k(\cdot, \cdot)$  in  $H_0^1(\mathcal{S}) \times H_0^1(\mathcal{S})$  corresponding to (8) and such that, for every  $\psi, \varphi \in H_0^1(\mathcal{S})$ ,

$$a_k(\psi, \varphi) := \sum_{j=1}^D (\partial_x \psi_j, \partial_x \varphi_j)_{L^2(0,L)} + k^- \sum_{j=D+1}^N (\partial_x \psi_j, \partial_x \varphi_j)_{L^2(0,L)}.$$

Denoted  $\langle \cdot, \cdot \rangle$  the duality product between  $H^{-1}(\mathcal{S})$  and  $H_0^1(\mathcal{S})$ , the result is ensured by proving that, for every  $f \in H^{-1}(\mathcal{S})$ , there exists  $\varphi \in H_0^1(\mathcal{S})$  such that

$$a_k(\varphi, \psi) = \langle f, \psi \rangle, \quad \forall \psi \in H_0^1(\mathcal{S}).$$

**1) Case  $|k^-| < D/(N - D)$ .** Let us introduce the operator

$$T_1 : (\psi_1, \dots, \psi_N) \in H_0^1(\mathcal{S}) \mapsto \left( \psi_1, \dots, \psi_D, -\psi_{D+1} + \frac{2}{D} \sum_{j=1}^D \psi_j, \dots, -\psi_N + \frac{2}{D} \sum_{j=1}^D \psi_j \right) \in H_0^1(\mathcal{S}).$$

Thanks to the young's inequality, for every  $\eta > 0$ ,

$$\begin{aligned} a_k(\psi, T_1 \psi) &= \sum_{l=1}^D \|\partial_x \psi_l\|_{L^2(0,L)}^2 - k^- \sum_{l=D+1}^N \|\partial_x \psi_l\|_{L^2(0,L)}^2 \\ &\quad + \frac{2k^-}{D} \sum_{l=D+1}^N \sum_{j=1}^D (\partial_x \psi_l, \partial_x \psi_j)_{L^2(0,L)} \\ &\geq \sum_{l=1}^D \|\partial_x \psi_l\|_{L^2(0,L)}^2 + |k^-| \sum_{l=D+1}^N \|\partial_x \psi_l\|_{L^2(0,L)}^2 \\ &\quad - \frac{|k^-|}{D} \sum_{l=D+1}^N \sum_{j=1}^D \left( \frac{1}{\eta} \|\partial_x \psi_l\|_{L^2(0,L)}^2 + \eta \|\partial_x \psi_j\|_{L^2(0,L)}^2 \right) \\ &\geq \left( 1 - \frac{|k^-|(N-D)}{D\eta} \right) \sum_{l=1}^D \|\partial_x \psi_l\|_{L^2(0,L)}^2 + |k^-|(1-\eta) \sum_{l=D+1}^N \|\partial_x \psi_l\|_{L^2(0,L)}^2. \end{aligned}$$

Now, for every  $|k^-| < D/(N - D)$ , we set  $\eta = \frac{|k^-|(N-D)}{D(1-\epsilon)}$  with  $\epsilon \in (0, 1 - \frac{|k^-|(N-D)}{D})$  which implies  $\eta \in (0, 1)$  and then

$$1 - \frac{|k^-|(N-D)}{D\eta} = \epsilon > 0, \quad 1 - \eta > 0.$$

Finally,  $T_1$  is a diffeomorphism from  $H_0^1(\mathcal{S})$  to itself such that  $T_1 \circ T_1 = Id$  and the bilinear form  $\tilde{a}_k(\cdot, \cdot) := a_k(\cdot, T_1 \cdot)$  is coercive over  $H_0^1(\mathcal{S}) \times H_0^1(\mathcal{S})$ . Fixed  $f \in H^{-1}(\mathcal{S})$ , we call  $l(\cdot) := \langle f, \cdot \rangle$  and  $l_1(\cdot) := l(T_1 \cdot)$  is a continuous linear form in  $H_0^1(\mathcal{S})$ . The Lax-Milgram's theorem yields the existence of a unique  $\varphi \in H_0^1(\mathcal{S})$  such that  $\tilde{a}_k(\varphi, \psi) = l_1(\psi)$  for every  $\psi \in H_0^1(\mathcal{S})$  continuously depending on the form  $l_1$ . Since  $T_1$  is an isomorphism, there exists a unique  $\varphi \in H_0^1(\mathcal{S})$  such that  $a_k(\varphi, \psi) = l(\psi)$  for every  $\psi \in H_0^1(\mathcal{S})$ . This fact concludes the proof when  $|k^-| < D/(N - D)$ .

**2) Case  $|k^-| > D/(N - D)$ .** Let us introduce the operator

$$T_2 : (\psi_1, \dots, \psi_N) \in H_0^1(\mathcal{S}) \mapsto \left( \psi_1 - \frac{2}{N-D} \sum_{j=D+1}^N \psi_j, \dots, \psi_D - \frac{2}{N-D} \sum_{j=D+1}^N \psi_j, -\psi_{D+1}, \dots, -\psi_N \right) \in H_0^1(\mathcal{S}).$$

Thanks to the young's inequality, for every  $\eta > 0$ ,

$$\begin{aligned} a_k(\psi, T_2\psi) &= \sum_{l=1}^D \|\partial_x \psi_l\|_{L^2(0,L)}^2 + |k^-| \sum_{l=D+1}^N \|\partial_x \psi_l\|_{L^2(0,L)}^2 \\ &\quad - \frac{2}{N-D} \sum_{j=1}^D \sum_{l=D+1}^N (\partial_x \psi_j, \partial_x \psi_l)_{L^2(0,L)} \\ &\geq \sum_{l=1}^D \|\partial_x \psi_l\|_{L^2(0,L)}^2 + |k^-| \sum_{l=D+1}^N \|\partial_x \psi_l\|_{L^2(0,L)}^2 \\ &\quad - \frac{1}{N-D} \sum_{l=D+1}^N \sum_{j=1}^D \left( \eta \|\partial_x \psi_l\|_{L^2(0,L)}^2 + \frac{1}{\eta} \|\partial_x \psi_j\|_{L^2(0,L)}^2 \right) \\ &\geq (1 - \eta) \sum_{l=1}^D \|\partial_x \psi_l\|_{L^2(0,L)}^2 + \left( |k^-| - \frac{D}{(N-D)\eta} \right) \sum_{l=D+1}^N \|\partial_x \psi_l\|_{L^2(0,L)}^2. \end{aligned}$$

Now, for every  $|k^-| > D/(N - D)$ , we set  $\eta = \frac{D}{(N-D)(|k^-| - \epsilon)}$  with  $\epsilon \in (0, |k^-| - \frac{D}{N-D})$  which implies  $\eta \in (0, 1)$  and then

$$|k^-| - \frac{D}{(N-D)\eta} = \epsilon > 0, \quad 1 - \eta > 0.$$

The claim is then proved as in the previous case thanks to the coercivity of the bilinear form  $a_k(\cdot, T_2\cdot)$ .  $\square$

We are finally ready to provide the statement in the general case where the star network is composed by edges of 2 different lengths and materials.

**Corollary 2.2.** *Let us consider the problem (8) with  $L_1 = \dots = L_D = L^+ \in \mathbb{R}_+^*$  for every  $1 \leq j \leq D$  with  $D \in \mathbb{N}^*$  such that  $D < N$  and  $L_{D+1} = \dots = L_N = L^- \in \mathbb{R}_+^*$ . Assume  $(k_1, \dots, k_N)$  be such that  $k_j = k^+ > 0$  for every  $1 \leq j \leq D$  and  $k_j = k^- < 0$  for every  $D + 1 \leq j \leq N$ . If*

$$\frac{|k^-|}{k^+} \neq \frac{D}{L^+} \frac{L^-}{N-D},$$

*then for every  $f \in H^{-1}(\mathcal{S})$ , there exists a unique solution  $\psi \in H_0^1(\mathcal{S})$  solving (8).*

*Proof.* The results follow from Theorem 2.1. First, we use the following change of variable on every edge of the graph

$$h_j : x \in (0, L_j) \mapsto y = \frac{x}{L_j}.$$



We substitute the parameters  $k^+$  with  $\tilde{k}^+ = \frac{k^+}{L^+}$  and  $\tilde{k}^- = \frac{k^-}{L^-}$ . We also replace  $f$  with  $\tilde{f} = (L_1 \tilde{f}_1, \dots, L_N \tilde{f}_N)$ . Second, we divide each equation of (8) with respect to  $\tilde{k}^+$  and we call  $\hat{f} = \frac{\tilde{f}}{\tilde{k}^+}$  and  $\hat{k}^- = \frac{\tilde{k}^-}{\tilde{k}^+}$ . These two transformations allow us to rewrite the problem (8) in an equivalent one defined on a star graph with edges of equal length. The new problem is still of the form of (8) where the parameters  $k_j$  are  $k_j = 1$  for every  $1 \leq j \leq D$  and  $k_j = \hat{k}^-$  with  $k^- < 0$  for every  $D + 1 \leq j \leq N$ , while  $f$  is substitute with  $\hat{f}$ . Finally, the result follows from Theorem 2.1.  $\square$

## 2.2 General multiphase star-graph networks

In the following theorem, we ensure well-posedness in the case of equilateral star graphs with non-constant conductivities. The general case where the edges may have different lengths is treated in the subsequent corollary.

**Theorem 2.3.** *Let us consider the problem (8) with  $L_1 = \dots = L_N = L \in \mathbb{R}$ . Assume  $(k_1, \dots, k_N)$  be such that  $k_j > 0$  for every  $1 \leq j \leq D$  with  $D \in \mathbb{N}^*$  such that  $D < N$  and  $k_j < 0$  for every  $D + 1 \leq j \leq N$ . If one of the following inequalities is satisfied*

$$\left( \max_{D+1 \leq l \leq N} S_l \right) \left( \sum_{l=1}^D \frac{1}{I_l} \right) < \frac{D^2}{N-D}, \quad \left( \max_{l \leq D} S_l \right) \left( \sum_{l=D+1}^N \frac{1}{I_l} \right) < \frac{(N-D)^2}{D},$$

then, for every  $f \in H^{-1}(\mathcal{S})$ , there exists a unique solution  $\psi \in H_0^1(\mathcal{S})$  solving (8).

*Proof.* The statement follows by the T-coercivity developed in Theorem 2.1. We consider again the bilinear form  $a_k(\cdot, \cdot)$  in  $H_0^1(\mathcal{S}) \times H_0^1(\mathcal{S})$  corresponding to (8) and such that, for every  $\psi, \varphi \in H_0^1(\mathcal{S})$ ,

$$a_k(\psi, \varphi) := - \sum_{j=1}^N (\partial_x \psi_j, k_j \partial_x \varphi_j)_{L^2(0,L)}.$$

**1) First case.** Let us consider the operator  $T_1 : (\psi_1, \dots, \psi_N) \in H_0^1(\mathcal{S}) \rightarrow H_0^1(\mathcal{S})$  introduced in the proof of Theorem 2.1. Thanks to the young's inequality, for every  $\eta_{j,l} > 0$  with  $l \in \{D + 1, \dots, N\}$  and  $j \in \{1, \dots, D\}$ ,

$$\begin{aligned} a_k(\psi, T_1 \psi) &= \sum_{l=1}^N (\partial_x \psi_l, |k_l| \partial_x \psi_l)_{L^2(0,L)} - \frac{2}{D} \sum_{l=D+1}^N \sum_{j=1}^D (\partial_x \psi_l, |k_l| \partial_x \psi_j)_{L^2(0,L)} \\ &\geq \sum_{l=1}^N \left\| \sqrt{|k_l|} \partial_x \psi_l \right\|_{L^2(0,L)}^2 \\ &\quad - \frac{1}{D} \sum_{l=D+1}^N \sum_{j=1}^D \left( \eta_{j,l} \left\| \sqrt{|k_l|} \partial_x \psi_l \right\|_{L^2(0,L)}^2 + \frac{1}{\eta_{j,l}} \left\| \sqrt{|k_l|} \partial_x \psi_j \right\|_{L^2(0,L)}^2 \right) \\ &\geq \sum_{l=1}^D \int_0^L \left( |k_l(x)| - \frac{1}{D} \sum_{j=D+1}^N \frac{|k_j(x)|}{\eta_{l,j}} \right) |\partial_x \psi_l(x)|^2 dx \\ &\quad + \sum_{l=D+1}^N \left( 1 - \frac{1}{D} \sum_{j=1}^D \eta_{j,l} \right) \left\| \sqrt{|k_l|} \partial_x \psi_l \right\|_{L^2(0,L)}^2. \end{aligned}$$

Let  $S^+ := \max_{D+1 \leq l \leq N} S_l$ . If  $K_1 := S^+ \sum_{l=1}^D \frac{1}{I_l} < \frac{D^2}{N-D}$ , then we set  $\eta_{j,l} = \frac{S_l(N-D)}{I_j D(1-\epsilon)}$  with  $\epsilon \in (0, 1 - \frac{K_1(N-D)}{D^2})$  and,

$$|k_l(x)| - \frac{1}{D} \sum_{j=D+1}^N \frac{|k_j(x)|}{\eta_{j,l}} \geq |k_l(x)| - I_l + I_l \epsilon \geq I_l \epsilon > 0, \quad \forall x \in (0, L), \quad \forall l \leq D,$$

$$1 - \frac{1}{D} \sum_{j=1}^D \eta_{j,l} \geq 1 - \frac{S^+(N-D)}{D^2} \sum_{j=1}^D \frac{1}{I_j(1-\epsilon)} > 1 - \frac{S^+}{K_1} \sum_{j=1}^D \frac{1}{I_j} = 0, \quad \forall D+1 \leq l \leq N.$$

The last relations yield that there exists  $C > 0$  such that  $a_k(\psi, T_1\psi) \geq C \|\partial_x \psi\|_{L^2}^2$  for every  $\psi \in H_0^1(\mathcal{S})$ . The well-posedness in this case is ensured as in the proof of Theorem 2.1.

**2) Second case.** Let us consider the operator  $T_2 : (\psi_1, \dots, \psi_N) \in H_0^1(\mathcal{S}) \rightarrow H_0^1(\mathcal{S})$  introduced in the proof of Theorem 2.1. Thanks to the Young's inequality, for every  $\eta_{j,l} > 0$  with  $l \in \{D+1, \dots, N\}$  and  $j \in \{1, \dots, D\}$ ,

$$\begin{aligned} a_k(\psi, T_2\psi) &= \sum_{l=1}^N (\partial_x \psi_l, |k_l| \partial_x \psi_l)_{L^2(0,L)} - \frac{2}{N-D} \sum_{l=1}^D \sum_{j=D+1}^N (\partial_x \psi_l, |k_l| \partial_x \psi_j)_{L^2(0,L)} \\ &\geq \sum_{l=1}^N \|\sqrt{|k_l|} \partial_x \psi_l\|_{L^2(0,L)}^2 \\ &\quad - \frac{1}{N-D} \sum_{l=1}^D \sum_{j=D+1}^N \left( \eta_{j,l} \|\sqrt{|k_l|} \partial_x \psi_l\|_{L^2(0,L)}^2 + \frac{1}{\eta_{j,l}} \|\sqrt{|k_l|} \partial_x \psi_j\|_{L^2(0,L)}^2 \right) \\ &\geq \sum_{l=1}^D \left( 1 - \frac{1}{N-D} \sum_{j=D+1}^N \eta_{j,l} \right) \|\sqrt{|k_l|} \partial_x \psi_l\|_{L^2(0,L)}^2 + \\ &\quad \sum_{l=D+1}^N \int_0^L \left( |k_l(x)| - \frac{1}{N-D} \sum_{j=1}^D \frac{|k_j(x)|}{\eta_{j,l}} \right) |\partial_x \psi_l(x)|^2 dx. \end{aligned}$$

Let  $S_+ := \max_{l \leq D} S_l$ . If  $K_2 := S_+ \sum_{l=D+1}^N \frac{1}{I_l} < \frac{(N-D)^2}{D}$ , then we set each  $\eta_{j,l} = \frac{S_j D}{I_l(N-D)(1-\epsilon)}$  with  $\epsilon \in \left(0, 1 - \frac{K_2 D}{(N-D)^2}\right)$  such that

$$|k_l(x)| - \frac{1}{N-D} \sum_{j=1}^D \frac{|k_j(x)|}{\eta_{j,l}} \geq |k_l(x)| - I_l + I_l \epsilon \geq I_l \epsilon > 0, \quad \forall x \in (0, L), \quad \forall D+1 \leq l \leq N,$$

$$1 - \frac{1}{N-D} \sum_{j=D+1}^N \eta_{j,l} \geq 1 - \frac{S_+ D}{(N-D)^2} \sum_{j=D+1}^N \frac{1}{I_j(1-\epsilon)} > 1 - \frac{S_+}{K_2} \sum_{j=D+1}^N \frac{1}{I_j} = 0, \quad \forall l \leq D.$$

The last relations yield that there exists  $C > 0$  such that  $a_k(\psi, T_2\psi) \geq C \|\partial_x \psi\|_{L^2}^2$  for every  $\psi \in H_0^1(\mathcal{S})$ . Finally, the proof is concluded as the one of Theorem 2.1.  $\square$

**Corollary 2.4.** *Let us consider the problem (8). Assume  $(k_1, \dots, k_N) \in \widehat{H}^1(\mathcal{S})$  be such that  $k_j > 0$  for every  $1 \leq j \leq D$  with  $D \in \mathbb{N}^*$  such that  $D < N$  and  $k_j < 0$  for every  $D + 1 \leq j \leq N$ . If one of the following inequalities is satisfied*

$$\left( \max_{D+1 \leq l \leq N} \frac{S_l}{L_l} \right) \left( \sum_{l=1}^D \frac{L_l}{I_l} \right) < \frac{D^2}{N-D}, \quad \left( \max_{l \leq D} \frac{S_l}{L_l} \right) \left( \sum_{l=D+1}^N \frac{L_l}{I_l} \right) < \frac{(N-D)^2}{D},$$

then for every  $f \in H^{-1}(\mathcal{S})$ , there exists a unique solution  $\psi \in H_0^1(\mathcal{S})$  solving (8).

*Proof.* The statement follows from Theorem 2.3 thanks to the arguments adopted in the proof of Corollary 2.2.  $\square$

### 3 Star-networks with Dirichlet and Neumann boundaries

In this section, we continue the study of the well-posedness of a system as (8) in the case we do not only consider Dirichlet boundary conditions on the external vertices but also Neumann-type conditions. In this case, we use a different approach from the T-coercivity adopted in the previous section and assume constant conductivities on each edge.

We denote  $N_d^+$  and  $N_n^+$  the numbers of dielectric edges equipped with Dirichlet and Neumann boundary conditions, respectively. Equivalently,  $N_d^-$  and  $N_n^-$  are the negative index edges respectively equipped with Dirichlet and Neumann boundary conditions. In details, consider  $(k_1, \dots, k_N) \in \mathbb{R}^N$  be such that

$$\begin{cases} k_j > 0, & \text{for } 1 \leq j \leq N_d^+ + N_n^+, \\ k_j < 0, & \text{for } N_d^+ + N_n^+ + 1 \leq j \leq N = N_d^+ + N_n^+ + N_d^- + N_n^-. \end{cases}$$

If we refer to the notation of the previous section  $D = N_d^+ + N_n^+$  and  $N - D = N_d^- + N_n^-$ . We also recall that

$$\widehat{H}^1(\mathcal{S}) := \prod_{j=1}^N H^1(e_j, \mathbb{R}).$$

Assume  $N_d^+ + N_d^- \geq 1$  (we refer to Remark 3.2 for the case  $N_d^+ + N_d^- = 0$ ). For  $f = (f_1, \dots, f_N) \in H^{-1}(\mathcal{S})$ , we study the existence and the unicity of a solution  $\psi = (\psi_1, \dots, \psi_N) \in \widehat{H}^1(\mathcal{S})$  verifying in a weak sense the following system

$$\begin{cases} -k_j \partial_x^2 \psi_j(x) = f_j(x), & x \in (0, L_j), 1 \leq j \leq N, \\ \psi_j(0) = 0, & 1 \leq j \leq N_d^+ \text{ and } N_d^+ + N_n^+ + 1 \leq j \leq N_d^+ + N_n^+ + N_d^-, \\ \partial_x \psi_j(0) = 0, & N_d^+ + 1 \leq j \leq N_d^+ + N_n^+ \text{ and } N_d^+ + N_n^+ + N_d^- + 1 \leq j \leq N, \\ \psi_l(L_l) = \psi_k(L_k), & 1 \leq l, k \leq N, \\ \sum_{l=1}^N k_l \partial_x \psi_l(L_l) = 0. \end{cases} \quad (9)$$

**Theorem 3.1.** *Let us consider the problem (9) with  $L_1 = \dots = L_N = 1$ . Assume  $(k_1, \dots, k_N)$  be such that  $k_j = k^+ > 0$  for  $1 \leq j \leq N_d^+$  and  $k_j = k^- < 0$  with  $k > 0$  for  $N_d^+ + N_n^+ + 1 \leq j \leq N_d^+ + N_n^+ + N_d^-$ . If*

$$\frac{|k^-|}{k^+} \neq \frac{N_d^+}{N_d^-},$$

then for every  $f \in H^{-1}(\mathcal{S})$ , there exists a unique solution  $\psi \in \widehat{H}^1(\mathcal{S})$  solving (9).

**Proof. Unicity of the solutions:** We show that, if  $\psi$  solves (9) with  $f = (0, \dots, 0)$ , then  $\psi = (0, \dots, 0)$ . Indeed, by integrating the equation, we obtain

$$\partial_x \psi_j(x) = \partial_x \psi_j(0), \quad x \in (0, L_j), \quad 1 \leq j \leq N.$$

The boundary conditions yields

$$\begin{aligned} \partial_x \psi_j(x) &= \partial_x \psi_j(0), & x \in (0, L_j), \quad 1 \leq j \leq N_d^+ \quad \text{and} \quad N_d^+ + N_n^+ + 1 \leq j \leq N_d^+ + N_n^+ + N_d^-, \\ \partial_x \psi_j(x) &= 0, & x \in (0, L_j), \quad N_d^+ + 1 \leq j \leq N_d^+ + N_n^+ \quad \text{and} \quad N_d^+ + N_n^+ + N_d^- + 1 \leq j \leq N, \end{aligned} \quad (10)$$

and then

$$\begin{aligned} \partial_x \psi_j(L_j) &= \partial_x \psi_j(0), & 1 \leq j \leq N_d^+ \quad \text{and} \quad N_d^+ + N_n^+ + 1 \leq j \leq N_d^+ + N_n^+ + N_d^-, \\ \partial_x \psi_j(L_j) &= 0, & N_d^+ + 1 \leq j \leq N_d^+ + N_n^+ \quad \text{and} \quad N_d^+ + N_n^+ + N_d^- + 1 \leq j \leq N. \end{aligned} \quad (11)$$

The last relations show that on the edges equipped with boundary Neumann boundary conditions,  $\psi$  can only be a constant function. We integrate again the equations (10) and we obtain

$$\begin{aligned} \psi_j(x) &= x \partial_x \psi_j(0) + \psi_j(0), & x \in (0, L_j), \quad 1 \leq j \leq N_d^+ \quad \text{and} \quad N_d^+ + N_n^+ + 1 \leq j \leq N_d^+ + N_n^+ + N_d^-, \\ \psi_j(x) &= \psi_j(0), & x \in (0, L_j), \quad N_d^+ + 1 \leq j \leq N_d^+ + N_n^+ \quad \text{and} \quad N_d^+ + N_n^+ + N_d^- + 1 \leq j \leq N, \end{aligned}$$

and thanks to the validity of the boundary conditions

$$\begin{aligned} \psi_j(x) &= x \partial_x \psi_j(0), & x \in (0, L_j), \quad 1 \leq j \leq N_d^+ \quad \text{and} \quad N_d^+ + N_n^+ + 1 \leq j \leq N_d^+ + N_n^+ + N_d^-, \\ \psi_j(x) &= \psi_j(0), & x \in (0, L_j), \quad N_d^+ + 1 \leq j \leq N_d^+ + N_n^+ \quad \text{and} \quad N_d^+ + N_n^+ + N_d^- + 1 \leq j \leq N. \end{aligned}$$

From the last relation and (11), we obtain

$$\begin{aligned} \frac{1}{L_j} \psi_j(L_j) &= \partial_x \psi_j(0) = \partial_x \psi_j(L_j), & 1 \leq j \leq N_d^+ \quad \text{and} \quad N_d^+ + N_n^+ + 1 \leq j \leq N_d^+ + N_n^+ + N_d^-, \\ \partial_x \psi_j(L_j) &= 0, & N_d^+ + 1 \leq j \leq N_d^+ + N_n^+ \quad \text{and} \quad N_d^+ + N_n^+ + N_d^- + 1 \leq j \leq N, \end{aligned}$$

We use the last identities in the boundary conditions in the internal vertex appearing in the problem (9) and we obtain the following linear system of equation

$$\begin{cases} \psi_l(L_l) = \psi_k(L_k), & 1 \leq l, k \leq N, \\ \sum_{l=1}^{N_d^+} \frac{k_l}{L_j} \psi_j(L_j) + \sum_{l=N_d^+ + N_n^+ + 1}^{N_d^+ + N_n^+ + N_d^-} \frac{k_l}{L_j} \psi_j(L_j) = 0. \end{cases} \quad (12)$$

We write (12) in a matrix form as  $M \cdot v = 0$  where  $v = (\psi_1(L_1), \dots, \psi_N(L_N))$  and

$$M = \begin{pmatrix} 1 & -1 & 0 & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 1 & -1 & 0 & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & 0 & -1 & 1 \\ \frac{k_1}{L_1} & \dots & \frac{k_{N_d^+}}{L_{N_d^+}} & 0 & \dots & 0 & \frac{k_{N_d^+ + N_n^+ + 1}}{L_{N_d^+ + N_n^+ + 1}} & \dots & \frac{k_{N_d^+ + N_n^+ + 1}}{L_{N_d^+ + N_n^+ + 1}} & 0 & \dots & 0 & 0 \end{pmatrix}. \quad (13)$$

Notice that  $\det(M) = \sum_{l=1}^{N_d^+} \frac{k_l}{L_j} + \sum_{l=N_d^++N_n^++N_d^-} \frac{k_l}{L_j}$  and  $M$  is invertible when

$$\sum_{l=1}^{N_d^+} \frac{k_l}{L_j} + \sum_{l=N_d^++N_n^++N_d^-} \frac{k_l}{L_j} \neq 0.$$

We recall the assumptions on the parameters  $k_j$  and the lengths  $L_j$  so that the last identity becomes

$$N_d^+ k^+ + N_d^- k^- \neq 0.$$

Finally, we have  $\psi = (0, \dots, 0)$  when  $\frac{-k^-}{k^+} \neq \frac{N_d^+}{N_d^-}$ , which implies the unicity of solutions.

**Existence of the solutions:** We solve the problem (9) by integrating the equation twice such as in the previous step. Here, we consider a general source term  $f$  and the boundary conditions verified in the internal vertex of  $\mathcal{S}$  correspond to a linear system that can be written in a matrix form as follows

$$Mv + F = 0. \quad (14)$$

The matrix  $M$  is the same introduced in (13), while  $F = (0, \dots, 0, F_N)$  is defined by

$$\begin{aligned} F_N = & \sum_{j=1}^{N_d^+} \int_0^1 \left( f_j(t) - \int_0^t f_j(s) ds \right) dt + \sum_{j=N_d^++1}^{N_n^+} \int_0^1 f_j(t) dt \\ & + \sum_{j=N_d^++N_n^++1}^{N_d^++N_n^++N_d^-} \int_0^1 \left( f_j(t) - \int_0^t f_j(s) ds \right) dt + \sum_{j=N_d^++N_n^++N_d^-+1}^N \int_0^1 f_j(t) dt. \end{aligned}$$

The equation (14) is solvable when  $M$  is invertible which is guaranteed when  $\frac{-k^-}{k^+} \neq \frac{N_d^+}{N_d^-}$ . □

**Remark 3.2.** Notice that when only Dirichlet boundary conditions are verified, we have  $N_n^+ = N_n^- = 0$  and the last theorem provides the same results of Corollary 2.2 since  $N_d^+ = D$  and  $N_d^- = N - D$ . On the other hand, when the graph is equipped with only Neumann boundary conditions and  $N_d^+ + N_d^- = 0$ , the well-posedness is guaranteed independently of the choice of the parameters  $k_j$ . However, the result is only valid for source terms  $f$  such that  $\int_{\mathcal{S}} f(x) dx = 0$ .

**Remark 3.3.** A simple but still interesting consequence of Theorem 3.1 is the well-posedness in the case of an interval. Let  $a \in (0, 1)$  and  $k_1, k_2 \in \mathbb{R}^*$ . We consider the problem in  $H^1(0, a) \times H^1(a, 1)$

$$\begin{cases} -k_1 \partial_x^2 \psi(x) = f(x), & x \in (0, a), \\ -k_2 \partial_x^2 \psi(x) = f(x), & x \in (a, 1), \end{cases}$$

with  $f \in H^{-1}(0, a) \times H^{-1}(a, 1)$  equipped with the boundary condition at the internal point

$$\psi(a^-) = \psi(a^+), \quad k_1 \partial_x \psi_l(a^-) = k_2 \partial_x \psi(a^+),$$

and at the external points

$$\psi(0) = \partial_x \psi(1) = 0 \quad \text{or} \quad \partial_x \psi(0) = \psi(1) = 0.$$

The existence and the unicity of the solution  $\psi$  of the problem is guaranteed independently of the constants  $k_1$  and  $k_2$  (also with different signs). Notice that the well-posedness is ensured only for  $k_1/k_2 \neq -1$  when we study the same problem with only Dirichlet boundary conditions on the external points. Here, the presence of one Neumann boundary condition yields the result for every  $k$  also negative. The same property is also true in the case only Neumann boundary conditions are verified when we choose a source term  $f$  such that  $\int_0^1 f(x)dx = 0$ .

**Corollary 3.4.** *Let us consider the problem (9). Assume  $(k_1, \dots, k_N)$  be such that*

$$\sum_{l=1}^{N_d^+} \frac{k_l}{L_j} + \sum_{l=N_d^++N_n^+}^{N_d^++N_n^++N_d^-} \frac{k_l}{L_j} \neq 0.$$

For every  $f \in H^{-1}(\mathcal{S})$ , there exists a unique solution  $\psi \in \widehat{H}^1(\mathcal{S})$  solving (9).

*Proof.* The result is a direct consequence of the proof of Theorem 3.1. □

## 4 Tadpole-shaped networks

### 4.1 A simple case: two-phase network

Consider a tadpole graph  $\mathcal{T}$  composed of two edges  $e_1$  and  $e_2$  of lengths  $L_1$  and  $L_2$ , respectively. The self-closing edge  $e_1$ , the “head” of the tadpole, is connected to another edge  $e_2$ , the tail, in the vertex  $v$ . The graph  $\mathcal{T}$  has a symmetry axis denoted by  $r$  which goes through  $e_2$  and crosses  $e_1$  in  $v$  and the center of  $e_1$ . The edge  $e_1$  is parameterized in the clockwise direction with a coordinate going from 0 to  $L_1$  (the length of  $e_1$ ). The edge  $e_2$  is a half-line equipped with a coordinate starting from 0 and arriving at  $L_2$  in  $v$  (see Figure 2 for further details).




Figure 2: The parametrization of the tadpole graph and its symmetry axis  $r$ .

We consider  $\mathcal{T}$  as domain of functions  $f := (f_1, f_2) : \mathcal{T} \rightarrow \mathbb{R}$ , such that  $f_j : e_j \rightarrow \mathbb{R}$  with  $j = 1, 2$ . Let  $L^2(\mathcal{T}) = L^2((0, L_1), \mathbb{R}) \times L^2((0, L_2), \mathbb{R})$  be the Hilbert space equipped with the norm  $\|\cdot\|$  induced by the scalar product

$$(\psi, \varphi)_{L^2} := \int_{e_1} \psi_1(x)\varphi_1(x)dx + \int_{e_2} \psi_2(x)\varphi_2(x)dx, \quad \forall \psi, \varphi \in L^2(\mathcal{T}).$$

Let  $\widehat{H}^1(\mathcal{T}) := H^1(e_1, \mathbb{R}) \times H^1(e_2, \mathbb{R})$ . By duality according to  $L^2(\mathcal{T}, \mathbb{R})$  scalar product, we define the space  $H^{-1}(\mathcal{T})$ . We consider  $k_1, k_2 \in \mathbb{R}^*$  and investigate the well-posedness of the following stationary problem

$$-k_j \partial_x^2 \psi_j(x) = f_j(x), \quad x \in (0, L_j), \quad j \in \{1, 2\}, \quad (15)$$

when the following boundary conditions are verified

$$\begin{aligned}\psi_2(0) &= 0, & \psi_1(0) &= \psi_1(L_1) = \psi_2(L_2), \\ k_1 \partial_x \psi_1(L_1) - k_1 \partial_x \psi_1(0) + k_1 \partial_x \psi_1(L_1) &= 0.\end{aligned}$$

**Theorem 4.1.** *Let us consider the problem (15) with  $f \in H^{-1}(\mathcal{T}, \mathbb{R})$ . Assume  $(k_1, k_2)$  be such that  $k_1$  and  $k_2$  have with different signs. There exists a unique solution  $\psi \in \widehat{H}^1(\mathcal{T})$  solving (15).*

*Proof.* Let us assume  $L_1 = 2$  and  $L_2 = 1$ . However, the same proof is also valid in the general case. We denote by  $R$  the reflection w.r.t. the symmetry axis  $r$  as the operator  $R$  acting on  $L^2(\mathcal{T})$  such that  $Rf = (f_1(1-x), f_2)$  for every  $f = (f_1, f_2) \in L^2(\mathcal{T})$ . The problem (15) is equivalent to study in the same space

$$\begin{cases} -k_1 \partial_x^2 \psi_1^a(x) - k_1(x) \partial_x^2 \psi_1^s(x) = f_1^a(x) + f_1^s(x), & x \in (0, 2), \\ -k_2 \partial_x^2 \psi_2^a(x) - k_2(x) \partial_x^2 \psi_2^s(x) = f_2^a(x) + f_2^s(x), & x \in (0, 1), \end{cases} \quad (16)$$

with

$$\begin{aligned}\psi^a &= (\psi_1^a, \psi_2^a) = \frac{\psi - R\psi}{2}, & \psi^s &= (\psi_1^s, \psi_2^s) = \frac{\psi + R\psi}{2}, \\ f^a &= (f_1^a, f_2^a) = \frac{f - Rf}{2}, & f^s &= (f_1^s, f_2^s) = \frac{f + Rf}{2}.\end{aligned}$$

We notice that  $\psi^a$  and  $f^a$  are antisymmetric w.r.t. the axis  $r$  and they vanish on  $e_2$ , while  $\psi^s$  and  $f^s$  are symmetric. This new formulation yields that studying the problem (15) can be done by considering  $\psi$  and  $f$  antisymmetric at first, and after symmetric.

First, the problem (15) with  $\psi \in \widehat{H}^1(\mathcal{T})$  antisymmetric is equivalent to the following problem  $f \in H^{-1}((0, 1), \mathbb{R})$

$$\begin{cases} -k_1 \partial_x^2 \psi_1(x) = f_1(x), & x \in (0, 1), \\ \psi(0) = \psi(1) = 0. \end{cases}$$

Indeed,  $\psi^a$  and  $f^a$  are equal to zero on  $e_2$  thanks to the definition of  $R$ . The last problem is clearly well-posed independently of the choice of  $k_1$  (and obviously of  $k_2$ ).

Second, we consider the trivial star-graph  $\mathcal{S}$  composed of two connected edges of length 1. The problem (8) with  $\psi \in \widehat{H}^1(\mathcal{T})$  symmetric w.r.t. the axis  $r$  is equivalent to studying the following one in  $H^1(0, 1) \times H^1(0, 1)$

$$\begin{cases} -k_1 \partial_x^2 \psi_1(x) = f_1(x), & x \in (0, 1), \\ -k_2 \partial_x^2 \psi_2(x) = f_2(x), & x \in (0, 1), \end{cases}$$

equipped with the boundary conditions

$$\partial_x \psi_1(0) = \psi_2(0) = 0, \quad \psi_1(L_1) = \psi(L_2), \quad 2k_1 \partial_x \psi_1(L_1) + k_2 \partial_x \psi_2(L_2) = 0.$$

However, if we denote  $\tilde{k}_1 = 2k_1$  and  $\tilde{f}_1 = 2f_1$ , then we obtain (9). We refer to Theorem 3.1 and, as explained in Remark 3.3, the existence and the unicity of solutions hold independently of the choice of  $k_1$  and  $k_2$ .  $\square$

**Remark 4.2.** *The well-posedness of (15) when the Neumann boundary condition is verified on the external vertex is also ensured independently of the choice of the parameters  $k_j$  when we consider the source term  $f$  such that  $\int_{\mathcal{T}} f(x) dx = 0$  thanks to Remark 3.2.*

## 4.2 A more complex case: tree-phase tadpole networks

Let us parametrize a tadpole graph  $\mathcal{T}$  differently from the previous chapter (see Figure 3). We use three edges  $\tilde{e}_1$ ,  $\tilde{e}_2$  and  $\tilde{e}_3$  so that  $|\tilde{e}_1| = \tilde{L}_1$ ,  $|\tilde{e}_2| = \tilde{L}_2$  and  $|\tilde{e}_3| = \tilde{L}_3$ , and we denote it  $\tilde{\mathcal{T}}$ . We consider the “head” of the tadpole ( $e_1$  in the previous subsection) as composed of two edges  $\tilde{e}_1$  and  $\tilde{e}_2$ . The two edges are connected on the left in a vertex  $\tilde{v}$  parametrized with the coordinate 0 while they are both connected on the right with the edge  $\tilde{e}_3$  at the vertex  $v$ . The coordinate 0 of the edge  $\tilde{e}_3$  is located at the external point.




Figure 3: The new parametrization of the tadpole graph.

We consider  $\tilde{\mathcal{T}}$  as domain of functions  $f := (f_1, f_2, f_3) : \tilde{\mathcal{T}} \rightarrow \mathbb{R}$ , such that  $f_j : \tilde{e}_j \rightarrow \mathbb{R}$  with  $j = 1, 2, 3$ . Let  $L^2(\tilde{\mathcal{T}}) = L^2(\tilde{e}_1, \mathbb{R}) \times L^2(\tilde{e}_2, \mathbb{R}) \times L^2(\tilde{e}_3, \mathbb{R})$  be the Hilbert space equipped with the corresponding norm and scalar product. Let  $\hat{H}^1(\tilde{\mathcal{T}}) := H^1(\tilde{e}_1, \mathbb{R}) \times H^1(\tilde{e}_2, \mathbb{R}) \times H^1(\tilde{e}_3, \mathbb{R})$ . By duality, we define the space  $H^{-1}(\tilde{\mathcal{T}})$ . We investigate the well-posedness of

$$\begin{cases} -\tilde{k}_1 \partial_x^2 \psi_1(x) = f_1(x), & x \in (0, \tilde{L}_1), \\ -\tilde{k}_2 \partial_x^2 \psi_2(x) = f_2(x), & x \in (0, \tilde{L}_2), \\ -\tilde{k}_3 \partial_x^2 \psi_3(x) = f_3(x), & x \in (0, \tilde{L}_3). \end{cases} \quad (17)$$

equipped with the boundary conditions

$$\begin{aligned} \psi_3(0) &= 0, \\ \psi_1(0) &= \psi_2(0), \end{aligned} \quad (18)$$

$$\tilde{k}_1 \partial_x \psi_1(0) + \tilde{k}_2 \partial_x \psi_2(0) = 0, \quad (19)$$

$$\psi_1(\tilde{L}_1) = \psi_2(\tilde{L}_2) = \psi_3(\tilde{L}_3), \quad (20)$$

$$\tilde{k}_1 \partial_x \psi_1(\tilde{L}_1) + \tilde{k}_2 \partial_x \psi_2(\tilde{L}_2) + \tilde{k}_3 \partial_x \psi_3(\tilde{L}_3) = 0. \quad (21)$$

**Theorem 4.3.** *Let us consider the problem (17) with  $\tilde{L}_1 = \tilde{L}_2 = \tilde{L}_3 = 1$ . Assume  $(\tilde{k}_1, \tilde{k}_2, \tilde{k}_3) = (k^-, k^+, k^+)$  be such that  $k^+ > 0$  and  $k^- < 0$ . If*

$$\frac{|k^-|}{k^+} \neq 1,$$

*then, for every  $f \in H^{-1}(\tilde{\mathcal{T}}, \mathbb{R})$ , there exists a unique solution  $\psi \in \hat{H}^1(\tilde{\mathcal{T}})$  solving (17).*

*Proof.* The proof follows from the same arguments adopted in the proof of Theorem 3.1. We integrate each component of the equation (17) and we impose the boundary conditions. As in the mentioned



proof, we study the linear system obtained by the boundary conditions on the two internal vertices  $v$  and  $\tilde{v}$ . In this case, we obtain a vector  $v$  such that

$$v = (\psi_1(1), \psi_2(1), \psi_3(1), \psi_1(0), \psi_2(0))$$

and a matrix  $M$ .

$$M = \begin{pmatrix} 1 & -1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 1 & -1 \\ \frac{\tilde{k}_1}{\tilde{L}_1} & \frac{\tilde{k}_2}{\tilde{L}_2} & 0 & -\frac{\tilde{k}_1}{\tilde{L}_1} & -\frac{\tilde{k}_2}{\tilde{L}_2} \\ \frac{\tilde{k}_1}{\tilde{L}_1} & \frac{\tilde{k}_2}{\tilde{L}_2} & \frac{\tilde{k}_3}{\tilde{L}_3} & -\frac{\tilde{k}_1}{\tilde{L}_1} & -\frac{\tilde{k}_2}{\tilde{L}_2} \end{pmatrix}.$$

Notice that the first two lines follow from the boundary conditions (20), the third from (18), the fourth from (19), and the fifth from (21). The invertibility of the matrix  $M$  is guaranteed when

$$\det(M) = \frac{\tilde{k}_1}{\tilde{L}_1} \frac{\tilde{k}_3}{\tilde{L}_3} + \frac{\tilde{k}_2}{\tilde{L}_2} \frac{\tilde{k}_3}{\tilde{L}_3} \neq 0.$$

We use the hypotheses on the conductivities  $\tilde{k}_j$  and the lengths  $\tilde{L}_j$ . The existence and unicity are again implied by the invertibility of  $M$  which is guaranteed when  $\det(M) = -k^+(k^+ + k^-) \neq 0$ .  $\square$

**Corollary 4.4.** *Let us consider the problem (17) If*

$$-\frac{\tilde{k}_1}{\tilde{k}_2} \neq \frac{\tilde{L}_1}{\tilde{L}_2},$$

*then, for every  $f \in H^{-1}(\tilde{\mathcal{T}}, \mathbb{R})$ , there exists a unique solution  $\psi \in \hat{H}^1(\tilde{\mathcal{T}})$  solving (17).*

*Proof.* The result is a direct consequence of the proof Theorem 4.3.  $\square$

**Remark 4.5.** *If we consider (17) and we change the boundary condition at the external vertex with a Neumann one, then the existence and unicity of solutions is ensured independently of the choice of the conductivities when  $f$  is such that  $\int_{\tilde{\mathcal{T}}} f(x) dx = 0$ .*

**Remark 4.6.** *The techniques adopted in the proof of Theorem 3.1 and Theorem 4.3 can be adopted to study the well-posedness of a problem as (3) defined on any network. It is sufficient to integrate the associated equations and rewrite the boundary conditions on the internal vertices as a linear system of the form  $M \cdot v = F$ . The matrix  $M$  keeps track of the structure of the network, and its invertibility infers the well-posedness of the problem (3). Clearly, this task can be really challenging when we study networks composed by several edges and with a complex structure.*

## 5 Spectral representation for star-networks with Dirichlet boundaries

This section aims to provide some spectral properties to the problem (8) defined on a star-graph  $\mathcal{S}$  when  $k_j = 1$  for  $j \leq D$ , while  $k_j = k^- < 0$  for  $D + 1 \leq k \leq N$ . Let us consider the following quadratic forms

$$(\psi, \varphi)_k = \sum_{j=1}^D \int_{e_j} \psi_j \varphi_j + \sum_{j=D+1}^N \int_{e_j} k^- \psi_j \varphi_j.$$

Notice that the quadratic form  $(\cdot, \cdot)_k$  is also a scalar product only when  $k^- > 0$ .

- A natural problem concerning (8) is the following eigenvalue problem: find  $\lambda$  and  $\psi \in H_0^1(\mathcal{S})$  such that, for all  $\varphi \in H_0^1(\mathcal{S})$ ,

$$a_k(\psi, \varphi) := \sum_{j=1}^D \int_{e_j} \psi'_j \varphi'_j + \sum_{j=D+1}^N \int_{e_j} k^- \psi'_j \varphi'_j = \lambda(\psi, \varphi). \quad (22)$$

It corresponds to the study of the eigenfunctions of the following operator in  $L^2(\mathcal{S})$ :

$$A\psi = \left( -\partial_x^2 \psi_1, \dots, -\partial_x^2 \psi_D, -k^- \partial_x^2 \psi_{D+1}, \dots, -k^- \partial_x^2 \psi_N \right) \quad \text{for } \psi \in D(A) \quad \text{with}$$

$$D(A) = \left\{ \psi \in \prod_{j=1}^N H^2(e_j, \mathbb{R}) : \psi_1(0) = \dots = \psi_N(0) = 0, \psi_1(|e_1|) = \dots = \psi_N(|e_N|), \right. \\ \left. \sum_{j=1}^D \partial_x \psi_j(|e_j|) + k^- \sum_{j=D+1}^N \partial_x \psi_j(|e_j|) = 0 \right\}.$$

**Proposition 5.1.** *The operator  $A$  is self-adjoint and admits compact resolvent when*

$$|k^-| \neq \frac{D}{N-D}.$$

*Proof.* The proof of the self-adjointness of the operator  $A$  is very standard. Indeed, the integration by parts in the one-dimensional integrals defining  $\langle \cdot, \cdot \rangle_{L^2}$  leads to the symmetry of the operator at first. Second, it is straightforward to show that  $D(A) = D(A^*)$ . Finally, the fact that  $A$  admits compact resolvent is a direct consequence of Theorem 2.1. □

- On the other hand, it can also be interesting to study the following “alternative” eigenvalue problem: find  $\lambda$  and  $\psi \in H_0^1(\mathcal{S})$  such that, for all  $\varphi \in H_0^1(\mathcal{S})$ ,

$$a_k(\psi, \varphi) = \lambda(\psi, \varphi)_k. \quad (23)$$

This problem can be studied by seeking for an eigenfunction  $\psi$  satisfying  $\tilde{A}\psi = \lambda\psi$ , where the operator  $\tilde{A}$  is

$$\tilde{A}\psi = \left( -\partial_x^2 \psi_1, \dots, -\partial_x^2 \psi_N \right) \quad \text{for } \psi \in D(\tilde{A}) \quad \text{with}$$

$$D(\tilde{A}) = \left\{ \psi \in \prod_{j=1}^N H^2(e_j, \mathbb{R}) : \psi_1(0) = \dots = \psi_N(0) = 0, \psi_1(|e_1|) = \dots = \psi_N(|e_N|), \right. \\ \left. \sum_{j=1}^D \partial_x \psi_j(|e_j|) + k^- \sum_{j=D+1}^N \partial_x \psi_j(|e_j|) = 0 \right\}.$$

**Remark 5.2.** *Notice that  $\tilde{A}$  is neither self-adjoint nor symmetric, on the contrary of  $A$  (Proposition 5.1). Indeed, there exist  $\psi, \varphi \in D(\tilde{A})$ , such that  $\langle \tilde{A}\psi, \varphi \rangle_{L^2} \neq \langle \psi, \tilde{A}\varphi \rangle_{L^2}$ .*

## 5.1 Standard spectral representation for equilateral star-networks

Let us focus on the spectral problem (22) when the length of all the edges is equal to  $L$ . To make things simple, we assume  $L = 1$ . The other cases can be treated in the same manner. To solve the spectral problem, we set  $\xi = is\mu$  with  $s = \sqrt{|k^-|}$ . The eigen-elements  $\psi$  and  $\xi$  equivalently satisfy

$$\psi = (\alpha_1 \sin(\xi x), \dots, \alpha_D \sin(\xi x), \beta_{D+1} \sin(\mu x), \dots, \beta_N \sin(\mu x)),$$

where the coefficients  $\alpha = (\alpha_1, \dots, \alpha_D)$  and  $\beta = (\beta_{D+1}, \dots, \beta_N)$  satisfy

$$\begin{cases} \alpha_j \sin(\xi) = \alpha_{j+1} \sin(\xi), & 1 \leq j \leq D-1, \\ \alpha_D \sin(\xi) = \beta_{D+1} \sin(\mu), \\ \beta_j \sin(\mu) = \beta_{j+1} \sin(\mu), & D+1 \leq j \leq N-1, \\ \sum_{j=1}^D \alpha_j \xi \cos(\xi) + k^- \sum_{j=D+1}^N \beta_j \mu \cos(\mu) = 0. \end{cases} \quad (24)$$

- Notice that when  $\sin(\xi) = 0$ , we find the eigenvalues  $\lambda$  of the form  $n^2\pi^2$  with  $n \in \mathbb{N}^*$ . In this case,  $\beta_j = 0$  with  $D+1 \leq j \leq N$ , thanks to the continuity condition in (24), and we can define the corresponding eigenfunctions of the form

$$(\alpha_1 \sin(n\pi x), \dots, \alpha_D \sin(n\pi x), 0, \dots, 0),$$

where  $\alpha_j$  are some normalizing constants such that

$$\sum_{j=1}^D \alpha_j = 0.$$

We observe that each eigenspace associated to such eigenvalues has dimension  $D-1$  and then the eigenvalues are multiple if  $D \geq 3$ . We denote by  $(\nu_j)_{j \in \mathbb{N}^*}$  such eigenvalues counted with their multiplicity, and we denote by  $(\varphi^j)_{j \in \mathbb{N}^*}$  some corresponding normalized eigenfunctions.

- Similarly to the previous case, when  $\sin(\mu) = 0$ , we find the eigenvalues  $\lambda$  of the form  $k^- n^2\pi^2$  with  $n \in \mathbb{N}^*$ . Here,  $\alpha_j = 0$  with  $1 \leq j \leq D$  and the corresponding eigenfunctions have the form

$$(0, \dots, 0, \beta_{D+1} \sin(n\pi x), \dots, \beta_N \sin(n\pi x)),$$

where  $\beta_j$  are normalizing constants such that

$$\sum_{j=D+1}^N \beta_j = 0.$$

Each eigenspace has dimension  $N-D-1$  and the eigenvalues are multiple if  $N-D \geq 3$ . We denote by  $(\theta_j)_{j \in \mathbb{N}^*}$  such eigenvalues counted with their multiplicity, and we denote by  $(\rho^j)_{j \in \mathbb{N}^*}$  some corresponding normalized eigenfunctions.

- When  $\sin(\xi), \sin(\mu) \neq 0$ , the conditions (24) can be rewritten as

$$\begin{pmatrix} 1 & -1 & 0 & \dots & \dots & \dots & \dots & \dots \\ 0 & 1 & -1 & 0 & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & 0 & \sin(\xi) & -\sin(\mu) & 0 & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & 0 & 1 & -1 \\ \xi \cos(\xi) & \dots & \xi \cos(\xi) & \xi \cos(\xi) & k^- \mu \cos(\mu) & k^- \mu \cos(\mu) & \dots & k^- \mu \cos(\mu) \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \end{pmatrix} = 0$$

and its determinant yields the dispersion relation

$$D\xi \cos(\xi) \sin(\mu) + (N - D)k^- \mu \sin(\xi) \cos(\mu) = 0.$$

We recall that  $\xi = is\mu$  and then

$$Ds \cosh(s\mu) \sin(\mu) + (N - D)k^- \sinh(s\mu) \cos(\mu) = 0.$$

Now, we notice that  $s = \sqrt{|k^-|}$  which implies  $s = -\frac{k^-}{s}$  and, with  $x = \frac{D}{s(N-D)}$ ,

$$\cos(\mu) \sinh(s\mu) - x \sin(\mu) \cosh(s\mu) = 0. \quad (25)$$

Assuming that  $\mu = a + ib$ , with  $a, b \in \mathbb{R}$ , this expression can be rewritten in the form

$$0 = [\cos(a) \cosh(b) - i \sin(a) \sinh(b)] [\sinh(sa) \cos(sb) + i \cosh(sa) \sin(sb)] - x [\sin(a) \cosh(b) + i \cos(a) \sinh(b)] [\cosh(sa) \cos(sb) + i \sinh(sa) \sin(sb)].$$

Taking the real and imaginary part of this relation yields two equations of the form

$$A \begin{pmatrix} x \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix},$$

where  $A$  is a  $2 \times 2$  matrix composed by the elements

$$\begin{aligned} A_{1,1} &= \cos(a) \sinh(b) \sinh(as) \sin(bs) - \sin(a) \cosh(b) \cosh(as) \cos(bs), \\ A_{1,2} &= \cos(a) \cosh(b) \sinh(as) \cos(bs) + \sin(a) \sinh(b) \cosh(as) \sin(bs), \\ A_{2,1} &= -\cos(a) \sinh(b) \cosh(as) \cos(bs) - \sin(a) \cosh(b) \sinh(as) \sin(bs), \\ A_{2,2} &= \cos(a) \cosh(b) \cosh(as) \sin(bs) - \sin(a) \sinh(b) \sinh(as) \cos(bs) \end{aligned}$$

and its determinant simplifies to

$$\begin{aligned} \det(A) &= \cosh(sa) \sinh(sa) \cosh(b) \sinh(b) - \sin(a) \cos(a) \sin(sb) \cos(sb) \\ &= \frac{1}{4} (\sinh(2sa) \sinh(2b) - \sin(2a) \sin(2sb)). \end{aligned}$$

Since  $\sin(x) < x < \sinh(x)$  for any  $x > 0$ , if  $ab > 0$  we see that  $\det(A) > 0$  and the above system cannot have a non-trivial solution. The same conclusion holds if  $ab < 0$  since  $\sin$  and

$\sinh$  are odd functions. We conclude that the dispersion relation (25) may have solutions only when  $\mu \in \mathbb{R}$  and when  $\mu \in i\mathbb{R}$ , *i.e.*

$$\begin{cases} \mu = a \in \mathbb{R} & \text{with} & \tan(a) - \frac{s(N-D)}{D} \tanh(sa) = 0, \\ \mu = ib/s \in i\mathbb{R} & \text{with} & \tan(b) - \frac{D}{s(N-D)} \tanh(b/s) = 0. \end{cases}$$

Recalling that  $s = \sqrt{|k^-|}$ , there is exactly one solution  $a_n \in ]n\pi, (n+1)\pi[$  to

$$\tan(a_n) = \frac{s(N-D)}{D} \tanh(sa_n)$$

and exactly one solution  $b_n \in ]n\pi, (n+1)\pi[$  to

$$\tan(b_n) = \frac{D}{s(N-D)} \tanh(b_n/s),$$

for  $n \geq 1$ . It follows that we can construct two families of eigenfunctions. The first corresponds to choosing  $\mu_n = a_n, n \geq 1$  and

$$\phi^{2n}(x) = c_{2n}(\phi_1^{2n}, \dots, \phi_N^{2n}),$$

$$\phi_j^{2n} = \begin{cases} \frac{\sin(a_n)}{\sinh(sa_n)} \sinh(sa_n x), & 1 \leq j \leq D, \\ \sin(a_n x), & D+1 \leq j \leq N, \end{cases}$$

where  $c_{2n}$  is a normalization factor. The second family corresponds to setting  $\mu_n = ib_n/s$  or equivalently  $\xi_n = b_n$  and the associated eigenfunctions are

$$\phi^{2n+1}(x) = c_{2n+1}(\phi_1^{2n+1}, \dots, \phi_N^{2n+1}),$$

$$\phi_j^{2n+1}(x) = \begin{cases} \sin(b_n x), & 1 \leq j \leq D, \\ \frac{\sin(b_n)}{\sinh\left(\frac{b_n}{s} x\right)} \sinh\left(\frac{b_n}{s} x\right), & D+1 \leq j \leq N, \end{cases}$$

with a normalization factor  $c_{2n+1}$ .

Finally, the eigenvalues  $(\lambda_n)_{n \in \mathbb{N}^*}$  of  $A$  are obtained by reordering the sequences

$$(\nu_n)_{n \in \mathbb{N}^*}, \quad (\theta_n)_{n \in \mathbb{N}^*}, \quad (k^- a_n^2)_{n \in \mathbb{N}^*}, \quad (b_n^2)_{n \in \mathbb{N}^*}.$$

We denote the corresponding eigenfunctions  $(\psi^n)_{n \in \mathbb{N}^*}$  obtained by reordering

$$(\varphi^n)_{n \in \mathbb{N}^*}, \quad (\rho^n)_{n \in \mathbb{N}^*}, \quad (\phi^n)_{n \in \mathbb{N}^*}.$$

**Proposition 5.3.** *The functions  $(\psi^n)_{n \geq 1}$  form a complete orthonormal system of  $L^2(\mathcal{S})$ .*

*Proof.* The result is a direct consequence of Theorem 5.1.  $\square$

**Remark 5.4.** Let us consider the following Schrödinger type equation in  $L^2(\mathcal{S}, \mathbb{C})$

$$\begin{cases} i\partial_t \varphi = A\varphi, & t > 0, \\ \varphi(t=0) = f \in L^2(\mathcal{S}). \end{cases} \quad (26)$$

If we extend our analysis leading to Proposition 5.3 for complex-valued function, then we can ensure the well-posedness of the problem (26) and express its solutions  $\varphi(t)$  for any initial data  $f \in L^2(\mathcal{S}, \mathbb{C})$  as follows:

$$S(t)f := \varphi(t) = \sum_{j=1}^{\infty} e^{-i\lambda_j t} \psi^j(\psi^j, f)_{L^2}.$$

We observe that  $(S(t))_{t \in \mathbb{R}}$  is a unitary group of  $L^2(\mathcal{S})$  generates by  $-iA$  and, for all  $f \in D(A)$ , the system (26) admits a unique solution

$$\varphi \in C([0, +\infty); D(A)) \cap C^1([0, +\infty); L^2(\mathcal{S}, \mathbb{C})), \quad \|\varphi(t)\|_{L^2} = \|f\|_{L^2}, \forall t \geq 0.$$

**Remark 5.5.** Now, consider the following heat-type equation

$$\begin{cases} \partial_t u + \tilde{A}u = 0, & t > 0, \\ u(t=0) = f \in L^2(\mathcal{S}, \mathbb{R}). \end{cases} \quad (27)$$

Due to the presence of negative eigenvalues, we can not define solutions of the dynamics (27) in the whole space  $L^2(\mathcal{S}, \mathbb{R})$ . Nevertheless, we can introduce the space

$$X = \{u \in L^2(\mathcal{S}, \mathbb{R}) : \exists N \in \mathbb{N} : \langle u, \phi^{2j} \rangle = \langle u, \rho^j \rangle = 0, \forall j > N\}$$

which is dense in  $L^2(\mathcal{S}, \mathbb{R})$ . The dynamics of (27) is well-defined in  $X$  and the solutions  $u(t)$ , for any initial data  $f \in X$ , are decomposed as follows with suitable  $N_f \in \mathbb{N}$ :

$$\begin{aligned} u(t) &= \sum_{j=1}^{\infty} e^{-\nu_j t} \varphi^j(\varphi^j, f)_{L^2} + \sum_{j=1}^{N_f} e^{-\theta_j t} \rho^j(\rho^j, f)_{L^2} \\ &\quad + \sum_{j=1}^{N_f} e^{k^- a_j^2 t} \phi^{2j}(\phi^{2j}, f)_{L^2} + \sum_{j=1}^{\infty} e^{-b_j^2 t} \phi^{2j+1}(\phi^{2j+1}, f)_{L^2}. \end{aligned}$$

## 5.2 Standard spectral representation for some non-equilateral star-networks

Let us consider the spectral problem (22) when the lengths  $L_j = |e_j|$  of the edges of the star-graph are such that  $L_j/L_k \notin \mathbb{Q}$  for every distinct  $j, k \leq N$ . As in the previous section, we study the eigenfunctions of the operator  $A$  and we set  $\xi = is\mu$  with  $s = \sqrt{|k^-|}$ . The eigen-elements  $\psi$  and  $\xi$  equivalently satisfy

$$\psi = (\alpha_1 \sin(\xi x), \dots, \alpha_D \sin(\xi x), \beta_{D+1} \sin(\mu x), \dots, \beta_N \sin(\mu x)),$$

where the coefficients  $\alpha = (\alpha_1, \dots, \alpha_D)$  and  $\beta = (\beta_{D+1}, \dots, \beta_N)$  satisfy

$$\begin{cases} \alpha_j \sin(\xi L_j) = \alpha_{j+1} \sin(\xi L_{j+1}), & 1 \leq j \leq D-1, \\ \alpha_D \sin(\xi L_D) = \beta_{D+1} \sin(\mu L_{D+1}), \\ \beta_j \sin(\mu L_j) = \beta_{j+1} \sin(\mu L_{j+1}), & D+1 \leq j \leq N-1, \\ \sum_{j=1}^D \alpha_j \xi \cos(\xi L_j) + k^- \sum_{j=D+1}^N \beta_j \mu \cos(\mu L_j) = 0. \end{cases} \quad (28)$$

Notice that, in contrast to the previous section, we have that  $\sin(\xi L_j), \sin(\mu L_j) \neq 0$  for every  $1 \leq j \leq D$  and  $D+1 \leq l \leq N$ . Indeed, if for instance  $\sin(\xi L_j) = 0$  for  $1 \leq j \leq D$ , then  $\beta_l = 0$  for all  $D+1 \leq l \leq N$  and there exists  $1 \leq m \leq D$  distinct from  $j$  such that

$$\sin(\xi L_j) = \sin(\xi L_m) = 0.$$

This identity implies that  $\xi$  is of the form

$$\frac{n_j \pi}{L_j} \quad \text{and} \quad \frac{n_m \pi}{L_m} \quad \text{with} \quad n_j, n_m \in \mathbb{Z}^*,$$

which is impossible since  $L_j/L_k \notin \mathbb{Q}$  for every distinct  $j, k \leq N$ . This argument leads to

$$\sin(\xi L_j), \sin(\mu L_l) \neq 0, \quad 1 \leq j \leq D, \quad D+1 \leq l \leq N,$$

and it also implies that the eigenfunctions of  $A$  can not have any vanishing components thanks to the continuity condition in (28). Now, the determinant corresponding to the linear system (28) yields

$$\sum_{j=1}^D \xi \cot(\xi L_j) + k^- \sum_{j=D+1}^N \mu \cot(\mu L_j) = 0. \quad (29)$$

We can observe that the eigenvalues are simple. Indeed, if  $\lambda$  is a multiple eigenvalue, then there exist at least two distinct eigenfunctions  $f$  and  $g$  both corresponding to  $\lambda$ . Such eigenfunctions do not have any vanishing component and their first components can be respectively written as

$$f_1 = \alpha_1^f \sin(\sqrt{\lambda} L_1), \quad g_1 = \alpha_1^g \sin(\sqrt{\lambda} L_1).$$

Now,  $h = \alpha_1^g f - \alpha_1^f g$  is another eigenfunction of  $A$  corresponding to the eigenvalue  $\lambda$  and its first component  $h_1 = \alpha_1^g \alpha_1^f \sin(\sqrt{\lambda} L_1) - \alpha_1^f \alpha_1^g \sin(\sqrt{\lambda} L_1) = 0$  which is absurd.

Finally, the eigenvalues  $(\lambda_n)_{n \in \mathbb{N}^*}$  of  $A$  are simple, and they obtained by solving the transcendental equation (29). In other words, since  $\mu = -i \frac{\xi}{s}$  with  $s = \sqrt{|k^-|}$ , they are the solutions of

$$\sum_{j=1}^D \cot(\sqrt{\lambda_n} L_j) + \sqrt{|k^-|} \sum_{j=D+1}^N \coth\left(\frac{\sqrt{\lambda_n}}{\sqrt{|k^-|}} L_j\right) = 0.$$

Some corresponding eigenfunctions  $(\psi^n)_{n \in \mathbb{N}^*}$  have the form

$$\begin{aligned} \psi^n = \alpha_n \left( \sin(\sqrt{\lambda_n} x), \frac{\sin(\sqrt{\lambda_n} L_1)}{\sin(\sqrt{\lambda_n} L_2)} \sin(\sqrt{\lambda_n} x), \dots, \frac{\sin(\sqrt{\lambda_n} L_1)}{\sin(\sqrt{\lambda_n} L_D)} \sin(\lambda_n x), \right. \\ \left. \frac{\sin(\sqrt{\lambda_n} L_1)}{\sinh\left(\frac{\sqrt{\lambda_n}}{\sqrt{|k^-|}} L_{D+1}\right)} \sinh\left(\frac{\sqrt{\lambda_n}}{\sqrt{|k^-|}} x\right), \dots, \frac{\sin(\sqrt{\lambda_n} L_1)}{\sinh\left(\frac{\sqrt{\lambda_n}}{\sqrt{|k^-|}} L_N\right)} \sinh\left(\frac{\sqrt{\lambda_n}}{\sqrt{|k^-|}} x\right) \right), \end{aligned}$$

with  $\alpha_n$  some normalization constants. As in the previous section, the operator  $A$  is self-adjoint with compact resolvent which implies the following result.

**Proposition 5.6.** *The functions  $(\psi^n)_{n \geq 1}$  form a complete orthonormal system of  $L^2(\mathcal{S})$ .*

### 5.3 Spectral representation via pseudo-eigenfunctions for equilateral star-networks

In the current section, we want to investigate the “alternative” spectral problem (23) when the edges of the star-graph have the same length. We assume for simplicity that  $|e_j| = 1$  for every  $1 \leq j \leq N$ . Notice that an eigenfunction of  $\tilde{A}$  has the form

$$\psi = (\psi_1, \dots, \psi_N), \quad \psi_j(x) = \alpha_j \sin(\sqrt{\lambda}x), \quad 1 \leq j \leq N, \quad (30)$$

where the coefficients  $\alpha_j$  satisfy

$$\begin{aligned} \alpha_j \sin(\sqrt{\lambda}) &= \alpha_N \sin(\sqrt{\lambda}), \quad 1 \leq j \leq N, \\ \sum_{j=1}^D \alpha_j \cos(\sqrt{\lambda}) + k^- \sum_{j=D+1}^N \alpha_j \cos(\sqrt{\lambda}) &= 0. \end{aligned}$$

One easily checks that the determinant of this linear system is equal to

$$\sin(\sqrt{\lambda})^{N-1} \cos(\sqrt{\lambda}) (-D - k^-(N-D))$$

and we recover the fact that the bilinear form  $a_k$  is an isomorphism under the condition appearing in Theorem 2.1 which is

$$-k^- \neq \frac{D}{N-D}.$$

We find two families of eigenpairs. The first one  $(\nu_n, \phi^n)_{n \in \mathbb{N}^*}$  is given by

$$\nu_n = \frac{(2n-1)^2 \pi^2}{4} \quad \text{and} \quad \phi^n = (\phi_1^n, \dots, \phi_N^n), \quad \phi_j^n = \sin\left(\frac{(2n-1)\pi}{2}x\right). \quad (31)$$

Notice that  $\cos(\sqrt{\nu_n}) = 0$  and the derivatives of  $\phi^n$  vanish at the endpoint  $x = 1$  of each edge. The second family of eigenpairs corresponds to eigenvalues of multiplicity  $N-1$ :

$$\gamma_n = n^2 \pi^2, \quad n \in \mathbb{N}^*.$$

The associated eigenspace to each  $\gamma_n$  corresponds to eigenfunctions of the form

$$\varphi = (\alpha_1 \sin(n\pi x), \dots, \alpha_N \sin(n\pi x)) \quad \text{with} \quad \sum_{j=1}^D \alpha_j + k^- \sum_{j=D+1}^N \alpha_j = 0.$$

Thus, we can construct the associated eigenfunctions as follows

$$\varphi = (k^- \tilde{\alpha}_1 \sin(n\pi x), \dots, k^- \tilde{\alpha}_D \sin(n\pi x), \tilde{\alpha}_{D+1} \sin(n\pi x), \dots, \tilde{\alpha}_N \sin(n\pi x)), \quad \text{with} \quad \sum_{j=1}^N \tilde{\alpha}_j = 0.$$



An example of basis for an eigenspace when  $D = 1$  and  $N = 3$  is the one formed by the two orthogonal functions

$$(0, -\sin(n\pi x), \sin(n\pi x)), \quad (-2k^- \sin(n\pi x), \sin(n\pi x), \sin(n\pi x)).$$

Finally, we call  $(\lambda_n)_{n \in \mathbb{N}^*}$  the sequence of eigenvalues of  $\tilde{A}$  obtained by reordering the sequences  $(\nu_n)_{n \in \mathbb{N}^*}$  and  $(\gamma_n)_{n \in \mathbb{N}^*}$  and then

$$\lambda_n = \frac{n^2 \pi^2}{4}.$$

Each eigenvalue  $\lambda_n$  with  $n \in 2\mathbb{N}^*$  has multiplicity  $N - 1$ , while the others are simple. We denote by  $(\psi^n)_{n \in \mathbb{N}^*}$  a sequence of orthonormalized eigenfunctions corresponding to  $(\lambda_n)_{n \in \mathbb{N}^*}$ .

**Proposition 5.7.** *The family of functions  $(\psi^n)_{n \in \mathbb{N}^*}$  forms a Riesz Basis of  $L^2(\mathcal{S})$ , i.e. it is isomorphic to an orthonormal system of  $L^2(\mathcal{S})$ .*

*Proof.* First, notice that the two families of eigenfunctions respectively corresponding to the eigenvalues  $(\nu_n)_{n \in \mathbb{N}^*}$  and  $(\gamma_n)_{n \in \mathbb{N}^*}$  are mutually orthogonal since

$$\left( \sin(n\pi x), \sin\left(\frac{2l-1}{2}\pi x\right) \right)_{L^2} = 0, \quad \forall n, l \in \mathbb{N}^*.$$

We denote by  $\mathcal{H}_1$  and  $\mathcal{H}_2$  the corresponding eigenspaces. Second, we introduce the linear mapping  $T : \mathcal{H}_1 + \mathcal{H}_2 \rightarrow L^2(\mathcal{S})$  such that  $T|_{\mathcal{H}_1} = Id_{\mathcal{H}_1}$  and  $T|_{\mathcal{H}_2}$  is defined as follows

$$T\psi = \left( \frac{1}{k^-} \psi_1, \dots, \frac{1}{k^-} \psi_D, \psi_{D+1}, \dots, \psi_N \right), \quad \psi \in \mathcal{H}_2.$$

We notice that  $T$  is an isomorphism from  $\mathcal{H}_1 + \mathcal{H}_2$  to  $T(\mathcal{H}_1 + \mathcal{H}_2)$ . Now,  $(\phi^n)_{n \in \mathbb{N}^*}$  with  $\phi^j = S(T\psi^j)$  are eigenfunctions of a standard Dirichlet Laplacian defined on a star-graph with edges of equal length (see for instance [15, Section 4.1]) which is self-adjoint with compact resolvent. Finally,  $(\phi^n)_{n \in \mathbb{N}^*}$  define a complete orthonormal system,  $\mathcal{H}_1 + \mathcal{H}_2 = L^2(\mathcal{S}, \mathbb{R})$  and  $(\psi^n)_{n \in \mathbb{N}^*}$  forms a Riesz Basis in  $L^2(\mathcal{S})$ .  $\square$

**Remark 5.8.** *Similarly to Remark 5.4, we can exploit Proposition 5.7 to characterize some evolution equations defined with the operator  $\tilde{A}$ . Consider for instance the following Schrödinger-type equation in  $L^2(\mathcal{S}, \mathbb{C})$*

$$\begin{cases} i\partial_t v = \tilde{A}v, & t > 0, \\ v(t=0) = f \in L^2(\mathcal{S}). \end{cases} \quad (32)$$

*Notice that our analysis leading to Proposition 5.7 is still valid for complex valued functions. First, we introduce the biorthogonal family in  $L^2(\mathcal{S}, \mathbb{C})$  of the Riesz basis  $(\psi^n)_{n \in \mathbb{N}^*}$  : this is the unique sequence of functions  $(\sigma^n)_{n \in \mathbb{N}^*}$  such that*

$$(\psi^n, \sigma^k)_{L^2} = \delta_{k,n}, \quad \forall k \in \mathbb{N}^*.$$

*The functions  $\sigma^n$  can be explicitly computed by using the orthogonal projectors  $\pi_k : L^2 \rightarrow \overline{\text{span}\{\psi^j : j \neq k\}}^{L^2}$  as follows:*

$$\sigma^k = \frac{\psi^k - \pi_k \psi^k}{\|\psi^k - \pi_k \psi^k\|_{L^2}^2}, \quad \forall k \in \mathbb{N}^*.$$

Second, we ensure the well-posedness of the evolution equation (32) by using the properties of the Riesz basis and its solutions  $v(t)$  for any initial data  $f \in L^2(\mathcal{S}, \mathbb{C})$  are of the form:

$$T(t)f := v(t) = \sum_{j=1}^{\infty} e^{-i\lambda_j t} \psi^j(\sigma^j, f)_{L^2}.$$

We have that  $(T(t))_{t \in \mathbb{R}}$  is a group of  $L^2(\mathcal{S}, \mathbb{C})$  generated by  $-i\tilde{A}$  and, for all  $f \in D(\tilde{A})$ , the system 32 admits a unique solution

$$v \in C([0, +\infty); D(\tilde{A})) \cap C^1([0, +\infty); L^2(\mathcal{S}, \mathbb{C})), \quad \|v(t)\|_{L^2} = \|f\|_{L^2}, \forall t \geq 0.$$

**Remark 5.9.** Consider now the following heat-type equation in  $L^2(\mathcal{S}, \mathbb{R})$

$$\begin{cases} \partial_t u + \tilde{A}u = 0, & t > 0, \\ u(t=0) = f \in L^2(\mathcal{S}, \mathbb{R}). \end{cases} \quad (33)$$

Equivalently to the complex case (Remark 5.8), we can define a family of eigenfunctions  $(\psi^n)_{n \in \mathbb{N}^*}$  of  $\tilde{A}$  forming a Riesz basis of  $L^2(\mathcal{S}, \mathbb{R})$  and we denote by  $(\sigma^n)_{n \in \mathbb{N}^*}$  its biorthogonal family. Now, the solutions  $u(t)$  of the problem (33) for any initial data  $f \in L^2(\mathcal{S}, \mathbb{R})$  are:

$$R(t)f := u(t) = \sum_{j=1}^{\infty} e^{-\lambda_j t} \psi^j(\sigma^j, f)_{L^2}.$$

This solution is a semigroup solution and, when  $f \in D(\tilde{A})$ ,

$$u \in C([0, +\infty); D(\tilde{A})) \cap C^1([0, +\infty); L^2(\mathcal{S})).$$

Notice that, on the contrary for the operator  $A$  considered in Remark 5.5, the eigenvalues of  $\tilde{A}$  are positive, which ensures the fact the solutions are well-defined.

#### 5.4 Spectral representation via pseudo-eigenfunctions for some non-equilateral star-networks

This section aims to study (23) when the lengths  $L_j = |e_j|$  of the edges of the star-graph are such that  $L_j/L_k \notin \mathbb{Q}$  for every distinct  $j, k \leq N$ . As in the previous section, we study the eigenfunctions of the operator  $\tilde{A}$  which have the form

$$\psi = (\psi_1, \dots, \psi_N), \quad \psi_j(x) = \alpha_j \sin(\sqrt{\lambda}x), \quad 1 \leq j \leq N, \quad (34)$$

where the coefficients  $\alpha_j$  satisfy

$$\begin{aligned} \alpha_j \sin(\sqrt{\lambda}L_j) &= \alpha_N \sin(\sqrt{\lambda}L_N), \quad 1 \leq j \leq N, \\ \sum_{j=1}^D \alpha_j \cos(\sqrt{\lambda}L_j) + k^- \sum_{j=D+1}^N \alpha_j \cos(\sqrt{\lambda}L_j) &= 0. \end{aligned}$$

On the contrary of the previous section, we can observe that the coefficients

$$\alpha_j \neq 0, \quad \forall 1 \leq j \leq N.$$

Indeed, if one of these coefficients vanishes, then there exist distinct  $l, m \leq N$  such that  $\alpha_l, \alpha_m \neq 0$ , and

$$\sin(\sqrt{\lambda}L_l) = \sin(\sqrt{\lambda}L_m) = 0.$$

This identity implies that  $\lambda$  is of the form

$$\frac{n_l^2 \pi^2}{L_l^2} \quad \text{and} \quad \frac{n_m^2 \pi^2}{L_m^2} \quad \text{with} \quad n_l, n_m \in \mathbb{Z}^*,$$

which is impossible since  $L_j/L_k \notin \mathbb{Q}$  for every distinct  $j, k \leq N$ . This argument not only yields that all the components of  $\psi$  are different from 0, but also that

$$\sin(\sqrt{\lambda}L_l) \neq 0, \quad \forall 1 \leq l \leq N,$$

and it can also be used to prove that

$$\cos(\sqrt{\lambda}L_l) \neq 0, \quad \forall 1 \leq l \leq N.$$

In addition, we have that each eigenvalue  $\lambda$  is simple since the existence of multiple eigenvalues would imply the existence of eigenfunctions vanishing in at least one edge which is impossible (as in Section 5.2).

Finally, we can define the eigenpairs  $(\lambda_n, \psi^n)_{n \in \mathbb{N}^*}$  where  $(\lambda_n)_{n \in \mathbb{N}^*}$  is a sequence of simple eigenvalues defined by the relation

$$\sum_{j=1}^D \cot(\sqrt{\lambda_n}L_j) + k^- \sum_{j=D+1}^N \cot(\sqrt{\lambda_n}L_j) = 0,$$

while  $(\psi^n)_{n \in \mathbb{N}^*}$  is a sequence of eigenfunctions such that

$$\psi^n = \alpha_n \left( \frac{\sin(\sqrt{\lambda_n}L_N)}{\sin(\sqrt{\lambda_n}L_1)} \sin(\lambda_n x), \dots, \frac{\sin(\sqrt{\lambda_n}L_N)}{\sin(\sqrt{\lambda_n}L_{N-1})} \sin(\sqrt{\lambda_n}x), \sin(\sqrt{\lambda_n}x) \right),$$

with  $\alpha_n$  some normalizing constants.

**Proposition 5.10.** *The family of functions  $(\psi^n)_{n \in \mathbb{N}^*}$  forms a Riesz Basis of  $L^2(\mathcal{S})$ , i.e. it is isomorphic to an orthonormal system of  $L^2(\mathcal{S})$ .*

*Proof.* The result is ensured equivalently to Proposition 5.7. Indeed, the functions  $\phi_j = T\psi_j$  are the eigenfunctions of the standard Dirichlet Laplacian defined on star graphs with edges of different lengths which is self-adjoint with compact resolvent (see again [15, Section 4.1]).  $\square$

## 6 Spectral representation for star-networks with mixed boundaries

This section aims to retrace the theory of Section 5 when some external vertices of the star-graph are equipped with Neumann boundary conditions. In detail, we consider the two spectral problems (22) and (23) and we study them with functions in  $\widehat{H}^1(\mathcal{S})$  when the boundary conditions in (9) are verified. To simplify the theory, we assume  $N_d^+ = N_n^+ = N_d^- = N_n^- = 2$ , but the same approach can be used in the general case.

## 6.1 Standard spectral representation for equilateral star-networks

Let us consider the spectral problem (22) when the length of all the edges is equal to  $L = 1$ . The same results can be directly extended to the case where all the edges have the same length.

- We find the eigenvalues  $\lambda$  of the form

$$n^2\pi^2, \quad \forall n \in \mathbb{N}^*$$

and corresponding to the eigenfunctions of the form

$$\psi = (\sin(n\pi x), -\sin(n\pi x), 0, 0, 0, 0, 0, 0).$$

Notice that such eigenvalues are simple due to the choice of  $N_d^+ = 2$ . In the general case, the same type of construction can be repeated and the multiplicity is  $N_d^+ - 1$ .

- We find the eigenvalues  $\lambda$  of the form

$$\frac{(2n-1)^2\pi^2}{4}, \quad \forall n \in \mathbb{N}^*$$

and corresponding to the eigenfunctions of the form

$$\psi = \left(0, 0, \cos\left(\frac{(2n-1)\pi}{2}x\right), -\cos\left(\frac{(2n-1)\pi}{2}x\right), 0, 0, 0, 0\right).$$

These eigenvalues are simple and, in the general case, their multiplicity is  $N_d^+ - 1$ .

- We find the eigenvalues  $\lambda$  of the form

$$k^- n^2\pi^2, \quad \forall n \in \mathbb{N}^*$$

and corresponding to the eigenfunctions of the form

$$\psi = (0, 0, 0, 0, \sin(n\pi x), -\sin(n\pi x), 0, 0),$$

As above, the eigenvalues are simple (in the general case the multiplicity is  $N_d^- - 1$ ).

- We find the eigenvalues  $\lambda$  of the form

$$k^- \frac{(2n-1)^2\pi^2}{4}, \quad \forall n \in \mathbb{N}^*$$

and corresponding to the eigenfunctions of the form

$$\psi = \left(0, 0, 0, 0, 0, 0, \cos\left(\frac{(2n-1)\pi}{2}x\right), -\cos\left(\frac{(2n-1)\pi}{2}x\right)\right),$$

The eigenvalues are simple (the multiplicity is  $N_d^- - 1$ ).

- The last family of eigenvalues  $\lambda \in \mathbb{R}$  are not of the form

$$\frac{n^2\pi^2}{4}, \quad k^- \frac{n^2\pi^2}{4}, \quad \forall n \in \mathbb{N}^*.$$

They are defined as the simple zeros of the equation

$$\begin{aligned} & N_d^+ \cot(\sqrt{\lambda}) - N_n^+ \tan(\sqrt{\lambda}) - \sqrt{|k^-|} N_d^- \coth\left(\frac{\sqrt{\lambda}}{\sqrt{|k^-|}}\right) - \sqrt{|k^-|} N_n^- \tanh\left(\frac{\sqrt{\lambda}}{\sqrt{|k^-|}}\right) \\ & = 2 \cot(\sqrt{\lambda}) - 2 \tan(\sqrt{\lambda}) - 2\sqrt{|k^-|} \coth\left(\frac{\sqrt{\lambda}}{\sqrt{|k^-|}}\right) - 2\sqrt{|k^-|} \tanh\left(\frac{\sqrt{\lambda}}{\sqrt{|k^-|}}\right) = 0. \end{aligned} \quad (35)$$

The corresponding eigenfunctions have the form

$$\psi(x) = (\psi_1, \dots, \psi_8),$$

$$\psi_j(x) = \begin{cases} \sin(\sqrt{\lambda}x) & 1 \leq j \leq 2, \\ \frac{\sin(\sqrt{\lambda})}{\cos(\sqrt{\lambda})} \cos(\sqrt{\lambda}x) & 3 \leq j \leq 4, \\ \frac{\sin(\sqrt{\lambda})}{\sinh\left(\frac{\sqrt{\lambda}}{\sqrt{|k^-|}}x\right)} \sinh\left(\frac{\sqrt{\lambda}}{\sqrt{|k^-|}}x\right) & 5 \leq j \leq 6, \\ \frac{\sin(\sqrt{\lambda})}{\cosh\left(\frac{\sqrt{\lambda}}{\sqrt{|k^-|}}x\right)} \cosh\left(\frac{\sqrt{\lambda}}{\sqrt{|k^-|}}x\right) & 7 \leq j \leq 8. \end{cases}$$

The definition of this eigenpairs can be directly extended to the general case of  $\mathbb{N}_d^+, \mathbb{N}_n^+, \mathbb{N}_d^-, \mathbb{N}_n^- \in \mathbb{N}^*$ .

## 6.2 Standard spectral representation for some non-equilateral star-networks

We consider the spectral problem (22) in the general case when  $L_j/L_k \notin \mathbb{Q}$  and we proceed as in Section 5.2. The eigenvalues  $\lambda \in \mathbb{R}$  in such a framework are simple, and they are defined as the zeros of the identity

$$\begin{aligned} 0 = & \sum_{j=1}^2 \cot(\sqrt{\lambda}L_j) - \sum_{j=3}^4 \tan(\sqrt{\lambda}L_j) - \\ & \sqrt{|k^-|} \sum_{j=5}^6 \coth\left(\frac{\sqrt{\lambda}}{\sqrt{|k^-|}}L_j\right) - \sqrt{|k^-|} \sum_{j=7}^8 \tanh\left(\frac{\sqrt{\lambda}}{\sqrt{|k^-|}}L_j\right). \end{aligned} \quad (36)$$

The corresponding eigenfunctions have the form

$$\psi(x) = (\psi_1, \dots, \psi_8),$$

$$\psi_j(x) = \begin{cases} \sin(\sqrt{\lambda}x) & 1 \leq j \leq 2, \\ \frac{\sin(\sqrt{\lambda})}{\cos(\sqrt{\lambda})} \cos(\sqrt{\lambda}x) & 3 \leq j \leq 4, \\ \frac{\sin(\sqrt{\lambda})}{\sinh\left(\frac{\sqrt{\lambda}}{\sqrt{|k^-|}}x\right)} \sinh\left(\frac{\sqrt{\lambda}}{\sqrt{|k^-|}}x\right) & 5 \leq j \leq 6, \\ \frac{\sin(\sqrt{\lambda})}{\cosh\left(\frac{\sqrt{\lambda}}{\sqrt{|k^-|}}x\right)} \cosh\left(\frac{\sqrt{\lambda}}{\sqrt{|k^-|}}x\right) & 7 \leq j \leq 8. \end{cases}$$

**Remark 6.1.** Notice that the equation (36) can be extended to the general case of  $N_d^+, N_n^+, N_d^-, N_n^- \in \mathbb{N}^*$  by changing the extremes of summation of each term. The construction of the eigenfunctions follows equivalently.

### 6.3 Spectral representation via pseudo-eigenfunctions for equilateral star-networks

In the current section, we study the spectral problem (23) when the edges of the star-graph have the same length  $L = 1$  and in the presence of some Neumann boundary conditions.

- We can define two families of eigenvalues. The first are the numbers  $\lambda$  of the form

$$\frac{(2n-1)^2\pi^2}{4}, \quad \forall n \in \mathbb{N}^*,$$

and with multiplicity 4. The corresponding eigenfunctions  $\psi$  have the following forms

$$\begin{aligned} & \left( \sin(\sqrt{\lambda}x), \sin(\sqrt{\lambda}x), 1, 1, \sin(\sqrt{\lambda}x), \sin(\sqrt{\lambda}x), 1, 1 \right), \\ & \left( 0, 0, \cos(\sqrt{\lambda}x), -\cos(\sqrt{\lambda}x), 0, 0, 0, 0 \right), \\ & \left( 0, 0, 0, 0, 0, 0, \cos(\sqrt{\lambda}x), -\cos(\sqrt{\lambda}x) \right), \\ & \left( 0, 0, -k^- \cos(\sqrt{\lambda}x), -k^- \cos(\sqrt{\lambda}x), 0, 0, \cos(\sqrt{\lambda}x), \cos(\sqrt{\lambda}x) \right). \end{aligned}$$

For general  $N_d^+, N_n^+, N_d^-, N_n^- \in \mathbb{N}^*$ , the 4 type of eigenfunctions can be defined equivalently: the first type is always simple, the second has multiplicity  $N_n^+ - 1$ , the third has multiplicity  $N_n^- - 1$ , and the fourth is always simple.

- The second family of eigenvalues is composed of numbers  $\lambda$  of the form:

$$n^2\pi^2, \quad n \in \mathbb{N}^*$$

with multiplicity 3 (in the general case  $N_d^+ + N_d^- - 1$ ). The corresponding eigenfunctions have the form

$$\begin{aligned} \psi &= (\psi^1, \psi^2, 0, 0, \psi^5, \psi^6, 0, 0), \quad \psi^j(x) = \alpha_j \sin(n\pi x) \\ &\text{with} \quad \alpha_1 + \alpha_2 + k^- \alpha_5 + k^- \alpha_6 = 0. \end{aligned}$$

An example is given by the following three orthogonal functions

$$\begin{aligned} & (-\sin(n\pi x), \sin(n\pi x), 0, 0, 0, 0, 0, 0), \quad (0, 0, 0, 0, -\sin(n\pi x), \sin(n\pi x), 0, 0), \\ & (-k^- \sin(n\pi x), -k^- \sin(n\pi x), 0, 0, \sin(n\pi x), \sin(n\pi x), 0, 0). \end{aligned}$$

## 6.4 Spectral representation via pseudo-eigenfunctions for some non-equilateral star-networks

Finally, we study (23) when  $L_j = |e_j|$  are such that  $L_j/L_k \notin \mathbb{Q}$  for every distinct  $j, k \leq N$ . In this case, the eigenvalues  $\lambda$  are simple (this also true for general  $N_d^+, N_n^+, N_d^-, N_n^-$ ) and they are the zeros of the equation

$$\sum_{j=1}^2 \cot(\sqrt{\lambda}L_j) - \sum_{j=3}^4 \tan(\sqrt{\lambda}L_j) + k^- \sum_{j=5}^6 \cot(\sqrt{\lambda}L_j) - k^- \sum_{j=7}^8 \tan(\sqrt{\lambda}L_j) = 0.$$

The corresponding eigenfunctions have the form

$$\psi(x) = (\psi_1, \dots, \psi_8),$$

$$\psi_j(x) = \begin{cases} \sin(\sqrt{\lambda}x) & 1 \leq j \leq 2, \\ \frac{\sin(\sqrt{\lambda})}{\cos(\sqrt{\lambda})} \cos(\sqrt{\lambda}x) & 3 \leq j \leq 4, \\ \frac{\sin(\sqrt{\lambda})}{\sinh\left(\frac{\sqrt{\lambda}}{\sqrt{|k^-|}}\right)} \sin\left(\frac{\sqrt{\lambda}}{\sqrt{|k^-|}}x\right) & 5 \leq j \leq 6, \\ \frac{\sin(\sqrt{\lambda})}{\cosh\left(\frac{\sqrt{\lambda}}{\sqrt{|k^-|}}\right)} \cos\left(\frac{\sqrt{\lambda}}{\sqrt{|k^-|}}x\right) & 7 \leq j \leq 8. \end{cases}$$

## 7 Spectral representation for two-phase tadpole networks

This section aims to study the spectral representations considered in Section 5 for the case of the tadpole graphs  $\mathcal{T}$  introduced in Section 4.1. The case in which the external vertex is not equipped with Dirichlet boundary conditions but with the Neumann one can be treated in the same way.

Assume that the lengths of the edges are  $|e_1| = L_1$  and  $|e_2| = L_2$ . We choose the constant conductivities:  $k_1 = 1$  and  $k_2 = k^- < 0$ . As in the previous section, we introduced the quadratic form

$$(\psi, \varphi)_k = \int_{e_1} \psi_1 \varphi_1 + \int_{e_2} k^- \psi_2 \varphi_2.$$

and, again, one can specify two different interesting spectral problems.

- We aim to find  $\lambda$  and  $\psi \in H_0^1(\mathcal{T})$  such that, for all  $\varphi \in H_0^1(\mathcal{T})$ ,

$$a_k(\psi, \varphi) := \int_{e_1} \psi_1' \varphi_1' + \int_{e_2} k^- \psi_2' \varphi_2' = \lambda(\psi, \varphi). \quad (37)$$

It corresponds to the study the eigenfunctions of the following operator in  $L^2(\mathcal{T})$ :

$$A\psi = \left( -\partial_x^2 \psi_1, -k^- \partial_x^2 \psi_2 \right) \quad \text{for } \psi \in D(A) \quad \text{with}$$

$$D(A) = \left\{ \psi \in H^2(e_1, \mathbb{R}) \times H^2(e_2, \mathbb{R}) : \psi_2(0) = 0, \psi_1(0) = \psi_1(|e_1|) = \psi_N(|e_2|), \right. \\ \left. \partial_x \psi_1(0) - \partial_x \psi_1(|e_1|) - k^- \partial_x \psi_2(|e_2|) = 0 \right\}.$$

**Proposition 7.1.** *The operator  $A$  is self-adjoint and admits compact resolvent for every  $k^- < 0$ .*

*Proof.* First, the fact that  $A$  is self-adjoint follows exactly as in the proof of Proposition 5.1. Second,  $A$  admits compact resolvent thanks to Theorem 4.1.  $\square$

- The second “alternative” eigenvalue problem is: find  $\lambda$  and  $\psi \in H_0^1(\mathcal{T})$  such that, for all  $\varphi \in H_0^1(\mathcal{T})$ ,

$$a_k(\psi, \varphi) = \lambda(\psi, \varphi)_k. \quad (38)$$

It corresponds to study the spectral problem associated with the operator:

$$\tilde{A}\psi = (-\partial_x^2 \psi_1, -\partial_x^2 \psi_2) \quad \text{for } \psi \in D(\tilde{A}) \quad \text{with}$$

$$D(\tilde{A}) = \left\{ \psi \in H^2(e_1, \mathbb{R}) \times H^2(e_2, \mathbb{R}) : \psi_2(0) = 0, \psi_1(0) = \psi_1(|e_1|) = \psi_N(|e_2|), \right. \\ \left. \partial_x \psi_1(0) - \partial_x \psi_1(|e_1|) - k^- \partial_x \psi_2(|e_2|) = 0 \right\}.$$

**Remark 7.2.** *As in Remark 5.2, the operator  $\tilde{A}$  is neither self-adjoint, nor symmetric.*

## 7.1 Standard spectral representation

Let us start by considering the spectral problem (37). We exploit the ideas developed in the proof of Theorem 4.1 and the eigenfunctions are either symmetric or antisymmetric w.r.t. symmetry axe  $r$ . This fact allows us to distinguish between two families of eigenpairs. The first  $(\nu_j, \varphi^j)_{j \in \mathbb{N}^*}$  is composed of antisymmetric functions:

$$\nu_j = \frac{4j^2 \pi^2}{L_1^2}, \quad \varphi^j = \left( \alpha_j \sin\left(\frac{2j\pi}{L_1} x\right), 0 \right).$$

with some normalizing  $\alpha_j$ . The second  $(\theta_j, \phi^j)_{j \in \mathbb{N}^*}$  is instead associated to symmetric eigenfunctions. Each eigenvalue, in this case, is associated to a function of form:

$$\left( \beta_1 \cos\left(\xi\left(x - \frac{L_1}{2}\right)\right), \beta_2 \sin(\mu x) \right),$$

with  $\xi = is\mu$  and  $s = \sqrt{|k^-|}$ . Now, the boundary conditions defining  $D(A)$  yield

$$\begin{cases} \beta_1 \cos\left(\xi \frac{L_1}{2}\right) = \beta_2 \sin(\mu L_2), \\ 2\beta_1 \xi \sin\left(\xi \frac{L_1}{2}\right) - k^- \beta_2 \mu \cos(\mu L_2) = 0. \end{cases} \quad (39)$$

These identities tell us that  $\beta_1, \beta_2 \neq 0$  and  $\cos(\xi \frac{L_1}{2}), \sin(\mu L_2) \neq 0$ . Thus, by recalling  $\mu = -i \frac{\xi}{\sqrt{|k^-|}}$ , the eigenvalues  $\theta_j$  are those numbers solving the following equation

$$2 \tan\left(\sqrt{\theta_j} \frac{L_1}{2}\right) + \sqrt{|k^-|} \coth\left(\frac{\sqrt{\theta_j}}{\sqrt{|k^-|}} L_2\right) = 0.$$



Notice that  $(\theta_j)_{j \in \mathbb{N}^*}$  are also real thanks to the self-adjointness of  $A$  and simple thanks to the same arguments adopted in Section 5.2. The eigenfunctions  $\phi^j$  can be defined via some normalizing constants  $\alpha_j$  as follows:

$$\phi^j = \alpha_j \left( \cos \left( \sqrt{\theta_j} \left( x - \frac{L_1}{2} \right) \right), \frac{\cos \left( \sqrt{\theta_j} \frac{L_1}{2} \right)}{\sinh \left( \frac{\sqrt{\theta_j}}{\sqrt{|k^-|}} L_2 \right)} \sinh \left( \frac{\sqrt{\theta_j}}{\sqrt{|k^-|}} x \right) \right).$$

Finally, the eigenvalues  $(\lambda_j)_{j \in \mathbb{N}^*}$  of  $A$  are obtained by reordering  $(\nu_j)_{j \in \mathbb{N}^*}$  and  $(\theta_j)_{j \in \mathbb{N}^*}$ . We denote by  $(\psi^j)_{j \in \mathbb{N}^*}$  the corresponding family of eigenfunctions obtained by reordering  $(\varphi^j)_{j \in \mathbb{N}^*}$  and  $(\phi^j)_{j \in \mathbb{N}^*}$ .

**Proposition 7.3.** *The functions  $(\psi^n)_{n \geq 1}$  form a complete orthonormal system of  $L^2(\mathcal{T})$ .*

## 7.2 Pseudo-eigenfunctions for a tadpole network

As above, we distinguish between two families of eigenpairs. The first  $(\nu_j, \varphi^j)_{j \in \mathbb{N}^*}$  is the same defined for the operator  $A$ . The second eigenpair instead  $(\tilde{\theta}_j, \tilde{\phi}^j)_{j \in \mathbb{N}^*}$  is associated to eigenfunctions of form

$$\left( \tilde{\beta}_1 \cos \left( \sqrt{\tilde{\theta}} \left( x - \frac{L_1}{2} \right) \right), \tilde{\beta}_2 \sin(\sqrt{\tilde{\theta}} x) \right).$$

such that

$$\begin{cases} \tilde{\beta}_1 \cos \left( \sqrt{\tilde{\theta}} \frac{L_1}{2} \right) = \tilde{\beta}_2 \sin(\sqrt{\tilde{\theta}} L_2), \\ 2\tilde{\beta}_1 \sin \left( \sqrt{\tilde{\theta}} \frac{L_1}{2} \right) - k^- \tilde{\beta}_2 \cos(\sqrt{\tilde{\theta}} L_2) = 0 \end{cases} \quad (40)$$

with  $\tilde{\beta}_1, \tilde{\beta}_2 \neq 0$  and. Notice that, if

$$\cos \left( \sqrt{\tilde{\theta}} \frac{L_1}{2} \right) = \sin(\sqrt{\tilde{\theta}} L_2) = 0, \quad (41)$$

then  $\tilde{\theta}$  has the form

$$\frac{(2n_1 - 1)^2 \pi^2}{L_1^2} \quad \text{and} \quad \frac{n_2^2 \pi^2}{L_2^2},$$

which is possible only when

$$\frac{L_1}{L_2} \in \mathbb{Q}.$$

Some corresponding eigenfunctions have the form, with  $\tilde{\alpha}$  suitable,

$$\tilde{\phi} = \left( \cos \left( \sqrt{\tilde{\theta}} \left( x - \frac{L_1}{2} \right) \right), \tilde{\alpha} \sin \left( \sqrt{\tilde{\theta}} x \right) \right).$$

When  $\cos \left( \sqrt{\tilde{\theta}} \frac{L_1}{2} \right) \neq 0$  and  $\sin(\sqrt{\tilde{\theta}} L_2) \neq 0$ , we can define the eigenvalues by the relation

$$2 \tan \left( \sqrt{\tilde{\theta}} \frac{L_1}{2} \right) - k^- \cot(\sqrt{\tilde{\theta}} L_2) = 0 \quad (42)$$

In this case, some corresponding eigenfunctions are defined by

$$\tilde{\phi} = \left( \cos \left( \sqrt{\tilde{\theta}} \left( x - \frac{L_1}{2} \right) \right), \frac{\cos \left( \sqrt{\tilde{\theta}} \frac{L_1}{2} \right)}{\sin(\sqrt{\tilde{\theta}} L_2)} \sin \left( \sqrt{\tilde{\theta}} x \right) \right).$$

Now, the sequence of eigenvalues  $(\tilde{\theta}_j)_{j \in \mathbb{N}^*}$  is defined by reordering the solutions of (41) and of (42) and we denote by  $(\tilde{\phi}^j)_{j \in \mathbb{N}^*}$  some corresponding normalized eigenfunctions.

Finally, the eigenvalues  $(\tilde{\lambda}_j)_{j \in \mathbb{N}^*}$  of  $\tilde{A}$  are obtained by reordering  $(\nu_j)_{j \in \mathbb{N}^*}$  and  $(\tilde{\theta}_j)_{j \in \mathbb{N}^*}$ . We denote by  $(\tilde{\psi}^j)_{j \in \mathbb{N}^*}$  the corresponding family of eigenfunctions obtained by reordering  $(\varphi^j)_{j \in \mathbb{N}^*}$  and  $(\tilde{\phi}^j)_{j \in \mathbb{N}^*}$ .

As done for the star graphs with Proposition 5.7 and Propositions 5.10, we can see now that  $(\tilde{\psi}^j)_{j \in \mathbb{N}^*}$  are isomorphic to the eigenfunctions of the Dirichlet Laplacian defined on a tadpole graph (see [15, Section 4.2]), which yields the following proposition

**Proposition 7.4.** *The family of functions  $(\tilde{\psi}^n)_{n \in \mathbb{N}^*}$  forms a Riesz Basis of  $L^2(\mathcal{T})$ , i.e. it is isomorphic to an orthonormal system of  $L^2(\mathcal{T})$ .*

## References

- [1] H. Ammari, G. Ciraolo, H. Kang, H. Lee and G. W. Milton. Spectral theory of a neumann–poincaré-type operator and analysis of cloaking due to anomalous localized resonance. *Archive for Rational Mechanics and Analysis*, 208(2):667–692, 2013.
- [2] H. Ammari, P. Millien, M. Ruiz and Hai Zhang. Mathematical analysis of plasmonic nanoparticles: the scalar case. *Archive on Rational Mechanics and Analysis*, 224(2): 597–658, 2017.
- [3] H. Ammari and P. Millien. Shape and size dependence of dipolar plasmonic resonance of nanoparticles. *J. Math. Pures Appl.*, 129(2):242–265, 2019.
- [4] H. Ammari and D. Bryn. *Metamaterial Analysis and Design: A Mathematical Treatment of Cochlea-inspired Sensors*, Volume 9 in the series De Gruyter Series in Applied and Numerical Mathematics, 2023.
- [5] A.-S. Bonnet-Ben Dhia, L. Chesnel and P. Ciarlet Jr. Two-dimensional Maxwell’s equations with sign-changing coefficients. *J. Applied Numerical Mathematics*, 79:29–41, 2014.
- [6] A.S. Bonnet-Ben Dhia, L. Chesnel, P. Ciarlet. T-coercivity for scalar interface problems between dielectrics and metamaterials. *ESAIM: Math. Model. Numer. Anal.* 46:1363–1387, 2012.
- [7] A. S. Bonnet-Ben Dhia, P. Ciarlet and C. M. Zwölf. Time harmonic wave diffraction problems in materials with sign-shifting coefficients. *J. Comput. Appl. Math.*, 234(6):1912–1919, 2010.
- [8] A.-S. Bonnet-Bendhia, M. Dauge and K. Ramdani. Analyse spectrale et singularités d’un problème de transmission non coercif. *C. R. Acad. Sci. Paris Sér. I Math.*, 328(8): 717–720, 1999.
- [9] E. Bonnetier and F. Triki. On the spectrum of the poincaré variational problem for two close-to-touching inclusions in 2d. *Archive for Rational Mechanics and Analysis*, 209(2):541–567, 2013.

- [10] E. Bonnetier, and H. Zhang. Characterization of the essential spectrum of the Neumann-Poincaré operator in 2D domains with corner via Weyl sequences. *Rev. Mat. Iber.*, 35(3):925–948, 2019.
- [11] E. Bonnetier, C. Dapogny, F. Triki, and H. Zhang. The plasmon resonances of a bowtie antenna. *Anal. Theory Appl.*, 235:85–116. 2019.
- [12] R. Bunoiu, L. Chesnel, K. Ramdani and M. Rihani. Homogenization of Maxwell’s equations and related scalar problems with sign-changing coefficients. *Annales de la Faculté des Sciences de Toulouse : Mathématiques, Ser. 6*, 30 (5):1075–1119, 2021.
- [13] R. Bunoiu and K. Ramdani. Homogenization of materials with sign changing coefficients. *Communications in Mathematical Sciences*, 14 (4):1137–1154, 2016.
- [14] R. Bunoiu, K. Ramdani and C Timofte. Homogenization of a transmission problem with sign-changing coefficients and interfacial flux jump. *Communications in Mathematical Sciences*, 21(7):2029–2049, 2023.
- [15] P. Cannarsa, A. Duca and C. Urbani. Exact controllability to eigensolutions of the bilinear heat equation on compact networks. *Discrete Contin. Dyn. Syst. Ser. S.*, 15(6):1377–1401, 2022.
- [16] P. Ciarlet Jr. T-coercivity: application to the discretization of Helmholtz-like problems. *Comput. Math. Appl.*, 64:22–34, 2012.
- [17] J. Behrndt and D. Krejcirik. An indefinite Laplacian on a rectangle. *J. Anal. Math.*, 134(2):501–522, 2018.
- [18] M. Costabel and E. Stephan. A direct boundary integral equation method for transmission problems. *Journal of Mathematical Analysis and Applications*, 106(2):367–413, 1985.
- [19] M. Costabel, M. Dauge and S. Nicaise. Corner Singularities and Analytic Regularity for Linear Elliptic Systems. Part I: Smooth Domains, Prépublication IRMAR, 10-09. hal-00453934, version 2.
- [20] M. Dauge and B. Texier. Non-coercive transmission problems in polygonal domains. *arXiv:1102.1409[math.NA]*.
- [21] D. Grieser. The plasmonic eigenvalue problem. *Rev. Math. Phys.*, 26:145000, 2014.
- [22] L. Grubisic, V. Kostykin, K. A. Makarov, and K. Veselic. Representation theorems for indefinite quadratic forms revisited. *Mathematika*, 59:169–189, 2013.
- [23] A. Hussein. Spectral theory of differential operators on finite metric graphs and on bounded domains. Ph.D thesis, University of Mainz, 2013.
- [24] A. Hussein. Sign-indefinite second order differential operators on finite metric graphs. *Rev. Math. Phys.*, **26** (2014), 1430003.
- [25] R. V. Kohn, J. Lu, B. Schweizer and M. I. Weinstein. A variational perspective on cloaking by anomalous localized resonance. *Communications in Mathematical Physics*, 328:1–27, 2014.
- [26] P. Kuchment. Quantum graphs. I. Some basic structures. *Special section on quantum graphs. Waves Random Media*, 14(1):S107–S128, 2004.

- [27] I. D. Mayergoyz. Plasmon resonances in nanoparticles, their applications to magnetics and relation to the Riemann hypothesis. *Physica B: Condensed Matter*, 407(9):1307–1313, 2012.
- [28] I. D. Mayergoyz, D. R. Fredkin and Z. Zhang. Electrostatic (plasmon) resonances in nanoparticles. *Phys. Rev. B*, 72:155412, 2005.
- [29] H. M. Nguyễn. Negative index materials and their applications: recent mathematics progress. *Chinese Annals of Mathematics, Series B*, 38:601–628, 2017.
- [30] S. Nicaise, J. Venel. A posteriori error estimates for a finite element approximation of transmission problems with sign changing coefficients. *Journal of Computational and Applied Mathematics*, 235(14):4272–4282, 2011.
- [31] D. R. Smith, J.B. Pendry and M. C. K. Wiltshire. Metamaterials and negative refractive index. *Science*, 305:788–792, 2004.
- [32] V.G. Veselago. The electrodynamics of substances with simultaneously negative values of  $\epsilon$  and  $\mu$ . *Sov. Phys. Usp*, 10:509–514, 1968.