



3D kinetic Monte-Carlo simulation of diamond growth

Audrey Valentin, Ovidiu Brinza, Samir Farhat, Fabien Benedic, Jocelyn Achard

► To cite this version:

Audrey Valentin, Ovidiu Brinza, Samir Farhat, Fabien Benedic, Jocelyn Achard. 3D kinetic Monte-Carlo simulation of diamond growth. SBDD XXVI, Mar 2022, Hasselt, Belgium. hal-04643844

HAL Id: hal-04643844

<https://hal.science/hal-04643844v1>

Submitted on 10 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

3D kinetic Monte-Carlo simulation of diamond growth

Audrey Valentin, Ovidiu Brinza, Samir Farhat, Fabien Bénédic, Jocelyn Achard


¹LSPM-CNRS, Laboratoire des Sciences des Procédés et des Matériaux, Université Sorbonne Paris Nord,
99 avenue Jean-Baptiste Clément, 93430 Villetaneuse, France

Context



The KMC model


Processes taken into account
[CH₃] and T taken from [6]



Process	Frequency	Ref
Adsorption	$\frac{v_{CH_3}}{4} [CH_3] \times \rho \times S \times R_H$	
Desorption	$\frac{k_B T}{h} \exp\left(-\frac{E_a}{RT}\right)$, $E_a = 200 \text{ kJ/mol}$	[4]
Etching	$\frac{k_B T}{h} \exp\left(-\frac{E_a}{RT}\right) \exp\left(-\frac{\epsilon N_a E_a}{RT}\right)$, $\epsilon = 0.365$	
Surface migration	$6.13 \times 10^{13} \times \exp\left(-\frac{18269}{T}\right) \times R_H$	[5]

Lattice geometry


- (1 0 0) surfaces
- Dimer rows (in green) taken into account




Calculated growth rates

Comparison to experimental data [7]

- Good reproduction at low methane concentration
- Unrelevant modeling of germination?




Conclusion & perspectives



Growth of a single layer


Effect of temperature

- [CH₄] = 4%
- Long islands due to dimer rows
 - Germination at low and high temperature



Effect of methane concentration

- T = 850 C
- Effect of methane concentration is not visible
 - Germination expected at high concentration




Growth on nanostructures

All simulation below performed with:

- [CH₄] = 4% and T = 850 C
- (1 0 0) growth model: new faces appear but their growth is not treated


(1 1 0)-oriented pillars

Appearance of (1 1 1) and (1 1 0) faces




(1 0 0)-oriented pillars

Appearance of (1 1 1) faces



(1 1 1)-oriented pyramids

Two atomic layers high steps



References

- G. Lombardi *et al.*, J. Appl. Phys. **98**, 053303 (2005).
- A. Valentin *et al.*, Diam. Relat. Mater. **123**, 108865 (2022).
- F. Silva *et al.*, Phys. Status Solidi A **203**, 3049 (2006).
- W. J. Rodgers *et al.*, J. Chem. Phys. **142**, 214707 (2015).
- A. Netto and M. Frenklach, Diam. Relat. Mater. **14**, 1630 (2005).
- F. Silva *et al.*, Diam. Relat. Mater. **18**, 683-697 (2009).
- O. Brinza *et al.*, Phys. Status Solidi A **205** 2114 (2008)